

## ORIGINAL RESEARCH

## Impact of baseline body mass index on the efficacy and safety of tofacitinib in patients with psoriatic arthritis

Jon T Giles,<sup>1</sup> Alexis Ogdie,<sup>2</sup> Juan J Gomez-Reino,<sup>3</sup> Philip S Helliwell ,<sup>4</sup> Rebecca Germino,<sup>5</sup> Lori Stockert,<sup>6</sup> Pamela Young,<sup>6</sup> Wael Joseph,<sup>6</sup> Rajiv Mundayat,<sup>5</sup> Daniela Graham,<sup>7</sup> Christopher Ritchlin <sup>8</sup>

**To cite:** Giles JT, Ogdie A, Gomez-Reino JJ, *et al.* Impact of baseline body mass index on the efficacy and safety of tofacitinib in patients with psoriatic arthritis. *RMD Open* 2021;7:e001486. doi:10.1136/rmdopen-2020-001486

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/rmdopen-2020-001486>).

Data in this manuscript were previously presented at the ACR/ARP 2019 Annual Meeting (Ritchlin C, Ogdie A, Giles J, *et al.* *Arthritis Rheumatol* 2019;71 [Suppl 10]:1513).

Received 16 October 2020  
Revised 21 December 2020  
Accepted 30 December 2020



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

**Correspondence to**

Dr Jon T Giles;  
jtg2122@cumc.columbia.edu

**ABSTRACT**

**Objectives** This post-hoc analysis explored the impact of body mass index (BMI) on tofacitinib efficacy/safety in patients with active psoriatic arthritis (PsA).

**Methods** Data were pooled from two phase 3 studies (NCT01877668; NCT01882439). Analyses included patients randomised to tofacitinib 5/10 mg two times a day or placebo, stratified by baseline BMI: <25 kg/m<sup>2</sup>, ≥25–<30 kg/m<sup>2</sup>, ≥30–<35 kg/m<sup>2</sup> or ≥35 kg/m<sup>2</sup>. Endpoints (month 3): American College of Rheumatology (ACR20/50/70), Health Assessment Questionnaire-Disability Index (HAQ-DI) and Psoriasis Area and Severity Index (PASI) 75 response rates; dactylitis/enthesitis resolution rates; changes from baseline Short Form-36 Health Survey version 2 (SF-36v2) Physical/Mental Component Summary (PCS/MCS) scores and Functional Assessment of Chronic Illness Therapy-Fatigue (FACIT-F) total score. Safety was also reported.

**Results** Analysis included 710 patients; 43.8% were obese (BMI ≥30 kg/m<sup>2</sup>). Tofacitinib demonstrated higher efficacy response rates at month 3, compared with placebo, regardless of baseline BMI. Generally, ACR20/50/70 and HAQ-DI response rates, enthesitis resolution rates and changes from baseline in SF-36v2 PCS score and FACIT-F total score (month 3) were reduced in patients with baseline BMI ≥35 kg/m<sup>2</sup> versus patients with lower BMIs. Elevated alanine aminotransferase/aspartate aminotransferase levels were reported in patients with baseline BMI ≥35 kg/m<sup>2</sup> receiving tofacitinib 5 mg but not 10 mg two times a day.

**Conclusion** Tofacitinib demonstrated greater efficacy than placebo in patients with PsA, regardless of baseline BMI. For all treatment arms, reduced efficacy was observed in patients with baseline BMI ≥35 kg/m<sup>2</sup>. Safety was generally comparable across BMI categories, although the effect of tofacitinib on liver enzymes in patients with baseline BMI ≥35 kg/m<sup>2</sup> was inconclusive.

**INTRODUCTION**

Obesity (body mass index (BMI) ≥30 kg/m<sup>2</sup>) poses a significant global health and economic burden.<sup>1</sup> Elevated BMI, a surrogate indicator of obesity,<sup>2</sup> is a key risk factor for numerous conditions, including cardiovascular disease,<sup>3</sup>

**Key messages****What is already known about this subject?**

- Obesity is prevalent in patients with psoriatic arthritis (PsA) and is associated with a reduced response to tumour necrosis factor inhibitors (TNFi).

**What does this study add?**

- The oral Janus kinase inhibitor tofacitinib demonstrated greater efficacy than placebo in patients with PsA, irrespective of baseline body mass index (BMI).
- Efficacy response rates to tofacitinib and placebo were generally reduced in patients with baseline BMI ≥35 kg/m<sup>2</sup>, compared with patients with a lower baseline BMI.
- The safety profile of tofacitinib was largely similar across baseline BMI categories; however, per cent changes from baseline alanine aminotransferase and aspartate aminotransferase levels were greater in patients treated with tofacitinib 5 mg but not 10 mg two times a day in the baseline BMI ≥35 kg/m<sup>2</sup> category compared with those in the other baseline BMI categories.

**How might this impact on clinical practice?**

- Our data suggest that tofacitinib can still be considered an effective treatment option for patients with PsA, regardless of their BMI status.

diabetes mellitus<sup>3</sup> and chronic autoimmune disorders such as psoriatic arthritis (PsA).<sup>4,5</sup>

PsA is a chronic, systemic inflammatory arthritis characterised by involvement of the joints, entheses, skin and nails, and the axial skeleton.<sup>6,7</sup> The prevalence of obesity is increased in patients with PsA compared with patients with psoriasis or rheumatoid arthritis (RA) and with the general population.<sup>8–10</sup> Furthermore, in a population-based study of ~2 million individuals, obesity was identified as a risk factor for developing PsA in patients with psoriasis and in the general population.<sup>4</sup>

Treatment options for PsA include: non-steroidal anti-inflammatory drugs; conventional synthetic disease-modifying antirheumatic drugs (csDMARDs), for example, methotrexate, sulfasalazine and leflunomide; biological DMARDs (bDMARDs), for example, tumour necrosis factor inhibitors (TNFi), interleukin (IL)-12/23 inhibitors and IL-17 inhibitors; and targeted synthetic DMARDs, for example, phosphodiesterase inhibitors and Janus kinase (JAK) inhibitors.<sup>11 12</sup> Intriguingly, therapeutic response to TNFi was decreased in patients with obesity with PsA compared with patients without obesity.<sup>13 14</sup> Moreover, in a meta-analysis of 19 372 patients with various autoimmune conditions, including PsA, patients with obesity were found to have 60% higher odds of failing TNFi therapy versus patients without obesity.<sup>15</sup> Weight loss ( $\geq 5\%$  from baseline) is associated with higher rates of achieving minimal disease activity (MDA) in overweight patients or patients with obesity with PsA treated with TNFi.<sup>16</sup> Accordingly, the Group for Research and Assessment of Psoriasis and Psoriatic Arthritis guidelines recommend maintaining a healthy body weight in patients with PsA to improve TNFi effectiveness.<sup>12</sup>

Tofacitinib is an oral JAK inhibitor for the treatment of PsA. The efficacy and safety of tofacitinib was demonstrated in phase 3 trials of up to 12 months' duration in patients with active PsA and an inadequate response to csDMARDs or TNFi therapy,<sup>17 18</sup> and for up to 36 months' duration in a long-term extension study.<sup>19</sup> Here, for the first time, we examined the impact of baseline BMI on the efficacy and safety of tofacitinib in patients with active PsA. Understanding the potential effect of BMI on treatment response in patients with PsA may inform therapeutic decision-making in the clinic.

## METHODS

### Patients

This post-hoc analysis pooled data from two randomised, placebo-controlled, double-blind, phase 3 studies (OPAL Broaden (NCT01877668); OPAL Beyond (NCT01882439)) in patients with active PsA.<sup>17 18</sup>

OPAL Broaden was a 12-month study in patients who had an inadequate response to  $\geq 1$  csDMARD and were TNFi-naïve. Patients were randomised 2:2:2:1:1 to receive oral tofacitinib 5 or 10 mg two times a day, adalimumab 40 mg subcutaneously every 2 weeks, or placebo progressing to tofacitinib 5 or 10 mg two times a day at 3 months.<sup>17</sup> OPAL Beyond was a 6-month study in patients who had an inadequate response to  $\geq 1$  TNFi and were randomised 2:2:1:1 to receive tofacitinib 5 or 10 mg two times a day, or placebo progressing to tofacitinib 5 or 10 mg two times a day at 3 months.<sup>18</sup> Throughout both studies, all patients were required to receive a stable dose of one csDMARD (eg, methotrexate, leflunomide or sulfasalazine). Full study details, including patient inclusion and exclusion criteria, have been published previously.<sup>17 18</sup> Briefly, eligible patients were aged  $\geq 18$  years, had signs and symptoms consistent with a diagnosis of

PsA  $\geq 6$  months prior to the study and met the Classification Criteria for Psoriatic Arthritis at screening.<sup>17 18</sup>

The current post-hoc analysis study included all patients from OPAL Broaden and OPAL Beyond who received tofacitinib 5 or 10 mg two times a day, or placebo. Patients were stratified into the following baseline BMI categories according to the WHO<sup>2</sup>: underweight/normal ( $< 25 \text{ kg/m}^2$ ), overweight ( $\geq 25$ – $< 30 \text{ kg/m}^2$ ), class 1 obesity ( $\geq 30$ – $< 35 \text{ kg/m}^2$ ), and class 2 and 3 obesity ( $\geq 35 \text{ kg/m}^2$ ).

Both OPAL Broaden and OPAL Beyond were conducted in accordance with the Declaration of Helsinki and Good Clinical Practice Guidelines of the International Conference on Harmonisation, and were approved by the relevant Institutional Review Board and/or Independent Ethics Committee of the investigational centres. All patients provided written, informed consent.

### Efficacy outcomes

Efficacy data were pooled at month 3 from OPAL Broaden and OPAL Beyond and stratified by baseline BMI category. Endpoints included: American College of Rheumatology (ACR) 20/50/70% (ACR20/50/70) improvement from baseline in tender and swollen joint counts, and in three of the five remaining ACR core domains (patient's and physician's global assessment of arthritis activity, patient's assessment of arthritis pain, measurement of disability and level of acute-phase reactants); Health Assessment Questionnaire-Disability Index (HAQ-DI) response rate (defined as a  $\geq 0.35$ -point decrease from baseline, which is considered to be the minimum clinically important difference)<sup>20</sup>; response rate in achieving  $\geq 75\%$  reduction from baseline Psoriasis Area and Severity Index (PASI75) in patients with baseline body surface area (BSA)  $\geq 3\%$  and PASI  $> 0$ ; dactylitis and enthesitis resolution rates (Dactylitis Severity Score (DSS)=0; Leeds Enthesitis Index (LEI)=0) in patients with baseline DSS  $> 0$  and LEI  $> 0$ , respectively; least squares (LS) mean changes from baseline Short Form-36 Health Survey version 2 (SF-36v2) Physical Component Summary (PCS) and Mental Component Summary (MCS) scores; LS mean change from baseline Functional Assessment of Chronic Illness Therapy-Fatigue (FACIT-F) total scores; and LS mean change from baseline HAQ-DI scores.

Patients achieved MDA response if they met  $\geq 5$  of the following criteria: tender joint count  $\leq 1$ ; swollen joint count  $\leq 1$ ; PASI score  $\leq 1$  or BSA  $\leq 3\%$ ; patient's assessment of arthritis pain visual analogue scale (VAS)  $\leq 15$  mm; patient's global assessment of arthritis activity VAS  $\leq 20$  mm; HAQ-DI score  $\leq 0.5$ ; or LEI  $\leq 1$ .

### Safety outcomes

Safety data were pooled up to month 3 from OPAL Broaden and OPAL Beyond and stratified by baseline BMI category. Endpoints included adverse events (AEs), changes from baseline in liver function tests and lipid panel parameters.

## Statistical analyses

All analyses were performed on the full analysis set, which included patients who were randomised and received  $\geq 1$  dose of study treatment.

For the binary efficacy endpoints (ie, response rates), non-responder imputation was used to account for missing values; patients with missing values were treated as non-responders. Observed data were used for continuous efficacy endpoints.

Multivariable analyses were conducted to evaluate the possible association between baseline BMI (as a continuous or categorical variable) and the efficacy endpoints: ACR20/50/70, HAQ-DI and PASI75 response rates, and change from baseline in HAQ-DI, LEI and DSS. For each tofacitinib treatment group, a multivariable logistic regression analysis for binary endpoints and multivariable regression analysis for continuous endpoints were conducted, to determine significant predictors for each efficacy endpoint based on a backward selection method using a 5% level of significance. Separate models were run using categorical BMI and continuous BMI as covariates. Other baseline covariates included: age, race, gender, region, weight, smoking status, disease duration, prior bDMARD use, FACIT-F total score, LEI, DSS, C-reactive protein (CRP), metabolic syndrome, corticosteroid use, prior TNFi use, pain VAS, HAQ-DI, SF-36v2 PCS and SF-36v2 MCS. No multiplicity adjustment was performed in this post-hoc analysis. Safety endpoints were analysed descriptively.

## RESULTS

### Patients

In total, 710 patients (OPAL Broaden,  $n=316$ ; OPAL Beyond,  $n=394$ ) were included in this post-hoc analysis, 311 (43.8%) of whom were obese. Of these, 186 had a BMI  $\geq 30$ – $<35$  kg/m<sup>2</sup>, and 125 had a BMI  $\geq 35$  kg/m<sup>2</sup>. The majority of patients were middle-aged, female and Caucasian (table 1). Slightly more female patients were randomised to the  $<25$  kg/m<sup>2</sup> and  $\geq 35$  kg/m<sup>2</sup> baseline BMI categories compared with patients in the other baseline BMI categories. Metabolic syndrome and elevated CRP levels ( $>2.87$  mg/L) were more frequent in patients with baseline BMI  $\geq 35$  kg/m<sup>2</sup> versus those with a lower baseline BMI (table 1). Higher HAQ-DI scores, tender and swollen joint counts and lower SF-36v2 PCS scores were observed in patients in the baseline BMI  $\geq 35$  kg/m<sup>2</sup> category, compared with patients in the other baseline BMI categories. Additionally, prior use of one bDMARD was more prevalent in patients with baseline BMI  $\geq 35$  kg/m<sup>2</sup> versus those with a lower baseline BMI.

Patient demographics and baseline disease characteristics by baseline BMI category and treatment are shown in online supplemental table 1.

### Efficacy outcomes by BMI category

Consistent with the main findings of OPAL Broaden and OPAL Beyond, ACR20/50/70, HAQ-DI and PASI75

response rates at month 3 were higher in tofacitinib-treated patients, compared with placebo-treated patients, across all baseline BMI categories (figure 1). In all treatment groups, response rates were generally lower in patients in the baseline BMI  $\geq 35$  kg/m<sup>2</sup> category, compared with patients in the other baseline BMI categories, particularly when treated with tofacitinib 10 mg two times a day (figure 1). In patients treated with placebo, the PASI75 response rate was higher in patients in the baseline BMI  $\geq 35$  kg/m<sup>2</sup> category, compared with patients in the other baseline BMI categories (figure 1E). It should be noted that PASI75 response rate was only assessed in patients with baseline BSA  $\geq 3\%$  and PASI  $>0$ , and patient numbers were lower than those for the ACR20/50/70 and HAQ-DI response rates.

Consistent with the main findings of OPAL Broaden and OPAL Beyond, dactylitis and enthesitis resolution rates at month 3 were higher in tofacitinib-treated patients, compared with placebo-treated patients, across all baseline BMI categories (figure 2). The dactylitis resolution rate was numerically higher in placebo-treated patients with baseline BMI  $\geq 35$  kg/m<sup>2</sup>, compared with placebo-treated patients in the other baseline BMI categories (figure 2A). Across all baseline BMI categories, the dactylitis resolution rate was increased in patients treated with tofacitinib 10 mg two times a day versus those treated with tofacitinib 5 mg two times a day or placebo (figure 2A). Patients with baseline BMI  $\geq 35$  kg/m<sup>2</sup> exhibited lower enthesitis resolution rates, compared with patients in the other baseline BMI categories, regardless of treatment (figure 2B).

Overall, patients treated with tofacitinib 5 and 10 mg two times a day exhibited greater mean changes from baseline SF36v2 PCS and MCS scores, FACIT-F total scores and HAQ-DI at month 3, compared with patients treated with placebo (figure 3). Patients in the baseline BMI  $\geq 35$  kg/m<sup>2</sup> category treated with tofacitinib 5 or 10 mg two times a day exhibited lower mean changes from baseline SF-36v2 PCS score and FACIT-F total score, compared with patients in the other baseline BMI categories (figure 3A,C).

Multivariable analyses with BMI as both a categorical and continuous variable demonstrated that BMI was not a significant predictor of response across most efficacy endpoints at month 3, and across both tofacitinib treatment groups (online supplemental table 2). However, in patients treated with tofacitinib 5 mg two times a day, BMI was a significant predictor of response for: ACR50 response (BMI categorical;  $p=0.035$ ), ACR70 response (BMI categorical ( $p=0.023$ ) and continuous ( $p=0.002$ )), MDA rate (BMI categorical ( $p=0.048$ ) and continuous ( $p=0.003$ )), and change from baseline at month 3 in HAQ-DI (BMI continuous;  $p=0.050$ ).

### Safety outcomes by BMI category

Online supplemental table 3 shows AEs reported in  $\geq 5\%$  of patients up to month 3 for each treatment group across baseline BMI categories. The most common

**Table 1** Patient demographics and baseline disease characteristics by baseline body mass index (BMI) category (pooled data from OPAL Broaden and OPAL Beyond)

	Baseline BMI category			
	<25 kg/m <sup>2</sup> (underweight/normal) N=161 (22.7%)	≥25–<30 kg/m <sup>2</sup> (overweight) N=238 (33.5%)	≥30–<35 kg/m <sup>2</sup> (class 1 obesity) N=186 (26.2%)	≥35 kg/m <sup>2</sup> (class 2 and 3 obesity) N=125 (17.6%)
Patient demographics				
Age, years, mean (SD)	44.5 (12.9)	50.0 (11.8)	50.5 (11.7)	51.2 (11.3)
Female, n (%)	101 (62.7)	118 (49.6)	92 (49.5)	82 (65.6)
White, n (%)	149 (92.5)	224 (94.1)	175 (94.1)	121 (96.8)
BMI, kg/m <sup>2</sup> , mean (SD)	22.5 (1.7)	27.5 (1.4)	32.3 (1.4)	39.6 (4.1)
Weight, kg, n (%)				
<60	55 (34.2)	0	1 (0.5)	1 (0.8)
≥60–<90	104 (64.6)	202 (84.9)	78 (41.9)	11 (8.8)
≥90	2 (1.2)	36 (15.1)	107 (57.5)	113 (90.4)
Waist circumference, cm, mean (SD)*	81.0 (9.0)	95.2 (10.5)	106.0 (9.3)	118.0 (13.2)
Metabolic syndrome, n (%)†	7 (4.3)	85 (35.7)	107 (57.5)	95 (76.0)
Never smoked, n (%)	110 (68.3)	147 (61.8)	108 (58.1)	72 (57.6)
Baseline disease characteristics				
PsA duration, years, mean (SD)	8.1 (7.7)	8.0 (7.8)	8.5 (7.1)	7.3 (6.3)
HAQ-DI, mean (SD)	1.1 (0.7)	1.1 (0.7)	1.2 (0.6)	1.5 (0.7)
SF-36v2 PCS score, mean (SD)	36.5 (8.7)	35.3 (8.5)	34.2 (7.9)	30.8 (8.2)
SF-36v2 MCS score, mean (SD)	38.6 (11.4)	41.9 (11.7)	39.9 (11.8)	39.3 (12.6)
FACIT-F total score, mean (SD)	27.5 (10.7)	28.9 (10.8)	27.3 (10.1)	24.1 (11.8)
CRP mg/L, mean (SD)	14.2 (27.5)	10.6 (18.8)	9.9 (17.3)	14.4 (19.2)
>2.87 mg/L, n (%)‡	79 (49.1)	143 (60.1)	117 (62.9)	105 (84.0)
PASI, mean (SD)	9.2 (8.5)	9.7 (8.3)	10.4 (9.5)	10.1 (8.1)
BSA ≥3%, n (%)	113 (70.2)	162 (68.1)	125 (67.2)	81 (64.8)
SJC, mean (SD)	9.9 (7.7)	12.1 (9.6)	11.7 (9.6)	14.7 (11.5)
TJC, mean (SD)	18.5 (13.0)	21.0 (14.2)	20.7 (13.6)	26.4 (16.8)
DSS, mean (SD) (N1)§	8.7 (7.7) (82)	8.9 (8.7) (133)	7.2 (6.5) (88)	9.6 (9.5) (70)
LEI, mean (SD) (N2)¶	2.6 (1.4) (101)	2.9 (1.6) (160)	3.1 (1.7) (123)	3.2 (1.7) (95)
Number of prior bDMARDs, n (%)				
0	70 (43.5)	114 (47.9)	79 (42.5)	42 (33.6)
1	55 (34.2)	79 (33.2)	59 (31.7)	53 (42.4)
≥2	36 (22.4)	45 (18.9)	48 (25.8)	30 (24.0)
Corticosteroid use at baseline, n (%)	35 (21.7)	47 (19.7)	40 (21.5)	31 (24.8)
NSAID use at baseline, n (%)	91 (56.5)	135 (56.7)	96 (51.6)	70 (56.0)
Concomitant methotrexate, n (%)	122 (75.8)	202 (84.9)	139 (74.7)	93 (74.4)
Dose, mg/week, mean (SD)	15.3 (4.2)	14.8 (4.3)	14.8 (4.6)	15.3 (4.2)
≤15 mg/week, n (%)	79 (49.1)	140 (58.8)	91 (48.9)	61 (48.8)
>15 mg/week, n (%)	43 (26.7)	62 (26.1)	48 (25.8)	23 (25.6)

\*Baseline waist circumference data were unavailable for one patient in the <25 kg/m<sup>2</sup> category and three patients in the ≥25–<30 kg/m<sup>2</sup> category.  
 †A patient was classified as having baseline metabolic syndrome if any three of the five metabolic syndrome risk factors at baseline were satisfied: hypertension (systolic ≥130 mm Hg and/or diastolic ≥85 mm Hg and/or concomitant antihypertensive medication), elevated triglycerides (≥150 mg/dL, 1.7 mmol/L and/or concomitant lipid-lowering medication), reduced high-density lipoprotein cholesterol (<40 mg/dL, 1.0 mmol/L in males; <50 mg/dL, 1.3 mmol/L in females), elevated waist circumference (population-specific and country-specific definitions) and elevated fasting glucose levels (≥100 mg/dL and/or concomitant anti-diabetic medication).<sup>45</sup>

‡Elevated level of CRP defined as >2.87 mg/L.


§Assessed only in patients with baseline DSS >0.

¶Assessed only in patients with baseline LEI >0.


bDMARD, biological disease-modifying antirheumatic drug; BSA, body surface area; CRP, C-reactive protein; DSS, Dactylitis Severity Score; FACIT-F, Functional Assessment of Chronic Illness Therapy-Fatigue; HAQ-DI, Health Assessment Questionnaire-Disability Index; LEI, Leeds Enthesitis Index; MCS, Mental Component Summary; N, number of patients randomised and treated; N1, number of patients with baseline DSS >0; N2, number of patients with baseline LEI >0; NSAID, non-steroidal anti-inflammatory drug; PASI, Psoriasis Area and Severity Index; PCS, Physical Component Summary; PsA, psoriatic arthritis; SF-36v2, Short Form-36 Health Survey version 2; SJC, swollen joint count; TJC, tender joint count.




■ BMI <25 kg/m<sup>2</sup> (Underweight/Normal) ■ BMI ≥25–<30 kg/m<sup>2</sup> (Overweight) ■ BMI ≥30–<35 kg/m<sup>2</sup> (Class 1 Obesity) ■ BMI ≥35 kg/m<sup>2</sup> (Class 2 & 3 Obesity)



**Figure 1** Response rates at month 3 for (A) ACR20,\* (B) ACR50,\* (C) ACR70,\* (D) HAQ-DI† and (E) PASI75‡ by baseline BMI category and treatment group (pooled data from OPAL Broaden and OPAL Beyond). N indicates the number of patients with non-missing BMI evaluable at month 3; n indicates the number of responders. All patients received a stable dose of one csDMARD throughout each study. \*ACR20/50/70 response rates were defined as the proportions of patients achieving a ≥20/50/70% improvement from baseline in tender and swollen joint counts and ≥20/50/70% improvement from baseline in three of the five remaining ACR core measures. †HAQ-DI response rate was defined as the proportion of patients achieving a reduction from baseline HAQ-DI of ≥0.35, considered the minimum clinically important difference. ‡PASI75 response rate was defined as the proportion of patients achieving a reduction from baseline PASI of ≥75%, assessed only in patients with baseline BSA ≥3% and a baseline PASI>0. ACR20/50/70, American College of Rheumatology≥20/50/70% response criteria; BID, two times a day; BMI, body mass index; BSA, body surface area; csDMARD, conventional synthetic disease-modifying antirheumatic drug; HAQ-DI, Health Assessment Questionnaire-Disability Index; PASI75, ≥75% Psoriasis Area and Severity Index improvement from baseline.



**Figure 2** Resolution rates at month 3 for (A) dactylitis\* and (B) enthesitis† by baseline BMI category and treatment group (pooled data from OPAL Broaden and OPAL Beyond). N indicates the number of patients with non-missing BMI evaluable at month 3; n indicates the number of responders. All patients received a stable dose of one csDMARD throughout each study. \*Dactylitis resolution rates, defined as the absence of dactylitis in all of the 20 assessed digits, assessed only in patients with baseline DSS >0. †Enthesitis resolution rates, defined as the absence of enthesitis in all of the six assessed sites, assessed only in patients with baseline LEI >0. BID, two times a day; BMI, body mass index; csDMARD, conventional synthetic disease-modifying antirheumatic drug; DSS, Dactylitis Severity Score; LEI, Leeds Enthesitis Index.



**Figure 3** LS mean (SE) changes from baseline at month 3 in (A) SF-36v2 PCS score, (B) SF-36v2 MCS score, (C) FACIT-F total score and (D) HAQ-DI by baseline BMI category and treatment group (pooled data from OPAL Broaden and OPAL Beyond). N indicates the number of patients with non-missing BMI evaluable at month 3; n indicates the number of responders. All patients received a stable dose of one csDMARD throughout each study.  $\Delta$ , change from baseline; BID, two times a day; BMI, body mass index; csDMARD, conventional synthetic disease-modifying antirheumatic drug; FACIT-F, Functional Assessment of Chronic Illness Therapy-Fatigue; HAQ-DI, Health Assessment Questionnaire-Disability Index; LS, least squares; MCS, Mental Component Summary; PCS, Physical Component Summary; SF-36v2, Short Form-36 Health Survey version 2.

AEs were similar to past reports<sup>17,18</sup>; falls, constipation, dizziness, bronchitis and dry skin were reported more frequently among patients with baseline BMI  $\geq 35$  kg/m<sup>2</sup>. Up to month 3, three cardiovascular events were reported: a non-fatal cerebrovascular accident and a transient ischaemic attack (both occurring in the same patient; tofacitinib 5 mg two times a day; baseline BMI  $\geq 30$ – $<35$  kg/m<sup>2</sup>) and a coronary artery revascularisation (patient received placebo; baseline BMI  $\geq 35$  kg/m<sup>2</sup>). One patient treated with tofacitinib 5 mg two times a day in the baseline BMI  $\geq 30$ – $<35$  kg/m<sup>2</sup> category had an arterial thromboembolic event. Those that experienced cardiovascular and thromboembolic events were found to have a history of hypertension, coronary heart disease, abnormal ECGs or angina. In this dataset, no deep vein thrombosis or pulmonary embolism events were reported up to month 3.

At month 3, per cent changes from baseline alanine aminotransferase (ALT) and aspartate aminotransferase (AST) levels were greater in patients treated with tofacitinib 5 and 10 mg two times a day, compared with patients treated with placebo, regardless of BMI category (table 2). Among patients treated with tofacitinib 5 mg two times a day, per cent changes in ALT and AST levels were notably greater in the baseline BMI  $\geq 35$  kg/m<sup>2</sup> category than in the other baseline BMI categories; among patients treated with tofacitinib 10 mg two times a day, per cent changes in ALT and AST levels were greater in the baseline BMI  $\geq 25$ – $<30$  kg/m<sup>2</sup> category than in the other baseline BMI categories. Generally, observations were similar for absolute liver enzyme levels (table 2).

For most BMI categories, per cent changes from baseline at month 3 in low-density lipoprotein, high-density lipoprotein, total cholesterol and triglyceride levels were greater in tofacitinib-treated patients, compared with placebo-treated patients. However, per cent changes from baseline triglyceride levels were higher in patients treated with placebo, compared with patients treated with tofacitinib 5 mg two times a day in the baseline BMI  $<25$  kg/m<sup>2</sup> category, and tofacitinib 5 and 10 mg two times a day in the baseline BMI  $\geq 30$ – $<35$  kg/m<sup>2</sup> category (table 2). Among patients treated with tofacitinib 5 mg two times a day, per cent change from baseline triglyceride levels was notably lower in the baseline BMI  $<25$  kg/m<sup>2</sup> category versus the other baseline BMI categories.

## DISCUSSION

This post-hoc analysis investigated the impact of baseline BMI on tofacitinib efficacy and safety in patients with active PsA. Tofacitinib 5 and 10 mg two times a day demonstrated higher efficacy response rates at month 3, compared with placebo, regardless of baseline BMI. Patients in the baseline BMI  $\geq 35$  kg/m<sup>2</sup> category had more active disease (eg, elevated CRP levels ( $>2.87$  mg/L) and higher tender/swollen joint counts) at baseline compared with patients in the other baseline BMI categories. For tofacitinib and placebo treatment

arms, efficacy response rates were generally reduced in patients with baseline BMI  $\geq 35$  kg/m<sup>2</sup> compared with patients with lower baseline BMIs. Additionally, baseline BMI was found to be a significant predictor of response for ACR50/70 and MDA rates, and change from baseline in HAQ-DI at month 3 in patients receiving tofacitinib 5 mg (but not 10 mg) two times a day. Up to month 3, there were no notable differences in the safety profile of tofacitinib between baseline BMI categories; however, the per cent changes from baseline ALT and AST levels were greater in patients treated with tofacitinib 5 mg (but not 10 mg) two times a day in the baseline BMI  $\geq 35$  kg/m<sup>2</sup> category, compared with patients in the other baseline BMI categories.

Elevated ALT and AST levels in patients with baseline BMI  $\geq 35$  kg/m<sup>2</sup> treated with tofacitinib 5 mg but not 10 mg two times a day provide limited evidence for an effect of tofacitinib on the liver enzymes of this BMI cohort. The small sample size within each tofacitinib arm (tofacitinib 5 mg two times a day, n=42; tofacitinib 10 mg two times a day, n=43) of the baseline BMI  $\geq 35$  kg/m<sup>2</sup> category may have led to this random finding, although there could be other putative explanations. For example, undiagnosed non-alcoholic fatty liver disease, which is prevalent in obesity,<sup>21</sup> might have contributed to the elevated ALT and AST levels in patients with baseline BMI  $\geq 35$  kg/m<sup>2</sup> treated with tofacitinib 5 mg two times a day; although baseline ALT and AST levels in patients in the BMI  $\geq 35$  kg/m<sup>2</sup> category were relatively similar to those in patients in the other baseline BMI categories. Furthermore, there were no notable differences in baseline disease characteristics or prior/baseline/concomitant treatments between BMI categories (or study treatment arms) that could account for the elevated liver enzyme levels observed in patients with baseline BMI  $\geq 35$  kg/m<sup>2</sup> receiving tofacitinib 5 mg two times a day.

Overall, for most lipid panel measures, the per cent changes from baseline at month 3 were greater in patients treated with tofacitinib than in those treated with placebo, regardless of baseline BMI. Small increases in cholesterol, low-density lipoprotein, high-density lipoprotein and triglyceride levels have been reported previously in patients with moderate to severe psoriasis treated with tofacitinib at week 16.<sup>22</sup> Accordingly, blockade of IL-6, a downstream target of tofacitinib,<sup>23</sup> is associated with increased low-density lipoprotein levels in patients with RA.<sup>24</sup> Although obesity is associated with dyslipidaemia,<sup>25</sup> we observed no notable relationship between BMI and per cent changes from baseline lipid panel measures in response to tofacitinib.

In obese individuals, adipocyte hypertrophy can lead to overproduction of proinflammatory cytokines (eg, TNF- $\alpha$  and IL-6) which may increase the risk and severity of inflammatory disorders.<sup>26–29</sup> Accordingly, there is an increased prevalence of obesity among patients with PsA.<sup>8–10</sup> Moreover, patients with obesity with PsA have been shown to exhibit higher Disease Activity Score 28 and pain scores, compared with patients without

**Table 2** Liver enzyme and lipid panel parameters at month 3, by baseline BMI category and treatment group

	Baseline BMI category			
	<25 kg/m <sup>2</sup> (underweight/normal)	≥25–<30 kg/m <sup>2</sup> (overweight)	≥30–<35 kg/m <sup>2</sup> (class 1 obesity)	≥35 kg/m <sup>2</sup> (class 2 and 3 obesity)
<b>Liver function test</b>				
ALT, % change from baseline, mean (SD) (N)				
Tofacitinib 5 mg two times a day	22.67 (73.7) (46)	12.70 (52.9) (84)	15.81 (50.5) (56)	34.13 (68.8) (40)
Tofacitinib 10 mg two times a day	19.91 (41.9) (43)	34.33 (123.4) (78)*	21.22 (41.2) (60)	27.30 (40.3) (41)
Placebo	16.54 (84.7) (58)	11.68 (58.4) (66)	7.23 (46.1) (62)	11.39 (48.0) (35)
ALT, IU/L, mean (SD) (N)				
Tofacitinib 5 mg two times a day	22.93 (16.71) (46)	26.42 (14.37) (84)	30.50 (22.19) (56)	31.88 (19.06) (40)
Tofacitinib 10 mg two times a day	20.19 (10.39) (43)	28.87 (15.68) (78)	31.83 (13.63) (60)	31.44 (13.93) (41)
Placebo	18.02 (9.97) (58)	26.33 (15.56) (66)	24.53 (12.09) (62)	27.17 (18.63) (35)
AST, % change from baseline, mean (SD) (N)				
Tofacitinib 5 mg two times a day	13.67 (36.0) (46)	13.97 (42.2) (83)	10.22 (30.8) (55)	32.52 (52.9) (40)
Tofacitinib 10 mg two times a day	18.07 (43.1) (43)	20.13 (43.1) (78)	16.25 (32.0) (59)	19.05 (33.5) (41)
Placebo	5.70 (31.9) (58)	6.32 (42.0) (66)	3.01 (29.7) (62)	1.17 (26.0) (35)
AST, IU/L, mean (SD) (N)				
Tofacitinib 5 mg two times a day	22.67 (7.89) (46)	24.23 (9.81) (83)	25.22 (10.02) (55)	28.20 (16.34) (40)
Tofacitinib 10 mg two times a day	22.70 (8.90) (43)	26.03 (8.86) (78)	26.29 (8.47) (59)	27.15 (8.17) (41)
Placebo	19.50 (5.48) (58)	24.85 (17.92) (66)	20.44 (6.09) (62)	20.69 (8.91) (35)
<b>Lipid panel†</b>				
LDL, % change from baseline, mean (SD) (N)				
Tofacitinib 5 mg two times a day	4.29 (15.1) (45)	12.22 (24.2) (73)	7.08 (18.3) (53)	12.11 (19.3) (37)
Tofacitinib 10 mg two times a day	16.88 (27.3) (40)	14.96 (24.8) (71)	13.03 (26.0) (57)	11.35 (26.8) (33)
Placebo	1.77 (21.0) (57)	7.27 (17.7) (61)	4.98 (21.2) (56)	1.98 (13.4) (31)
HDL, % change from baseline, mean (SD) (N)				
Tofacitinib 5 mg two times a day	6.64 (13.4) (45)	13.27 (24.9) (78)	5.33 (14.3) (53)	11.80 (21.9) (38)
Tofacitinib 10 mg two times a day	20.05 (21.0) (40)	9.63 (18.9) (72)	15.59 (17.0) (58)	14.88 (18.0) (36)
Placebo	-2.16 (18.4) (57)	0.87 (15.5) (64)	-1.15 (13.8) (58)	0.24 (16.6) (32)
Total cholesterol, % change from baseline, mean (SD) (N)				
Tofacitinib 5 mg two times a day	3.70 (11.2) (45)	11.42 (17.4) (79)	6.27 (13.0) (53)	10.41 (9.3) (38)
Tofacitinib 10 mg two times a day	15.90 (20.7) (40)	12.16 (17.3) (73)	12.94 (18.1) (58)	7.62 (18.8) (37)
Placebo	-0.56 (12.6) (57)	4.35 (14.0) (64)	3.50 (15.1) (58)	-0.57 (9.4) (32)
Triglycerides, % change from baseline, mean (SD) (N)				
Tofacitinib 5 mg two times a day	0.37 (29.6) (45)	9.87 (42.9) (79)	14.72 (47.6) (53)	11.20 (32.7) (38)
Tofacitinib 10 mg two times a day	12.12 (59.2) (40)	23.21 (57.8) (72)	16.86 (45.9) (58)	9.84 (62.3) (36)
Placebo	8.53 (38.1) (57)	8.76 (49.1) (64)	17.54 (53.8) (58)	0.12 (31.8) (32)

N indicates the number of patients assessed at baseline and month 3. All patients received a stable dose of one csDMARD throughout each study.

\*High SD value due to a per cent change in ALT level of 1037.50 reported for one patient. Note that this value was not considered to be clinically significant, and had decreased at retesting.

†Data on lipids for patients in fasting state only.

ALT, alanine aminotransferase; AST, aspartate aminotransferase; BMI, body mass index; csDMARD, conventional synthetic disease-modifying antirheumatic drug; HDL, high-density lipoprotein; IU/L, international units/litre; LDL, low-density lipoprotein.

obesity.<sup>14</sup> The ongoing PsABio study (NCT02627768), including patients with PsA from eight European countries, recently reported an association between elevated BMI and higher clinical Disease Activity in Psoriatic

Arthritis, Psoriatic Impact of Disease-12 and HAQ-DI scores.<sup>30</sup> Here, increased baseline BMI (≥35 kg/m<sup>2</sup>) was generally associated with increased tender and swollen joint counts, PASI75 and HAQ-DI scores, elevated CRP



(>2.87 mg/L) and decreased SF-36v2 PCS scores at baseline. Additionally, among placebo-treated patients, those with a baseline BMI  $\geq 35$  kg/m<sup>2</sup> typically exhibited reduced efficacy response rates at month 3, compared with those in lower baseline BMI categories.

Tofacitinib inhibits the downstream signalling of several JAK-dependent cytokines, including TNF- $\alpha$  and IL-6<sup>23</sup>; it is, therefore, possible that the therapeutic effectiveness of tofacitinib is attenuated by the elevated expression of proinflammatory cytokines in patients with obesity. Accordingly, among patients treated with tofacitinib 5 mg and 10 mg two times a day, those with a baseline BMI  $\geq 35$  kg/m<sup>2</sup> generally experienced reduced efficacy response rates at month 3, versus those with lower baseline BMIs. Interestingly, an analysis of pooled data from two phase 3 studies in patients with RA treated with the JAK inhibitor baricitinib revealed reduced ACR20/50/70 response rates after 12 weeks in patients with baseline BMI  $>30.7$  kg/m<sup>2</sup>, compared with those with lower baseline BMIs.<sup>31</sup> Additionally, patients with obesity (BMI  $\geq 30$  kg/m<sup>2</sup>) with RA were reported to have reduced rates of achieving low disease activity and remission compared with normal weight (BMI  $<25$  kg/m<sup>2</sup>) and overweight (BMI  $\geq 25$ – $<30$  kg/m<sup>2</sup>) patients at 6 months after commencing baricitinib or tofacitinib treatment.<sup>32</sup> It should be noted that tofacitinib has been shown to be efficacious versus placebo, in patients with PsA in the current study, and in patients with RA,<sup>33</sup> regardless of BMI.

It is also noteworthy that pooled analyses of phase 3 studies in patients with RA<sup>33</sup> and ulcerative colitis<sup>34</sup> receiving tofacitinib 5 or 10 mg two times a day, or placebo, revealed similar efficacy outcomes across all treatment groups, regardless of baseline BMI category; however, these analyses included three BMI categories ( $<25$  kg/m<sup>2</sup>,  $25$ – $<30$  kg/m<sup>2</sup> and  $\geq 30$  kg/m<sup>2</sup>) and the current analysis included a fourth BMI category ( $\geq 35$  kg/m<sup>2</sup>), so a direct comparison with this analysis is limited.

Tofacitinib has moderate lipophilicity. Analysis of pharmacokinetic parameters for tofacitinib demonstrated that while increased body weight results in lower peak ( $C_{\max}$ ) and higher trough ( $C_{\min}$ ) plasma concentrations, it has no clinically relevant impact on total drug exposure (AUC); therefore, no dose adjustment is recommended for tofacitinib based on patients' body weight.<sup>35</sup> The reduced efficacy response to tofacitinib in patients with baseline BMI  $\geq 35$  kg/m<sup>2</sup> could be due to a larger volume of distribution, which can lower plasma concentrations. Body weight has been shown to influence the PK profile of biological agents, including TNFi, in various immune disorders.<sup>36–38</sup> In a population PK study in patients with chronic inflammatory diseases, including PsA, treated with the TNFi infliximab, drug volume of distribution and clearance increased with body weight.<sup>37</sup>

In patients with obesity, increased immunogenicity and elevated drug clearance may account for the effect of high body weight on decreased efficacy of TNFi and other biological agents.<sup>15 39</sup> A study in patients with

PsA, RA and controls without rheumatic disease (n=89) reported increased adipose expression of inflammatory genes in patients with PsA, compared with patients with RA or controls, across all BMI categories. Additionally, patients treated with biologics expressed high adipose levels of inflammatory genes regardless of baseline BMI.<sup>40</sup>

Interestingly, although obesity is associated with a slight attenuated therapeutic response to TNFi and JAK inhibitors in patients with PsA and RA,<sup>13–15 31</sup> the response to the IL-6 inhibitor tocilizumab in patients with RA was reported to not be affected by high baseline BMI.<sup>41</sup> However, IL-6 inhibitors are not indicated for the treatment of PsA. The reasons why TNFi and JAK inhibitors have reduced effectiveness in patients with obesity compared with other drugs are poorly understood; however, PK properties, volume of distribution and lipophilicity may be contributing factors.<sup>42</sup> It is also possible that, due to the inflammatory processes associated with obesity,<sup>26–29</sup> higher doses of TNFi or JAK inhibitors are required to treat individuals with an elevated BMI, compared with those with lower BMIs.

Current ACR guidelines for the treatment of PsA recommend weight loss in overweight patients or patients with obesity to potentially improve pharmacologic responses.<sup>43</sup> Correspondingly, in patients with obesity with PsA, weight loss has been reported to improve disease activity in a dose-dependent manner.<sup>44</sup>

A limitation of this research is that the analysis was post-hoc, and comparisons between tofacitinib and placebo treatment groups were limited to the 3-month, placebo-controlled portion of the OPAL Broaden and OPAL Beyond studies; further research is required to evaluate the long-term impact of BMI on treatment response to tofacitinib in patients with PsA in the real world. No adjustments for multiple comparisons were performed for this analysis; therefore, chance findings cannot be ruled out. In addition, small numbers of patients in the treatment and BMI strata limit the breadth of the conclusions that can be made from these analyses. Measures of peripheral proinflammatory molecules, such as IL-6 or leptin were not part of the study design and may have provided additional insight into the study results.

## CONCLUSIONS

Across all baseline BMI categories, tofacitinib demonstrated higher efficacy in patients with PsA at month 3, compared with placebo. However, similar to TNFi therapy,<sup>13–15</sup> reduced efficacy was generally observed in tofacitinib-treated and placebo-treated patients with baseline BMI  $\geq 35$  kg/m<sup>2</sup>, compared with patients in the other baseline BMI categories. In general, tofacitinib safety appeared similar across all baseline BMI categories up to month 3, although the effect of tofacitinib on liver enzymes in patients with baseline BMI  $\geq 35$  kg/m<sup>2</sup> was inconclusive.

### Author affiliations

<sup>1</sup>Department of Medicine/Division of Rheumatology, Columbia University Vagelos College of Physicians and Surgeons, New York, New York, USA

<sup>2</sup>Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania, USA

<sup>3</sup>Fundación del Instituto de Investigación Sanitaria, Hospital Clínico Universitario, Santiago de Compostela, Spain

<sup>4</sup>Leeds Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds, Leeds, UK

<sup>5</sup>Pfizer Inc, New York, New York, USA

<sup>6</sup>Pfizer Inc, Collegeville, Pennsylvania, USA

<sup>7</sup>Pfizer Inc, Groton, Connecticut, USA

<sup>8</sup>Department of Medicine/Division of Allergy, Immunology and Rheumatology, University of Rochester Medical Center, Rochester, New York, USA

**Contributors** All authors provided critical revision of the manuscript. All authors interpreted the results, approved the final draft, and had the final decision to submit the manuscript for publication. Pfizer did not control the analysis or interpretation of the study results. Publication of this article was not contingent upon approval by Pfizer.

**Funding** This study was sponsored by Pfizer. Medical writing support, under the guidance of the authors, was provided by Dominic Singson, MD, and Emma Mitchell, PhD, CMC Connect, McCann Health Medical Communications and was funded by Pfizer, New York, New York, USA in accordance with Good Publication Practice (GPP3) guidelines (Ann Intern Med 2015;163:461–4).

**Competing interests** JTG has received research grants from Pfizer; and consultancy fees from AbbVie, Bristol-Myers Squibb, Eli Lilly, Genentech, Gilead and UCB. AO has received research grants from Amgen (to Forward/National Databank), Novartis (to the University of Pennsylvania) and Pfizer Inc (to the University of Pennsylvania); and consultancy fees from AbbVie, Amgen, Bristol-Myers Squibb, Celgene, Corrona, Eli Lilly, Janssen, Novartis, Pfizer, Takeda Pharmaceutical Company and UCB. AO's husband has received royalties from Novartis. JGG-R has received funding from Pfizer. PSH has received research grants from AbbVie, Janssen and Novartis; and honoraria from AbbVie, Amgen, Celgene, Galapagos, Janssen, Pfizer and UCB. RG, LS, PY, WJ, RM and DG are employees and shareholders of Pfizer. CR has received research grants from AbbVie, Amgen and UCB; and consultancy fees from AbbVie, Amgen, Bristol-Myers Squibb, Eli Lilly, Janssen, Novartis, Pfizer and UCB.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request. Upon request, and subject to certain criteria, conditions and exceptions (see <https://www.pfizer.com/science/clinical-trials/trial-data-and-results> for more information), Pfizer will provide access to individual deidentified participant data from Pfizer-sponsored global interventional clinical studies conducted for medicines, vaccines and medical devices (1) for indications that have been approved in the US and/or EU, or (2) in programmes that have been terminated (ie, development for all indications has been discontinued). Pfizer will also consider requests for the protocol, data dictionary and statistical analysis plan. Data may be requested from Pfizer trials 24 months after study completion. The deidentified participant data will be made available to researchers whose proposals meet the research criteria and other conditions, and for which an exception does not apply, via a secure portal. To gain access, data requestors must enter into a data access agreement with Pfizer.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

### ORCID iDs

Philip S Helliwell <http://orcid.org/0000-0002-4155-9105>

Christopher Ritchlin <http://orcid.org/0000-0002-2602-1219>

## REFERENCES

- Chooi YC, Ding C, Magkos F. The epidemiology of obesity. *Metabolism* 2019;92:6–10.
- World Health Organization. Global health Observatory (GHO) data, 2020. Available: [https://www.who.int/gho/ncd/risk\\_factors/bmi\\_text/en/](https://www.who.int/gho/ncd/risk_factors/bmi_text/en/) [Accessed 21 May 2020].
- Singh I, Rajagopalan S, Srinivasan A, *et al*. Preoperative statin therapy is associated with lower requirement of renal replacement therapy in patients undergoing cardiac surgery: a meta-analysis of observational studies. *Interact Cardiovasc Thorac Surg* 2013;17:345–52.
- Love TJ, Zhu Y, Zhang Y, *et al*. Obesity and the risk of psoriatic arthritis: a population-based study. *Ann Rheum Dis* 2012;71:1273–7.
- Harpsoe MC, Basit S, Andersson M, *et al*. Body mass index and risk of autoimmune diseases: a study within the Danish national birth cohort. *Int J Epidemiol* 2014;43:843–55.
- Gladman DD, Antoni C, Mease P, *et al*. Psoriatic arthritis: epidemiology, clinical features, course, and outcome. *Ann Rheum Dis* 2005;64(Suppl 2):ii14–17.
- Kang EJ, Kavanaugh A. Psoriatic arthritis: latest treatments and their place in therapy. *Ther Adv Chronic Dis* 2015;6:194–203.
- Labitigan M, Bahçe-Altuntas A, Kremer JM, *et al*. Higher rates and clustering of abnormal lipids, obesity, and diabetes mellitus in psoriatic arthritis compared with rheumatoid arthritis. *Arthritis Care Res* 2014;66:600–7.
- Jafri K, Bartels CM, Shin D, *et al*. Incidence and management of cardiovascular risk factors in psoriatic arthritis and rheumatoid arthritis: a population-based study. *Arthritis Care Res* 2017;69:51–7.
- Ogdie A, Eder L. Improving cardiovascular health and metabolic comorbidities in patients with psoriatic arthritis. *Int J Clin Rheumatol* 2015;10:451–9.
- Gossec L, Smolen JS, Ramiro S, *et al*. European League against rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2015 update. *Ann Rheum Dis* 2016;75:499–510.
- Coates LC, Kavanaugh A, Mease PJ, *et al*. Group for research and assessment of psoriasis and psoriatic arthritis 2015 treatment recommendations for psoriatic arthritis. *Arthritis Rheumatol* 2016;68:1060–71.
- di Minno MND, Peluso R, Iervolino S, *et al*. Obesity and the prediction of minimal disease activity: a prospective study in psoriatic arthritis. *Arthritis Care Res* 2013;65:141–7.
- Højgaard P, Grintborg B, Kristensen LE, *et al*. The influence of obesity on response to tumour necrosis factor- $\alpha$  inhibitors in psoriatic arthritis: results from the DANBIO and ICEBIO registries. *Rheumatology* 2016;55:2191–9.
- Singh S, Facciorusso A, Singh AG, *et al*. Obesity and response to anti-tumor necrosis factor- $\alpha$  agents in patients with select immune-mediated inflammatory diseases: a systematic review and meta-analysis. *PLoS One* 2018;13:e0195123.
- Di Minno MND, Peluso R, Iervolino S, *et al*. Weight loss and achievement of minimal disease activity in patients with psoriatic arthritis starting treatment with tumour necrosis factor  $\alpha$  blockers. *Ann Rheum Dis* 2014;73:1157–62.
- Mease P, Hall S, FitzGerald O, *et al*. Tofacitinib or adalimumab versus placebo for psoriatic arthritis. *N Engl J Med* 2017;377:1537–50.
- Gladman D, Rigby W, Azevedo VF, *et al*. Tofacitinib for psoriatic arthritis in patients with an inadequate response to TNF inhibitors. *N Engl J Med* 2017;377:1525–36.
- Nash P, Coates LC, Kivitz AJ, *et al*. SAT0293 Safety and efficacy of tofacitinib, an oral Janus kinase inhibitor, up to 36 months in patients with active psoriatic arthritis: data from the third interim analysis of OPAL Balance, an open-label long-term extension study [abstract]. *Ann Rheum Dis* 2018;77:SAT0293.
- Mease PJ, Woolley JM, Bitman B, *et al*. Minimally important difference of health assessment questionnaire in psoriatic arthritis: relating thresholds of improvement in functional ability to patient-rated importance and satisfaction. *J Rheumatol* 2011;38:2461–5.
- Milić S, Lulić D, Štimac D. Non-alcoholic fatty liver disease and obesity: biochemical, metabolic and clinical presentations. *World J Gastroenterol* 2014;20:9330–7.
- Wolk R, Armstrong EJ, Hansen PR, *et al*. Effect of tofacitinib on lipid levels and lipid-related parameters in patients with moderate to severe psoriasis. *J Clin Lipidol* 2017;11:1243–56.
- Hodge JA, Kawabata TT, Krishnaswami S, *et al*. The mechanism of action of tofacitinib - an oral Janus kinase inhibitor for the treatment of rheumatoid arthritis. *Clin Exp Rheumatol* 2016;34:318–28.
- Robertson J, Porter D, Sattar N, *et al*. Interleukin-6 blockade raises LDL via reduced catabolism rather than via increased synthesis: a cytokine-specific mechanism for cholesterol changes in rheumatoid arthritis. *Ann Rheum Dis* 2017;76:1949–52.
- Klop B, Elte JWF, Cabezas MC. Dyslipidemia in obesity: mechanisms and potential targets. *Nutrients* 2013;5:1218–40.
- Goyal R, Faizy AF, Siddiqui SS, *et al*. Evaluation of TNF- $\alpha$  and IL-6 levels in obese and non-obese diabetics: pre- and Postinsulin effects. *N Am J Med Sci* 2012;4:180–4.

- 27 Popko K, Gorska E, Stelmaszczyk-Emmel A, *et al.* Proinflammatory cytokines IL-6 and TNF- $\alpha$  and the development of inflammation in obese subjects. *Eur J Med Res* 2010;15(Suppl 2):120–2.
- 28 Eder K, Baffy N, Falus A, *et al.* The major inflammatory mediator interleukin-6 and obesity. *Inflamm Res* 2009;58:727–36.
- 29 Hamminga EA, van der Lely AJ, Neumann HAM, *et al.* Chronic inflammation in psoriasis and obesity: implications for therapy. *Med Hypotheses* 2006;67:768–73.
- 30 Siebert S, Bergmans P, de Vlam K. OP0007 High body mass index (BMI) in psoriatic arthritis (PSA) is associated with higher disease activity in joints and skin, impaired quality of life and more disability: results from the PSABIO study [abstract]. *Ann Rheum Dis* 2019;78:OP0007.
- 31 Zerbini CAF, Muram D, Arora VK, *et al.* Effect of BMI on baricitinib efficacy: pooled analysis from two phase 3 rheumatoid arthritis clinical trials [abstract]. *Arthritis Rheumatol* 2016;68:1640 <https://acrabstracts.org/abstract/effect-of-bmi-on-baricitinib-efficacy-pooled-analysis-from-two-phase-3-rheumatoid-arthritis-clinical-trials>
- 32 Meißner Y, Baganz L, Schneider M, *et al.* Baricitinib and tofacitinib in real life - does obesity impact response to Janus kinase inhibitor therapy in rheumatoid arthritis? [abstract]. *Arthritis Rheumatol* 2018;70:1518 <https://acrabstracts.org/abstract/baricitinib-and-tofacitinib-in-real-life-does-obesity-impact-response-to-janus-kinase-inhibitor-therapy-in-rheumatoid-arthritis/>
- 33 Dikranian A, Gonzalez-Gay MA, Wellborne F, *et al.* The efficacy of tofacitinib in patients with rheumatoid arthritis stratified by baseline body mass index [abstract]. *Arthritis Rheumatol* 2018;69:2371 <https://acrabstracts.org/abstract/the-efficacy-of-tofacitinib-in-patients-with-rheumatoid-arthritis-stratified-by-baseline-body-mass-index/>
- 34 Farraye FA, Qazi T, Kotze PG, *et al.* Tu1753 – analysis of the impact of body mass index on efficacy and safety in the tofacitinib Octave ulcerative colitis program. *Gastroenterology* 2019;156:S-1111–2.
- 35 Pfizer Inc. Xeljanz® (tofacitinib): highlights of prescribing information, 2015. Available: <http://labeling.pfizer.com/ShowLabeling.aspx?id=959> [Accessed 21 Nov 2020].
- 36 Mostafa NM, Nader AM, Noertersheuser P, *et al.* Impact of immunogenicity on pharmacokinetics, efficacy and safety of adalimumab in adult patients with moderate to severe chronic plaque psoriasis. *J Eur Acad Dermatol Venereol* 2017;31:490–7.
- 37 Passot C, Mulleman D, Bejan-Angoulvant T, *et al.* The underlying inflammatory chronic disease influences infliximab pharmacokinetics. *MAbs* 2016;8:1407–16.
- 38 Wade JR, Parker G, Kosutic G, *et al.* Population pharmacokinetic analysis of certolizumab pegol in patients with Crohn's disease. *J Clin Pharmacol* 2015;55:866–74.
- 39 Keizer RJ, Huitema ADR, Schellens JHM, *et al.* Clinical pharmacokinetics of therapeutic monoclonal antibodies. *Clin Pharmacokinet* 2010;49:493–507.
- 40 Yun J-H, Winchester R, Zhang H-Z, *et al.* Adipose tissue Inflammation in psoriatic arthritis: overexpression of a wide array of inflammatory mediators and associations with disease and treatment characteristics [abstract]. *Arthritis Rheumatol* 2018;70:1618 <https://acrabstracts.org/abstract/adipose-tissue-inflammation-in-psoriatic-arthritis-overexpression-of-a-wide-array-of-inflammatory-mediators-and-associations-with-disease-and-treatment-characteristics/>
- 41 Pers Y-M, Godfrin-Valnet M, Lambert J, *et al.* Response to tocilizumab in rheumatoid arthritis is not influenced by the body mass index of the patient. *J Rheumatol* 2015;42:580–4.
- 42 Toussiro E. The interrelations between biological and targeted synthetic agents used in inflammatory joint diseases, and obesity or body composition. *Metabolites* 2020;10:107.
- 43 Singh JA, Guyatt G, Ogdie A, *et al.* Special article: 2018 American College of Rheumatology/National psoriasis Foundation guideline for the treatment of psoriatic arthritis. *Arthritis Rheumatol* 2019;71:5–32.
- 44 Klingberg E, Bilberg A, Björkman S, *et al.* Weight loss improves disease activity in patients with psoriatic arthritis and obesity: an interventional study. *Arthritis Res Ther* 2019;21:17.
- 45 Alberti KGMM, Eckel RH, Grundy SM, *et al.* Harmonizing the metabolic syndrome: a joint interim statement of the International diabetes Federation Task force on epidemiology and prevention; National heart, lung, and blood Institute; American heart association; world heart Federation; international atherosclerosis Society; and international association for the study of obesity. *Circulation* 2009;120:1640–5.