

Attitudes and preferences of consumers toward food allergy labeling practices by diagnosis of food allergies

Se-young Ju¹, Jong-Hwan Park², Tong-Kyoung Kwak^{2S} and Kyu-earn Kim³

¹Department of Food Science and Nutrition, Dankook University, Yongin, Gyeonggi 448-701, Korea

²Department of Food Science and Nutrition, Yonsei University, 50 Yonsei-ro, Seoul 120-749, Korea

³Department of pediatrics, Yonsei University College of Medicine, Seoul 120-752, Korea

BACKGROUND/OBJECTIVES: The objective of this study was to investigate food allergens and prevalence rates of food allergies, followed by comparison of consumer attitudes and preferences regarding food allergy labeling by diagnosis of food allergies.

SUBJECTS/METHODS: A total of 543 individuals living in Seoul and Gyeonggi area participated in the survey from October 15 to 22 in 2013.

RESULTS: The results show that the prevalence of doctor-diagnosed food allergies was 17.5%, whereas 6.4% of respondents self-reported food allergies. The most common allergens of doctor-diagnosed and self-reported food allergy respondents were peaches (30.3%) and eggs (33.3%), respectively, followed by peanuts, cow's milk, and crab. Regarding consumer attitudes toward food labeling, checking food allergens as an item was only significantly different between allergic and non-allergic respondents among all five items ($P < 0.001$). All respondents reported that all six items (bold font, font color, box frame, warning statement, front label, and addition of potential allergens) were necessary for an improved food allergen labeling system. PLSR analysis determined that the doctor-diagnosed group and checking of food allergens were positively correlated, whereas the non-allergy group was more concerned with checking product brands.

CONCLUSIONS: An effective food labeling system is very important for health protection of allergic consumers. Additionally, government agencies must develop policies regarding prevalence of food allergies in Korea. Based on this information, the food industry and government agencies should provide clear and accurate food labeling practices for consumers.

Nutrition Research and Practice 2015;9(5):517-522; doi:10.4162/nrp.2015.9.5.517; pISSN 1976-1457 eISSN 2005-6168

Keywords: Food allergy, food labeling, self-reported allergy, attitudes, preferences

INTRODUCTION

The prevalence of food allergies is 3-5% in adults and 3-8% in children worldwide [1,2]. Furthermore, the rate of self-reported food allergies has risen up to 25% in the United States. The prevalence of food allergies has increased, resulting in a greater frequency of public health problems. A previous study in 2007 reported an overall prevalence of self-reported food allergies from 1.7% to 17%, although there were limitations in determining accurate incidences such as rapid evolution of food allergies, lack of confirmative diagnostic tests, and numerous potential factors [2-4]. A 23 large-scale meta-analysis of the EuroPrevall program reported overall prevalence rates of 12% for self-reported food allergies in infants and children as well as 13% across all ages [3]. Lee *et al.* [5] reported prevalence rates of 12.6% for experienced food allergies in students along with self-reported food allergy rates of 5.8% among 27,679 children and adolescents in Korea.

Food allergies are associated with a variety of symptoms,

ranging from mild to severe or life-threatening, affecting the skin (urticaria, dermatitis, eczema, angioedema, and itching), gastrointestinal tract (nausea, vomiting, diarrhea, and abdominal cramping), and respiratory tract (rhinitis, asthma, and laryngeal edema), in addition to anaphylactic shock [6]. Anaphylactic shock is the most frightening food-allergic symptom and develops very rapidly after ingestion of allergenic foods, leading to death if prompt treatment is not provided [7].

There are more than 160 foods confirmed to cause food allergies [6]. However, the most common allergenic foods are eggs, cow's milk, soy, and peanuts in infant and children. For adults, wheat, shellfish, peanuts, and tree nuts are the most common allergenic foods [8]. The only consistent preventive method is complete avoidance of all foods containing the causative allergen. Accordingly, complete and accurate allergen labeling is essential to successful avoidance of problematic allergens by food-allergic consumers [9]. Food labeling legislation was passed in Australia and New Zealand, followed by the European Commission and United States in 2003-2004 [10]. In

This study was supported by Ministry of Food and Drug Safety(13052.901).

^S Corresponding Author: Tong-Kyoung Kwak, Tel. 82-10-5259-4090, Email. Kwaktk@yonsei.ac.kr

Received: March 2, 2015, Revised: May 20, 2015, Accepted: May 24, 2015

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Korea, allergenic labeling regulations introduced in 2004 require mandatory labeling of 12 allergenic foods: eggs, cow's milk, buckwheat, peanuts, soybeans, wheat, mackerel, crab, shrimp, pork, peaches, and tomatoes as well as ingredients from those foods [8]. Although allergenic food labeling regulations were introduced, there is still room for improvements to current allergenic labeling, which provides insufficient information and confusing terminology [11]. Several studies have investigated consumer preferences and satisfaction regarding allergenic food labeling. In a report by Lee *et al.* [12], 993 consumers who visited seven university hospitals were not satisfied with existing labeling practices. Cornelisse-Vermaat *et al.* [13] also showed that consumers with food allergies in Greece and Netherlands indicate problems of readability and accessibility to allergen information on labels. Therefore, the objective of this study was to investigate food allergens and prevalence rates of food allergies, followed by comparison of consumer attitudes and preferences regarding food labeling by diagnosis of food allergies.

SUBJECTS AND METHODS

Study population and study design

A total of 543 individuals living in Seoul and Gyeonggi area participated in the survey from October 15 to 22 in 2013. The participants were grouped by age; teenagers (elementary, middle, and high school students), twenties, thirties, forties, fifties, and sixties. The questionnaire consisted of four sections based on a review of the literature related to food allergies [5,13-16]. The study was approved by the institutional review board of Yonsei University (1040917-201311-SB-125-02), Seoul, Korea.

The first section contained three questions that identified subjects' food allergy information, including diagnosis by a doctor. The second section contained six items that assessed subjects' attitudes toward food labeling, including brand, nutrition facts, origin, food allergens, quality, and overall checking, on a 5-point Likert scale ranging from 1 "Never check" to 5 "Always check" for checking labels. The third section consisted of six items, including bold font, font color, box frame, warning statement, front label for allergens, and addition of potential allergens for improvement of labeling information, on a using 5-point Likert scale ranging from 1 "Strongly disagree" to 5 "Strongly agree". The final section consisted of general characteristic questions such as residential area, gender, age, job, and education level.

Statistical analysis

The data were analyzed using the Statistical Package for Social Science (Ver. 21.0, SPSS, Inc., Chicago, IL). For the categorical variables, frequency and percentage were calculated using frequency analysis. For the continuous variables, descriptive analysis was generated for mean and standard deviation. Significant testing was conducted by One-way Anova. When a significant difference was observed, Duncan's multiple comparison was used to examine differences between the doctor-diagnosed group and self-reported group at $P < 0.05$. Partial Least Square Regression (PLSR) Analysis was performed for correlation between attitudes of food labeling and improve-

ment of food allergen labeling system based on whether or not respondents had diagnosed food allergies.

RESULTS

General characteristics of respondents

A total of 543 participants returned the questionnaire from 558, resulting in a response rate of 97.3%. Table 1 shows the general characteristics of the respondents. The respondents' age range was teenagers (27.3%), twenties (12.9%), thirties (14.9%), forties (14.7%), fifties (15.5%), and sixties and more (14.7%). Approximately 50% of respondents were women. The majority of respondents lived in a large city (92.1%), and the highest education level of respondents was 68.8% high school or higher.

Food allergy prevalence rates of respondents

Table 2 presents food allergy prevalence rates of respondents. The number of respondents who had food allergies was 130

Table 1. General characteristics of respondents (n = 543)

	n	%
Gender		
Male	271	49.9
Female	272	50.1
Age (yrs)		
< 19	149	27.4
20-29	67	12.3
30-39	83	15.3
40-49	78	14.4
50-59	83	15.3
≥60	83	15.3
Region		
Large city	500	92.1
Middle and small city	43	7.9
Job		
Professional	81	15.0
Production, technical and self-employed	63	11.7
Service, and sales	45	8.3
Housewife	101	18.7
University student	65	12.0
Teenagers (elementary, middle, and high school students)	145	26.9
Unemployed	40	7.4
Education level		
6 yrs and less	91	16.9
9 yrs	77	14.3
12 yrs	203	37.6
over 12 yrs	169	31.2
Total	543	100.0

Table 2. Food allergy prevalence rates of respondents (n = 543)

Item	n	%
Doctor diagnosed food allergy group	95	17.5
Self-reported food allergy group	35	6.4
Non-allergy group	413	76.1
Total	543	100.0

(23.9%), which included doctor-diagnosed (n = 95, 17.5%) and self-reported respondents (n = 35, 6.4%). The other 413 respondents out of 543 did not report food allergies.

Types of food allergens reported by doctor-diagnosed and self-reported respondents

Table 3 shows food allergens reported by doctor-diagnosed and self-reported respondents. For allergens reported by doctor-diagnosed respondents, peaches (30.3%) was reported as the highest food allergen, followed by peanuts (21.3%), eggs (20.2%), and crab (13.8%). For allergens reported by self-reported respondents, eggs (33.3%) was reported as the most common allergen, followed by peaches (30.3%), milk (27.3%), peanuts (18.2%), and crab (18.2%). In addition, other allergenic foods such as kiwi, pineapple, taro, hazelnut, pine nuts, pistachio, walnut, and mushroom showed a rate of 11.2% among food allergenic respondents.

Comparison of attitudes regarding improvement of food allergen labeling and checking food labels according to food allergy status

Table 4 shows attitudes regarding improvement of food allergen labeling system based on food allergy status of the respondents. All respondents reported that all six items (bold font, font color, box frame, warning statement, front label, and addition of potential allergens) were necessary for an improved food allergen labeling system. However, there were no significant differences among doctor-diagnosed, self-reported, and non-allergenic groups. Table 5 presents differences in food labeling attitudes according to food allergy status of respondents. On

Table 3. Types of food allergens reported by doctor-diagnosed and self-reported respondents

Allergen	Doctor-diagnosed ¹⁾		Self-reported ²⁾		Total	
	n	%	n	%	n	%
Eggs	19	20.2	11	33.3	30	13.5
Cow's milk	15	16.0	9	27.3	24	10.8
Buckwheat	3	3.2	2	6.1	5	2.2
Peanuts	20	21.3	6	18.2	26	11.7
Soybeans	4	4.3	3	9.1	7	3.1
Wheat	1	1.1	1	3.0	2	.9
Mackerel	10	10.6	5	15.2	15	6.7
Crab	13	13.8	6	18.2	19	8.5
Shrimp	9	9.6	1	3.0	10	4.5
Pork	8	8.5	2	6.1	10	4.5
Peaches	29	30.9	10	30.3	39	17.5
Tomatoes	5	5.3	2	6.1	7	3.1
Sulfite	1	1.1	3	9.1	4	1.8
Other foods	21	22.3	4	12.1	25	11.2
Total	94	100.0	33	100.0	127	100.0

¹⁾ Non response (n = 1)

²⁾ Non response (n = 2)

average, all respondents gave middle (3/5 points) scores for all items, except food allergen label check item. For checking food allergen labels, there was a significant difference between allergic and non-allergic respondents ($P < 0.001$).

Table 4. Comparison of attitudes toward food allergen labeling improvement by food allergy status

	Doctor-diagnosed (n = 95)	Self-reported (n = 35)	Non-allergy (n = 413)	Total (n = 543)	P-value ³⁾
	Mean ¹⁾ ± SD ²⁾	Mean ± SD	Mean ± SD	Mean ± SD	
Bold font	4.03 ± 0.84	3.89 ± 0.93	4.07 ± 0.84	4.05 ± 0.85	0.449
Font color	4.05 ± 0.79	3.94 ± 0.80	4.07 ± 0.82	4.06 ± 0.81	0.669
Box frame	4.06 ± 0.77	3.83 ± 0.86	4.00 ± 0.85	4.00 ± 0.83	0.364
Warning statement	3.87 ± 0.89	3.51 ± 0.95	3.79 ± 0.95	3.79 ± 0.94	0.153
Front label	4.05 ± 0.79	3.89 ± 0.83	4.12 ± 2.17	4.09 ± 1.93	0.778
Addition of potential allergen	3.95 ± 0.90	3.74 ± 0.95	3.85 ± 0.87	3.86 ± 0.88	0.453

¹⁾ Mean response score (1: strongly unnecessary, 5: strongly necessary)

²⁾ SD = standard deviation

³⁾ P-value by One-way Anova

Table 5. Comparison of practice levels on food labels by food allergy status

	Doctor-diagnosed (n = 95)	Self-reported (n = 35)	Non-allergy (n = 413)	Total (n = 543)	P-value ³⁾
	Mean ¹⁾ ± SD ²⁾	Mean ± SD	Mean ± SD	Mean ± SD	
Brand	3.41 ± 1.28	3.71 ± 1.20	3.67 ± 1.32	3.63 ± 1.31	0.204
Nutrition facts	3.05 ± 1.25	2.94 ± 1.26	3.09 ± 1.30	3.08 ± 1.29	0.792
Origin	3.29 ± 1.22	3.69 ± 1.02	3.30 ± 1.30	3.33 ± 1.27	0.223
Allergenic ingredients	3.04 ± 1.34 ^a	3.03 ± 1.29 ^a	2.22 ± 1.20 ^b	2.41 ± 1.28	< 0.001
Food quality	3.01 ± 1.17	3.29 ± 1.20	3.01 ± 1.31	3.03 ± 1.28	0.476
Overall check	3.16 ± 0.97	3.33 ± 0.85	3.06 ± 0.94	3.10 ± 0.94	0.206

¹⁾ Mean response score (1: never check, 5: always check)

²⁾ SD = standard deviation

³⁾ P-value by One-way Anova

^{ab} Different superscript letters mean significantly different among groups at $\alpha = 0.05$ level by Duncan's multiple range test.

Table 6. VIP (Variable importance in the projection) values for each attribute by PLSR (Partial Least Square Regression) by correlating attitudes toward food allergen labeling improvement and practice levels on food labels by food allergy status.

Variable	PLS 1	PLS 2
Addition of potential allergen	0.848	1.116
Front label	1.041	1.066
Bold font	1.053	1.001
font color	1.057	0.960
Box frame	0.984	0.930
Warning statement	1.001	0.910




Fig. 1. PLSR analysis of food allergen labeling improvement and labeling attitudes

Partial Least Square Regression (PLSR) Analysis between food allergen labeling improvement and labeling attitudes of respondents

In Fig. 1, PLSR analysis was used to illustrate the correlation between food allergen labeling improvement and labeling attitudes of respondents. Partial Least Square Regression (PLSR) analysis analyzes the correlation between two data points and provides a visual summary [17-19]. The result shows that the doctor-diagnosed group and checking food allergens were positively correlated, whereas the non-allergy group was more concerned with checking brand of products. Self-reported group checked overall labeling information and especially origin and food quality. In PLSR analysis, all respondents answered that improvement of allergen labeling system is necessary. Regarding the effects of food allergen labeling improvement on labeling attitudes of respondents, variable importance in the projection (VIP) was calculated by PLSR analysis (Table 6). Food allergen labeling improvement with a VIP value greater than 1 had a large impact on labeling attitudes of respondents [20,21]. The factors of food allergen labeling improvement in

PLS 1 for affecting labeling attitudes of respondents were 'Front label', 'Bold font', 'Font color', and 'Warning statement', whereas the factors in PLS 2 affecting labeling attitudes of respondents were 'Addition of potential allergen', 'Front label', and 'Bold font'.

DISCUSSION

This study compared food labeling attitudes and food allergen labeling improvement by diagnosis of food allergies. The results show that prevalence of doctor-diagnosed food allergies was 17.5%, and 6.4% of respondents self-reported food allergies. The most common allergens of doctor-diagnosed and self-reported food allergy respondents were peaches (30.3%) and eggs (33.3%), respectively, followed by peanuts, cow's milk, and crab. For consumers' attitudes toward food labeling, the rate of checking food allergens was only significantly different between allergic and non-allergic respondents among all five items ($P < 0.001$). All respondents reported that all six items (bold font, font color, box frame, warning statement, front label, and addition of potential allergens) were necessary for an improved food allergen labeling system. PLSR analysis showed that the doctor-diagnosed group and checking food allergens were positively correlated, whereas the non-allergy group was more concerned with checking product brand.

Shah *et al.* [22] reported that the prevalence of self-reported food allergies ranges from 3 to 35% in America. On the other hand, the prevalence of self-reported food allergies is 20-25% of the population in Korea [23]. Lee *et al.* [5] surveyed 27,679 elementary, middle, and high school students in Korea and found that the prevalence of food allergies in students was 12.6% with doctor-diagnosed food allergies at 6.8%. Peaches (17.5%) was found to be one of the most common food allergens. Our results show a higher prevalence (23.9%) of food allergies than other studies [5,24]. The higher prevalence of food allergies in our study may be attributed to several factors. First, 27% of respondents were children and adolescents aged 19 or younger, who usually show higher prevalence of food allergies than adults. Second, some respondents might have been confused about food allergy/food intolerance due to misconceptions concerning accurate food allergy knowledge, self-diagnosis, and parental diagnosis. Third, some respondents may have been unaware of tolerance to food allergens (eggs and cow's milk) after diagnosis of food allergies at an earlier age. Forth, up to 60% of allergies from fruits and vegetables may actually be oral allergy syndrome instead of anaphylaxis shock in accordance with a previous study [4,6,25-28].

In our results, the most common allergens of doctor-diagnosed and self-reported food allergy respondents were peaches (30.3%) and eggs (33.3%). Lee *et al.* [12] surveyed 993 parents with allergic children and adults and investigated the prevalence of allergenic foods. It was found that peaches (18.6%) was the most common allergenic food, followed by eggs (14.2%), cow's milk and products (9.7%), and mackerel (8%) among adults. On the other hand, eggs (27%) was the most common allergenic food, followed by cow's milk and products (18.3%), peaches (10.4%), and shrimp (8.6%) among children. Lee *et al.*'s study [5] reported that the most common allergenic

foods were cow's milk and shrimp (7.0%), followed by eggs and mackerel (5.8%), peanuts and tomatoes (3.5%), wheat and crab (2.3%), and pork, peaches, and soybean (1.2%).

Over 160 allergenic foods have been reported worldwide. In addition, the variety of allergenic foods is usually determined by age, ethnicity, and country. For children, the most common allergenic foods are eggs, cow's milk, peanuts, seafood, and tree nuts. Among adults, peanuts are probably the most common allergenic food, followed by tree nuts, seafood, and crustaceans (shrimp, crab, and lobster) [6,8]. Allergies toward cow's milk or eggs are usually outgrown during childhood or adolescence [29]. A study on the prevalence and main characteristics of food allergenic school children in France reported that 62 children (25.4%) among 182 food allergenic respondents outgrew their food allergies [30]. Peanut allergy is more common in America than in other countries due to the popularity of peanut butter. Likewise, in Southeast Asian countries such as Japan and Korea, buckwheat allergy is more common than in other parts of the world since buckwheat noodles and products are popular. This means that specific foods causing food allergies may be affected by cultural dietary habits [6,8,31]. Our results also demonstrate other specific foods such as kiwi, pineapple, taro, hazelnut, pine nuts, pistachio, walnut, and mushroom as prevalent causes of food allergies among the Korean population. Therefore, clinical research and allergenic food labeling improvement should be performed to investigate specific foods that generate less frequent but severe reactions [6].

In our results, respondents reported that current allergenic labeling practices such as readability (bold font and font color) and visibility of information (box frame and label on front of product) need to be improved. A prior study on public awareness of food allergenic labeling system investigated 993 subjects (337 food allergic and 656 non-allergic consumers) from seven university hospitals in Korea. The results showed that 23.5% of respondents checked food labels, and 46.6% of doctor-diagnosed and 31.8% of self-reported allergic consumers checked allergenic ingredients on labels. Moreover, 73.1% of respondents reported that they were not satisfied with current food allergic labeling practices. Common complaints were 'unclear and too technical ingredient list and allergen information', 'too small font size or undistinguishable color contrast of label for allergen ingredients', and 'inappropriate and insufficient information' [12]. In another study [13], a total of 40 food allergic consumers who suffered from one or more common allergens such as milk, eggs, and peanuts/tree nuts in the Netherlands and Greece were asked for their preferences for current food labeling practices and information. The results showed that the current food labeling system does not meet the needs of food allergic consumers. The main problems were 'label appearance for readability' and 'contents of the ingredient list on the label'. They also reported that some ingredient lists include too much information as well as too technical terms.

In conclusion, an effective food labeling system is very important for health protection of food-allergic consumers. Additionally, government agencies must develop policies regarding prevalence of food allergies in Korea. Based on this information, the food industry and government agencies should provide clear and accurate food labeling practices for consumers.

REFERENCES

- Gupta RS, Springston EE, Warrier MR, Smith B, Kumar R, Pongracic J, Holl JL. The prevalence, severity, and distribution of childhood food allergy in the United States. *Pediatrics* 2011;128:e9-17.
- Sicherer SH. Epidemiology of food allergy. *J Allergy Clin Immunol* 2011;127:594-602.
- Rona RJ, Keil T, Summers C, Gislason D, Zuidmeer L, Sodergren E, Sigurdardottir ST, Lindner T, Goldhahn K, Dahlstrom J, McBride D, Madsen C. The prevalence of food allergy: a meta-analysis. *J Allergy Clin Immunol* 2007;120:638-46.
- Soller L, Ben-Shoshan M, Harrington DW, Fragapane J, Joseph L, St Pierre Y, Godefroy SB, La Vieille S, Elliott SJ, Clarke AE. Overall prevalence of self-reported food allergy in Canada. *J Allergy Clin Immunol* 2012;130:986-8.
- Lee AH, Kim KE, Lee KE, Kim SH, Wang TW, Kim KW, Kwak TK. Prevalence of food allergy and perceptions on food allergen labeling in school foodservice among Korean students. *Allergy Asthma Respir Dis* 2013;1:227-34.
- Taylor SL, Hefle SL. Food allergies and other food sensitivities. *Food Technol* 2001;55:68-83.
- Sampson HA, Muñoz-Furlong A, Campbell RL, Adkinson NF Jr, Bock SA, Branum A, Brown SG, Camargo CA Jr, Cydulka R, Galli SJ, Gidudu J, Gruchalla RS, Harlor AD Jr, Hepner DL, Lewis LM, Lieberman PL, Metcalfe DD, O'Connor R, Muraro A, Rudman A, Schmitt C, Scherrer D, Simons FE, Thomas S, Wood JP, Decker WW. Second symposium on the definition and management of anaphylaxis: summary report—second National Institute of Allergy and Infectious Disease/Food Allergy and Anaphylaxis Network symposium. *Ann Emerg Med* 2006;47:373-80.
- Lee SY. IgE mediated food allergy in Korean children: focused on plant food allergy. *Asia Pac Allergy* 2013;3:15-22.
- Gendel SM. Comparison of international food allergen labeling regulations. *Regul Toxicol Pharmacol* 2012;63:279-85.
- Zurzolo GA, Mathai ML, Koplin JJ, Allen KJ. Precautionary allergen labelling following new labelling practice in Australia. *J Paediatr Child Health* 2013;49:E306-10.
- Van Hengel AJ. Declaration of allergens on the label of food products purchased on the European market. *Trends Food Sci Technol* 2007;18:96-100.
- Lee SY, Kim KW, Ahn K, Kim HH, Pyun BY, Park YM, Kim KE. Consumer's use and satisfaction of allergic food labels. *Pediatr Allergy Respir Dis* 2011;21:294-301.
- Cornelisse-Vermaat JR, Voordouw J, Yiakoumaki V, Theodoridis G, Frewer LJ. Food-allergic consumers' labelling preferences: a cross-cultural comparison. *Eur J Public Health* 2008;18:115-20.
- Son KY, Park KS, Hwang HH, Yun BS, Lee SJ, Kim MA, Park JY, Kim KE, Jang KC. Prevalence of allergic diseases among primary school children in Ilsan, Gyeonggi and changes of symptoms after environmental control in 2005. *Pediatr Allergy Respir Dis* 2007;17:384-93.
- Ajala AR, Cruz AG, Faria JA, Walter EH, Granato D, Sant' Ana AS. Food allergens: knowledge and practices of food handlers in restaurants. *Food Control* 2010;21:1318-21.
- Jee HM, Kim KW, Kim CS, Sohn MH, Shin DC, Kim KE. Prevalence of asthma, rhinitis and eczema in Korean children using the International Study of Asthma and Allergies in Childhood (ISAAC) Questionnaires. *Pediatr Allergy Respir Dis* 2009;19:165-72.
- Chung SJ, Heymann H, Grün IU. Application of GPA and PLSR in

- correlating sensory and chemical data sets. *Food Qual Prefer* 2003;14:485-95.
18. Chung L, Chung SJ. Understanding the factors affecting the acceptance for fermented soybean products. *Food Sci Biotechnol* 2008;17:144-50.
 19. Gwak MJ, Chung SJ, Kim Y. Sensory drivers of liking for adlay (*Coix lacryma-jobi*) tea. *Korean J Food Cult* 2012;27:512-20.
 20. Azizan KA, Baharum SN, Resson HW, Noor NM. GC-MS analysis and PLS-DA validation of the trimethyl silyl-derivatization techniques. *Am J Appl Sci* 2012;9:1124-36.
 21. Kwak HS, Oh YJ, Kang HB, Kim TH. Descriptive profile and liking/disliking factors for aseptic-packaged rice porridge. *J Korean Soc Food Sci Nutr* 2013;42:1878-85.
 22. Shah SS, Parker CL, Davis CM. Improvement of teacher food allergy knowledge in socioeconomically diverse schools after educational intervention. *Clin Pediatr (Phila)* 2013;52:812-20.
 23. Seo WH, Jang EY, Han YS, Ahn KM, Jung JT. Management of food allergies in young children at a child care center and hospital in Korean. *Pediatr Allergy Respir Dis* 2011;21:32-8.
 24. Jung YH, Ko H, Kim HY, Seo JH, Kwon JW, Kim BJ, Kim HB, Lee SY, Jang GC, Song DJ, Kim WK, Shim JY, Hong SJ. Prevalence and risk factors of food allergy in preschool children in Seoul. *Korean J Asthma Allergy Clin Immunol* 2011;31:177-83.
 25. Bock SA, Lee WY, Remigio LK, May CD. Studies of hypersensitivity reactions to foods in infants and children. *J Allergy Clin Immunol* 1978;62:327-34.
 26. Sloan AE, Powers ME. A perspective on popular perceptions of adverse reactions to foods. *J Allergy Clin Immunol* 1986;78:127-33.
 27. Burks AW, Jones SM, Boyce JA, Sicherer SH, Wood RA, Assa'ad A, Sampson HA. NIAID-sponsored 2010 guidelines for managing food allergy: applications in the pediatric population. *Pediatrics* 2011;128:955-65.
 28. Zuidmeer L, Goldhahn K, Rona RJ, Gislason D, Madsen C, Summers C, Sodergren E, Dahlstrom J, Lindner T, Sigurdardottir ST, McBride D, Keil T. The prevalence of plant food allergies: a systematic review. *J Allergy Clin Immunol* 2008;121:1210-1218.e4.
 29. Skypala I. Adverse food reactions--an emerging issue for adults. *J Am Diet Assoc* 2011;111:1877-91.
 30. Rancé F, Grandmottet X, Grandjean H. Prevalence and main characteristics of schoolchildren diagnosed with food allergies in France. *Clin Exp Allergy* 2005;35:167-72.
 31. Wieslander G, Norbäck D. Buckwheat allergy. *Allergy* 2001;56:703-4.