

## RESEARCH ARTICLE

# Who is seeking help for psychological distress associated with the COVID-19 pandemic? Characterization of risk factors in 1269 participants accessing low-threshold psychological help

Kevin Hilbert <sup>\*</sup>, Ole Boeken, Julia Asbrand, Sophia Seemann, Till Langhammer, Berit Praxl, Leonore Horváth, Andrea Ertle, Ulrike Lueken

Department of Psychology, Humboldt-Universität zu Berlin, Berlin, Germany

\* [Kevin.Hilbert@hu-berlin.de](mailto:Kevin.Hilbert@hu-berlin.de)



## OPEN ACCESS

**Citation:** Hilbert K, Boeken O, Asbrand J, Seemann S, Langhammer T, Praxl B, et al. (2022) Who is seeking help for psychological distress associated with the COVID-19 pandemic? Characterization of risk factors in 1269 participants accessing low-threshold psychological help. PLoS ONE 17(7): e0271468. <https://doi.org/10.1371/journal.pone.0271468>

**Editor:** Yann Benetreau, Public Library of Science, UNITED STATES

**Received:** November 4, 2021

**Accepted:** June 29, 2022

**Published:** July 18, 2022

**Copyright:** © 2022 Hilbert et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** Data cannot be shared publicly as this is not included in the informed consent by participants and the mental health data is particularly sensitive. However, deidentified participant data with annotations will be made available to other researchers upon reasonable request (towards the study administration at [corona-stressfrei.psychologie@hu-berlin.de](mailto:corona-stressfrei.psychologie@hu-berlin.de)). Additionally, a registered study protocol is available at the German

## Abstract

### Background

The COVID-19 pandemic and accompanying restrictions are associated with substantial psychological distress. However, it is unclear how this increased strain translates into help-seeking behavior. Here, we aim to characterize those individuals who seek help for COVID-19 related psychological distress, and examine which factors are associated with their levels of distress in order to better characterize vulnerable groups.

### Methods

We report data from 1269 help-seeking participants subscribing to a stepped-care program targeted at mental health problems due to the COVID-19 pandemic. Sample characteristics were compared to population data, and linear regression analyses were used to examine which risk factors and stressors were associated with current symptom levels.

### Results

Seeking for help for COVID-19 related psychological distress was characterized by female gender, younger age, and better education compared to the general population. The majority reported mental health problems already before the pandemic. 74.5% of this help-seeking sample also exceeded clinical thresholds for depression, anxiety, or somatization. Higher individual symptom levels were associated with higher overall levels of pandemic stress, younger age, and pre-existing mental health problems, but were buffered by functional emotion regulation strategies.

### Conclusions

Results suggest a considerable increase in demand for mental-healthcare in the pandemic aftermath. Comparisons with the general population indicate diverging patterns in help-

Clinical Trial Register (DRKS00023220; <https://www.drks.de>), a methods paper describing the overall research project is also available (16).

**Funding:** The authors received no specific funding for this work.

**Competing interests:** I have read the journal's policy and the authors of this manuscript have the following competing interests: KH, OB, SS, TL, LH, AE, JA and UL declare no competing interests. BP declares having performed paid freelance work for the Mental Tech GmbH and having stocks or stock options for the same company. The Mental Tech GmbH provided the chatbot program 'Aury' that was offered in step one of the stepped-care program (data not included in this manuscript). This does not alter our adherence to PLOS ONE policies on sharing data and materials.

seeking behavior: while some individuals seek help themselves, others should be addressed directly. Individuals that are young, have pre-existing mental health problems and experience a high level of pandemic stress are particularly at-risk for considerable symptom load. Mental-healthcare providers should use these results to prepare for the significant increase in demand during the broader aftermath of the COVID-19 pandemic as well as allocate limited resources more effectively.

## Introduction

Soon after the emergence of COVID-19 and its pandemic spreading, first reports indicated that this disease would not only cause a medical but a mental health crisis as well [1, 2]. A subsequent, considerable corpus of work demonstrated an increase in general psychological distress and a wide range of symptoms including anxiety and depression due to the pandemic's burden (as reviewed in [3]). Overall, a recent meta-regression estimated a global increase of 27.6% and 25.6% for major depressive disorder and anxiety disorders, respectively [4].

It is crucial to point out, that many mental health problems will occur with considerable delay and a substantial share of affected individuals may experience a relatively chronic course [5–7]. This indicates an increase in the need for mental healthcare in the near future and indeed, first reports seem to confirm this notion: in Germany, a survey conducted by a professional body of psychotherapists reported an increase in demand for an initial psychotherapy session of 40.8% in January 2021 compared to January 2020 [8].

Following diathesis-stress models, individual burden and symptoms emerges by the interplay between situational stressors and pre-existing vulnerability which can be modified by the ability to successfully regulate emotions [9]. For the COVID-19 pandemic, first results suggest that duration of quarantine and—more broadly—social isolation, financial difficulties as well as uncertainty may be central stressors [10, 11]. Further, female sex, younger age, low education level and unemployment, and pre-existing mental disorders may be personal dispositions signaling increased risk [11–15]. However, it remains unclear who will seek help for mental problems associated with the pandemic. Therefore, reports on generally increased distress and mental health problems due to the COVID-19 pandemic need to be complemented by data describing which individuals actually seek help. A characterization of such help-seeking samples, including a comparison to representative samples, might shed light on specifics of help-seeking individuals. Furthermore, an examination of which factors help-seeking individuals associate with their levels of distress may be informative on further characterizing vulnerability factors.

Such information will help to allocate limited resources in mental healthcare efficiently and to plan for upcoming treatment-demand. Accordingly, we used baseline data from an help-seeking sample accessing a stepped-care program designed to alleviate psychological distress due to the COVID-19 pandemic. We examined sociodemographic characteristics, the presence of a range of risk-factors as well as stressors. In addition, we compared core characteristics with population scores from government data and representative studies in Germany in order to characterize this help-seeking sample. We examined which characteristics were associated with symptom severity at the start of this program in order to examine which factors they associate with their levels of distress. We hypothesized that the help-seeking sample will be predominantly female and well-educated. We also expected that their level of distress will be associated with pre-existing psychopathology and that they have had more exposure to

adverse experiences in the pandemic. In contrast, high levels of social support and good emotion regulation abilities were expected to be buffering factors.

## Materials and methods

### Trial design

Here, we report cross-sectional data from a sample of help-seeking individuals participating in a stepped-care program against psychological distress due to the COVID-19 pandemic at study-entry ('baseline'), before any interventions were conducted.

The overall study was planned as a longitudinal observational study following participants over the course of two years [16]. A stepped-care program including a first online intervention and, if necessary, a second face-to-face digital group program was offered. After completing a baseline assessment at study entry, participants were invited to use the online intervention (chatbot program 'Aury') which was specifically adapted to the pandemic. Aury provided information and low-threshold interventions for symptoms (e.g. sleeping disorders, depression, anxiety, rumination, interpersonal conflicts) and activated individual resources. After a four week-waiting period, participants with lasting symptoms entered the second step of a 6-week group-based prevention program conducted by cognitive behavioral therapists. Participants were invited for 6 and 12 month follow-ups (still ongoing). All procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2013. The study protocol was registered at the German Clinical Trial Register (DRKS00023220; <https://www.drks.de>) and approved by the ethics committee of the Department of Psychology at Humboldt-Universität zu Berlin (#2020–35).

### Sample

The stepped-care program ("Stressfrei nach Corona: ein psychologisches Hilfsprogramm der Humboldt-Universität zu Berlin": [www.corona-stressfrei.de](http://www.corona-stressfrei.de)) was promoted via official press releases and mailing lists of the Humboldt-Universität zu Berlin, social media (Instagram, Facebook). We further dispatched information material to mental health care providers and email contact to Corporate Health Management offices in the Berlin metropolitan area. Additionally, the program was promoted on local and nationwide newspaper, radio and TV interviews.

We examined the baseline data of adults aged 18 or older registering for the stepped-care program from the beginning of the program, September 1, 2020 until the end of the recruitment phase, April 15, 2021. There were  $n = 1752$  registrations of which  $n = 421$  were excluded as incomplete.  $N = 62$  were excluded as double registrations, resulting in a final analysis sample of  $n = 1269$  participants. For those who started or completed the baseline assessment more than once, we selected the earliest complete dataset. We followed the principles of the Declaration of Helsinki, whereby all subjects gave their written informed consent for participation, data collection, analysis and publication.

### Materials and procedures

Participants signed up to the study by providing their email-address to a double-opt-in procedure, and in turn received a link to the baseline assessment. The email-address was used to re-contact participants over the course of the study but was kept separate of any other information. During the baseline assessment, participants generated an individual code for data pseudonymization. They provided sociodemographic information such as age, sex, children and

education. Additionally, participants completed the following questionnaires before receiving a link to the first step of the stepped-care program.

**Patient Health questionnaire (PHQ).** Symptom severity was assessed using three subscales from the PHQ [17]: the PHQ-9 for depressive symptoms, the GAD-7 for anxiety symptoms and the PHQ-15 for somatization. These scales have demonstrated good psychometric properties [17–20], higher scores equal higher symptom load.

**Short-form-8 (SF-8).** The SF-8 [21] was used to assess health-related quality of life as a secondary outcome. It assesses impairment due to and satisfaction with mental and physical wellbeing with good psychometric properties [21]. Higher scores denote more impairment and lower quality of life.

**COVID-19 pandemic events questionnaire.** A customized collection of items on potential psychological distress associated with the COVID-19 pandemic was developed in-house after the beginning of the pandemic. Items were created based on literature on psychological distress due to natural disasters, previous pandemics, social isolation and quarantine. Items cover distress due to fear and worry about somatic health, due to social isolation and contact restrictions, due to financial or work-related burdens, due to increased child-care or home-schooling, and due to previous mental health problems. Items focused on the first wave of the COVID-19 pandemic in Germany. Additionally, the presence or absence of a variety of emotional states was checked for the time of the first wave of the COVID-19 pandemic in Germany and for the time of the baseline assessment. All items are given in [S1 Table](#).

**Emotion Regulation Questionnaire (ERQ).** The ERQ [22] was used to assess emotion regulation as a form of coping. It assesses the strategies reappraisal and suppression and showed good psychometric properties.

## Analysis

Data were screened for improbable scores and outliers, both of which were set to missing values, some variables were recoded (details are given in [S1 File](#)). We calculated a sum score on the overall level of pandemic stress based on individual items on situation stressors with higher scores indicating more stress (for details see also [S1 File](#)). Descriptive statistics were used to examine the sample in detail. We used McNemar's test to examine whether affective states were currently being reported more frequently than during the first wave of the COVID-19 pandemic. Results were Bonferroni-corrected to account for the high number of affective states.

Comparisons of sample characteristics to population data were done using one-sample t-tests and one-sample chi-square tests as appropriate. Population data was drawn from census data, census extrapolations and representative surveys provided by the German Federal Statistical Office, from statistics by the German Federal Employment Agency and the Robert Koch Institute, which is the German body for monitoring the COVID-19 outbreak (see [S3 Table](#) for a detailed description of this data and all data sources).

Separate multiple stepwise linear regression analyses were used in order to examine which risk factors and stressors were associated with mental health problems as indicated by the PHQ-9, GAD-7 and PHQ-15 scales. We used forward selection and decided on a threshold of  $p < 0.01$  for inclusion given the high number of potentially associated variables. All analyses were conducted in SPSS 26 (IBM, Armonk, NY, USA), with the level of significance set at  $p < 0.05$ .

## Results

### Overall description of help-seeking sample

[S2 Table](#) provides a detailed overview on the raw data for all sociodemographic characteristics and responses to all potential stressors. The help-seeking sample of participants was mainly

female (79.4%), well educated (87.1% with highest secondary school or better) and only rarely unemployed (6.6%). Slightly more than half of the participants were in a relationship and slightly less than half of the participants had a child or children. When examining responses to the potential COVID-19 related stressors, marked differences between mean ratings appeared: while participants were less concerned about their individual COVID-19 related risk, they worried substantially about potential threat to friends and relatives. Contact restrictions were perceived as the strongest stressor, while experiencing a burden at work because of the pandemic was common as well. On the contrary, financial worries were rather uncommon in the sample.

Regarding their mental health, the majority of participants (54.5%) reported having mental health problems before the pandemic already. [S1 Fig](#) shows the results from coding the previous mental health problems participants reported in their free responses, indicating that sub-clinical and clinical anxiety and depression were by far the most common. Crucially, the vast majority of participants with previous mental health problems reported a further deterioration of their mental health due to the pandemic, only few participants reported benefits or no changes ([Fig 1](#)). Moreover, a majority of 59.3% of participants reported new mental health problems caused by stress due to the pandemic. For both old and new mental health problems, the lack of support was more common than the availability of support.


The severity of depression and somatization was rated higher than the severity of worry symptoms (means and standard deviations: PHQ-9: 11.3 (5.7), GAD-7: 9.7 (5.0), PHQ-15: 11.3 (5.4)). On all three scales, substantial proportions of participants scored ten or above, which is commonly used as cut off [17]: PHQ-9: 58.2%, GAD-7: 47.7%, PHQ-15: 58.3% ([Fig 1](#)). 74.5% scored ten or above in at least one of these scales. Impaired quality of life as measured by the SF-8 was substantial (mean and standard deviation: 23.7 (5.9)). A variety of additional symptoms were reported, with worry, rumination, physical arousal, restlessness and sleep problems being the most frequent ones ([Fig 1](#)). Positive experiences such as life slowing down were rare in the sample. All symptoms and experiences were more frequent during the first wave of the pandemic compared to the current assessment which took place during the second to third wave.

### Comparison of help-seeking sample with population data

The participating sample differed in many characteristics from population averages. Participants were considerably more female, younger, and better educated than the general population (see [Table 1](#)). No differences to the population emerged when examining the average size of the flat or house. Being unmarried and separated was more frequent, being married, divorced or widowed less frequent in the participating sample compared to the population. More participants reported on having two children or children in general at home and fewer reported on having no children at home, compared to the general population. Substantially more participants worked in the health sector than common for the population, while fewer participants reported a verified COVID-19 infection.

### Contribution of vulnerability factors and pandemic-related stressors on symptom severity

Multiple stepwise linear regression analyses on each PHQ subscale explained substantial proportions of the outcome variance (PHQ-9:  $R^2_{\text{corr}} = 0.40$ , GAD-7:  $R^2_{\text{corr}} = 0.34$ , PHQ-15:  $R^2_{\text{corr}} = 0.30$ ). [Table 2](#) presents the associated variables for each regression with their standardized coefficients. General and specific patterns emerged for the associated variables. The overall level of pandemic stress and pre-existing mental health problems were the most important



**Fig 1. Ratings of psychological distress and clinical scores.** Change in mental health problems due to the pandemic in the sample (valid answers only, without N/As; upper). Percentage of participants scoring ten or above across the primary outcomes (middle). Percentage of affective states reported as occurring during the first wave of the COVID-19 pandemic and currently (all categories negative except for the right outer two, lower). Here, all comparisons between both time-points were highly significant (\*\*\*)  $p < 0.001$ , all Bonferroni-corrected).

<https://doi.org/10.1371/journal.pone.0271468.g001>

variables associated with increasing symptom severity for all primary outcomes. Moreover, the development of new mental health problems due to the pandemic was associated with higher symptom severity as well. Pre-existing depressiveness and pre-existing depressive disorders were associated with PHQ-9 severity, while the presence of multiple pre-existing mental health problems and younger age were associated with PHQ-9 severity and GAD-7 severity. Worries about not getting adequate medical care for other conditions were associated with PHQ somatization. Interestingly, higher reappraisal in emotion regulation as assessed by the ERQ was associated with lower PHQ-9 and GAD-7 scores.

### Discussion

Plenty of studies reported increased psychological distress and mental health problems associated with the COVID-19 pandemic for considerable proportions of the population. Yet, it is unclear which individuals will subsequently seek help for these problems and which factors are central drivers of distress in such individuals. Examining a help-seeking sample of 1269 participants from an easily-accessible stepped-care program against COVID-19-induced psychological distress, the extent of depression and anxiety symptoms in our help-seeking sample was substantial with 74.5% of participants scoring above at least one of the cutoff scores, thus indicating a clinically relevant symptom load. We found support for our hypothesis that



Table 1. Comparison of sample characteristics to population values, numbers per category except where noted.

	Sample		Population		Comparison	
	mean / N	sd / %	mean / N	sd / %	Chi-square / t	p
<i>sex</i>						
					456.348	<0.001
female	1008	(79.4)	42,052,500	(50.7)		
male	238	(18.8)	40,966,700	(49.3)		
<i>age</i>						
	42.1	(13.3)	44.4	N/A	-6.099	<0.001
<i>education status</i>						
					1631.960	<0.001
lowest secondary school	19	(1.5)	N/A <sup>1</sup>	(29.6)		
inter-mediate secondary school	144	(11.3)	N/A <sup>1</sup>	(29.9)		
highest secondary school (including university degree and PhD)	1083	(85.3)	N/A <sup>1</sup>	(32.5)		
other	23	(1.8)	N/A <sup>1</sup>	(7.8)		
<i>relationship status</i>						
					141.151	<0.001
unmarried	692	(55.4)	34,661,000	(41.9)		
married	433	(34.1)	35,351,000	(42.7)		
separated	44	(3.5)	1,725,000	(2.1)		
divorced	61	(4.8)	5,727,000	(6.9)		
widowed	18	(1.4)	5,321,000	(6.4)		
<i>working in the health sector</i>						
					35.598	<0.001
yes	193	(16.2)	3,657,135	(10.8)		
no	1001	(83.8)	30,134,730	(89.2)		
<i>number of children at home</i>						
					35.224	<0.001
none	853	(67.3)	N/A <sup>1</sup>	(72.1)		
one	182	(14.4)	N/A <sup>1</sup>	(14.2)		
two	192	(15.2)	N/A <sup>1</sup>	(10.2)		
three or more	40	(3.2)	N/A <sup>1</sup>	(3.5)		
<i>housing space (in m<sup>2</sup>)</i>						
	93.4	(48.8)	94.1	N/A	-0.547	0.585
<i>COVID-19 status</i>						
					5.078	0.024
without COVID-19 diagnosis	1217	(97.5)	80,081,589	(96.3)		
with COVID-19 diagnosis	31	(2.5)	3,073,442	(3.7)		

<sup>1</sup> Absolute numbers for the population were not available, but percentage data was. Chi-square tests were done by comparing the actual number of participants per category in the sample with the expected number of participants given the population percentages. Categories are not raw data, but have been modified in order to match the categories for which population data was available. This means that some categories were dropped for this comparison (such as the *diverse* category for *gender*) while others have been recoded (such as the *highest secondary school* category, which here includes also the *university degree* and *PhD* entries that had their own categories in the raw data). More detail is available in supplement D.

<https://doi.org/10.1371/journal.pone.0271468.t001>

particularly well-educated women seek psychological support, however, the help-seeking sample also differed from the general population on additional dimensions such as being younger, less frequently married and more having children at home. Partly confirming our second hypotheses, individual psychological distress in this help-seeking sample was associated with pre-existing psychopathology and more exposure to adverse experiences in the pandemic, while emotion regulation (reappraisal) showed a buffering effect.

In line with our expectations, help-seeking individuals as identified by participating in our program were considerably more often female and well-educated. Beyond our hypotheses, we also found the study participants to be younger, more often unmarried and with children at home, more often working in healthcare and less often personally affected by a verified COVID-19 infection than in the general population. Results on sex and age match previous reports on groups who are at increased risk for psychological distress due to the pandemic [11–13]. However, other results stand in contrast to these reports: high education status, for

**Table 2. Final stepwise regression models on the PHQ scores.**

Variable	Coefficient			Modell	
	Standardized Beta	T	p	R	R <sup>2</sup> <sub>corr</sub>
<i>PHQ-9 Depression</i>					
Overall level of pandemic stress	0.243	7.138	<0.001		
Suffering from mental health problems before the pandemic	0.169	5.693	<0.001		
ERQ: reappraisal sum score	-0.154	-6.855	<0.001		
Developed mental health problems due to pandemic: yes	0.145	6.181	<0.001		
Mental-health problems before pandemic: Depressive Disorder	0.140	5.306	<0.001		
Found work meaningful or relevant during the pandemic	-0.099	-4.135	<0.001		
ERQ: suppression sum score	0.109	4.884	<0.001		
Age	-0.101	-4.366	<0.001		
Impression that social relationships have suffered long-term damage	0.116	4.365	<0.001		
Relationship status: married	-0.065	-2.818	0.005		
Mental-health problems before pandemic: depressiveness	0.088	3.786	<0.001		
Relationship status: separated	0.069	3.090	0.002		
Mental health problems have changed since the pandemic*	-0.081	-2.834	0.005		
Worried about financial situation	0.132	3.820	<0.001		
Financial losses due to the pandemic	-0.090	-2.798	0.005		
Verified COVID-19 illness: yes	0.058	2.630	0.009		
Mental-health problems before pandemic: multiple problems	0.065	2.591	0.010		
				0.639	0.401
<i>PHQ GAD-7</i>					
Overall level of pandemic stress	0.356	11.515	<0.001		
Suffering from mental health problems before the pandemic	0.197	8.245	<0.001		
Developed mental health problems due to pandemic: no	-0.113	-3.265	0.001		
ERQ: reappraisal sum score	-0.176	-7.536	<0.001		
Age	-0.117	-5.053	<0.001		
Mental-health problems before pandemic: multiple problems	0.096	4.078	<0.001		
Developed mental health problems due to pandemic: yes	0.126	3.655	<0.001		
Burden felt at work because of the pandemic	-0.091	-3.050	0.002		
				0.586	0.339
<i>PHQ-15 Somatoform</i>					
Overall level of pandemic stress	0.341	11.717	<0.001		
Suffering from mental health problems before the pandemic	0.140	5.530	<0.001		
Had to work from home	-0.081	-3.188	0.001		
Developed mental health problems due to pandemic: yes	0.112	4.514	<0.001		
Worried about not getting adequate medical care for other conditions	0.108	3.987	<0.001		
Occupational status: self-employed	-0.080	-3.332	<0.001		
Sex: male	-0.100	-4.100	<0.001		
Mental-health problems before pandemic: Depressive Disorder	0.084	3.355	<0.001		
Education: PhD	-0.096	-3.905	<0.001		
Worried about having Covid-19, but no test was done or not done in time: no, not worried	-0.071	-2.973	0.003		
Education: university degree	-0.068	-2.691	0.007		
ERQ: suppression sum score	0.064	2.620	0.009		
				0.555	0.308

Total n = 1261.

PHQ-9: Patient Health Questionnaire-9 depression scale; GAD-7: Generalized Anxiety Disorder Scale-7; PHQ-15: Patient Health Questionnaire-15 somatic symptom scale; ERQ: Emotion Regulation Questionnaire.

\*Change to the worse coded with lower numbers and change to the better coded with higher numbers, i.e. subjective change to the worse predicted elevated PHQ-9 scores.

<https://doi.org/10.1371/journal.pone.0271468.t002>



instance, has rather been reported as a protective factor. Note that these results are likely also influenced by diverging patterns in help-seeking behaviors. For instance, younger, female and better educated individuals are also overrepresented in routine psychotherapy compared to their proportion in the general population of Germany. An important implication of this finding is that those individuals who seek help for their COVID-19 related psychological distress by themselves are most likely only a portion of all affected individuals. Individuals from certain groups such as those with low educational status may be particularly hard to reach. As such, they may benefit from a broader public health approach addressing them more directly in order to mitigate the individual burden and societal cost of the pandemic's effect on mental health. Overall, it is important to note that three quarters of our help-seeking sample showed symptoms of anxiety and depression to clinically relevant degree. This considerably exceeds the percentages above cut-offs in the general population (ca. 6.6% for the PHQ-9, 5.1% for the GAD-7, and 9.3% for the PHQ-15; [23–25]). These figures underscore the unmet need for people suffering from mental health problems in the aftermath of the pandemic. Expecting a certain amount of these persons to request professional help is in line with reports from German professional organizations of a more than 40% increase in demand for regular psychotherapy [8]. In summary, this data suggests considerable unmet needs in mental-healthcare during the aftermath of the COVID-19 pandemic. Therefore, a call for low-threshold and scalable psychological interventions for indicated prevention or early intervention in this target group seems warranted. A number of risk factors and stressors were associated with individual symptom severity within this sample of help-seeking individuals. The results indicate that diathesis-stress models of psychological distress also apply to the unique pandemic situation: symptom levels were strongly influenced by the overall level of pandemic stress and by the occurrence of relevant risk factors. Some findings are particularly notable. In agreement with previous research [14, 15], our data indicates that individuals with pre-existing mental health problems are a major high-risk group for psychological distress due to the pandemic and therefore should be monitored closely if possible. This could occur for instance during general practitioner or psychiatrist visits. Second, reappraisal emotion regulation abilities were found to be important buffers of symptom levels, although this was only true for depression and anxiety but not for somatization. As expected, reappraisal as a functional emotion regulation strategy was associated with lower symptom levels, contrary to suppression. This finding may be exploited in order to mitigate the psychological distress due to this pandemic and future similar events by encouraging the use of low-level (digital) interventions targeting individual emotion regulation abilities, which are becoming more and more available and are tested for efficacy [26]. Third, contrary to our hypotheses, the availability of social support for mental health problems was not associated with symptom levels. However, being married was associated with low symptom levels and being separated was associated with elevated symptom levels for depression. This finding may signify the lack of availability of some sort of support in close relationships. Thus, relationship status may be another sociodemographic factor associated with increased risk for high psychological distress from the disorder. In addition, young age was associated with elevated symptom levels across all PHQ subscales except somatization.

The current investigation complements previous surveys with data from a help-seeking sample of substantial size. The sample is well characterized regarding a variety of sociodemographic characteristics and pandemic-related stressors, enabling both a comparison to representative population data and the explanation of considerable proportions of variance in individual symptoms levels. However, there are also limitations. While the study was advertised on national newspapers and television occasionally, recruitment largely focused on the Berlin metropolitan area. Although we did not record the place of residence for our participants, it seems likely that ours is a relatively urban sample. This may influence some findings

such as the low average age of the sample. Additionally, the assessment of pandemic-related stressors focused on the circumstances during the first major lockdown in Germany. It is possible that further important factors affecting individual symptom levels occurred afterwards, or that participants deviated from this focus in their answers. Finally, when comparing the sample to representative population data, both sources used different data collection methods and response categories. For instance, 'diverse' was not available as a response category for sex in the population data. As a result, we had to compare data which was not completely similar. However, given that significant differences to the population data were very pronounced, this limitation appears to not have had strong consequences for the interpretation of results.

In conclusion, we here report the characteristics from a help-seeking sample including more than one-thousand participants applying for a stepped-care program targeting psychological distress due to the COVID-19 pandemic. Supplementing previous research, we found that help-seeking behavior was associated with both risk-factors for pandemic-related psychological distress and established patterns in help-seeking behaviors. This indicates that only a subset of individuals with psychological distress will proactively seek help by themselves, while other vulnerable groups (e.g. low socio-economic status, low education background) may need to be addressed more directly. Individual symptom levels were significantly higher among this sample compared to the general population, suggesting an emerging and possibly increasing gap between supply and demand for mental-healthcare in the pandemic aftermath. Moreover, individual symptom levels clearly follow the diathesis-stress approach: individuals of younger age, who have pre-existing mental health problems and experience a high level of pandemic stress are particularly at-risk for a considerable symptom load. This may in turn manifest in either relapse or new incidence of mental disorders if untreated. In summary, these data suggest considerable unmet needs in mental-healthcare in the aftermath of the COVID-19 pandemic and call for low-threshold and scalable psychological interventions for indicated prevention or early intervention in this target group. Providers of mental-healthcare should use these results to prepare for a significant increase in demand of psychological help in the broader aftermath of the COVID-19 pandemic and to allocate limited resources most effectively.

## Supporting information

**S1 Table. Items and response options for the in-house developed 'stressors for mental health associated with the COVID-19 pandemic' questionnaire.**

(DOCX)

**S2 Table. Raw response data for the in-house developed 'stressors for mental health associated with the COVID-19 pandemic' questionnaire (except free texts).**

(DOCX)

**S3 Table. Data sources and analytic approach for comparison with population data.**

(DOCX)

**S1 Fig. Previous mental health problems in the sample (valid answers only, without N/As).**

(DOCX)

**S1 File. Outlier screening and recoding.**

(DOCX)

## Acknowledgments

The authors would like to thank Noah McCuen-Koytek for his support with the spell and grammar check of the revised manuscript.

## Author Contributions

**Conceptualization:** Kevin Hilbert, Julia Asbrand, Ulrike Lueken.

**Data curation:** Kevin Hilbert, Sophia Seemann, Leonore Horváth.

**Formal analysis:** Kevin Hilbert, Ole Boeken, Sophia Seemann.

**Investigation:** Kevin Hilbert, Sophia Seemann, Leonore Horváth.

**Methodology:** Kevin Hilbert, Ole Boeken, Sophia Seemann, Ulrike Lueken.

**Project administration:** Kevin Hilbert, Julia Asbrand, Leonore Horváth, Ulrike Lueken.

**Resources:** Kevin Hilbert, Julia Asbrand, Sophia Seemann, Till Langhammer, Berit Praxl, Leonore Horváth, Andrea Ertle, Ulrike Lueken.

**Software:** Kevin Hilbert.

**Supervision:** Julia Asbrand, Ulrike Lueken.

**Validation:** Till Langhammer.

**Visualization:** Kevin Hilbert.

**Writing – original draft:** Kevin Hilbert.

**Writing – review & editing:** Kevin Hilbert, Ole Boeken, Julia Asbrand, Sophia Seemann, Till Langhammer, Berit Praxl, Leonore Horváth, Andrea Ertle, Ulrike Lueken.

## References

1. Qiu J, Shen B, Zhao M, Wang Z, Xie B, Xu Y. A nationwide survey of psychological distress among Chinese people in the COVID-19 epidemic: implications and policy recommendations. *General Psychiatry*. 2020; 33(2):e100213. <https://doi.org/10.1136/gpsych-2020-100213> PMID: 32215365
2. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, et al. Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the General Population in China. *Int J Environ Res Public Health*. 2020; 17(5):1729. <https://doi.org/10.3390/ijerph17051729> PMID: 32155789
3. Torales J O'Higgins M, Castaldelli-Maia JM, Ventriglio A. The outbreak of COVID-19 coronavirus and its impact on global mental health. *Int J Soc Psychiatry*. 2020; 66(4):317–20. <https://doi.org/10.1177/0020764020915212> PMID: 32233719
4. Collaborators C-MD. Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the COVID-19 pandemic. *Lancet*. 2021; 398(10312):1700–12. [https://doi.org/10.1016/S0140-6736\(21\)02143-7](https://doi.org/10.1016/S0140-6736(21)02143-7) PMID: 34634250
5. Meewisse ML, Olf M, Kleber R, Kitchiner NJ, Gersons BP. The course of mental health disorders after a disaster: predictors and comorbidity. *J Trauma Stress*. 2011; 24(4):405–13. <https://doi.org/10.1002/jts.20663> PMID: 21815216
6. Lee AM, Wong JG, McAlonan GM, Cheung V, Cheung C, Sham PC, et al. Stress and Psychological Distress among SARS Survivors 1 Year after the Outbreak. *Canadian journal of psychiatry Revue canadienne de psychiatrie*. 2007; 52(4):233–40. <https://doi.org/10.1177/070674370705200405> PMID: 17500304
7. Katz CL, Pellegrino L, Pandya A, Ng A, DeLisi LE. Research on psychiatric outcomes and interventions subsequent to disasters: a review of the literature. *Psychiatry Res*. 2002; 110(3):201–17. [https://doi.org/10.1016/s0165-1781\(02\)00110-5](https://doi.org/10.1016/s0165-1781(02)00110-5) PMID: 12127471
8. PsychotherapeutenVereinigung D. Patients requesting therapy during the corona-pandemic [Patientenanfragen während der Corona-Pandemie]. 2021.
9. Braet C, Van Vlierberghe L, Vandevivere E, Theuwis L, Bosmans G. Depression in early, middle and late adolescence: differential evidence for the cognitive diathesis-stress model. *Clin Psychol Psychother*. 2013; 20(5):369–83. <https://doi.org/10.1002/cpp.1789> PMID: 22473794
10. Kuehner C, Schultz K, Gass P, Meyer-Lindenberg A, Dressing H. [Mental Health Status in the Community During the COVID-19-Pandemic]. *Psychiatr Prax*. 2020; 47(7):361–9. <https://doi.org/10.1055/a-1222-9067> PMID: 32823293

11. Pfefferbaum B, North CS. Mental Health and the Covid-19 Pandemic. *N Engl J Med*. 2020; 383(6):510–2. <https://doi.org/10.1056/NEJMp2008017> PMID: 32283003
12. Vindegaard N, Benros ME. COVID-19 pandemic and mental health consequences: Systematic review of the current evidence. *Brain Behav Immun*. 2020; 89:531–42. <https://doi.org/10.1016/j.bbi.2020.05.048> PMID: 32485289
13. Xiong J, Lipsitz O, Nasri F, Lui LMW, Gill H, Phan L, et al. Impact of COVID-19 pandemic on mental health in the general population: A systematic review. *J Affect Disord*. 2020; 277:55–64. <https://doi.org/10.1016/j.jad.2020.08.001> PMID: 32799105
14. Bendau A, Kunas SL, Wyka S, Petzold MB, Plag J, Asselmann E, et al. Longitudinal changes of anxiety and depressive symptoms during the COVID-19 pandemic in Germany: The role of pre-existing anxiety, depressive, and other mental disorders. *J Anxiety Disord*. 2021; 79:102377. <https://doi.org/10.1016/j.janxdis.2021.102377> PMID: 33662702
15. O'Connor RC, Wetherall K, Cleare S, McClelland H, Melson AJ, Niedzwiedz CL, et al. Mental health and well-being during the COVID-19 pandemic: longitudinal analyses of adults in the UK COVID-19 Mental Health & Wellbeing study. *Br J Psychiatry*. 2020:1–8.
16. Langhammer T, Hilbert K, Praxl B, Kirschbaum C, Ertle A, Asbrand J, et al. Mental health trajectories of individuals and families following the COVID-19 pandemic: Study protocol of a longitudinal investigation and prevention program. *Ment Health Prev*. 2021; 24:200221. <https://doi.org/10.1016/j.mhp.2021.200221> PMID: 34608431
17. Kroenke K, Spitzer RL, Williams JB, Lowe B. The Patient Health Questionnaire Somatic, Anxiety, and Depressive Symptom Scales: a systematic review. *Gen Hosp Psychiatry*. 2010; 32(4):345–59. <https://doi.org/10.1016/j.genhosppsy.2010.03.006> PMID: 20633738
18. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *Journal of general internal medicine*. 2001; 16(9):606–13. <https://doi.org/10.1046/j.1525-1497.2001.016009606.x> PMID: 11556941
19. Kroenke K, Spitzer RL, Williams JB. The PHQ-15: validity of a new measure for evaluating the severity of somatic symptoms. *Psychosom Med*. 2002; 64(2):258–66. <https://doi.org/10.1097/00006842-200203000-00008> PMID: 11914441
20. Spitzer RL, Kroenke K, Williams JB, Lowe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med*. 2006; 166(10):1092–7. <https://doi.org/10.1001/archinte.166.10.1092> PMID: 16717171
21. Ware JE, GlaxoSmithKline. How to score and interpret single-item health status measures: a manual for users of the of the SF-8 health survey: (with a supplement on the SF-6 health survey). Lincoln, RI; Boston, MA: QualityMetric, Inc. ; Health Assessment Lab; 2001.
22. Gross JJ, John OP. Individual differences in two emotion regulation processes: implications for affect, relationships, and well-being. *J Pers Soc Psychol*. 2003; 85(2):348–62. <https://doi.org/10.1037/0022-3514.85.2.348> PMID: 12916575
23. Kocalevent RD, Hinz A, Brahler E. Standardization of a screening instrument (PHQ-15) for somatization syndromes in the general population. *Bmc Psychiatry*. 2013; 13:91. <https://doi.org/10.1186/1471-244X-13-91> PMID: 23514436
24. Lowe B, Decker O, Muller S, Brahler E, Schellberg D, Herzog W, et al. Validation and standardization of the Generalized Anxiety Disorder Screener (GAD-7) in the general population. *Med Care*. 2008; 46(3):266–74. <https://doi.org/10.1097/MLR.0b013e318160d093> PMID: 18388841
25. Hinz A, Mehnert A, Kocalevent RD, Brahler E, Forkmann T, Singer S, et al. Assessment of depression severity with the PHQ-9 in cancer patients and in the general population. *Bmc Psychiatry*. 2016; 16:22. <https://doi.org/10.1186/s12888-016-0728-6> PMID: 26831145
26. Hides L, Dingle G, Quinn C, Stoyanov SR, Zelenko O, Tjondronegoro D, et al. Efficacy and Outcomes of a Music-Based Emotion Regulation Mobile App in Distressed Young People: Randomized Controlled Trial. *JMIR Mhealth Uhealth*. 2019; 7(1):e11482. <https://doi.org/10.2196/11482> PMID: 30664457