

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

cis-Triaqua[1,1'-(propane-1,3-diyl)-bis(pyridin-1-ium-4-carboxylato)- κ O]-bis(thiocyanato- κ N)manganese(II) dihydrate

Qing-Hua Tan and Yan-Qin Wang*

College of Chemistry and Chemical Engineering, Department of Chemistry, Inner Mongolia University, Huhhot 010021, People's Republic of China

Correspondence e-mail: yqwang_chem@imu.edu.cn


Received 8 January 2014; accepted 11 January 2014

Key indicators: single-crystal X-ray study; $T = 273$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.045; wR factor = 0.104; data-to-parameter ratio = 16.6.

In the title compound, $[\text{Mn}(\text{NCS})_2(\text{C}_{15}\text{H}_{14}\text{N}_2\text{O}_4)(\text{H}_2\text{O})_3] \cdot 2\text{H}_2\text{O}$, the metal ion is octahedrally coordinated by three water molecules, one carboxylate O atom from a 1,1'-(propane-1,3-diyl)bis(pyridinium-4-carboxylate) ligand and two N atoms from two thiocyanate anions in *cis* positions, forming a mononuclear complex molecule. In the crystal, molecules are connected into a three-dimensional architecture through $\text{O}-\text{H} \cdots \text{O}$ hydrogen bonds involving water molecules and carboxylate groups.

Related literature

For related literature concerning the ligand, see: Jiang & Li (2006); Li *et al.* (2007); Wu *et al.* (2006); Zhang *et al.* (2002).



Experimental

Crystal data

 $[\text{Mn}(\text{NCS})_2(\text{C}_{15}\text{H}_{14}\text{N}_2\text{O}_4)(\text{H}_2\text{O})_3] \cdot 2\text{H}_2\text{O}$ $M_r = 547.46$ Monoclinic, $P2_1/c$ $a = 17.056$ (2) Å $b = 11.7514$ (16) Å $c = 11.8962$ (16) Å $\beta = 92.984$ (2)° $V = 2381.1$ (6) Å³ $Z = 4$ Mo $K\alpha$ radiation
 $\mu = 0.78$ mm⁻¹ $T = 273$ K
 $0.10 \times 0.08 \times 0.05$ mm

Data collection

Bruker SMART CCD area-detector
diffractometerAbsorption correction: multi-scan
(SADABS; Bruker, 1999) $T_{\min} = 0.926$, $T_{\max} = 0.962$ 14572 measured reflections
5451 independent reflections
3409 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.049$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.045$ $wR(F^2) = 0.104$ $S = 1.04$

5451 reflections

328 parameters

6 restraints

H atoms treated by a mixture of
independent and constrained
refinement $\Delta\rho_{\max} = 0.34$ e Å⁻³ $\Delta\rho_{\min} = -0.33$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{O5}-\text{H5WA} \cdots \text{O3}^{\text{i}}$	0.89 (4)	1.84 (4)	2.707 (3)	164 (4)
$\text{O5}-\text{H5WB} \cdots \text{O3}^{\text{ii}}$	0.81 (4)	1.88 (4)	2.682 (3)	174 (4)
$\text{O6}-\text{H6WA} \cdots \text{O9}^{\text{iii}}$	0.78 (3)	2.09 (3)	2.862 (3)	168 (4)
$\text{O6}-\text{H6WB} \cdots \text{O8}^{\text{iv}}$	0.82 (3)	2.02 (3)	2.837 (3)	179 (4)
$\text{O7}-\text{H7WA} \cdots \text{O1}^{\text{iii}}$	0.88 (3)	1.91 (3)	2.780 (3)	169 (3)
$\text{O7}-\text{H7WB} \cdots \text{O8}$	0.82 (3)	1.90 (3)	2.710 (3)	169 (3)
$\text{O8}-\text{H8WA} \cdots \text{O9}^{\text{v}}$	0.84 (2)	1.93 (2)	2.767 (3)	175 (3)
$\text{O8}-\text{H8WB} \cdots \text{O4}^{\text{vi}}$	0.87 (2)	1.91 (2)	2.755 (3)	164 (3)
$\text{O9}-\text{H9WA} \cdots \text{O2}$	0.85 (2)	2.05 (2)	2.830 (3)	152 (3)
$\text{O9}-\text{H9WB} \cdots \text{O4}^{\text{ii}}$	0.83 (2)	2.03 (2)	2.862 (3)	174 (3)

Symmetry codes: (i) $-x+1, y-\frac{1}{2}, -z+\frac{3}{2}$; (ii) $x-1, -y+\frac{3}{2}, z-\frac{1}{2}$; (iii) $x, -y+\frac{3}{2}, z-\frac{1}{2}$; (iv) $x, -y+\frac{3}{2}, z+\frac{1}{2}$; (v) $-x, -y+1, -z+1$; (vi) $x-1, y, z-1$.

Data collection: *SMART* (Bruker, 1999); cell refinement: *SAINT-Plus* (Bruker, 1999); data reduction: *SAINT-Plus*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and local programs.

We are thankful for financial support from the NSFC (21301087), SPH-IMU-30105-125135, and the Inner Mongolia autonomous region Natural Science Fund project (2013MS0206).

Supporting information for this paper is available from the IUCr electronic archives (Reference: BT6955).

References

- Bruker (1999). *SMART*, *SAINT-Plus* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Jiang, N.-Y. & Li, S.-L. (2006). *Chin. J. Struct. Chem.* **25**, 957–964.
- Li, Y., Li, G.-Q., Zou, W.-Q., Zheng, F.-K., Zou, J.-P., Guo, G.-C., Lu, C.-Z. & Huang, J.-S. (2007). *J. Mol. Struct.* **837**, 231–236.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wu, A.-Q., Li, Y., Zheng, F.-K., Guo, G.-C. & Huang, J.-S. (2006). *Cryst. Growth Des.* **6**, 444–450.
- Zhang, X., Guo, G.-C., Zheng, F.-K., Zhou, G.-W., Mao, J.-G., Dong, Z.-C., Huang, J.-S. & Mak, T. C. W. (2002). *J. Chem. Soc. Dalton Trans.* pp. 1344–1349.

supplementary materials

Acta Cryst. (2014). E70, m42 [doi:10.1107/S1600536814000701]

***cis*-Triaqua[1,1'-(propane-1,3-diyl)bis(pyridin-1-ium-4-carboxylato)- κ O]bis(thiocyanato- κ N)manganese(II) dihydrate**

Qing-Hua Tan and Yan-Qin Wang

1. Comment

The double betaine 1,3-bis(pyridinium-4-carboxylato)propane is a neutral dicarboxylate ligand. In recent years, an increasing number of coordination compounds with this ligand have been reported (Jiang *et al.*, 2006; Li *et al.*, 2007; Wu *et al.*, 2006; Zhang *et al.*, 2002). The structure of the title compound consists of the mononuclear complex molecules $[\text{Mn}(\text{C}_{15}\text{O}_4\text{N}_2\text{H}_{14})(\text{NCS})_2(\text{H}_2\text{O})_3]$ and lattice water molecules. The metal ion adopts an octahedral coordination geometry completed by three water molecules (O5, O6 and O7), a carboxylato O atom (O1) from the betaine ligand, and two N atoms (N1 and N2) from two thiocyanato anions in *cis* position (Fig. 1). There is a lot of intermolecular O—H $\cdots$ O hydrogen bonds involving the carboxylate groups and water molecules. The O5 water molecule forms two hydrogen bonds with uncoordinated carboxylate oxygen atoms (O3) from different complex molecules, the O6 water molecule contributes its hydrogen atoms to two lattice water molecules (O8 and O9), and the O6 water molecule forms two hydrogen bonds with the coordinated O1 atom and a lattice water molecule (O8). The lattice water molecules are four-coordinated: they accept hydrogen atoms from coordinated water molecules and contribute their hydrogen atoms to carboxylate oxygen atoms (O4 and O2) or water molecules. These hydrogen bonds lead to a three-dimensional architecture (Fig. 2). In addition, there is a short S $\cdots$ S contact [$\text{S1}\cdots\text{S2}(1-x, 1-y, 1-z)$, 3.625 (1) Å] between the thiocyanato ions from neighboring complex molecules.

2. Experimental

The preparation of the compound: 1,3-bis(pyridinium-4-carboxylato)propane (0.029 g, 0.10 mmol) and Potassium thiocyanate (0.039 g, 0.40 mmol) dissolved in ethanol (3 ml) and water (5 ml) were added into the ethanol solution of manganese(II) dichloride hexahydrate (0.040 g, 0.2 mmol), and the resulting light yellow solution was stirred for several minutes, filtered. The filtrate was kept at room temperature, and a few days later, light yellow single crystals suitable for X-ray diffraction was obtained.

3. Refinement

All hydrogen atoms attached to carbon atoms were placed geometrically at calculated positions and refined using the riding model with secondary C—H = 0.97 Å and aromatic C—H = 0.93 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. H atoms bonded to O atoms have been refined isotropically with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. In the two non-coordinating water molecules the O—H distances were restrained to 0.85 (2) Å and the H $\cdots$ H distances to 1.38 (2) Å.

Computing details

Data collection: *SMART* (Bruker, 1999); cell refinement: *SAINT-Plus* (Bruker, 1999); data reduction: *SAINT-Plus* (Bruker, 1999); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure:

SHELXTL (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008) and local programs.




Figure 1

The structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. Hydrogen atoms are omitted.




Figure 2

View of the intermolecular interactions in the title compound. Hydrogen bonds and S...S contacts are drawn with dashed lines. Hydrogen atoms attached to C atoms are omitted.

***cis*-Triaqua[1,1'-(propane-1,3-diyl)bis(pyridin-1-ium-4-carboxylato)- κ O]bis(thiocyanato- κ N)manganese(II) dihydrate**

Crystal data

$[\text{Mn}(\text{NCS})_2(\text{C}_{15}\text{H}_{14}\text{N}_2\text{O}_4)(\text{H}_2\text{O})_3] \cdot 2\text{H}_2\text{O}$

$M_r = 547.46$

Monoclinic, $P2_1/c$

$a = 17.056(2) \text{ \AA}$

$b = 11.7514(16) \text{ \AA}$

$c = 11.8962(16) \text{ \AA}$

$\beta = 92.984(2)^\circ$

$V = 2381.1(6) \text{ \AA}^3$

$Z = 4$

$F(000) = 1132$

$D_x = 1.527 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2306 reflections

$\theta = 2.4\text{--}23.7^\circ$

$\mu = 0.78 \text{ mm}^{-1}$
 $T = 273 \text{ K}$

Rectangular, yellow
 $0.10 \times 0.08 \times 0.05 \text{ mm}$

Data collection

Bruker SMART CCD area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 phi and ω scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 1999)
 $T_{\min} = 0.926$, $T_{\max} = 0.962$

14572 measured reflections
 5451 independent reflections
 3409 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.049$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.1^\circ$
 $h = -20 \rightarrow 22$
 $k = -9 \rightarrow 15$
 $l = -12 \rightarrow 15$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.104$
 $S = 1.04$
 5451 reflections
 328 parameters
 6 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0366P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.34 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.33 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Mn1	0.19350 (2)	0.66835 (4)	0.46440 (3)	0.02899 (13)
C1	0.91446 (16)	0.7847 (3)	1.0248 (3)	0.0356 (7)
C2	0.85603 (15)	0.6997 (2)	0.9725 (2)	0.0307 (6)
C3	0.83380 (19)	0.7061 (3)	0.8598 (3)	0.0515 (9)
H3A	0.8537	0.7634	0.8155	0.062*
C4	0.82470 (16)	0.6147 (2)	1.0353 (2)	0.0344 (7)
H4A	0.8381	0.6094	1.1119	0.041*
C5	0.78248 (19)	0.6283 (3)	0.8131 (3)	0.0547 (10)
H5A	0.7680	0.6327	0.7368	0.066*
C6	0.77363 (15)	0.5376 (2)	0.9852 (2)	0.0348 (7)
H6A	0.7534	0.4792	1.0278	0.042*
C7	0.69666 (14)	0.4628 (2)	0.8224 (2)	0.0332 (7)
H7B	0.6962	0.3943	0.8678	0.040*
H7A	0.7141	0.4423	0.7488	0.040*

C8	0.61420 (14)	0.5110 (2)	0.8099 (2)	0.0346 (7)
H8B	0.6153	0.5865	0.7772	0.042*
H8A	0.5918	0.5165	0.8830	0.042*
C9	0.56506 (15)	0.4320 (3)	0.7341 (2)	0.0374 (7)
H9B	0.5863	0.4306	0.6600	0.045*
H9A	0.5681	0.3554	0.7644	0.045*
C10	0.46197 (16)	0.5602 (3)	0.6618 (2)	0.0376 (7)
H10A	0.5003	0.5987	0.6239	0.045*
C11	0.42645 (16)	0.4099 (3)	0.7759 (2)	0.0374 (7)
H11A	0.4405	0.3459	0.8183	0.045*
C12	0.38622 (15)	0.5987 (2)	0.6539 (2)	0.0338 (7)
H12A	0.3736	0.6639	0.6126	0.041*
C13	0.34983 (16)	0.4439 (2)	0.7678 (2)	0.0349 (7)
H13A	0.3119	0.4019	0.8029	0.042*
C14	0.32846 (15)	0.5407 (2)	0.7073 (2)	0.0278 (6)
C15	0.24337 (16)	0.5814 (3)	0.6987 (2)	0.0324 (7)
C16	0.36520 (17)	0.7912 (2)	0.4339 (2)	0.0315 (6)
C17	0.29664 (16)	0.4229 (3)	0.4638 (2)	0.0319 (6)
N1	0.25535 (14)	0.5006 (2)	0.45556 (19)	0.0418 (6)
N2	0.29939 (14)	0.7665 (2)	0.4249 (2)	0.0422 (6)
N3	0.75267 (12)	0.5455 (2)	0.87586 (18)	0.0312 (5)
N4	0.48197 (12)	0.4677 (2)	0.72342 (18)	0.0322 (5)
O1	0.23085 (10)	0.67351 (16)	0.64340 (14)	0.0335 (5)
O2	0.19488 (11)	0.52474 (19)	0.74602 (17)	0.0492 (6)
O3	0.93988 (12)	0.85658 (18)	0.95846 (18)	0.0455 (5)
O4	0.93195 (12)	0.7747 (2)	1.12736 (17)	0.0525 (6)
O5	0.08954 (12)	0.5776 (2)	0.4957 (2)	0.0585 (7)
H5WB	0.046 (2)	0.600 (3)	0.480 (3)	0.088*
H5WA	0.089 (2)	0.503 (4)	0.508 (3)	0.088*
O6	0.11913 (13)	0.82309 (19)	0.47446 (18)	0.0421 (6)
H6WB	0.095 (2)	0.846 (3)	0.527 (3)	0.063*
H6WA	0.095 (2)	0.835 (3)	0.418 (3)	0.063*
O7	0.16305 (12)	0.67069 (19)	0.28495 (16)	0.0372 (5)
H7WB	0.1212 (19)	0.651 (3)	0.255 (3)	0.056*
H7WA	0.1806 (18)	0.727 (3)	0.244 (3)	0.056*
O8	0.03603 (12)	0.5969 (2)	0.15924 (19)	0.0514 (6)
H8WB	−0.0005 (17)	0.648 (2)	0.161 (3)	0.077*
H8WA	0.0157 (18)	0.5357 (17)	0.181 (3)	0.077*
O9	0.03881 (12)	0.5981 (2)	0.77207 (17)	0.0453 (6)
H9WB	0.0092 (15)	0.633 (3)	0.726 (2)	0.068*
H9WA	0.0805 (13)	0.578 (3)	0.741 (2)	0.068*
S1	0.35705 (5)	0.31522 (7)	0.47422 (7)	0.0519 (2)
S2	0.45791 (5)	0.82371 (8)	0.45024 (7)	0.0530 (3)

Atomic displacement parameters ($\text{\AA}^2$)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mn1	0.0255 (2)	0.0327 (3)	0.0285 (2)	0.00140 (18)	−0.00004 (17)	0.00147 (19)
C1	0.0278 (15)	0.0328 (17)	0.0461 (19)	0.0018 (13)	0.0012 (14)	−0.0026 (14)
C2	0.0262 (15)	0.0340 (17)	0.0316 (16)	−0.0023 (12)	−0.0011 (12)	0.0016 (12)

C3	0.058 (2)	0.061 (2)	0.0353 (18)	−0.0301 (18)	−0.0047 (15)	0.0120 (16)
C4	0.0382 (16)	0.0374 (17)	0.0271 (15)	−0.0058 (13)	−0.0035 (12)	0.0000 (13)
C5	0.060 (2)	0.075 (3)	0.0286 (17)	−0.0327 (19)	−0.0078 (15)	0.0132 (17)
C6	0.0377 (16)	0.0349 (18)	0.0318 (16)	−0.0026 (13)	0.0002 (13)	0.0072 (13)
C7	0.0270 (14)	0.0346 (17)	0.0378 (16)	−0.0027 (12)	0.0001 (12)	−0.0048 (13)
C8	0.0278 (15)	0.0332 (17)	0.0427 (17)	−0.0001 (13)	0.0005 (13)	−0.0046 (14)
C9	0.0258 (15)	0.0373 (18)	0.0484 (18)	0.0022 (13)	−0.0039 (13)	−0.0076 (14)
C10	0.0310 (16)	0.0431 (19)	0.0388 (17)	−0.0070 (14)	0.0003 (13)	0.0057 (15)
C11	0.0383 (17)	0.0330 (18)	0.0401 (17)	−0.0016 (14)	−0.0050 (13)	0.0041 (14)
C12	0.0321 (15)	0.0382 (18)	0.0307 (15)	−0.0030 (13)	−0.0013 (12)	0.0089 (13)
C13	0.0309 (15)	0.0367 (18)	0.0371 (17)	−0.0036 (13)	0.0019 (13)	0.0064 (14)
C14	0.0291 (14)	0.0335 (16)	0.0203 (13)	0.0009 (12)	−0.0037 (11)	−0.0013 (12)
C15	0.0331 (15)	0.0409 (19)	0.0232 (15)	0.0017 (13)	0.0007 (12)	−0.0001 (13)
C16	0.0367 (17)	0.0302 (16)	0.0277 (15)	0.0013 (13)	0.0023 (13)	0.0057 (12)
C17	0.0336 (15)	0.0372 (18)	0.0245 (15)	−0.0003 (13)	−0.0006 (12)	−0.0033 (13)
N1	0.0423 (15)	0.0457 (17)	0.0366 (14)	0.0102 (13)	−0.0048 (12)	−0.0057 (12)
N2	0.0329 (14)	0.0525 (18)	0.0411 (15)	−0.0043 (12)	0.0002 (12)	0.0073 (13)
N3	0.0279 (12)	0.0370 (15)	0.0285 (13)	−0.0037 (10)	−0.0008 (10)	−0.0009 (11)
N4	0.0281 (12)	0.0339 (14)	0.0341 (13)	−0.0001 (11)	−0.0037 (10)	−0.0039 (11)
O1	0.0367 (11)	0.0362 (12)	0.0275 (10)	0.0079 (9)	0.0001 (8)	0.0023 (9)
O2	0.0328 (11)	0.0633 (16)	0.0522 (13)	0.0026 (11)	0.0087 (10)	0.0228 (12)
O3	0.0421 (12)	0.0357 (13)	0.0583 (14)	−0.0128 (10)	−0.0005 (10)	0.0030 (11)
O4	0.0566 (14)	0.0599 (16)	0.0394 (13)	−0.0165 (12)	−0.0121 (11)	−0.0053 (11)
O5	0.0266 (11)	0.0391 (14)	0.110 (2)	0.0006 (11)	0.0038 (13)	0.0183 (14)
O6	0.0475 (14)	0.0417 (13)	0.0371 (13)	0.0119 (11)	0.0029 (10)	0.0027 (11)
O7	0.0348 (12)	0.0452 (14)	0.0307 (12)	−0.0068 (10)	−0.0047 (9)	0.0059 (10)
O8	0.0427 (13)	0.0564 (17)	0.0543 (14)	−0.0137 (11)	−0.0044 (11)	0.0116 (13)
O9	0.0388 (13)	0.0530 (15)	0.0442 (13)	0.0006 (11)	0.0019 (10)	0.0046 (11)
S1	0.0514 (5)	0.0444 (5)	0.0606 (6)	0.0178 (4)	0.0094 (4)	0.0025 (4)
S2	0.0334 (4)	0.0735 (7)	0.0513 (5)	−0.0135 (4)	−0.0067 (4)	0.0118 (5)

Geometric parameters (Å, °)

Mn1—O5	2.118 (2)	C10—N4	1.345 (3)
Mn1—O7	2.171 (2)	C10—C12	1.367 (4)
Mn1—O1	2.1914 (18)	C10—H10A	0.9300
Mn1—N2	2.214 (2)	C11—N4	1.346 (3)
Mn1—O6	2.224 (2)	C11—C13	1.365 (4)
Mn1—N1	2.241 (3)	C11—H11A	0.9300
C1—O4	1.246 (3)	C12—C14	1.379 (3)
C1—O3	1.249 (3)	C12—H12A	0.9300
C1—C2	1.521 (4)	C13—C14	1.385 (4)
C2—C4	1.373 (4)	C13—H13A	0.9300
C2—C3	1.375 (4)	C14—C15	1.526 (4)
C3—C5	1.364 (4)	C15—O2	1.222 (3)
C3—H3A	0.9300	C15—O1	1.278 (3)
C4—C6	1.372 (4)	C16—N2	1.159 (3)
C4—H4A	0.9300	C16—S2	1.628 (3)
C5—N3	1.342 (4)	C17—N1	1.154 (3)
C5—H5A	0.9300	C17—S1	1.632 (3)

C6—N3	1.334 (3)	O5—H5WB	0.81 (4)
C6—H6A	0.9300	O5—H5WA	0.89 (4)
C7—N3	1.482 (3)	O6—H6WB	0.82 (3)
C7—C8	1.516 (3)	O6—H6WA	0.78 (3)
C7—H7B	0.9700	O7—H7WB	0.82 (3)
C7—H7A	0.9700	O7—H7WA	0.88 (3)
C8—C9	1.516 (4)	O8—H8WB	0.867 (17)
C8—H8B	0.9700	O8—H8WA	0.844 (17)
C8—H8A	0.9700	O9—H9WB	0.834 (16)
C9—N4	1.477 (3)	O9—H9WA	0.854 (16)
C9—H9B	0.9700	S1—S2 ⁱ	3.6247 (12)
C9—H9A	0.9700		
O5—Mn1—O7	91.13 (9)	N4—C9—H9B	109.2
O5—Mn1—O1	92.68 (9)	C8—C9—H9B	109.2
O7—Mn1—O1	176.17 (8)	N4—C9—H9A	109.2
O5—Mn1—N2	177.44 (9)	C8—C9—H9A	109.2
O7—Mn1—N2	86.61 (8)	H9B—C9—H9A	107.9
O1—Mn1—N2	89.57 (8)	N4—C10—C12	121.1 (3)
O5—Mn1—O6	85.24 (9)	N4—C10—H10A	119.5
O7—Mn1—O6	86.25 (8)	C12—C10—H10A	119.5
O1—Mn1—O6	93.60 (7)	N4—C11—C13	120.9 (3)
N2—Mn1—O6	93.40 (9)	N4—C11—H11A	119.6
O5—Mn1—N1	88.05 (9)	C13—C11—H11A	119.6
O7—Mn1—N1	93.10 (8)	C10—C12—C14	120.0 (3)
O1—Mn1—N1	87.49 (8)	C10—C12—H12A	120.0
N2—Mn1—N1	93.28 (9)	C14—C12—H12A	120.0
O6—Mn1—N1	173.24 (9)	C11—C13—C14	120.2 (3)
O4—C1—O3	127.5 (3)	C11—C13—H13A	119.9
O4—C1—C2	117.2 (3)	C14—C13—H13A	119.9
O3—C1—C2	115.3 (3)	C12—C14—C13	118.0 (3)
C4—C2—C3	118.4 (3)	C12—C14—C15	120.9 (2)
C4—C2—C1	121.3 (3)	C13—C14—C15	121.1 (2)
C3—C2—C1	120.3 (3)	O2—C15—O1	126.8 (3)
C5—C3—C2	119.9 (3)	O2—C15—C14	117.5 (3)
C5—C3—H3A	120.0	O1—C15—C14	115.6 (2)
C2—C3—H3A	120.0	N2—C16—S2	178.2 (3)
C6—C4—C2	120.0 (3)	N1—C17—S1	178.4 (3)
C6—C4—H4A	120.0	C17—N1—Mn1	168.1 (2)
C2—C4—H4A	120.0	C16—N2—Mn1	155.2 (2)
N3—C5—C3	120.9 (3)	C6—N3—C5	120.2 (2)
N3—C5—H5A	119.6	C6—N3—C7	120.4 (2)
C3—C5—H5A	119.6	C5—N3—C7	119.4 (2)
N3—C6—C4	120.6 (3)	C10—N4—C11	119.8 (2)
N3—C6—H6A	119.7	C10—N4—C9	119.4 (2)
C4—C6—H6A	119.7	C11—N4—C9	120.8 (2)
N3—C7—C8	111.8 (2)	C15—O1—Mn1	120.58 (17)
N3—C7—H7B	109.3	Mn1—O5—H5WB	125 (3)
C8—C7—H7B	109.3	Mn1—O5—H5WA	123 (2)

N3—C7—H7A	109.3	H5WB—O5—H5WA	110 (3)
C8—C7—H7A	109.3	Mn1—O6—H6WB	128 (2)
H7B—C7—H7A	107.9	Mn1—O6—H6WA	113 (3)
C7—C8—C9	108.1 (2)	H6WB—O6—H6WA	109 (4)
C7—C8—H8B	110.1	Mn1—O7—H7WB	126 (2)
C9—C8—H8B	110.1	Mn1—O7—H7WA	119 (2)
C7—C8—H8A	110.1	H7WB—O7—H7WA	107 (3)
C9—C8—H8A	110.1	H8WB—O8—H8WA	106 (2)
H8B—C8—H8A	108.4	H9WB—O9—H9WA	110 (2)
N4—C9—C8	112.0 (2)	C17—S1—S2 ⁱ	154.96 (11)

Symmetry code: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry ($\text{\AA}$, $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O5—H5WA $\cdots$ O3 ⁱⁱ	0.89 (4)	1.84 (4)	2.707 (3)	164 (4)
O5—H5WB $\cdots$ O3 ⁱⁱⁱ	0.81 (4)	1.88 (4)	2.682 (3)	174 (4)
O6—H6WA $\cdots$ O9 ^{iv}	0.78 (3)	2.09 (3)	2.862 (3)	168 (4)
O6—H6WB $\cdots$ O8 ^v	0.82 (3)	2.02 (3)	2.837 (3)	179 (4)
O7—H7WA $\cdots$ O1 ^{iv}	0.88 (3)	1.91 (3)	2.780 (3)	169 (3)
O7—H7WB $\cdots$ O8	0.82 (3)	1.90 (3)	2.710 (3)	169 (3)
O8—H8WA $\cdots$ O9 ^{vi}	0.84 (2)	1.93 (2)	2.767 (3)	175 (3)
O8—H8WB $\cdots$ O4 ^{vii}	0.87 (2)	1.91 (2)	2.755 (3)	164 (3)
O9—H9WA $\cdots$ O2	0.85 (2)	2.05 (2)	2.830 (3)	152 (3)
O9—H9WB $\cdots$ O4 ⁱⁱⁱ	0.83 (2)	2.03 (2)	2.862 (3)	174 (3)

Symmetry codes: (ii) $-x+1, y-1/2, -z+3/2$; (iii) $x-1, -y+3/2, z-1/2$; (iv) $x, -y+3/2, z-1/2$; (v) $x, -y+3/2, z+1/2$; (vi) $-x, -y+1, -z+1$; (vii) $x-1, y, z-1$.