

Original Article  
Medicine General &  
Social Medicine



# The Prevalence of Obesity and Metabolic Syndrome in the Korean Military Compared with the General Population

Jung Hwan Lee ,<sup>1,2\*</sup> Da Hea Seo ,<sup>2\*</sup> Min Jung Nam ,<sup>1</sup> Geon Hui Lee ,<sup>1</sup>  
Dong Hee Yang ,<sup>1</sup> Min Joo Lee ,<sup>1</sup> Ung-Rim Choi ,<sup>1</sup> and Seongbin Hong <sup>2</sup>

<sup>1</sup>The 5th Division the Medical Battalion of the Republic of Korea Armed Forces, Yeoncheon, Korea

<sup>2</sup>Division of Endocrinology and Metabolism, Department of Internal Medicine, Inha University Hospital, Inha University School of Medicine, Incheon, Korea



Received: Jan 17, 2018

Accepted: Apr 13, 2018

**Address for Correspondence:**

Seongbin Hong, MD

Division of Endocrinology and Metabolism,  
Department of Internal Medicine, Inha  
University Hospital, Inha University School of  
Medicine, 27 Inhang-ro, Jung-gu,  
Incheon 22332, Republic of Korea.  
E-mail: sbhongmd@inha.ac.kr

\*Jung Hwan Lee and Da Hea Seo contributed  
equally to this study as first author.

© 2018 The Korean Academy of Medical  
Sciences.

This is an Open Access article distributed  
under the terms of the Creative Commons  
Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>)  
which permits unrestricted non-commercial  
use, distribution, and reproduction in any  
medium, provided the original work is properly  
cited.

**ORCID IDs**

Jung Hwan Lee   
<https://orcid.org/0000-0001-7567-0664>

Da Hea Seo   
<https://orcid.org/0000-0003-2767-0293>

Min Jung Nam   
<https://orcid.org/0000-0001-6546-8921>

Geon Hui Lee   
<https://orcid.org/0000-0002-9290-6541>

Dong Hee Yang   
<https://orcid.org/0000-0002-8923-7738>

Min Joo Lee   
<https://orcid.org/0000-0003-3510-139X>

## ABSTRACT

**Background:** Obesity and related metabolic disorders are growing health challenges worldwide and individuals at military service are not exceptions. The purpose of this study was to examine the prevalence of obesity and metabolic syndrome (MS) in the Korean military and to compare with the general population.

**Methods:** This was a cross-sectional study of 4,803 young military participants who underwent a corporal health-screening program between October 2013 and October 2014. The National Cholesterol Education Program Adult Treatment Panel III criteria was used to identify MS. We also sampled 1,108 men aged 19–29 years from the Korea National Health and Nutritional Examination Survey from 2010 to 2013 to compare with their military counterparts.

**Results:** The mean age of military participants was  $20.8 \pm 1.1$  years, and 20.6% ( $n = 988$ ) were obese. The prevalence of MS was 0.8% in military participants, while 7.9% in general population. The risk factors of MS were less prominent among military participants relative to civilians, with the exception of high blood pressure, of which prevalence was higher among military participants (21.5% vs. 18.2%, respectively). In multiple logistic analysis, high physical activity conferred lower odds of MS and obesity in military participants (odds ratios, 0.19 and 0.81, respectively). Age older than 25 years increased risk of most components of MS among civilians.

**Conclusion:** The prevalence of obesity and MS is lower in military participants compared with civilians of similar age. Monitoring of high blood pressure and proper stress management are warranted in those at military service.

**Keywords:** Metabolic Syndrome; Obesity; Military; Cross-sectional Survey

## INTRODUCTION

Metabolic syndrome (MS) is defined as a cluster of cardiovascular disease (CVD) risk factors that typically include central obesity, elevated blood pressure (BP), impaired glucose metabolism, and dyslipidemia.<sup>1</sup> Because obesity is a key component of the syndrome, the prevalence of MS in adults continues to rise as the prevalence of obesity has increased

Ung-Rim Choi   
<https://orcid.org/0000-0001-9698-4996>  
Seongbin Hong   
<https://orcid.org/0000-0002-8189-395X>

### Funding

This research was supported by Korean Military Medical Research Program (Grant number: 2014-KMMRP-015) and Inha University Research Grant.

### Disclosure

The authors have no potential conflicts of interest to disclose.

### Author Contributions

Conceptualization: Lee JH. Formal analysis: Hong S. Data curation: Lee GH, L Yang DH, Lee MJ, Choi UR, Nam MJ. Investigation: Hong S, Seo DH. Methodology: Lee JH. Software: Lee JH. Validation: Lee JH. Writing - original draft: Lee JH, Seo DH. Writing - review & editing: Hong S.

dramatically.<sup>2</sup> MS is not only associated with an increased risk of type 2 diabetes but also with increased risk of CVD morbidity and mortality.<sup>3-5</sup>

The prevalence of MS is increasing worldwide and this increasing trend also has been observed in Asian countries. According to Korea National Health and Nutritional Examination Survey (KHANES) 1998–2007, the prevalence of MS in Korea has increased from 24.9% to 31.3%.<sup>6</sup> In particular, the prevalence of MS at younger age has increased significantly during this period. Young adults with MS have higher risk of developing CVD and type 2 diabetes. Prevention and early intervention of MS among younger individuals will be very important to decrease morbidity and mortality of CVD, diabetes, and its complications.

Most Korean military service consists of young soldiers, recruited at the average age of early 20s. Regular physical trainings, restricted dietary patterns, and the army lifestyle may have a positive impact on the overall health status but there is paucity of data regarding the prevalence of MS in the Korean military. Understanding the prevalence of MS and risk factors among young soldiers would be very important, as it is required for most Korean males to serve in the military.

A corporal health-screening program (CHSP) has been implemented throughout the Korean military since 2012. CHSP includes a basic physical examination consisting of anthropometric measurements and blood sampling for routine chemistry. We had access to a health database of approximately 5,000 young soldiers during the study period, including waist circumference which is a good indicator for abdominal obesity, a well-known risk factor for insulin resistance, which further predisposes individuals to type 2 diabetes.

The objectives of this study were to: 1) estimate the prevalence of obesity, and MS, and investigate common components of MS among Korean military members, using the 2005 revised National Cholesterol Education Program Adult Treatment Panels III (NCEP-ATP III) criteria, as proposed by the American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement (AHA/NHLB),<sup>7</sup> and 2) make a comparison with Korean young adults in the general population using KHANES data.

## METHODS

### Study population

We conducted a retrospective, cross-sectional study of 4,803 young military participants who underwent a CHSP at one division medical battalion in the Republic of Korea (ROK) armed forces between October 2013 and October 2014. A CHSP is required for all soldiers promoting to at least a rank of corporal in the ROK armed forces. We excluded 37 subjects who had missing data for any of the individual MS criteria necessary to diagnosis MS.

We also had access to data from the KHANES, a nation-wide, population-based, cross-sectional health survey conducted by the Korea Centers for Disease Control and Prevention. A stratified, multistage probability sampling design was used to produce representative data. We obtained a sample of 1,108 young adults with age between 19–29 years from the fifth KNHNS conducted in 2010–2013.

### Measurements

The participants in the CHSP underwent a routine physical examination that included measurement of height, weight, BP, and overnight fasting blood sampling. Standard height was measured without shoes to the nearest 0.1 cm. Weight was determined without shoes, wearing light clothes. Waist circumference was measured midway between the lower limit of the rib cage and the iliac crest. BP was measured using a standard mercury sphygmomanometer after being seated for at least 5 minutes. Blood was sampled after 12 hours of fasting, and venous blood glucose, total cholesterol (TC), triglycerides (TGs), and high- and low-density lipoprotein cholesterol (HDL-C and LDL-C, respectively) were measured.

A questionnaire on lifestyle was conducted among participants and the questionnaire included the following: how many times have you performed strenuous activity for more than 10 minutes that caused you to breathe heavily during the past one week? For smoking status, participants were asked whether they were smoking at the time of survey interview (yes/no). A similar survey was extracted from KHANES to match with the questionnaire conducted during CHSP.

### Definitions of MS and obesity

The revised NCEP-ATP III criteria was used to identify those individuals with MS<sup>7</sup> and the following were considered risk factors for MS: 1) abdominal obesity  $\geq 90$  cm according to World Health Organization Asian specific cut off value,<sup>8</sup> 2) hypertriglyceridemia, defined as TG  $\geq 150$  mg/dL, or taking TG-lowering medication, 3) low HDL-C, defined as HDL-C  $< 40$  mg/dL in men, 4) high BP, defined as systolic BP/diastolic BP  $\geq 130/85$  mmHg or taking antihypertensive medication, and 5) high fasting plasma glucose (FPG)  $\geq 100$  mg/dL or taking antidiabetic medication (insulin or oral agents).

### Statistical analysis

Data was presented as means  $\pm$  standard deviation and prevalence of obesity, MS, and all components of MS were provided in percent (%). The  $\chi^2$  test was used for the comparison of prevalence of MS in each group, and two sample t-tests were used for the comparison of continuous variables. Multivariate logistic regression analysis was used to identify independent risk factors for development of MS, obesity and each components of MS.  $P < 0.05$  was considered statistically significant. Statistical analyses were performed using SPSS version 19.0 (SPSS Inc., Chicago, IL, USA).

### Ethics statement

The study was approved by the Institutional Review Board of the Armed Forces Medical Command, Seoul, Korea (approval number: AFMC-14-IRB-006). Informed consent was waived due to the retrospective nature of the study.

## RESULTS

**Table 1** shows the baseline characteristics of the 4,803 military personnel compared with 1,108 civilians aged between 19–29 years old from KHANES. The mean age of military participants was  $20.8 \pm 1.1$  years, which was lower than that of the civilian participants. Among anthropometric and laboratory measurements, mean body mass index (BMI), waist circumference, diastolic BP, and FPG values were lower in the military group than in the civilian

**Table 1.** Comparison of basic characteristics between Korean military personnel and civilians

| Variables                         | Military (n = 4,803) | Civilians aged 19–29 years (n = 1,108) | P value |
|-----------------------------------|----------------------|--|---------|
| Age, yr                           | 20.8 ± 1.1           | 24.4 ± 3.1                             | < 0.001 |
| BMI, kg/m <sup>2</sup>            | 23.0 ± 2.81          | 23.7 ± 3.8                             | < 0.001 |
| Waist circumference, cm           | 76.6 ± 7.5           | 80.8 ± 9.8                             | < 0.001 |
| Systolic BP, mmHg                 | 118.2 ± 19.9         | 114.2 ± 10.7                           | < 0.001 |
| Diastolic BP, mmHg                | 68.6 ± 11.6          | 74.3 ± 9.7                             | < 0.001 |
| TGs, mg/dL                        | 77.1 ± 41.0          | 118.0 ± 120.0                          | < 0.001 |
| Fasting glucose, mg/dL            | 85.4 ± 8.6           | 89.0 ± 9.8                             | < 0.001 |
| Smoking, %                        | 54.3                 | 40.6                                   | < 0.001 |
| Physical activity, % <sup>a</sup> | 18.5                 | 8.8                                    | < 0.001 |

Data are presented as means ± standard deviation (range) or number (%).

BMI = body mass index, BP = blood pressure, TG = triglyceride.

<sup>a</sup>At least 10 minutes of vigorous physical activity 5 or more times a week.

**Table 2.** Prevalence of MS and each component abnormality between Korean military personnel and civilian men aged 19–29 years

| Groups              | Military |                  | Civilians |    | P value |
|---------------------|----------|------------------|-----------|----|---------|
|                     | %        | OR               | %         | OR |         |
| Obesity             | 20.6     | 0.63 (0.54–0.73) | 29.2      | 1  | < 0.001 |
| Waist circumference | 5.8      | 0.34 (0.28–0.42) | 15.2      | 1  | < 0.001 |
| TGs                 | 3.9      | 0.17 (0.14–0.21) | 19.6      | 1  | < 0.001 |
| HDL-C               | 5.2      | 0.20 (0.17–0.24) | 21.7      | 1  | < 0.001 |
| Hyperglycemia       | 1.7      | 0.20 (0.15–0.27) | 7.9       | 1  | < 0.001 |
| High BP             | 21.5     | 1.23 (1.05–1.45) | 18.2      | 1  | 0.016   |
| MS                  | 0.8      | 0.10 (0.07–0.14) | 7.9       | 1  | < 0.001 |

Data are presented as number (%) and OR with 95% confidence interval.

MS = metabolic syndrome, OR = odds ratio, TG = triglyceride, HDL-C = high-density lipoprotein cholesterol, BP = blood pressure.

group (all,  $P < 0.001$ ). Regarding lifestyle factors, 54.3% of military participants were smokers and 18.5% of military participants were engaged in regular, vigorous physical activity.

The prevalence of MS and obesity was much lower in the military group than in the civilian group (0.8% vs. 7.9% and 20.6% vs. 29.2%, MS and obesity, respectively; **Table 2**). Furthermore, those components to fulfill the diagnosis of MS were different between the two groups. Compared with civilian participants, military participants reported lower prevalence for most clinical variables such as BMI (20.6% vs. 29.2%), abdominal obesity (5.8% vs. 15.2%), hyperlipidemia (5.2% vs. 21.7%), high blood TG (3.9% vs. 19.6%) and hyperglycemia (1.7% vs. 7.9%). Conversely, higher prevalence of high BP was observed among military participants than civilians (21.5% vs. 18.2%).

Age older than 25 years was the most significant risk factor associated with development of MS and obesity in the civilian group (MS, odds ratio [OR], 3.17; 95% confidence interval [CI], 1.91–5.26 and obesity, OR, 1.61; 95% CI, 1.24–2.11) in multiple logistic analysis (**Table 3**). For the military group, the effect of age was not demonstrated as no participants were above 25 years. However, smoking lowered risk of obesity (OR, 0.81; 95% CI, 0.70–0.94), and high intensity physical activity showed protective effect against MS (OR, 0.19; 95% CI, 0.04–0.78) and obesity (OR, 0.81; 95% CI, 0.66–0.99) in the military group.

We also performed other analyses to further explore risk factors for each component of MS in both groups (**Table 4**). In the civilian group, age greater than 25 years old again was one of the strongest risk factors for development of most components of MS except low HDL-C level; abdominal obesity (OR, 1.77; 95% CI, 1.26–2.49), high BP (OR, 1.91; 95% CI, 1.39–2.63), hypertriglyceridemia (OR, 2.31; 95% CI, 1.68–3.19), and hyperglycemia (OR, 2.41; 95% CI,

**Table 3.** Multiple logistic analysis of MS and obesity among Korean military personnel and civilian men aged 19–29 years

| Variables           | MS        |                   |           | Obesity    |                   |            |
|---------------------|-----------|-------------------|-----------|------------|-------------------|------------|
|                     | No. (%)   | OR                | 95% CI    | No. (%)    | OR                | 95% CI     |
| <b>Military</b>     |           |                   |           |            |                   |            |
| Age, yr             |           |                   |           |            |                   |            |
| > 25                | 0 (0.0)   | 0.00              |           | 20 (27.4)  | 1.52              | 0.90–2.55  |
| ≤ 25                | 40 (0.9)  | 1                 |           | 869 (20.0) | 1                 |            |
| Smoking             |           |                   |           |            |                   |            |
| Yes                 | 19 (0.7)  | 0.78              | 0.41–1.48 | 497 (19.1) | 0.81 <sup>b</sup> | 0.70–0.94  |
| No                  | 21 (1.0)  | 1                 |           | 491 (22.4) | 1                 |            |
| Physical activity   |           |                   |           |            |                   |            |
| Low (0–2/week)      | 30 (1.1)  | 1                 |           | 577 (21.5) | 1                 |            |
| Moderate (3–4/week) | 8 (0.7)   | 0.54              | 0.25–1.20 | 244 (20.2) | 0.94              | 0.79–1.13  |
| High (≥ 5/week)     | 2 (0.2)   | 0.19 <sup>a</sup> | 0.04–0.78 | 163 (18.4) | 0.81 <sup>a</sup> | 0.66–0.99  |
| Branch of army      |           |                   |           |            |                   |            |
| Artillery, armored  | 17 (0.7)  | 2.15              | 0.98–4.74 | 190 (23.3) | 1.23 <sup>a</sup> | 1.01–14.98 |
| Support             | 11 (1.3)  | 1.62              | 0.77–3.41 | 246 (20.0) | 1.04              | 0.87–1.24  |
| Infantry            | 12 (1.0)  | 1                 |           | 502 (19.2) | 1                 |            |
| <b>Civilians</b>    |           |                   |           |            |                   |            |
| Age, yr             |           |                   |           |            |                   |            |
| > 25                | 62 (11.4) | 3.17 <sup>b</sup> | 1.91–5.26 | 184 (33.8) | 1.61 <sup>b</sup> | 1.24–2.11  |
| ≤ 25                | 25 (4.5)  | 1                 |           | 139 (24.7) | 1                 |            |
| Smoking             |           |                   |           |            |                   |            |
| Yes                 | 38 (7.6)  | 1.15              | 0.73–1.81 | 139 (28.2) | 1.12              | 0.85–1.46  |
| No                  | 49 (8.7)  | 1                 |           | 183 (31.4) | 1                 |            |
| Physical activity   |           |                   |           |            |                   |            |
| Low (0–2/week)      | 67 (7.9)  | 1                 |           | 251 (29.4) | 1                 |            |
| Moderate (3–4/week) | 11 (8.7)  | 1.20              | 0.61–2.36 | 38 (29.9)  | 1.05              | 0.69–1.58  |
| High (≥ 5/week)     | 5 (5.3)   | 0.69              | 0.27–1.78 | 26 (27.7)  | 0.95              | 0.59–1.54  |

MS = metabolic syndrome, OR = odds ratio, CI = confidence interval.

<sup>a</sup>P value < 0.05; <sup>b</sup>P value < 0.001.

1.49–3.86). Moreover, increased intensity of physical activities only showed the trend of lowering risk on each component of MS but did not meet statistical significance as only a small portion of civilians (8.8%) were engaged in regular physical activity. Conversely, higher intensity of physical activity showed protective effect on lowering of abdominal obesity (OR, 0.54; 95% CI, 0.37–0.79), high BP (OR, 0.79; 95% CI, 0.65–0.97) and hypertriglyceridemia (OR, 0.58; 95% CI, 0.37–0.91) in the military group. Interestingly, current smokers showed lower risk of having high BP in the military group.

Moreover, there were some differences among branches of the army in the military group. Compared with the infantry branch, the support branch appeared to have lower risk of high BP (OR, 0.76; 95% CI, 0.66–0.94) but higher risk of hypertriglyceridemia (OR, 1.46; 95% CI, 1.04–2.04).

## DISCUSSION

In this study, we have identified the prevalence and risk factors for MS, obesity and its components in a large military group and compared it with the civilians of similar age. The overall baseline health status was better in the military group compared with the general population, except high BP. As expected with greater portion of military participants engaged in physical activities, the prevalence of MS and obesity was lower in the military group compared with civilians and physical activity attenuated risk of having abdominal obesity, high BP and hypertriglyceridemia in military group. As demonstrated in previous studies,

**Table 4.** Multiple logistic analysis of the factors associated with MS among Korean military personnel and civilian men aged 19–29 years

| Variables           | Abdominal obesity |                   |           | High BP    |                   |           | Hypertriglyceridemia |                   |           | Low HDL-C  |                   |           | Hyperglycemia |                   |           |
|---------------------|-------------------|-------------------|-----------|------------|-------------------|-----------|----------------------|-------------------|-----------|------------|-------------------|-----------|---------------|-------------------|-----------|
|                     | No. (%)           | HR                | 95% CI    | No. (%)    | HR                | 95% CI    | No. (%)              | HR                | 95% CI    | No. (%)    | HR                | 95% CI    | No. (%)       | HR                | 95% CI    |
| <b>Military</b>     |                   |                   |           |            |                   |           |                      |                   |           |            |                   |           |               |                   |           |
| Age, yr             |                   |                   |           |            |                   |           |                      |                   |           |            |                   |           |               |                   |           |
| > 25                | 4 (5.5)           | 0.92              | 0.33–2.55 | 16 (21.9)  | 1.04              | 0.60–1.84 | 5 (6.8)              | 1.73              | 0.69–4.36 | 5 (6.8)    | 1.31              | 0.52–3.29 | 2 (2.7)       | 1.63              | 0.39–6.77 |
| ≤ 25                | 254 (5.9)         | 1                 |           | 933 (21.5) | 1                 |           | 175 (4.0)            | 1                 |           | 233 (5.4)  | 1                 |           | 72 (1.7)      | 1                 |           |
| Smoking             |                   |                   |           |            |                   |           |                      |                   |           |            |                   |           |               |                   |           |
| Yes                 | 140 (5.4)         | 0.84              | 0.65–1.08 | 461 (17.7) | 0.61 <sup>b</sup> | 0.53–0.70 | 107 (4.1)            | 1.11              | 0.82–1.50 | 141 (5.4)  | 1.06              | 0.81–1.38 | 43 (1.6)      | 0.77              | 0.48–1.22 |
| No                  | 137 (6.2)         | 1                 |           | 571 (26.0) | 1                 |           | 82 (3.7)             | 1                 |           | 111 (5.1)  | 1                 |           | 39 (1.8)      | 1                 |           |
| Physical activity   |                   |                   |           |            |                   |           |                      |                   |           |            |                   |           |               |                   |           |
| Low (0–2/week)      | 186 (6.9)         | 1                 |           | 624 (23.2) | 1                 |           | 119 (4.4)            | 1                 |           | 133 (4.9)  | 1                 |           | 50 (1.9)      | 1                 |           |
| Moderate (3–4/week) | 57 (4.7)          | 0.66 <sup>a</sup> | 0.48–0.90 | 236 (19.5) | 0.81 <sup>a</sup> | 0.68–0.96 | 45 (3.7)             | 0.83              | 0.58–1.18 | 80 (6.6)   | 1.32              | 0.98–1.77 | 16 (1.3)      | 0.68              | 0.38–1.22 |
| High (≥ 5/week)     | 34 (3.8)          | 0.54 <sup>a</sup> | 0.37–0.79 | 169 (19.1) | 0.79 <sup>a</sup> | 0.65–0.97 | 24 (2.7)             | 0.58 <sup>a</sup> | 0.37–0.91 | 39 (4.4)   | 0.84              | 0.58–1.22 | 16 (1.8)      | 0.86              | 0.47–1.59 |
| Branch of army      |                   |                   |           |            |                   |           |                      |                   |           |            |                   |           |               |                   |           |
| Artillery, armored  | 57 (7.0)          | 1.35              | 0.96–1.89 | 179 (22.0) | 0.96              | 0.79–1.18 | 31 (3.8)             | 1.16              | 0.76–1.77 | 64 (7.9)   | 1.66 <sup>b</sup> | 1.20–2.32 | 15 (1.8)      | 1.37              | 0.75–2.51 |
| Support             | 78 (6.4)          | 1.28              | 0.95–1.71 | 230 (18.7) | 0.76 <sup>a</sup> | 0.66–0.94 | 64 (5.2)             | 1.46 <sup>a</sup> | 1.04–2.04 | 67 (5.5)   | 1.20              | 0.88–1.64 | 21 (1.7)      | 1.21              | 0.70–2.07 |
| Infantry            | 135 (5.2)         | 1                 |           | 589 (22.6) | 1                 |           | 91 (3.5)             | 1                 |           | 118 (4.5)  | 1                 |           | 39 (1.5)      | 1                 |           |
| <b>Civilians</b>    |                   |                   |           |            |                   |           |                      |                   |           |            |                   |           |               |                   |           |
| Age, yr             |                   |                   |           |            |                   |           |                      |                   |           |            |                   |           |               |                   |           |
| > 25                | 100 (18.3)        | 1.77 <sup>b</sup> | 1.26–2.49 | 124 (18.8) | 1.91 <sup>b</sup> | 1.39–2.63 | 142 (15.1)           | 2.31 <sup>b</sup> | 1.68–3.19 | 120 (21.8) | 1.06              | 0.79–1.43 | 59 (9.4)      | 2.40 <sup>b</sup> | 1.49–3.86 |
| ≤ 25                | 68 (12.1)         | 1                 |           | 78 (16.9)  | 1                 |           | 75 (26.2)            | 1                 |           | 119 (21.6) | 1                 |           | 29 (5.6)      | 1                 |           |
| Smoking             |                   |                   |           |            |                   |           |                      |                   |           |            |                   |           |               |                   |           |
| Yes                 | 65 (15.9)         | 0.89              | 0.63–1.25 | 75 (38.1)  | 0.84              | 0.61–1.16 | 116 (54.2)           | 1.98 <sup>b</sup> | 1.45–2.71 | 94 (40.0)  | 1.03              | 0.76–1.39 | 25 (29.1)     | 0.54 <sup>a</sup> | 0.33–0.88 |
| No                  | 103 (14.7)        | 1                 |           | 122 (61.9) | 1                 |           | 98 (45.8)            | 1                 |           | 141 (60.0) | 1                 |           | 61 (70.9)     | 1                 |           |
| Physical activity   |                   |                   |           |            |                   |           |                      |                   |           |            |                   |           |               |                   |           |
| Low (0–2/week)      | 134 (15.7)        | 1                 |           | 166 (19.4) | 1                 |           | 169 (19.8)           | 1                 |           | 182 (21.5) | 1                 |           | 67 (7.8)      | 1                 |           |
| Moderate (3–4/week) | 15 (11.8)         | 0.73              | 0.41–1.30 | 22 (17.3)  | 0.93              | 0.57–1.52 | 23 (18.1)            | 0.94              | 0.58–1.55 | 24 (19.0)  | 0.87              | 0.54–1.40 | 11 (8.7)      | 1.19              | 0.61–2.34 |
| High (≥ 5/week)     | 15 (16.0)         | 1.06              | 0.59–1.91 | 11 (11.7)  | 0.59              | 0.31–1.14 | 16 (17.0)            | 0.81              | 0.45–1.46 | 23 (24.5)  | 1.23              | 0.75–2.03 | 8 (8.5)       | 1.01              | 0.45–2.23 |

MS = metabolic syndrome, BP = blood pressure, HDL-C = high-density lipoprotein cholesterol, HR = hazard ratio, CI = confidence interval.

<sup>a</sup>p value < 0.05; <sup>b</sup>p value < 0.001.



smoking again showed the inverse association with obesity in the military group.<sup>9-11</sup> In the civilian group, age was the most important risk factor affecting the overall health status.

Previous studies have investigated the prevalence of obesity and MS in the military in other countries.<sup>12-20</sup> A similar tendency was observed in the comparison between US military and civilians. The prevalence of MS in military personnel was significantly lower than that of the general population.<sup>21</sup> Soldiers tend to have higher BP and smoke more but have better lipid profile, fasting glucose and lower BMI than the general public.<sup>17</sup> However, the prevalence of MS in military personnel in the current study was significantly lower than other studies.<sup>12-14,18,20</sup> The organization of the ROK armed forces is quite unique from armed forces in other countries. Korea has a mandatory draft system and most Korean men serve in the military in their early 20s. Therefore, participants in the current study are limited to young soldiers with mean age of 20.8 years old, which is significantly younger than military members of other countries.

Moreover, there are other factors contributing to low prevalence of MS in Korean military. When young Korean men are initially recruited, all of them undergo physical examinations and those individuals with underlying medical conditions including extreme BMI or high BP are either being exempt from military service or being placed in public service. Therefore, those military personnel included in the current study may have superior health status relative to the general population. However, this is not the decisive reason contributing to the health status of army members, because most of the men in Korea are recruited in the army and CHSP was performed for those who were about one-year post-recruitment. In addition, the military creates special circumstances where members are exposed to regular, high intensity physical activities. According to previous studies, military life and training had positive impact on overall health status.<sup>19,20,22-24</sup> Furthermore, this finding may imply that the prevalence of MS differs due to ethnic groups. In general, Asians have lower risk of obesity and MS than people from western countries although its prevalence in Asian countries continues to rise since the introduction of westernized diet.<sup>25</sup>

Our study analyzed the risk factors associated with MS, obesity and each component of MS in both groups using multiple logistic regression analysis. The prevalence of high BP among military members was higher than that of civilians. Stressful situations in the army may have contributed to development of high BP which was also demonstrated in previous studies.<sup>12,17</sup> Moreover, stress in the army was considered as a risk factor for newly reported hypertension after combat deployments in US army.<sup>26,27</sup> Similarly, high BP was more prevalent among active duty service soldiers in the current study. In the analysis of military branches, the infantry branch had higher risk of high BP than the artillery and armored groups. The infantry is considered to be the most active branch among the armed forces and this finding also reinforces the notion that stressful conditions may increase the risk of high BP. Therefore, proper stress management in the army will have positive effect on prevention and management of high BP. As expected, regular physical activity attenuated the risk of MS and obesity in the military group. This is in accord with previous studies where they also found that physical activity was effective in decreasing the risk of MS in both the military and general population.<sup>12,14,28-30</sup> However, physical activity did not demonstrate the positive impact against MS among civilians in this study and we think that this is because the civilian group may have been influenced by multiple factors that are beyond the scope of the current study, while the military group is under controlled circumstances where confounders may have minimal effect.

Although smoking is a well-known risk factor for high BP as well as MS, the prevalence of high BP in the current study was lower in smokers than non-smokers in the military group. This is contrary to previous findings where they showed smoking was a risk factor for hypertension and MS.<sup>31-33</sup> We thought BMI rather than smoking may have affected BP in young adults.<sup>34-37</sup> It also reflects the fact that smokers have lower BMI.<sup>38,39</sup> Our study also supports this concept by the fact fewer obese soldiers were found in the smoking group. Interestingly, age older than 25 years was a key factor affecting the prevalence of obesity, MS, and each component of MS in civilians. This finding can be explained by the fact that the lifestyle of 20-year-old men varies greatly after 25 years of age. Many men tend to start their career with more sedentary lifestyle after 25 years of age and this change in their lifestyles may have significant impact on overall health status.

Our study has several strengths. First, this study included 4,803 military personnel to estimate the prevalence of MS, which is the largest study so far. Second, the baseline characteristics of the military were compared with general population of similar age, thus well demonstrating the effect of active duty in military. Third, we analyzed the risk factors of MS, obesity and each component of MS in both military and civilian groups.

The current study also has several limitations. First, this was a retrospective, cross-sectional study which may not demonstrate the direct causal effect. Second, the age distribution between the two groups was different, which might have affected the results of this study. While the age distribution in the military was concentrated in the early 20s, data from KHANES showed even distribution throughout ages 20–29 years. However, when we performed subgroup analysis, limiting to individuals under 25 years of age, we obtained similar results (data not shown). Third, the data were absent on other well-documented risk factors for MS and obesity, such as diet, family history or other medical conditions.

This study reported that the prevalence of MS and obesity in the Korean military are lower than civilians of the same age group. However, the prevalence of high BP and current smokers are higher in the military population, emphasizing the importance of proper stress management and/or preventative programs in the military. Future, prospective clinical studies which include variables related to lifestyle interventions such as physical activity and nutrition are needed in parallel to the assessment of MS and obesity in the military.

## ACKNOWLEDGMENTS

We would like to thank the previous 5th Division Commander, Chang-Hwan Joo, and all members of the 5th Division Medical Battalion for supporting our study.

## REFERENCES

1. Eckel RH, Grundy SM, Zimmet PZ. The metabolic syndrome. *Lancet* 2005;365(9468):1415-28.  
[PUBMED](#) | [CROSSREF](#)
2. Zimmet P, Alberti KG, Shaw J. Global and societal implications of the diabetes epidemic. *Nature* 2001;414(6865):782-7.  
[PUBMED](#) | [CROSSREF](#)
3. Suh S, Baek J, Bae JC, Kim KN, Park MK, Kim DK, et al. Sex factors in the metabolic syndrome as a predictor of cardiovascular disease. *Endocrinol Metab (Seoul)* 2014;29(4):522-9.  
[PUBMED](#) | [CROSSREF](#)



4. Mottillo S, Filion KB, Genest J, Joseph L, Pilote L, Poirier P, et al. The metabolic syndrome and cardiovascular risk a systematic review and meta-analysis. *J Am Coll Cardiol* 2010;56(14):1113-32.  
[PUBMED](#) | [CROSSREF](#)
5. Suh S, Lee MK. Metabolic syndrome and cardiovascular diseases in Korea. *J Atheroscler Thromb* 2014;21 Suppl 1:S31-5.  
[PUBMED](#) | [CROSSREF](#)
6. Lim S, Shin H, Song JH, Kwak SH, Kang SM, Won Yoon J, et al. Increasing prevalence of metabolic syndrome in Korea: the Korean National Health and Nutrition Examination Survey for 1998–2007. *Diabetes Care* 2011;34(6):1323-8.  
[PUBMED](#) | [CROSSREF](#)
7. Grundy SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, et al. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute scientific statement. *Curr Opin Cardiol* 2006;21(1):1-6.  
[PUBMED](#) | [CROSSREF](#)
8. Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, et al. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* 2009;120(16):1640-5.  
[PUBMED](#) | [CROSSREF](#)
9. Albanes D, Jones DY, Micozzi MS, Mattson ME. Associations between smoking and body weight in the US population: analysis of NHANES II. *Am J Public Health* 1987;77(4):439-44.  
[PUBMED](#) | [CROSSREF](#)
10. Flegal KM, Troiano RP, Pamuk ER, Kuczmarski RJ, Campbell SM. The influence of smoking cessation on the prevalence of overweight in the United States. *N Engl J Med* 1995;333(18):1165-70.  
[PUBMED](#) | [CROSSREF](#)
11. Whitlock G, Lewington S, Sherliker P, Clarke R, Emberson J, Halsey J, et al. Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet* 2009;373(9669):1083-96.  
[PUBMED](#) | [CROSSREF](#)
12. Bauduceau B, Baigts F, Bordier L, Burnat P, Ceppa F, Dumenil V, et al. Epidemiology of the metabolic syndrome in 2045 French military personnel (EPIMIL study). *Diabetes Metab* 2005;31(4 Pt 1):353-9.  
[PUBMED](#) | [CROSSREF](#)
13. Ceppa F, Merens A, Burnat P, Mayaudon H, Bauduceau B. Military community: a privileged site for clinical research: epidemiological study of metabolic syndrome risk factors in the military environment. *Mil Med* 2008;173(10):960-7.  
[PUBMED](#) | [CROSSREF](#)
14. Costa FF, Montenegro VB, Lopes TJ, Costa EC. Combination of risk factors for metabolic syndrome in the military personnel of the Brazilian Navy. *Arq Bras Cardiol* 2011;97(6):485-92.  
[PUBMED](#) | [CROSSREF](#)
15. Kalf KG, Maya-Pelzer P, Andexer A, Deuber HJ. Prevalence of the metabolic syndrome in military and civilian flying personnel. *Aviat Space Environ Med* 1999;70(12):1223-6.  
[PUBMED](#)
16. Khazale NS, Haddad F. Prevalence and characteristics of metabolic syndrome in 111 Royal Jordanian Air Force pilots. *Aviat Space Environ Med* 2007;78(10):968-72.  
[PUBMED](#) | [CROSSREF](#)
17. McGraw LK, Turner BS, Stotts NA, Dracup KA. A review of cardiovascular risk factors in US military personnel. *J Cardiovasc Nurs* 2008;23(4):338-44.  
[PUBMED](#) | [CROSSREF](#)
18. Al-Qahtani DA, Imtiaz ML. Prevalence of metabolic syndrome in Saudi adult soldiers. *Saudi Med J* 2005;26(9):1360-6.  
[PUBMED](#)
19. Filho RT, D'Oliveira A Jr. The prevalence of metabolic syndrome among soldiers of the military police of Bahia State, Brazil. *Am J Men Health* 2014;8(4):310-5.  
[PUBMED](#) | [CROSSREF](#)
20. Payab M, Hasani-Ranjbar S, Merati Y, Esteghamati A, Qorbani M, Hematabadi M, et al. The prevalence of metabolic syndrome and different obesity phenotype in Iranian male military personnel. *Am J Men Health* 2017;11(2):404-13.  
[PUBMED](#) | [CROSSREF](#)

21. Herzog CM, Chao SY, Eilerman PA, Luce BK, Carnahan DH. Metabolic syndrome in the Military Health System based on electronic health data, 2009–2012. *Mil Med* 2015;180(1):83-90.  
[PUBMED](#) | [CROSSREF](#)
22. Blacker SD, Horner FL, Brown PI, Linnane DM, Wilkinson DM, Wright A, et al. Health, fitness, and responses to military training of officer cadets in a Gulf Cooperation Council country. *Mil Med* 2011;176(12):1376-81.  
[PUBMED](#) | [CROSSREF](#)
23. Williams AG. Effects of basic training in the British Army on regular and reserve army personnel. *J Strength Cond Res* 2005;19(2):254-9.  
[PUBMED](#)
24. Williams AG, Rayson MP, Jones DA. Effects of basic training on material handling ability and physical fitness of British Army recruits. *Ergonomics* 1999;42(8):1114-24.  
[PUBMED](#) | [CROSSREF](#)
25. Park J, Mendoza JA, O'Neil CE, Hilmers DC, Liu Y, Nicklas TA. A comparison of the prevalence of the metabolic syndrome in the United States (US) and Korea in young adults aged 20 to 39 years. *Asia Pac J Clin Nutr* 2008;17(3):471-82.  
[PUBMED](#)
26. Granado NS, Smith TC, Swanson GM, Harris RB, Shahar E, Smith B, et al. Newly reported hypertension after military combat deployment in a large population-based study. *Hypertension* 2009;54(5):966-73.  
[PUBMED](#) | [CROSSREF](#)
27. Stewart IJ, Sosnov JA, Snow BD, Batou A, Howard JT, Janak JC, et al. Hypertension after injury among burned combat veterans: A retrospective cohort study. *Burns* 2017;43(2):290-6.  
[PUBMED](#) | [CROSSREF](#)
28. Park YW, Zhu S, Palaniappan L, Heshka S, Carnethon MR, Heymsfield SB. The metabolic syndrome: prevalence and associated risk factor findings in the US population from the Third National Health and Nutrition Examination Survey, 1988–1994. *Arch Intern Med* 2003;163(4):427-36.  
[PUBMED](#) | [CROSSREF](#)
29. Kim J, Tanabe K, Yokoyama N, Zempo H, Kuno S. Association between physical activity and metabolic syndrome in middle-aged Japanese: a cross-sectional study. *BMC Public Health* 2011;11(1):624.  
[PUBMED](#) | [CROSSREF](#)
30. Stensvold D, Nauman J, Nilsen TI, Wisløff U, Slørdahl SA, Vatten L. Even low level of physical activity is associated with reduced mortality among people with metabolic syndrome, a population based study (the HUNT 2 study, Norway). *BMC Med* 2011;9(1):109.  
[PUBMED](#) | [CROSSREF](#)
31. Chen CC, Li TC, Chang PC, Liu CS, Lin WY, Wu MT, et al. Association among cigarette smoking, metabolic syndrome, and its individual components: the metabolic syndrome study in Taiwan. *Metabolism* 2008;57(4):544-8.  
[PUBMED](#) | [CROSSREF](#)
32. Ishizaka N, Ishizaka Y, Toda E, Hashimoto H, Nagai R, Yamakado M. Association between cigarette smoking, metabolic syndrome, and carotid arteriosclerosis in Japanese individuals. *Atherosclerosis* 2005;181(2):381-8.  
[PUBMED](#) | [CROSSREF](#)
33. Virdis A, Giannarelli C, Neves MF, Taddei S, Ghiadoni L. Cigarette smoking and hypertension. *Curr Pharm Des* 2010;16(23):2518-25.  
[PUBMED](#) | [CROSSREF](#)
34. Brown CD, Higgins M, Donato KA, Rohde FC, Garrison R, Obarzanek E, et al. Body mass index and the prevalence of hypertension and dyslipidemia. *Obes Res* 2000;8(9):605-19.  
[PUBMED](#) | [CROSSREF](#)
35. Shihab HM, Meoni LA, Chu AY, Wang NY, Ford DE, Liang KY, et al. Body mass index and risk of incident hypertension over the life course: the Johns Hopkins Precursors Study. *Circulation* 2012;126(25):2983-9.  
[PUBMED](#) | [CROSSREF](#)
36. Papatheanasiou G, Zerva E, Zacharis I, Papandreou M, Papageorgiou E, Tzima C, et al. Association of high blood pressure with body mass index, smoking and physical activity in healthy young adults. *Open Cardiovasc Med J* 2015;9(1):5-17.  
[PUBMED](#) | [CROSSREF](#)
37. John U, Meyer C, Hanke M, Völzke H, Schumann A. Smoking status, obesity and hypertension in a general population sample: a cross-sectional study. *QJM* 2006;99(6):407-15.  
[PUBMED](#) | [CROSSREF](#)

38. Sneve M, Jorde R. Cross-sectional study on the relationship between body mass index and smoking, and longitudinal changes in body mass index in relation to change in smoking status: the Tromso Study. *Scand J Public Health* 2008;36(4):397-407.  
[PUBMED](#) | [CROSSREF](#)
39. Chiolero A, Faeh D, Paccaud F, Cornuz J. Consequences of smoking for body weight, body fat distribution, and insulin resistance. *Am J Clin Nutr* 2008;87(4):801-9.  
[PUBMED](#) | [CROSSREF](#)