


# Postprandial Plasma Amino Acid Responses Between Standard Whey Protein Isolate and Whey Protein Isolate Plus Novel Technology

Nutrition and Metabolic Insights  
Volume 12: 1–6  
© The Author(s) 2019  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/1178638819827970



Matthew H Sharp , Matthew W Stefan, Ryan P Lowery and Jacob M Wilson

Applied Science & Performance Institute, Department of Research, Tampa, FL, USA.

## ABSTRACT

**BACKGROUND:** Muscle mass is an important determinant of metabolic health and physical function. It has previously been demonstrated that the postprandial rise in circulating essential amino acids (EAA) acts as the main stimulus for muscle protein synthesis (MPS). This study investigated postprandial plasma amino acid (AA) responses of 2 different forms of whey protein isolate (WPI) with iso-caloric and iso-nitrogenous profiles to investigate plasma concentrations of EAA.

**METHODS:** In all, 12 healthy men ( $n = 12$ ) between 19 and 32 years of age were recruited for a randomized, cross-over design, which involved consumption of protein supplements on 2 testing days separated by a 6-day washout period between conditions. On each testing day, subjects consumed either 29.6 g of WPI or WPI + io (whey protein isolate plus Ingredient Optimized Protein<sup>®</sup>) mixed with 236 mL of water. Plasma EAA and branch chain amino acid (BCAA) concentrations were assessed from whole body donated by subjects at pre-consumption and 30, 60, 90, 120, and 180 minutes post consumption.

**RESULTS:** Plasma levels of total EAA concentration was significantly greater in WPI + io at 30, 60, 90, and 120 minutes post consumption ( $P < .01$ ,  $P < .001$ ,  $P < .01$ , and  $P < .01$ , respectively). Plasma levels of total BCAA concentration was significantly greater in WPI + io at 30, 60, 90, and 120 minutes post consumption ( $P < .01$ ,  $P < .001$ ,  $P < .01$ , and  $P < .05$ , respectively) compared with WPI. For leucine, only WPI + io had elevated levels compared with pre-test at 90 minutes post consumption ( $P < .001$ ).

**DISCUSSION:** Both conditions significantly elevated EAA, BCAA, and leucine from basal levels. However, we conclude that the consumption of the treated WPI significantly raises plasma EAA, BCAA, and leucine to a greater extent compared with WPI with no treatment. Thus, supplementation with WPI that has undergone Ingredient Optimized<sup>®</sup> technology may be highly beneficial for those who partake in regular exercise, elderly individuals, or those affected by a reduced sensitivity to amino acids.

**KEYWORDS:** Amino Acids, Whey Proteins, Leucine, Bioavailability, Postprandial Period

**RECEIVED:** October 16, 2018. **ACCEPTED:** October 21, 2018.

**TYPE:** Original Research

**FUNDING:** The author(s) disclosed receipt of the following financial support for the research, authorship and/or publication of this article: Corr-Jensen Inc financially supported this work and supplied the tested conditions.

**DECLARATION OF CONFLICTING INTERESTS:** The author(s) declared the following potential conflicts of interest with respect to the research, authorship, and/or publication of this article: Corr-Jensen Inc utilizes the trademark ingredient tested in this study. This does not alter the authors' adherence to all policies set forth by this journal, *Nutrition and Metabolic Insights*, as detailed online in the submission guidelines.

**CORRESPONDING AUTHOR:** Matthew H Sharp, Applied Science & Performance Institute, Tampa, FL 33607, USA. Email: msharp@theaspi.com

## Introduction

Muscle mass is an important determinant of metabolic health and physical function.<sup>1</sup> As such, muscle mass profiles have been indicated to positively impact performance in athletic events<sup>2</sup> and activities of daily living,<sup>3,4</sup> muscle strength,<sup>5</sup> and quality of life.<sup>6</sup> Thus, the maintenance or addition of muscle mass is highly sought. The loss of muscle mass and function is underpinned by increased rates of muscle protein breakdown or a reduced muscle protein synthetic response to protein feeding.<sup>7</sup> However, the ability of protein to impact muscle health is highly dependent on (1) the amount of protein ingested,<sup>8,9</sup> (2) the quality (ie amino acid composition) of the protein source,<sup>10</sup> and (3) absorption kinetics of amino acids.<sup>11–13</sup>

Increases in muscle protein synthesis (MPS) have been attributed to the postprandial rise in circulating essential amino acids (EAA).<sup>14</sup> Consequently, protein quality has been defined as the capacity of a protein to provide EAA.<sup>15</sup> However, a subgroup of EAA known as branched chain amino acids (BCAA; valine, isoleucine, and leucine) have also been shown

to be important regulators of protein anabolism.<sup>16</sup> Leucine, in particular, has been suggested to be a direct indicator of protein quality as it has demonstrated the ability to independently stimulate MPS protein synthesis alone.<sup>17,18</sup> Therefore, BCAA and leucine content of protein sources should be recognized when considering protein quality. Whey protein (WP) is considered to be a high-quality protein source due to its high concentrations of EAA, BCAA, and leucine.<sup>11,13,19</sup> As a result, WP has become a popular supplement among those looking to maintain or increase muscle mass.<sup>20</sup>

Recently, the use of atmospheric plasma has been implemented in powdered whey protein isolate (WPI). Plasma-altered WPI powders have exhibited increased surface area<sup>21</sup> as well as positive impacts to solubility and dispersibility,<sup>22</sup> which serve as potential benefits for beverage production. Plasma modification has further been shown to alter the taste and perceived mixability of powdered protein.<sup>23</sup> Furthermore, plasma modification has demonstrated ability to alter protein structure in such a way to expose the hydrophobic pockets of a protein.<sup>24</sup> These structural



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

**Table 1.** Subject characteristics.

VARIABLE	MEAN $\pm$ SD
Age (years)	25 $\pm$ 4
Height (cm)	181.3 $\pm$ 6.7
Fat mass (kg)	18.18 $\pm$ 6.83
Fat-free mass (kg)	73.70 $\pm$ 7.0
Total mass (kg)	91.88 $\pm$ 12.60
Body fat (%)	19.3 $\pm$ 4.5

alterations have been confirmed by using a protein thermal shift, which showed an improved ability for dye to bind to the protein.<sup>25</sup> Improving protein powder's hydrophobicity enhances enzymatic degradation and ultimately promotes increased digestibility, as has been demonstrated using other protein modification methods.<sup>26</sup>

To date, no study has looked at the aggregate improvements of atmospheric plasma and protein on the blood plasma amino acid response. Therefore, the purpose of this study was to investigate the postprandial plasma amino acid (AA) responses of 2 different forms of whey protein isolate (WPI) with iso-caloric and iso-nitrogenous profiles. In a randomized, crossover trial, one serving of WPI and one serving of WPI + treatment of atmospheric plasma (WPI + io) were used to investigate postprandial response of EAA, BCAA, and leucine plasma concentrations.

## Methods

### *Study population*

In all, 12 healthy men between 19 and 32 years of age who regularly engage in whole body resistance training (2-3 days/week) were recruited and considered eligible for the study. Subjects were screened to ensure that they met and would adhere to the following criteria prior to entry into the study: (1) maintain a diet consisting of 15%-20% protein, 45%-55% carbohydrate, and 25%-30% fat according to a 3-day dietary food recall; (2) not taking performance-enhancing supplements for the previous 6 weeks; (3) non-smokers; (4) not taking amino acid supplements; (5) not using anabolic or catabolic hormones; (6) not on medication or supplements known to influence any of the variables measured in the study; and (7) free of metabolic diseases. Written informed consent was obtained from all study participants, and the protocol was approved by an external Institutional Review Board (IntegReview IRB, Austin, TX; Protocol #1101). Characteristics of the subject pool are presented in Table 1. Anthropometric data were assessed from a whole-body, dual-energy X-ray absorptiometry scan (DXA; Hologic Inc., Bedford, MA, USA).

### *Study design and protocol*

The cross-over study (n=12) involved consumption of protein supplements on 2 testing days separated by a 6-day washout

period between conditions to evaluate plasma amino acid profiles following ingestion of supplement shakes. The investigated supplements (Corr-Jensen Inc., Denver, CO, USA) were WPI and WPI + treatment (WPI + io [ioProtein®]). Both WPI and WPI + io were sourced from a single 1 kg container to ensure that both conditions were from the same supplier, batch, had the same production date, and were stored in the same manner. On preparation of samples, half was designated for WPI and half was treated (WPI + io). The treatment condition was exposed to cold atmospheric plasma to incite functional and structural changes in the protein peptide to more readily expose binding sites for enzymatic cleavage.

A total of 12 subjects reported to the laboratory between 07:00 and 08:00 after an overnight fast ( $\geq 10$  hours), and a catheter (Introcan® Safety IV Catheter; Braun Medical Inc., Bethlehem, PA, USA) was inserted into an antecubital vein and a resting blood sample was drawn at time zero (pre). Immediately thereafter, subjects ingested a bolus of 29.6 g of one of the testing conditions mixed with 236 mL of water. Following ingestion of the supplement, subjects were not allowed to consume any food products until the 3-hour time course was completed. In addition, subjects were not allowed to consume water 1 hour before or 1 hour after consumption of investigational product. Serial blood samples were collected at time 0 (immediately prior to ingestion of the protein) and at 30, 60, 90, 120, and 180 minutes after consuming the study product. Immediately after collection, blood samples were stored in ice and centrifuged for 15 minutes at 2500  $\times g$  at 4°C. The resulting plasma was stored at -80°C until analysis of amino acid concentration. This process of overnight fasting, consumption of test shake (ie 29.6 g of either WPI or WPI + io mixed with 236 mL of water), and sequential blood draws were applied to both testing days.

### *Preparation of plasma samples*

For amino acid assessment, 900  $\mu\text{L}$  serum was added to 90  $\mu\text{L}$  of a 50% trichloroacetic acid (TCA) solution, vortexed, and snap frozen in liquid nitrogen and stored at -80°C until analysis. Sample preparation for liquid chromatography with tandem mass spectrometry (LC-MS/MS) analyses was as follows: an aliquot of 25  $\mu\text{L}$  of the TCA deproteinized serum was pipetted into a glass vial (1 mL) and diluted with 125  $\mu\text{L}$  water. A solid-phase extraction tip was attached to a 1.5 mL syringe and the diluted serum sample was pulled slowly through the SPE tip by moving the piston of the syringe. Thereafter, 200  $\mu\text{L}$  of the washing solution (solution 2) was added to the glass vial and pulled through the SPE tip to remove urea and other matrix components. The syringe was removed leaving the SPE tip inside the glass vial. Then, 100  $\mu\text{L}$  of a freshly prepared elution/extraction (solution 3A/3B: 3/2% (v/v)) buffer solution was pipetted into the vial, a syringe of 0.6 mL was attached to the pipette tip, and the piston was pulled up approximately 1 cm and attached to the SPE tip. The elution/extraction buffer was drawn into the SPE tip, and the solid-phase sorbent

material was then expelled into the glass vial by pushing the piston down. This step was repeated until all SPE sorbent material from the pipette tip was expelled.

Derivatization agent (solution 4:50  $\mu\text{L}$ ) was added and the vial was vortexed vigorously for 10 seconds. The derivatization agent was allowed to react for 1 minutes, and 100  $\mu\text{L}$  of organic extraction solvent (solution 5: propyl chloroformate in chloroform) was added and the vial was vortexed vigorously for 20 seconds and was allowed to stand for 1 minutes for organic solvent phase separation. In case the organic solvent phase separation was not complete, 100  $\mu\text{L}$  of a saturated sodium chloride solution was added, and the glass vial was vortexed one more time for 10 seconds, and after 1 minute of organic solvent phase separation, an aliquot of 100  $\mu\text{L}$  of the upper organic phase was transferred into a new glass vial, and the organic solvent was evaporated until dryness at ambient temperature with high-purity nitrogen gas. The residue was dissolved with 100  $\mu\text{L}$  of a mixture of methanol/water (62/38) and pipetted into a plastic spring-loaded micro-insert and placed into an autosampler vial with a septum cap.

#### *Liquid chromatography with tandem mass spectrometry*

The liquid chromatographic separation was performed on the Phenomenex LC AAA-MS column (250 mm  $\times$  3 mm ID, 4  $\mu\text{m}$  particles) using a Waters 2695 HPLC system (Milford, MA, USA) with integrated autosampler and sample chiller set at a temperature of 10°C and a Waters column heater module controlled by a Waters temperature control module (TCM). Separation of the amino acid derivatives was achieved by gradient elution using the following gradient: 0 minutes 38% A, 13 minutes 17% A, 13.01 minutes 38% A, and 20 minutes 38% A at a flow of 0.5 mL/min at a column temperature of 35°C. Eluent A consisted of 100% water containing 10 mM ammonium formate and B consisted of 100% methanol. The sample injection volume was 10  $\mu\text{L}$ .

A Thermo Quest TSQ triple quadrupole mass spectrometer (San Jose, CA, USA) equipped with an API 2 electro spray ionization (ESI) probe was employed for analyses of the amino acid enrichments. The mass spectrometer conditions were 1 second scan time; heated capillary temperature: 325°C; spray voltage: 4.5 kV; conversion dynode: 15 kV; electron multiplier voltage: 1600 V; and sheath gas pressure: 0.62 MPa. Collision-induced dissociation (CID) spectra of the (stable isotope labeled) amino acid derivatives were obtained at an argon collision cell pressure of  $8.27 \times 10^{-6}$  MPa.

#### *Statistical analysis*

Results were obtained for plasma concentrations and were EAAs (valine, leucine, isoleucine, threonine, methionine, tryptophan, phenylalanine, and lysine), BCAAs (valine, leucine, and isoleucine), and leucine alone. Classification of amino acid

essentiality is in accordance to previous literature.<sup>27</sup> Two-way repeated measures analysis of variance (ANOVA) was used to assess plasma amino acid concentrations assuming condition (WPI and WPI + io) and time (0 minutes [pre], 30 minutes, 60 minutes, 90 minutes, 120 minutes, and 180 minutes post supplementation) as fixed factors. Whenever a significant *F*-value was obtained, a post hoc test with a Tukey adjustment was performed for multiple-comparison purposes. Incremental area under the curve (iAUC) was calculated by subtracting the baseline (ie pre) concentration from each subsequent timepoint (ie 30, 60, 90, 120, and 180 minutes) and then applying the linear trapezoidal rule to the resulting concentration. The total iAUC of the 2 conditions were analyzed by paired-sample *t*-test. The significance level was previously set at  $P < .05$ . Results are expressed as mean  $\pm$  standard error.

## Results


### *Plasma amino acid concentrations*

The results for plasma AA concentrations are displayed in Figure 1, and the raw data are represented in Table 2. Both WPI and WPI + io stimulated a significant rise in EAA, BCAA, and leucine concentration by 30 minutes post ingestion ( $P < .001$ ); however, WPI + io demonstrated a more pronounced aminoacidemia than WPI ( $P < .001$ ). For plasma EAA, WPI + io demonstrated significantly higher concentrations than WPI at 30, 60, 90 ( $P < .001$ ), and 120 minutes post ingestion ( $P < .01$ ). In addition, WPI + io produced significantly higher BCAA concentrations at 60 and 90 minutes post ingestion and higher leucine concentration at 60 minutes post ingestion compared with WPI ( $P < .001$ ). The iAUC for plasma EAA, BCAA, and leucine concentration was significantly greater in WPI + io by 55.2%, 52.6%, and 50.8%, respectively ( $P < .001$ ; Table 3).

## Discussion

This study investigated postprandial plasma concentration of EAA, BCAA, and leucine in response to ingestion of 2 isocaloric and iso-nitrogenous WPI supplements in healthy men. Despite similar rates of AA appearance in plasma, WPI + io demonstrated significantly greater EAA, BCAA, and leucine availability compared with WPI as measured by plasma concentrations responses and iAUC. In addition, both conditions significantly elevated plasma EAA, BCAA, and leucine concentrations at 30 and 60 minutes. However, only in WPI + io were EAA and BCAA concentrations significantly elevated at the 90 and 120 minutes post consumption time points, respectively. Leucine concentration for WPI + io remained significantly elevated at the 90 minutes post consumption. In addition, plasma levels of total EAA and BCAA concentration were significantly greater in WPI + io at 30, 60, 90, and 120 minutes post ingestion compared with WPI.

This study is in agreement with previous literature that the ingestion of a fast absorbing, high-quality protein source



**Figure 1.** 3-hour time course response of plasma concentrations of (a) EAA, (b) BCAA, and (c) leucine. BCAA indicates branch chain amino acid; EAA, essential amino acids; a, b, and c indicate difference in A versus B condition at a given time point ( $P < .05$ ,  $P < .01$ ,  $P < .001$ ); † and ‡ indicate difference from 0 min ( $P < .01$ ,  $P < .001$ ).

**Table 2.** Plasma concentrations of leucine,  $\Sigma$ BCAA, and  $\Sigma$ EAA for WPI and WPI + io.

	PRE	30 MINUTES	60 MINUTES	90 MINUTES	120 MINUTES	180 MINUTES
Leucine (nmol/mL)						
WPI	121.13 $\pm$ 7.92	239.24 $\pm$ 21.22 <sup>#</sup>	193.23 $\pm$ 16.39 <sup>^</sup>	163.01 $\pm$ 12.58 <sup>#</sup>	136.48 $\pm$ 11.33	127.74 $\pm$ 8.33
WPI + io	139.29 $\pm$ 13.05	304.19 $\pm$ 25.46 <sup>#,a</sup>	310.03 $\pm$ 22.07 <sup>#,c</sup>	235.95 $\pm$ 15.40 <sup>#,b</sup>	192.08 $\pm$ 13.22	160.13 $\pm$ 7.57
$\Sigma$ BCAA (nmol/mL)						
WPI	411.08 $\pm$ 26.30	724.34 $\pm$ 58.34 <sup>#</sup>	608.32 $\pm$ 44.17 <sup>#</sup>	518.08 $\pm$ 31.65	446.33 $\pm$ 34.39	413.70 $\pm$ 24.52
WPI + io	476.98 $\pm$ 43.94	919.04 $\pm$ 58.22 <sup>#,c</sup>	932.51 $\pm$ 53.44 <sup>#,c</sup>	732.56 $\pm$ 46.88 <sup>#,b</sup>	621.88 $\pm$ 43.90 <sup>*,a</sup>	528.47 $\pm$ 23.48
$\Sigma$ EAA (nmol/mL)						
WPI	788.77 $\pm$ 37.14	1295.61 $\pm$ 95.70 <sup>#</sup>	1111.57 $\pm$ 55.75 <sup>#</sup>	987.09 $\pm$ 38.09	837.52 $\pm$ 47.65	804.57 $\pm$ 31.21
WPI + io	902.35 $\pm$ 76.95	1650.96 $\pm$ 100.05 <sup>#,b</sup>	1643.44 $\pm$ 78.88 <sup>#,c</sup>	1347.98 $\pm$ 83.08 <sup>#,b</sup>	1186.24 $\pm$ 88.04 <sup>^,b</sup>	1033.14 $\pm$ 44.69

Abbreviations: BCAA, branch chain amino acid; EAA, essential amino acids; WPI, whey protein isolate; WPI + io, whey protein isolate plus Ingredient Optimized Protein®. Data are expressed as mean  $\pm$  standard error.

a, b, c Significant differences between conditions at a given time point ( $P < .05$ ,  $P < .01$ , and  $P < .001$ ).

\*, ^, # Difference from PRE  $P < .05$ ,  $P < .01$ , and  $P < .001$ .

results in a robust rise in plasma amino acid levels in as early as 30 minutes.<sup>13,28</sup> For instance, Burke et al<sup>28</sup> demonstrated that postprandial AA levels of a fast absorbing protein (ie skim milk) peaked at 50 minutes, compared with

100 minutes for slower digesting proteins. In addition, the authors of this study found marked differences in plasma AA availability, despite both conditions being iso-nitrogenous, thus suggesting a difference in absorption kinetics between

**Table 3.** Plasma leucine, BCAA, and EAA expressed as iAUC.

	WPI	WPI + IO
iAUC × 10 <sup>2</sup> (min·nmol/mL)		
Leucine	78.52 ± 0.31	159.72 ± 0.47*
ΣBCAA	201.89 ± 0.74	426.61 ± 1.23*
ΣEAA	335.07 ± 1.16	747.58 ± 2.03*

Abbreviations: BCAA, branch chain amino acid; EAA, essential amino acids; iAUC, incremental area under the curve; WPI, whey protein isolate; WPI + io, whey protein isolate plus Ingredient Optimized Protein®.

Data are expressed as mean ± standard error.

\*Significantly greater than WPI ( $P < .001$ ).

two forms of fast absorbing, iso-caloric and iso-nitrogenous protein sources.

Both WPI and WPI + io of this study were sourced from the same product batch and therefore had similar AA profiles. However, the rise in EAA, BCAA, and leucine was of greater amplitude and lasted longer in WPI + io compared with WPI. These differences in circulating amino acids are likely due to WPI + io being treated with cold atmospheric plasma to provoke structural changes in protein peptides to more readily expose binding sites for enzymatic cleavage. This treatment has been shown to expose hydrophobic pockets of protein and increase protein surface hydrophobicity by as much as 20%.<sup>24</sup> Bioavailability of a protein depends on its ability to cross the intestinal mucosa and enter systemic circulation.<sup>29</sup> Proteins with a more hydrophobic surface can permeate the epithelial barrier with more efficiency than proteins with a hydrophilic surface,<sup>30</sup> thereby increasing bioavailability. Furthermore, it has been demonstrated that increasing hydrophobicity of WPI enhances enzymatic degradation, ultimately promoting greater bioavailability.<sup>26</sup>

Previous research showing divergent MPS responses to the same dose of protein suggest that the speed at which AA appear in the blood, and the leucine content of the protein, are the primary factors that determine the magnitude of the MPS response.<sup>31,32</sup> Furthermore, in a rested state, plasma availability of leucine may be more important than the rate of aminoacidemia in determining the MPS response.<sup>8,33</sup> As such, the similar rapid aminoacidemia resulting from both conditions may be more relevant to post exercise feeding, in which the muscle is sensitized to protein feeding.<sup>11,32</sup>

## Conclusions

In summary, we report that the ingestion of the treated WPI significantly raises plasma EAA, BCAA, and leucine compared with WPI with no treatment. Furthermore, the rise in EAA, BCAA, and leucine was extended to a larger degree in the treated group compared with the non-treated group. Thus, the technical application of treating WPI with plasma surface modification to further expose hydrophobic pockets and

increase enzymatic degradation appears to promote greater concentrations of circulating EAA, BCAA, and leucine. Future research should consider larger sample sizes to denote greater levels of significance between conditions. Furthermore, a similar design could be carried out in an exercising, not resting, condition to investigate the impact of these different proteins on aminoacidemia.

## Author Contributions

JMW and RPL were involved in conceptualizing the study and structuring the study design. MHS and MWS were involved in carrying out study procedures and data collection. MHS performed statistical analysis of study data. All authors assisted in writing the manuscript for submission and approved the final version of the manuscript. The results provided in this manuscript do not constitute endorsement of the product by the authors.

## ORCID iD

Matthew H Sharp  <https://orcid.org/0000-0003-2018-9229>

## REFERENCES

- von Haehling S, Morley JE, Anker SD. An overview of sarcopenia: facts and numbers on prevalence and clinical impact. *J Cachexia Sarcopenia Muscle*. 2010;1:129-133. doi:10.1007/s13539-010-0014-2.
- Perez-Gomez J, Rodriguez GV, Ara I, et al. Role of muscle mass on sprint performance: gender differences? *Eur J Appl Physiol*. 2008;102:685-694. doi:10.1007/s00421-007-0648-8.
- Freeman W. The relationship between habitual dietary protein intake and dual task performance in sedentary, recreationally active, and masters athlete older adults. *Health, human performance and recreation undergraduate honors theses*. May, 2017. <http://scholarworks.uark.edu/hhpruht/51>.
- Visser M, Kritchevsky SB, Goodpaster BH, et al. Leg muscle mass and composition in relation to lower extremity performance in men and women aged 70 to 79: the health, aging and body composition study. *J Am Geriatr Soc*. 2002;50:897-904. doi:10.1046/j.1532-5415.2002.50217.x.
- Folland JP, Williams AG. The adaptations to strength training: morphological and neurological contributions to increased strength. *Sports Med*. 2007;37:145-168. doi:10.2165/00007256-200737020-00004.
- Trombetti A, Reid KF, Hars M, et al. Age-associated declines in muscle mass, strength, power, and physical performance: impact on fear of falling and quality of life. *Osteoporos Int*. 2016;27:463-471. doi:10.1007/s00198-015-3236-5.
- Churchward-Venne TA, Breen L, Donato D, et al. Leucine supplementation of a low-protein mixed macronutrient beverage enhances myofibrillar protein synthesis in young men: a double-blind, randomized trial. *Am J Clin Nutr*. 2014;99:276-286. doi:10.3945/ajcn.113.068775.
- Mitchell WK, Phillips BE, Williams JP, et al. A dose-rather than delivery profile-dependent mechanism regulates the "muscle-full" effect in response to oral essential amino acid intake in young men. *J Nutr*. 2015;145:207-214. doi:10.3945/jn.114.199604.
- Witard OC, Jackman SR, Breen L, Smith K, Selby A, Tipton KD. Myofibrillar muscle protein synthesis rates subsequent to a meal in response to increasing doses of whey protein at rest and after resistance exercise. *Am J Clin Nutr*. 2014;99:86-95. doi:10.3945/ajcn.112.055517.
- Atherton PJ, Kumar V, Selby AL, et al. Enriching a protein drink with leucine augments muscle protein synthesis after resistance exercise in young and older men. *Clin Nutr*. 2017;36:888-895.
- Burd NA, Yang Y, Moore DR, Tang JE, Tarnopolsky MA, Phillips SM. Greater stimulation of myofibrillar protein synthesis with ingestion of whey protein isolate v. micellar casein at rest and after resistance exercise in elderly men. *Br J Nutr*. 2012;108:958-962.
- Pennings B, Boirie Y, Senden JMG, Gijsen AP, Kuipers H, van Loon LJC. Whey protein stimulates postprandial muscle protein accretion more effectively

- than do casein and casein hydrolysate in older men. *Am J Clin Nutr*. 2011;93:997–1005. doi:10.3945/ajcn.110.008102.
13. Tang JE, Moore DR, Kujbida GW, Tarnopolsky MA, Phillips SM. Ingestion of whey hydrolysate, casein, or soy protein isolate: effects on mixed muscle protein synthesis at rest and following resistance exercise in young men. *J Appl Physiol*. 2009;107:987–992. doi:10.1152/jappphysiol.00076.2009.
  14. Fujita S, Dreyer HC, Drummond MJ, et al. Nutrient signalling in the regulation of human muscle protein synthesis. *J Physiol*. 2007;582:987–992. doi:10.1113/jphysiol.2007.134593.
  15. Campbell B, Kreider RB, Ziegenfuss T, et al. International Society of Sports Nutrition position stand: protein and exercise. *J Int Soc Sports Nutr*. 2007;4:8. doi:10.1186/1550-2783-4-8.
  16. Kimball SR, Jefferson LS. Signaling pathways and molecular mechanisms through which branched-chain amino acids mediate translational control of protein synthesis. *J Nutr*. 2006;136:227S–231S. doi:10.1093/jn/136.1.227S.
  17. Suryawan A, Jeyapalan AS, Orellana RA, Wilson FA, Nguyen HV, Davis TA. Leucine stimulates protein synthesis in skeletal muscle of neonatal pigs by enhancing mTORC1 activation. *Am J Physiol Endocrinol Metab*. 2008;295:E868–E875. doi:10.1152/ajpendo.90314.2008.
  18. Norton LE, Layman DK, Bunpo P, Anthony TG, Brana DV, Garlick PJ. The leucine content of a complete meal directs peak activation but not duration of skeletal muscle protein synthesis and mammalian target of rapamycin signaling in rats. *J Nutr*. 2009;139:1103–1109. doi:10.3945/jn.108.103853.
  19. Bucci L, Unlu L. Proteins and amino acid supplements in exercise and sport. In: Driskell JA, Wolinsky I eds. *Energy-Yielding Macronutrients and Energy Metabolism in Sports Nutrition*. Boca Raton, FL: CRC Press; 2000:191–212.
  20. Mitchell CJ, D'Souza RF, Fanning AC, Poppitt SD, Cameron-Smith D. Short communication: muscle protein synthetic response to microparticulated whey protein in middle-aged men. *J Dairy Sci*. 2017;100:4230–4234. doi:10.3168/jds.2016-12287.
  21. Motosko C, Zakhem G. Effect of atmospheric plasma on the surface area of powdered whey protein isolate. *Matters*. 2017;3. doi:10.19185/matters.201707000006. <https://sciencematters.io/articles/201707000006>.
  22. Lightfoot H. The effect of atmospheric plasma on the solubility and dispersibility of powdered whey protein isolate. May, 2018. doi:10.31232/osf.io/rh8un. <https://osf.io/preprints/nutrixiv/rh8un/>.
  23. Lightfoot H. The effect of atmospheric plasma on the perception of taste and mixability of powdered whey protein isolate. May, 2018. doi:10.31232/osf.io/nyt9e. <https://osf.io/preprints/nutrixiv/nyt9e/>.
  24. Lightfoot H. The effect of atmospheric plasma on the hydrophobicity of powdered whey protein isolate. May, 2018. doi:10.31232/osf.io/hx75f. <https://osf.io/preprints/nutrixiv/hx75f/>.
  25. Aguilar D. The effect of atmospheric plasma on a protein thermal shift assay of powdered whey protein isolate. May, 2018. doi:10.31232/osf.io/pxdqb. <https://osf.io/preprints/nutrixiv/pxdqb/>.
  26. del Castillo-Santaella T, Sanmartín E, Cabrerizo-Vilchez MA, Arboleya JC, Maldonado-Valderrama J. Improved digestibility of  $\beta$ -lactoglobulin by pulsed light processing: a dilatational and shear study. *Soft Matter*. 2014;10:9702–9714. doi:10.1039/c4sm01667j.
  27. Grimble GK. Essential and conditionally-essential nutrients in clinical nutrition. *Nutr Res Rev*. 1993;6:97–119. doi:10.1079/NRR19930008.
  28. Burke LM, Winter JA, Cameron-Smith D, Ensen M, Farnfield M, Decombaz J. Effect of intake of different dietary protein sources on plasma amino acid profiles at rest and after exercise. *Int J Sport Nutr Exerc Metab*. 2012;22:452–462.
  29. Kwan KC. Oral bioavailability and first-pass effects. *Drug Metab Dispos*. 1997;25:1329–1336.
  30. Miklavžin A, Cegnar M, Kerč J, Kristl J. Effect of surface hydrophobicity of therapeutic protein loaded in polyelectrolyte nanoparticles on transepithelial permeability. *Acta Pharmaceut*. 2018;68:275–293. doi:10.2478/acph-2018-0032.
  31. Katsanos CS, Kobayashi H, Sheffield-Moore M, Aarsland A, Wolfe RR. A high proportion of leucine is required for optimal stimulation of the rate of muscle protein synthesis by essential amino acids in the elderly. *Am J Physiol Endocrinol Metab*. 2006;291:E381–E387.
  32. West DWD, Burd NA, Coffey VG, et al. Rapid aminoacidemia enhances myofibrillar protein synthesis and anabolic intramuscular signaling responses after resistance exercise. *Am J Clin Nutr*. 2011;94:795–803. doi:10.3945/ajcn.111.013722.
  33. Mitchell CJ, McGregor RA, D'Souza RF, et al. Consumption of milk protein or whey protein results in a similar increase in muscle protein synthesis in middle aged men. *Nutrients*. 2015;7:8685–8699. doi:10.3390/nu7105420.