

RESEARCH

Open Access



# Birth and Regulation of head Extension to Guide Manual perineal Assistance (BREGMA) study: a prospective cohort study

Abdelrahman Abouelhasan<sup>1,2,3,6</sup>, Vladimir Kalis<sup>3,4,5</sup>, Marc Schüssler<sup>2</sup>, Hassan Awwad<sup>1</sup>, Rania H. M. Ahmed<sup>1</sup> and Khaled M. Ismail<sup>3,5\*</sup>

## Abstract

**Background** The exact positioning of the dominant hand at the time of manual perineal protection (MPP) is not clearly specified. The main aim of this work was to identify the place on the perineum where pressure needs to be applied to achieve optimal forward fetal head displacement away from the anal sphincters using the bregma to posterior fourchette distance (BFD) and the perineal body length (PBL).

**Methods** This was a two-center prospective cohort study. Term cephalic singleton nulliparous women having spontaneous vaginal delivery were considered eligible for recruitment into the study. Once crowning was diagnosed and just prior the initiation of manual perineal protection (MPP) or cutting an episiotomy, the BFD and PBL were measured using a standardized measurement protocol.

**Results and discussion** A total of 100 women (50 women in each center) were recruited into the study. The overall mean BFD was  $2.8 \pm 0.5$  cm. The overall mean PBL was  $4.4 \pm 0.8$  cm. There were no statistically significant differences between the measurements taken in both units regarding BFD or PBL ( $p = 0.81$  and  $0.10$  respectively). There was a weak correlation between both measurements. Based on our measured parameters, it seems that the most effective point to apply perineal pressure to displace the head away from the anus is approximately 1 – 1.5 cm anterior to the anal margin.

**Conclusion** Measurements generated by this study will form the bases of future biomechanical studies to confirm their validity.

**Keywords** Manual perineal protection, Perineal mapping, Perineal body, Perineal tension

\*Correspondence:

Khaled M. Ismail  
khaled.ismail@lfp.cuni.cz

<sup>1</sup> Department of Obstetrics and Gynecology, Faculty of Medicine, Ain Shams University, 38 Abbassia, Next to the Al-Nour Mosque, Cairo 1181, Egypt

<sup>2</sup> Department of Obstetrics & Gynecology, Agaplesion General Hospital Hagen (Allgemeines Krankenhaus Hagen), Hagen, Germany

<sup>3</sup> Department of Obstetrics and Gynecology, Faculty of Medicine in Pilsen, Charles University, Alej Svobody 76, Pilsen 304 60, Czech Republic

<sup>4</sup> Department of Obstetrics and Gynecology, University Hospital, Pilsen, Czech Republic

<sup>5</sup> Biomedical Center, Faculty of Medicine in Pilsen, Charles University, Pilsen, Czech Republic

<sup>6</sup> Present Address: Department of Obstetrics & Gynecology, Feldkirch Academic Teaching Hospital, Carinagasse 47, Feldkirch 6800, Austria



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

## Background

Studies investigating perineal trauma at the time of vaginal birth found that primiparity, forceps delivery, midline episiotomy, persistent occipitoposterior position, and shoulder dystocia were significant risk factors [1]. Manual perineal protection (MPP) is an intrapartum intervention that is used to reduce the risk of severe perineal trauma and its consequences [2–4]. However, there is lack of complete understanding of how to accurately execute the technique and the intended value to be achieved by each of its components [5–9]. We believe that these factors, and the lack of objective evidence of how the technique relates to the dynamic changes occurring at the fetal head and the perineum at the time of crowning, are important limitations to the wide adoption of MPP amongst professionals and might account for the variation in the reproducibility of the technique's effectiveness in reducing perineal trauma.

Biomechanical analysis of a variety of MPP techniques that reduce the tension at the midpoint of the perineum identified the Finnish and Viennese maneuvers to be the most effective [4, 10–12]. Stereo-photogrammetric analysis and biomechanical studies have optimized the positioning of the thumb and finger(s) of the dominant hand to minimize tension at the midpoint of the perineum [10–13]. However, in addition to the reduction in perineal tension, both techniques also aim to regulate fetal head extension, while pushing it away from the anal sphincter complex, using the flexed middle finger or the ulnar border of the dominant hand respectively [14–17]. This component is considered to be important for the effectiveness of MPP [18].

In an occipital-anterior position, the bregma is the most posterior fetal head landmark in relation to the maternal perineum at the time of crowning. Therefore, to achieve the required anterior displacement of the head away from the anal sphincters, the pressure needs to be applied at a point on the perineum between the bregma and the anal margin. However, the normal range and variation of the position of the bregma in relation to the distended perineum at the time of crowning is not known.

## Objectives

The primary aim of this work was to identify the optimal place on the perineum where pressure needs to be applied during MPP to achieve optimal forward fetal head displacement away from the anal sphincters. We proposed that this could be achieved by collecting data regarding the position of the bregma in relation to the perineum at the time of crowning using the bregma to the posterior fourchette distance (BFD) and the perineal body length (PBL) as objective surrogate measures. As a secondary aim we wanted to assess the degree of


variability between these measurements in a diversified population.

## Methods

This is a prospective cohort study conducted in 2 maternity units in Hagen, Germany (unit 1) and in Cairo, Egypt (unit 2). Women having their first spontaneous vaginal birth of a singleton term baby in vertex presentation and occipitoanterior position were considered eligible to participate in the study. Suspected fetal distress or fetal anomalies, occipitoposterior position, operative vaginal delivery, and history of vaginal or perineal operations were reasons for exclusion. Women meeting the inclusion criteria were informed about the study on admission to the labor ward during the first stage of labor. Favorable ethical approvals were granted to the study by the Hospital Board of Management at Allgemeines Hospital Hagen on 02.01.2020 and the Research Ethics Committee at Ain Shams University on 05.01.2021 (Ref. FMASU MS 704/2020/2021) (supplementary information). All study participants provided a valid written informed consent.

Intrapartum care was undertaken based on the participating unit's protocol. One assessor (AA) performed all the measurements in both units. Measurements were performed while the woman was in a semi-recumbent lithotomy position with the legs flexed at the hip joints at an angle between 90° and 100° [19, 20]. Disposable sterile surgical skin marker and tape measure kits were used for the assessments (Viscot Medical, LLC; East Hanover, NJ 07936). The fixed reference points were the posterior fourchette (the posterior margin of the vestibular (navicular) fossa), and the most anterior point on the anal epithelium at 12 o'clock from the center of the anal orifice [21]. PBL was defined as the distance between the posterior fourchette and the anterior margin of the anal opening [21]. We used the anterior margin of the anal opening rather than the center of the anus, as proposed by the International Continence Society, to mitigate inaccuracies resulting from the varying degrees of anal dilatation that may occur at the time of fetal head crowning.

Measurements were taken once crowning was diagnosed and just prior the initiation of MPP or cutting an episiotomy, whichever came first. The index finger was inserted through the vaginal introitus in the midline between the fetal head and perineum to palpate the fetal bregma. Using a sterile skin marker, a point was marked externally in the midline on the perineal skin corresponding to the point at which the center of the fetal bregma was palpated. Using a sterile tape measure, the distance between this marked point on the perineum and posterior fourchette was measured (BFD). Using the same tape measure, the distance between the posterior fourchette and the anterior margin of the anal opening, (at 12



**Fig. 1** Simulation of how the BFD and PBL measurements were assessed

o'clock) was then measured (PBL) (Fig. 1). Additionally, we collected data on maternal age, BMI, duration of second stage of labor, birth weight, neonatal head circumference and any associated perineal trauma including episiotomy.

A convenience sample of 50 women per unit was set a priori for the study. Statistical analysis was performed using SAS 9.4 (SAS Institute Inc., Cary, NC, USA) statistical software. Difference in distribution of ranked data between groups was carried out with non-parametric ANOVA (2-sample Wilcoxon test or 2-sample Median test), while categorical variables were assessed by the  $\chi^2$  test or Fisher's exact test. The Pearson correlation and linear regression were used to assess the relation between PBL and BFD. A  $p < 0.05$  was considered statistically significant.

## Results

The study was conducted between January 2, 2020, and August 30, 2020, at unit 1 and between January 5, 2021, and May 30, 2021, at unit 2. A total of 100 primiparous women were recruited and contributed full data into the study in the two units. The number of patients


approached and recruited to achieve the target sample is presented in Fig. 2. The demographic details and obstetrics variables are presented in Table 1. There was a statistically significant difference between the two centers regarding duration of the 2nd stage of labor ( $p = 0.004$ ) and neonatal weight ( $p = 0.02$ ). The overall mean BFD was  $2.8 \pm 0.5$  cm ( $2.8 \pm 0.6$  cm in unit 1 and  $2.8 \pm 0.5$  cm in unit 2). The overall mean PBL was  $4.4 \pm 0.8$  cm ( $4.5 \pm 0.8$  cm in unit 1 and  $4.3 \pm 0.7$  cm in unit 2). There were no statistically significant differences between both units regarding BFD or PBL ( $p = 0.81$ , and  $0.10$  respectively).

On assessing the relationship between BFD and PBL using Pearson coefficient, there was low correlation between both measurements ( $r = 0.4820$ ) (Fig. 3).

## Discussion

### Summary of findings

To our knowledge, this is the first study to generate data for BFD and PBL at time of initiation of MPP. Our study demonstrated that there were no statistically significant differences between these measurements in two cohorts of women recruited in a unit in Germany and another one in Egypt despite some significant differences in



\* Not feasible to take measurements because the participant was too distressed, labor progressed too fast or the assessor was not available at time of birth.

Fig. 2 Flowchart of patient flow into the study

participants’ demographics. This observation suggests that ethnicity does not seem to affect these measurements; nonetheless, this finding may be limited by our sample size. There was no statistically significant difference regarding maternal age between the two groups. However, there was a statistically significant difference in neonatal weight where the mean birthweight was higher in Germany compared to Egypt. Exploring the reasons for this difference is beyond the scope of our work, however, it is plausible that constitutional and / or nutritional factors have contributed to such difference.

When comparing the duration of second stage of labor between the two centers, there was a statistically significant difference between the two groups. This could be due to variations in the management protocols adopted by the two centers. Indeed, at Ain Shams University, getting women to commence active pushing as soon as full cervical dilatation is confirmed is the preferred policy to shorten the second stage of labor as much as possible to be able to manage the large number of women admitted in labor per day. Moreover, in the absence of a midwifery service in Egypt, labor is totally managed by a medical

team, which might have had an impact on the observed difference.


**Comparison of findings to other studies**

Although our study group was not the first one to measure the PBL, to our knowledge, we are the first to measure it at the time of initiation of MPP and the first to propose and measure the BFD. Rizk et al. studied the relation between perineal length and position of the anus at the time of vaginal delivery in primigravidae. They measured PBL and the distance between the fourchette and the inferior margin of the coccyx to calculate the anal position index. Unlike our study, these measurements were obtained at the first stage of labor. In their study a short perineum and anterior displacement of the anus were associated with a traumatic vaginal delivery [22]. Interestingly, Rizk and associates’ reported mean PBL measurement of 4.6 cm is comparable to the mean PBL in our study despite the difference in the stage in labor when the measurements were taken. In contrast, the mean PBL measured in our study was shorter than that reported by Meriwhether et al. where their mean reported PBL increase from 3.7 cm

**Table 1** The demographic data and obstetrics variables in the study group

Obstetric variable	Cohort	Mean ±SD	Median (IQR)	p-value
Maternal age, years	overall	28.0 ± 5.9	27.5 (24.0 – 32.0)	0.46 *
	unit 1	28.4 ± 6.2	27.5 (24.0 – 33.0)	
	unit 2	27.5 ± 5.6	27.5 (24.0 – 31.0)	
Body mass index, kg/m <sup>2</sup>	overall	29.3 ± 4.2	29.0 (26.0 – 32.0)	0.63 *
	unit 1	29.4 ± 4.3	30.0 (26.0 – 32.4)	
	unit 2	29.2 ± 4.0	29.0 (26.0 – 32.0)	
Duration of the 2nd stage of labor, min	overall	82.3 ± 24.9	83.5 (65.0 – 98.0)	0.004 *
	unit 1	89.7 ± 27.0	92.0 (70.0 – 103.0)	
	unit 2	74.8 ± 20.3	75.0 (60.0 – 90.0)	
Birth weight, g	overall	3446 ± 403	3440 (3225—3670)	0.02 *
	unit 1	3531 ± 463	3535 (3300—3820)	
	unit 2	3361 ± 314	3345 (3150—3550)	
Neonatal head circumference, cm	overall	35.3 ± 1.5	35.0 (34.0 – 36.5)	0.38 *
	unit 1	35.5 ± 1.6	35.0 (34.0 – 37.0)	
	unit 2	35.1 ± 1.4	35.5 (34.0 – 36.0)	
Bregma – fourchette distance, cm	overall	2.8 ± 0.5	2.8 (2.5 – 3.2)	0.81 *
	unit 1	2.8 ± 0.6	2.9 (2.4 – 3.2)	
	unit 2	2.8 ± 0.5	2.8 (2.5 – 3.1)	
Perineal body length, cm	overall	4.4 ± 0.8	4.2 (3.9 – 4.9)	0.10 *
	unit 1	4.5 ± 0.8	4.4 (3.9 – 5.1)	
	unit 2	4.3 ± 0.7	4.1 (3.8 – 4.6)	

\* Wilcoxon Two Sample Test



**Fig. 3** Relationship between BFP and PBL using Pearson coefficient

during the first stage to a maximum of 6.1 cm during the second stage [23]. Nonetheless, in Rizk et al. and Meriwhether et al. studies PBL was defined as the distance between the posterior fourchette and the center of the anal orifice [22, 23]. However, we opted to use the anterior anal margin rather than the center of the anus to mitigate the potential inaccuracies caused by the varying degrees of anal dilatation at the later stages of labor [21]. Hence, it is plausible that, not only the difference in the stage of labor at which measurements were taken but also the reference point used for measurement could have contributed to some of the unexpected comparisons. Furthermore, Pihl and colleagues, studied the relation between the anovaginal distance (AVD) measured by transperineal ultrasound and the occurrence of external anal sphincter injury before suturing the perineal laceration. They reported that a short AVD could be a warning sign and should increase the awareness of accoucheurs of possible external sphincter injury before suturing a perineal tear [24]. Nevertheless, they measured the AVD after the vaginal birth and before suturing a perineal laceration, which does not reflect the actual perineal length at the time of crowning.

### Optimizing MPP dominant hand placement

Jansova et al. investigated the role of the thumb and index finger in manual perineal protection and concluded that the optimal placement of the thumb and index finger is 2 cm anterior of the fourchette and 12 cm apart while the most effective distance and direction of movement is to approximate these digits by 1 cm on either side without changing their antero-posterior orientation to the fourchette [10–12]. We believe that the results of this study provide important additional clinical information for the optimization of the placement of the flexed middle finger, in the Finnish technique, or the ulnar border of the hand, in the Viennese technique, to facilitate the delivery of the fetal head through the perineum with the least possible head circumference. Indeed, the lack of such measurements to objectively describe the degree of flexion of the fetal head at the time of instigating MPP, has been a major limitation for studies investigating this intervention. Based on our findings and the difference between the BFD and the PBL, it seems that for the pressure, by the middle finger or the ulnar border of the hand depending on the technique used, to achieve effective displacement of the fetal head at the time of crowning, it needs to be applied approximately 1 – 1.5 cm anterior to the anal margin.

### Strengths and limitations

We appreciate that our study has some limitations. The lack of availability of data on BFD and PBL at the time of crowning hindered our ability to perform a priori formal sample size calculation for this work. However, we were able to recruit a well-defined cohort of women from different ethnic backgrounds and were able to generate a fairly homogenous dataset for these measurements. Additionally, it could be argued that having one assessor collecting all the data could be a potential source for bias. Nevertheless, this design has negated the possibility of inter-rater variability.

### Conclusion

This is the first study to generate data on the bregma to fourchette distance and the perineal body length just prior to fetal head expulsion. Based on our measured parameters the point at which perineal pressure at the time of manual perineal protection could be most effective seems to be approximately 1 – 1.5 cm anterior to the anal margin. This information will form the basis of future *in silico* trials on birthing virtual twins to confirm their mechanistic validity.

### Abbreviations

BFD	Bregma to posterior fourchette distance
PBL	Perineal body length
MPP	Manual perineal protection

### Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12884-025-07428-z>.

Supplementary Material 1.

### Acknowledgements

Not applicable.

### Authors' contributions

AA performed literature search, developed protocol, collected, and analyzed data, and prepared the first draft of manuscript. VK conceived the idea, performed literature search, contributed to data analysis and interpretation, edited, and revised the manuscript. MS contributed to data interpretation. HA contributed to data interpretation. RA contributed to data interpretation. KI developed protocol, performed literature search, contributed to data analysis and interpretation, edited and revised the manuscript.

### Funding

This work was supported by the Integration of biomedical research and health care in the Pilsen metropolitan area"; reg. no. CZ.02.01.01/00/23\_021/0008828) - co-funded by the European Union and by the State Budget of the Czech Republic and by the Cooperatio program, Charles University. Funders were not involved in the design, analysis, or the reporting of this work. The funders did not have a role in the collection, analysis, and interpretation of data and in the writing of the manuscript.

### Data availability

The datasets used and analysed during the current study are available from the corresponding author on reasonable request.



## Declarations

### Ethics approval and consent to participate

Favorable ethical approvals were granted to the study by the Hospital Board of Management at Allgemeines Hospital Hagen on 02.01.2020 and the Research Ethics Committee at Ain Shams University on 05.01.2021 (Ref. FMASU MS 704/2020/2021). All study participants provided a valid written informed consent. All methods were performed in accordance with the relevant guidelines and regulations.

### Consent for publication

Not applicable.

### Competing interests

The authors declare no competing interests.

Received: 5 March 2023 Accepted: 5 March 2025

Published online: 21 March 2025

## References

- Third- and Fourth-degree Perineal Tears, Management (Green-top Guideline No. 29). <https://www.rcog.org.uk/en/guidelines-research-services/guidelines/gtg29/>. Accessed 9 Jan 2022.
- Aglijaloro M. The contribution of the Salerno School to the historical evolution of protection and suturing of the perineum in labor. *Minerva Med.* 1970;61:5201–3.
- Deweese WB. Relaxation and Management of the Perineum During Parturition. Read before the Golden Belt District Medical Society of Kansas, July 4, 1889. *JAMA.* 1889;XIII(24):841–8.
- Ismail KMK, Paschetta E, Papoutsis D, Freeman RM. Perineal support and risk of obstetric anal sphincter injuries: A Delphi survey. *Acta Obstetrica et Gynecologica Scandinavica.* 2015;94(2):165–74.
- Kleprlikova H, Kalis V, Lucovnik M, Rusavy Z, Blaganje M, Thakar R, et al. Manual perineal protection: The know-how and the know-why. *Acta Obstetrica et Gynecologica Scandinavica.* 2020;99(4):445–50.
- Kalis V, Rusavy Z, Havelkova L, Zitka T, et al. Metrics of perineal support (MOPS) study. *BMC Pregnancy Childbirth.* 2020;20:1–6.
- Manresa M, Kalis V, de Tayrac R, de Leeuw JW, Laine K, Räisänen S, et al. Hands up if you do not understand hands on. *Midwifery.* 2020;90:102836.
- Ali-Masri H, Hassan S, Fosse E, Zimmo KM, Zimmo M, Ismail KMK, et al. Impact of electronic and blended learning programs for manual perineal support on incidence of obstetric anal sphincter injuries: A prospective interventional study. *BMC Med Educ.* 2018;18(1):258.
- Ali HY, Vikanes Å, Anti M, Hassan S, Ismail KM, Zimmo K, et al. Evaluation of an animated instructional video as a training tool for manual perineum support during vaginal delivery. *Int J Gynecol Obstet.* 2017;137(2):213–9.
- Jansova M, Kalis V, Lobovsky L, Hyncik L, Karbanova J, Rusavy Z. The role of thumb and index finger placement in manual perineal protection. *Int Urogynecol J.* 2014;25(11):1533–40.
- Jansova M, Kalis V, Rusavy Z, Zemčik R, Lobovsky L, Laine K. Modeling manual perineal protection during vaginal delivery. *Int Urogynecol J.* 2014;25(1):65–71.
- Jansova M, Kalis V, Rusavy Z, Räisänen S, Lobovsky L, Laine K. Fetal head size and effect of manual perineal protection. *PLoS One.* 2017;12(12):e0189842.
- Zemčik R, Karbanova J, Kalis V, Lobovský L, Jansová M, Rusavy Z. Stereophotogrammetry of the perineum during vaginal delivery. *Int J Gynecol Obstet.* 2012;119(1):76–80.
- Laine K, Pirhonen T, Rolland R, Pirhonen J. Decreasing the incidence of anal sphincter tears during delivery. *Obstet Gynecol.* 2008;111(5):1053–7.
- Laine K, Skjeldestad FE, Sandvik L, Staff AC. Incidence of obstetric anal sphincter injuries after training to protect the perineum: Cohort study. *BMJ Open.* 2012;2(5):e001649.
- Poulsen MO, Madsen ML, Skriver-Møller AC, Overgaard C. Does the Finnish intervention prevent obstetric anal sphincter injuries? A systematic review of the literature. *BMJ Open.* 2015;5(9):e008346.
- Kalis V, Jansova M, Rusavy Z. Perineal Mapping. In: Ismail K. (eds) *Perineal Trauma at Childbirth*. Cham: Springer; 2016. [https://doi.org/10.1007/978-3-319-14860-1\\_3](https://doi.org/10.1007/978-3-319-14860-1_3).
- Cechova H, Kalis V, Havelkova L, Rusavy Z, Fiala P, Rybarova M, et al. Finite element modeling of maximum stress in pelvic floor structures during the head expulsion (FINESSE) study. *Int Urogynecol J.* 2021;32(7):1997–2003.
- Kalis V, Karbanova J, Horak M, Lobovsky L, Kralickova M, Rokyta Z. The incision angle of mediolateral episiotomy before delivery and after repair. *Int J Gynecol Obstet.* 2008;103(1):5–8.
- Kalis V, Landsmanova J, Bednarova B, Karbanova J, Laine K, Rokyta Z. Evaluation of the incision angle of mediolateral episiotomy at 60 degrees. *Int J Gynecol Obstet.* 2011;112(3):220–4.
- Kalis V, Karbanova J, Bukacova Z, Bednarova B, Rokyta Z, Kralickova M. Anal dilation during labor. *Int J Gynecol Obstet.* 2010;109(2):136–9.
- Rizk DEE, Thomas L. Relationship between the length of the perineum and position of the anus and vaginal delivery in primigravidae. *Int Urogynecol J.* 2000;11(2):79–83.
- Meriwether K V, Rogers RG, Dunivan GC, Alldredge JK, Qualls C, Migliaccio L, et al. Perineal body stretch during labor does not predict perineal laceration, postpartum incontinence, or postpartum sexual function: a cohort study. *Int Urogynecol J.* 2016;27(8):1193–200.
- Pihl S, Uustal E, Blomberg M. Anovaginal distance and obstetric anal sphincter injury: a prospective observational study. *Int Urogynecol J.* 2019;30(6):939–944.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.