

microRNA-124 Inhibits Migration and Invasion by Down-Regulating ROCK1 in Glioma

Liwen An¹*, Yongjun Liu²*, Anhua Wu³, Yifu Guan¹*

1 Key Laboratory of Medical Cell Biology, Ministry of Education, Department of Biochemistry and Molecular Biology, China Medical University, Shenyang, China, **2** Department of Neurosurgery, Fourth Affiliated Hospital, China Medical University, Shenyang, China, **3** Department of Neurosurgery, First Affiliated Hospital, China Medical University, Shenyang, China

Abstract

Background: The extraordinary invasive capability is a major cause of treatment failure and tumor recurrence in glioma, however, the molecular and cellular mechanisms governing glioma invasion remain poorly understood. Evidence in other cell systems has implicated the regulatory role of microRNA in cell motility and invasion, which promotes us to investigate the biological functions of miR-124 in glioma in this regard.

Results: We have found that miR-124 is dramatically downregulated in clinical specimen of glioma and is negatively correlated with the tumor pathological grading in the current study. The cells transfected by miR-124 expression vector have demonstrated retarded cell mobility. Using a bioinformatics analysis approach, rho-associated coiled-coil containing protein kinase 1 (ROCK1), a well-known cell mobility-related gene, has been identified as the target of miR-124. A dual-luciferase reporter assay was used to confirm that miR-124 targeted directly the 3'UTR of ROCK1 gene and repressed the ROCK1 expression in U87MG human glioma cell line. Furthermore, experiments have shown that the decreased cell mobility was due to the actin cytoskeleton rearrangements and the reduced cell surface ruffle in U87MG glioma cells. These results are similar to the cellular responses of U87MG glioma cells to the treatment of Y-27632, an inhibitor of ROCK protein. Moreover, a constitutively active ROCK1 in miR-124 over-expressed glioma cells reversed the effects of miR-124. Our results revealed a novel mechanism that miR-124 inhibits glioma cells migration and invasion via ROCK1 downregulation.

Conclusions: These results suggest that miR-124 may function as anti-migration and anti-invasion influence in glioma and provides a potential approach for developing miR-124-based therapeutic strategies for malignant glioma therapy.

Citation: An L, Liu Y, Wu A, Guan Y (2013) microRNA-124 Inhibits Migration and Invasion by Down-Regulating ROCK1 in Glioma. PLoS ONE 8(7): e69478. doi:10.1371/journal.pone.0069478

Editor: Svetlana Kotliarova, NIH/NCI, United States of America

Received: February 1, 2013; **Accepted:** June 10, 2013; **Published:** July 23, 2013

Copyright: © 2013 An et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by National Natural Science Foundation of China (no.31070705 and 31000390). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: yfgcmu@yahoo.com

† These authors contributed equally to this work.

Introduction

Brain tumors account for ~90% of all primary central nervous system tumors. In the United States, it is estimated 22,910 new cases of brain and related 13,700 deaths in 2012 [1]. Gliomas are the most common type of malignant primary brain tumor, accounting for 80% of malignant case [2,3]. Due to its high invasive neoplasm infiltrating diffusely into regions of normal brain, glioma is extremely difficult to be cured by total surgical resection or radiotherapy, leading to a high recurrences and poor prognosis. Despite of multi-modality treatment, the median survival of patients suffering from malignant glioma such as glioblastoma multiforme (GBM) is only 12 to 15 months [4]. Hence, it is urgently needed to understand the mechanisms of glioma cell's migration and invasion and develop more effective curative therapies.

MicroRNAs (miRNAs) are endogenous non-coding RNAs of approximately 21–23 nucleotides long. Being expressed in a tissue-specific manner during development of organisms, they regulate the gene expression by interacting specifically with 3'-untranslated

regions (3'UTR) of mRNA, reducing the stability of mRNAs and leading to reduced expression of protein [5]. Since miRNA may have many targets, they play key roles to regulate many biological processes such as embryonic development [6], differentiation [7], proliferation [8], cell death [9] and autophagy [10]. Emerging evidence has strongly suggested that aberrant miRNA expression is a common feature of many human cancers, functioning as either tumor suppressors or oncogenes [11–16].

Previous researches have shown that miRNAs have a very close relationship with glioma development [17–24]. microRNA-124 (miR-124) is abundantly expressed in normal brain tissue [25], necessary for embryonic neuronal differentiation which has been widely investigated in physiological neural development [26] and is highly conserved across species. It regulates some proliferation-related genes such as cyclin-dependent kinase 6 [27,28], aryl hydrocarbon receptor (AHR) [29], sphingosine kinase 1 (SPHK1) [30], androgen receptor (AR) [31], and solute carrier family 16, member 1 (SLC16A1) [32]. miR-124 has been documented as a tumor suppressor since low expression of miR-124 was observed in

several types of human cancers [27–35]. However, biological impacts of miR-124 on glioma cell migration and invasion have seldom been published.

In the current study, we have observed that miR-124 was downregulated in malignant glioma and its expression was correlated negatively with the pathological grading of glioma. Furthermore, we have identified that miR-124 regulated the ROCK1 gene, and ROCK1 protein expression caused actin cytoskeleton rearrangements, reduced cell surface ruffle, and suppressed glioma cell invasion. A constitutively active ROCK1 in miR-124 over-expressed glioma cells reversed the effects of miR-124. Our findings uncovered an important role of miR-124 in glioma morphology, motility and invasion via ROCK1 for the first time. These experimental results facilitate our understanding about the mechanism of the invasive ability of human glioma cells and pinpointed therapeutic options for glioma treatment.

Results

miR-124 is Significantly Downregulated in Human Glioma Tissues

To determine the expression level of miR-124 in clinical tissue specimens, we collected sixteen primary glioma tissue samples and extracted their total RNA. These sixteen primary glioma samples were classified into four grades according to WHO standard [2]: three WHO grade I, five WHO grade II, five WHO grade III and three WHO grade IV. Quantitative real-time PCR (qRT-PCR) was performed to determine the expression of miR-124 with respect to the internal standard RNU6-2.

Considering the fact that the our clinical samples were obtained from elderly patients and the research reports that anaplastic astrocytoma (WHO grade III) shares the similar biomolecular expression pattern and poor outcome of glioblastomas in the elderly patients [36,37], we divided clinical samples into two groups: low grade gliomas (grade I–II, $n = 8$) and high grade gliomas (grade III–IV, $n = 8$).

The qRT-PCR results (Figure 1) showed that the expression level of miR-124 was significantly lower in high grade gliomas than that in low grade gliomas, demonstrating a negatively correlation of the endogenous miR-124 expression with the WHO grade (Spearman $r = -0.5423$, $p < 0.01$). These data exhibits a good consistence with previous studies [38,39] and indicates a putative tumor suppressor role for miR-124 in glioma.

The Overexpressed miR-124 Inhibits the Cell Motility

To decipher the biological function of miR-124 in glioma cells, we constructed a miR-124 expression vector (named as pcDNA3.1-miR-124) and transiently transfected into HEK293ET and human glioma cells U87MG and U251, respectively, to create a gain-of-function behavior in cell lines. Forty-eight hours later, the expression levels of miR-124 in all cell lines were examined using qRT-PCR. As shown in Figure 2A, the expression of miR-124 was enhanced significantly (by a factor of hundreds) in HEK293ET cells transfected with pcDNA3.1-miR-124 vector when compared with cells transfected with pcDNA3.1. The transfected U87MG cells (Figure 2A) and U251 cells (data not shown) demonstrated similar trend. All these results proved a success and effectiveness of miR-124 expression vector construction and transfection.

The migration ability of the transfected cells was further tested using the wound-healing assay. In this study, two controls were used: lipofectamine2000 blank (referred to as L2K) and lipofectamine2000 blank fused with pcDNA3.1 (referred to as




Figure 1. qPCR assays of miR-124 expression levels in glioma tissue samples. The expression level of miR-124 was downregulated significantly in high grade human glioma tissues (five grade III and three grade IV) than that in low grade human glioma tissues (three grade I and five grade II) determined using qRT-PCR. doi:10.1371/journal.pone.0069478.g001

L2K+pcDNA3.1). L2K+pcDNA3.1-miR-124 was used as the miR-124 expression vector.

The wound-healing process was monitored at 0, 6, 24 and 48 hours after scrape for U87MG cells. Images in Figure 2B and 2C clearly showed that the cells transfected with vector pcDNA3.1-miR-124 have a retarded mobility in comparison with other two controls, and similar results were also observed in U251 cells (Figure S1). These results indicated the anti-migration effect of miR-124 in glioma cells.

miR-124 Interacts Specifically with the 3'UTR Region of ROCK1

After observing the altered cell mobility by miR-124, we started to search the potential genes involved in regulating the cell motility. Using TargetScan (Release 4.2) and miRanda (August 2010 Release) online searching programs, we have identified cell motility-related gene (ROCK1) as the potential target of miR-124. A 100% matched sequence was found at the nts 930–937 region of ROCK1 mRNA 3'UTR (NM_005406) (Figure 3A). The free energy (ΔG) was calculated about -26.4 kcal/mol for hybrid of ROCK1 3'UTR region and miR-124 by BibiServ analysis (Figure 3B). In addition, homology analyses have shown that the nucleotide sequences of 3'UTR of the ROCK1 gene targeted by miR-124 seed sequence were highly conserved among different species (Figure 3C).

To confirm the hypothesis that miR-124 targets the 3'UTR region of ROCK1, we cloned the vector that the entire 3'UTR region of ROCK1 was connected at the downstream of a modified luciferase reporter gene (named as pGL3M-ROCK1-3'UTR), and co-transfected the HEK293ET cells with this vector along with either the miR-124 expression vector or its negative control. In the case of pGL3M, the luciferase activities of cells transfected with either pcDNA3.1 or pcDNA3.1-miR-124 showed a slightly difference (by $< 16\%$). We found that co-transfection of miR-124 expression vector along with the full-length 3'UTR of ROCK1 caused a significant decrease by over 50% in luciferase units compared to controls (Figure 3D). These results demonstrated unambiguously that miR-124 targeted specifically the 3'UTR region of ROCK1.

Moreover, we transfected U87MG cells with either control or pcDNA3.1-miR-124, and determined the endogenous expression




Figure 2. Evaluation of biological functions of miR-124 in transfected HEK293ET and U87MG cells. (A) The expression level of miR-124 was measured using qRT-PCR after miR-124 transfected HEK293ET and U87MG cells for 48 h. Data presented are mean values of three independent experimental results and compared with the level of miR-124 obtained in mock control (Lipofectamine2000 blank) that is normalized to 1. (B) Wound-healing assay of U87MG glioma cells transfected with either control or the miR-124 expression vector, respectively. doi:10.1371/journal.pone.0069478.g002

of ROCK1 at both protein and mRNA levels, respectively. qRT-PCR results revealed that although the mRNA level of ROCK1 was attenuated, it was not significantly affected by miR-124 in comparison with the control after statistics calculation (Figure 3E). However, the ROCK1 protein was remarkably reduced after transfection (Figure 3F). These data provided strong evidence that miR-124 suppressed ROCK1 gene expression by regulating 3'UTR at the post-transcription level.

miR-124 Overexpression Suppresses the Glioma Cell Invasion by Affecting the Actin Cytoskeleton Rearrangements and Reducing Cell Surface Ruffles

Based on the inhibitory effects of miR-124 on ROCK1 protein expression and glioma cell locomotion, we reasoned that miR-124 may have an impact on cell invasive capacity.

Matrigel invasion assays have been employed to assess the correlation of the miR-124 and the cell invasion. Forty-eight hours after transfection, a cell suspension was added into the upper chamber of matrigel-coated inserts. The lower chamber was filled with 500 μ l of media containing 20% FBS and cultured for another 24 hours. Invasive cells were stained and the average number of cells was counted. Images of the stained cells demonstrated that ectopic overexpression of miR-124 significantly reduced the invasiveness of U87MG cells (lower left image, Figure 4A); the number of invasive cells able to digest the extracellular matrix and migrate through pores in the membrane was reduced by over 50%. Interestingly, this result was consistent with the result of Y-27632-treated cells (lower right image, Figure 4A), which has been known as a ROCK inhibitor.

It has been demonstrated that the Rho/ROCK pathway participates in regulating cytoskeletal signaling events and is

crucial for cell motility. Moreover, cytoskeletal reorganization exemplified by the formation of stress fibre bundling arrays is essential for the contractile motion of cancer cells [40,41]. We thus examined the status of stress fibre formation and polymerised actin in U87MG cells. Figure 4B showed the cytoskeleton changes inside cells. The actin filaments were stained with rhodamine phalloidin and examined by fluorescence microscopic. In the mock control or negative control U87MG cells (upper images, Figure 4B), the actin filaments were distribution presented as many dot-like and cluster-like actin assemblies spanning the length of the cells. However, the actin cytoskeletons of cells were completely different in the pcDNA3.1-miR-124 treated U87MG cells that they showed a decrease in the length and number of actin fibers in cells (lower left image, Figure 4B); do not have the same extent like the control groups. Similar phenotypic changes could be visualized in U87MG cells treated with 10 μ M Y-27632, showed a loss of actin stress fibers, which has been reported with 50 μ M Y-27632 [42].

Furthermore, filopodia and lamellipodia on cell membrane surfaces have been identified as dynamic cellular features, requiring actin polymerization/depolymerization for cancer cell invasion. We investigated the impact of miR-124-mediated cell morphology alternation in U87MG cells. Under scanning electron microscope, the negative control U87MG cells presented the elongated and fibroblast-like morphology with many filopodia and lamellipodia on the cell surfaces, probably for cell protrusion (upper images, Figure 4C). In contrast, higher miR-124 expression led to remarkable morphological changes in that cells became a round-like and shrunken form, and there was an obvious reduction of long and thin protrusions on the surface of cells transfected with pcDNA3.1-miR-124 (lower left image, Figure 4C).




Figure 4. Identification of biological function of miR-124 in U87MG cells. (A) In vitro invasion assay. Invasive cells were stained and the average number of cells was counted at random six fields of vision. The data was an average value of three independent experiments. (Magnification: 100 \times ; scale bars: 100 μ m) (B) Stress fiber staining in U87MG glioma cells (Scale bars: 20 μ m). (C) Scanning electron microscopy of U87MG cells treated with miR-124 expression vector or control and Y-27632 (Scale bars: 10 μ m). doi:10.1371/journal.pone.0069478.g004

our in vitro observations in miR-124 overexpressed cells, indicating a potential role of ROCK1 in glioma cell invasion.

ROCK1 Partially Rescues the miR-124-induced Invasion Inhibition

To further understand the role of ROCK1 in miR-124-mediated anti-invasion, we treated U87MG cells with pCAG-myc-p160^{ROCK} Δ 3 [43] (a constitutively active ROCK1, p160^{ROCK} Δ 3) or control pCAG-myc and co-transfected with pcDNA3.1-miR-124 followed by functional assays. Expression of these constructs was confirmed by western blot analysis for the myc-tagged antigen. As expected, ectopic expression of ROCK1 (without 3'UTR) significantly abrogated the miR-124-mediated anti-invasion by affecting the actin cytoskeleton rearrangements and reducing cell surface ruffles. The F-actin filaments stained with rhodamine phalloidin showed that ectopically expressing ROCK1 significantly increased the dot-like and cluster-like actin assemblies spanning the length of the cells (Figure 5A). Furthermore, the impact of miR-124-mediated cell morphology alternation in U87MG cells was restored compared to control pCAG-myc-treated cells, as observed after miR-124 suppression under scanning electron microscope (Figure 5B). Moreover, treatment with p160^{ROCK} Δ 3 also led to increased invasion of U87MG cells

(Figure 5C). These results suggest that the anti-invasion effects of miR-124 are in part facilitated by ROCK1 downregulation.

Discussion

Accumulating evidence has suggested that miRNAs are able to regulate multiple genes involving almost all aspects of cancer biology [13]. Investigation of these miRNAs would expand our view to better understand carcinogenesis by analyzing mRNA target associated and miRNA-mediated pathways. Several independent research groups have reported the correlation of alterations of miRNA expression with the gliomagenesis, patient diagnosing, potential prognosis and therapeutic tool [17,44–45]. In this study, we focused on miR-124, a putative tumor suppressor in several human cancers. A variety of targets have been found to be regulated by miR-124, including proliferation-related genes [27–30], invasion/metastasis-related genes [33–35] and so on. It has been known that invasion, one of the most important hallmarks of malignant tumors [46,47], is the incurable factor for human glioma. Study by Fowler et al [33] has reported that transfection of commercialized miR-124 precursor in GBM cell line A172 resulted in diminished cell migration and invasion as well as downregulated three targeted genes: Ras GTPase activating protein 1 (IQGAP1), cytoskeletal proteins laminin c1




Figure 5. Reintroduction of ROCK1 rescues the miR-124-induced invasion inhibition. (A) Stress fiber staining in U87MG glioma cells treated with miR-124 expression vector along with p160^{ROCK} Δ 3 or control pCAG-myc (Scale bars: 20 μ m). (B) Scanning electron microscopy of U87MG cells. (C) In vitro invasion assay. Invasive cells were stained and the average number of cells was counted at random six fields of vision. The data was an average value of three independent experiments. (Magnification: 100 \times ; scale bars: 100 μ m). doi:10.1371/journal.pone.0069478.g005

(LAMC1) and integrin β 1 (ITGB1). However, the median survival was not significantly different between the high and low miR-124 expression of GBM patients, which might be due to the extraordinary high malignancy of GBM. Xia et al [39] have reported the down-regulation of miR-124 in a larger patient clinical specimen, which included human glioma tissue samples ($n = 27$) and non-glioma patients samples ($n = 20$, two non-tumor brain tissues). They identified SNAI2, a member of the Snail family of zinc finger transcription factors - because it has been implicated in epithelial-mesenchymal transition (EMT) [48,49] and tumor metastasis, as a direct functional target of miR-124. The enhanced miR-124 expression significantly inhibited glioma cell invasion using matrigel invasion assay and tumor xenografts in nude mice.

In this study, we identified that low expression of miR-124 was closely associated with a more aggressive tumor phenotype. Furthermore, the level of endogenous miR-124 is negatively correlated with the tumor pathological grading, indicating an association with the progression of glioma. Therefore, it is possible to be developed into a biomarker for diagnosis.

We performed functional analysis to examine the function of miR-124. Reintroduction of miR-124 through an expression vector dramatically repressed glioma cell migration and invasion in vitro. These findings suggest that miR-124 plays a critical role in the invasive potential of glioma. We also observed that the number of miR-124 transfected cells was reduced slightly in the wound-healing assay (Figure 2B). It could be due to either the serum depletion during the longer experimental duration or some unclassified functions of miR-124 in regulating other proliferation-related genes. In addition, by transfecting EGFP plasmid in the

two glioma cell lines U87MG and U251, we identified that U87MG cells have a higher transfection efficiency than U251 (data not shown). Accordingly, we chose U87MG cells as the objective of the experiment.

As a next step, we used the on line websites to identify target genes of miR-124, interestingly, all of which are closely related to tumor migration and invasion. In general, cell movement is affected through a combination of protrusive and contractile events. Non-muscle cells contain stress fibres – bundles of approximately 10–30 actin filaments [50]. A large number of signaling proteins, scaffolding proteins and actin-binding proteins (ABPs) participate this very complicated biological process in a temporal-specific and spatial-specific manner, as well as in the cooperative fashion [51]. ROCKs have been shown to be a central player in the formation of stress fibers via phosphorylation of myosin light chain [40,52–53]. Two ROCK isoforms have been identified: ROCK1 and ROCK2. Sequence analyses have shown a 65% sequence homology between these two ROCK isoforms, and in the kinase domains, their sequence similarity could be as high as 92% [54]. The ROCK2 transcript is highly expressed in muscle and brain tissues, whereas the ROCK1 is localized in non-nerves tissues [53]. Deregulation of Rho/ROCK signaling pathway has been reported across diverse tumors types [55,56]. Several preclinical and clinical studies have utilized inhibitors of Rho/ROCK signaling pathway for anticancer therapeutics in prostate, lung, melanoma, and many other tumor types with remarkable success [57–59]. Using bioinformatics and experimental methods, we assessed ROCK1 as potential functional targets of miR-124. We performed a 3'UTR luciferase assay and observed that luciferase activity was decreased after co-transfection of the

miR-124 expression vector and a 3'UTR vector containing the ROCK1/miR-124 target sequence. ROCK1 protein expression was also significantly downregulated in U87MG cells that were transfected with the miR-124 expression vector. All these data clearly indicate that ROCK1 is a direct target of miR-124.

It has been reported that as a cell moves in a designated direction, cooperation of continuous actin polymerisation and depolymerisation occurs simultaneously, which leads the cells to protrude at its anterior front. Concurrently, the cell undergoes consecutive actomyosin contractions and separates from the posterior end, and directional cell movement occurs. In addition, formation of the stress fibres, filopodia and lamellipodia also requires actin polymerisation. In this study, our results show that exogenous expression of miR-124 in U87MG cells substantially suppressed the formation of the stress fibres and cell protrusions. Previous studies have reported that downregulation of ROCK1 reduced markedly cell spanning F-actin fibers but enhanced the production of cortical fibers. On the other hand, downregulation of ROCK2 promoted the production of shorter intracellular fibers and destabilized the cortical F-actin fibers, leading to the formation of invaginations in human-derived renal proximal tubular cell line HKC-8 peripheral cells [60]. The changes in the F-actin localization in glioma cells treated with miR-124 in this study showed a good agreement (Figure 4B).

In addition to the changes of stress fibres, the invasion behavior of glioma was inhibited by up-regulation of miR-124. This phenomenon is consistent with the effect of ROCK inhibitor Y-27632 reported previously using GBM cell lines T98G and 8401 [61,62]. These observations suggested that miR-124 retard the glioma cell migration and invasion by inhibiting the formation of the stress fibres via ROCK1 regulation. Notably, a constitutively active ROCK1 in miR-124 over-expressed glioma cells reversed the effects of miR-124, suggesting the biological role of preventive invasion of miR-124 due mainly to the ROCK1 down-regulation. However, contradictory evidence showed that the invasive and migratory properties of astrocytoma were enhanced by inhibition of ROCK [42,63]. The inconsistency among these studies might be due to either the differences of cell lines, or the inhibitor dosages or the model systems used in these different studies. More systematic studies, thus, have been proposed to be performed by following the standard experimental approaches to identify the true functions of ROCK proteins in migratory and invasive phenotypes of glioma cells.

To extend these findings to glioma tissues, we also measured ROCK1 mRNA expression by qRT-PCR in the same clinical samples for miR-124 expression. Analyses confirmed that this difference was statistically significant between the low grade gliomas and the high grade gliomas (Figure S2). However, the correlation between the endogenous ROCK1 mRNA expression of clinical samples and the pathological grading was not established since we were unable to extract enough protein for the measurement. A study by Oellers et al [64] reported that ROCK was implicated in the migration of glioma cells on myelinated axons, a process that is still poorly understood. They also showed another important finding that ROCK1 was much less expressed while ROCK2 was highly expressed in the normal brain tissue. In contrast, ROCK1 was highly expressed and ROCK2 expression was almost undetectable in three WHO grade III and three WHO grade IV gliomas using immunohistochemistry and western blotting. These data indicated that at least the ROCK1 isoform is expressed in malignant human gliomas with a high potential for migration and invasion although this conclusion needs to be confirmed via a large number of clinical samples. Various compounds such as Y-27632 have been found to inhibit

ROCK and inhibition of ROCK by fasudil leads to beneficial effects in patients with cardiovascular disease clinically [65]. The question is that the inhibitors are non-isoform-selective. Thus, designing an inhibitor that specifically blocks ROCK1 and using a siRNA approach would further help to unravel its role in primary gliomas.

Conclusions

In this study, we have demonstrated that the endogenous expression of miR-124 was negatively correlated with the tumor pathological grading of clinical glioma samples, and bioinformatic analyses have identified the target gene of miR-124. Further in vitro experiments have showed that miR-124 reduced migration and invasion, affected actin cytoskeleton rearrangements and reduced cell surface protrusion by suppressing the expression of ROCK1 protein. Moreover, a constitutively active ROCK1 in miR-124 over-expressed glioma cells rescued the effects of miR-124. Based on our observations and results reported by other groups, we have proposed a model to elucidate the potential roles of miR-124 in glioma (Figure 6). The experimental data and conclusions in the present study furnish valuable information regarding the biological functions of miR-124 and the possible mechanism of the migration and invasion of glioma tumor. Thus, further studies will be focused on the molecular network involved in the ROCK1 regulation on a large quantity of clinical sample basis.

Materials and Methods

Cells Lines and Reagents

HEK293ET and glioma cell lines U87MG and U251 (human) were purchased from the American Type Culture Collection (Manassas, VA, USA). All cell lines were cultured at 37°C with 5% CO₂ in DMEM supplemented with 10% fetal bovine serum (FBS).

Original stock solutions of Y-27632 (Calbiochem, Nottingham, Germany) at a concentration of 5 mM was stored at -20°C and freshly dissolved in culture medium before use.

Clinical Glioma Samples

Human glioma samples from surgery were collected from the neurosurgery department of the First Affiliated Hospital of China Medical University from adult patients, freshly resected during surgery. The samples were snap-frozen in liquid nitrogen and stored at -80°C for subsequent total RNA extraction. All human materials used were approved by the Ethical Committee in China Medical University. All participants have provided their written consent to participate in this study.

RNA Isolation, Reverse Transcription and Quantitative Real-time PCR of mRNA and miRNA

Total RNA, including miRNAs, was isolated by TRIzol reagent (Invitrogen, Carlsbad, USA) following the manufacturer's protocol. qRT-PCR were performed using ImProm-IITM reverse transcriptase (Promega, Madison, USA) and SYBR[®] Premix Ex TaqTM II (TaKaRa, Dalian, China), and detected with the ABI7500 Real-time PCR system instrument (Applied Biosystems, Foster City, CA, USA). For ROCK1 quantification with the internal control GAPDH, the primers for ROCK1 were AACCATGTGACTGAGTGCCC and TCAGTGTGTTGTGCCAAAGC. Primers for GAPDH were AATGGG-CAGCCGTTAGGAAA and TGAAGGGGTCATTGATGG-CA.




Figure 6. Proposed model of miR-124 function in glioma development and progression. The target of miR-124 in bold fonts was confirmed in this study.

doi:10.1371/journal.pone.0069478.g006

For quantification of miR-124, 10ng of RNA was used as a template and cDNA was synthesized with miRNA-specific primers, performed as described [66,67], the miR-124 level was analyzed with internal control RNU6-2. Specific stem-loop reverse transcription primer and forward primer of PCR for miR-124 were CTCAACTGGTGTCTGGAGTCGGCTACTAAGT-TGGCGAGATTCA and ACACTCCAAGGGCTGTAACG-GGTGCCGAA. Primers for RNU6-2 were CTCAAC-TGGTGTCTGGAGTCGGCAATTGACAAGTTGAAATA-TG and ACACTCCAAGGGCTGTAACGGGTGCCGAA.

The expression level of ROCK1 and miR-124 was calculated by using $2^{-\Delta\Delta Ct}$ analysis method [68], normalized to the control group.

Vector Construction

The human pre-miR-124 sequence was amplified and cloned into pcDNA3.1-hisA constructs (Invitrogen) to generate pcDNA3.1-miR-124 expression vector. The set of primers were GAGAATTCTTGCATCTCTAAGCCCTGT and TCTCTA-GAGCGCGCTTTTTATTTCTTT.

The full-length 3'UTR of ROCK1 was amplified using cDNA from U87MG cells with following primers: TCTCTA-GATTGTTTCGTGCTTCCC and TCGAATTCATCAGTG-CGGCTTTC. The 3'UTR was double-digested with XbaI/EcoRI and cloned downstream of firefly luciferase coding region sites of a modified pGL3-control plasmid.

Transfection

The HEK293ET cells, U87MG cells and U251 cells were transfected with plasmids using Lipofectamine 2000 reagent (Invitrogen) at 24h after plating. Transfection complexes were prepared according to the manufacturer's instructions. The transfection medium was replaced at 4h post-transfection.

Bioinformatics Analysis

The target gene information of miR-124 was analyzed using miRanda (<http://microrna.sanger.ac.uk/sequences/>) and the Human microRNA targets can (<http://www.microrna.org/mammalian/index.html>). The minimum free energy predicted for hybridization was determined by BibiServ analysis (<http://bibiserv.techfak.uni-bielefeld.de/genefisher2/>).

Luciferase Reporter Assay

HEK293ET cells were co-transfected using lipofectamine 2000 reagent with 100ng of firefly luciferase construct and 300ng of control-pcDNA3.1 or pcDNA3.1-miR-124 expression vector. Ten ng of pRL-CMV (Promega) was co-transfected as a normalization control. Reporter assays were performed 48h post-transfection using the Dual-luciferase assay system (Promega), normalized for

transfection efficiency by co-transfected Renilla luciferase. Cells were transfected in duplicated wells and such experiments were repeated three times on different days.

Western Blot

Total protein samples were collected from U87MG cells 72h post-transfection with cell lysis buffer (Beyotime, Jiangsu, China). The protein concentration was determined by bicinchoninic acid protein assay kit (Beyotime). Heat-denatured protein samples (30 μ g per lane) were resolved by 6%~12% SDS-polyacrylamide gel electrophoresis and transferred to Hybond-nitrocellulose membranes (Amersham, Buckinghamshire, UK). The membrane was incubated for 2h in TBS containing 0.1% Tween 20 and 5% BSA to block nonspecific binding, followed by incubation for 12 h at 4°C with primary mouse monoclonal anti-ROCK1 antibody (Abcam, Cambridge, UK) (at 1:1000 dilution). As a loading control, the GAPDH expression level was measured using mouse monoclonal anti-GAPDH antibody (Zsbio, Beijing, China), at 1:1000 dilution. The membrane was then incubated with goat anti-mouse secondary antibody (Zsbio) using at 1:5000 for 2h, then detected with SuperSignal West Pico Chemiluminescent Substrate (Thermo Scientific, Waltham, MA, USA) and visualized in an MF-ChemiBis 3.2 Bioimaging system (DNR, Jerusalem, Israel). Expression levels were quantified using ImageJ 1.44 software (National Institute of Health, Bethesda, MA, USA) and normalized to loading controls.

Wound-healing Assay

Cells were plated at 80% confluence in DMEM supplemented with 10% FBS. At 24h after seeding, the monolayers were wounded by scoring with a sterile plastic 200 μ l micropipette tip, washed, and then incubated in DMEM in the absence of serum. At vary hours, cells were photographed using a low-magnification fluorescence microscope (Olympus IX71, Miami, USA). The widths of the wound lines were measured by AlphaEase FC (Version 4.0, Alpha Innotech Corp.). The data are presented as percentages of the control.

Invasion Assay

Cell invasion was assessed by the invasion of the cells through Matrigel-coated Transwell inserts. Briefly, Transwell inserts with 8 μ m pore size were coated with 0.1ml of final concentration of 200 μ g/ml Matrigel in cold serum-free medium. U87MG cells transfected were harvested 48 hours after treatment and trypsinized, then the serum-deprived cell suspension (200 μ l; 0.5×10^5 cells/ml) was added in triplicate to wells. After 24h incubation, cells that invaded the Matrigel and passed through the filter were stained with crystal violet and photographed using fluorescence microscope (Olympus IX71). The number of invaded cells was quantified by counting them in at least six random fields with total

magnification of 100× per filter. In certain experiments, Y-27632 (10 μmol/L) was applied to the upper chamber.

Fluorescence Microscopy

U87MG cells treated in 6-well plates were fixed in 4% formaldehyde for 10min and then washed three times for 10min each in PBS. The cells were then permeabilized with 0.1% Triton X-100 for 10min. After blocking with 5% BSA in PBS, the cells were incubated with TRITC-labeled phalloidin (Sigma-Aldrich) to stain the actin filaments. The nuclei were stained with DAPI. The coverslips were mounted and analyzed using fluorescence microscope (Olympus BX51).

Scanning Electron Microscopy

U87MG cells were grown to sub confluence on 20mm square cover slip in a six-well plate. The cells were fixed with 2.5% glutaraldehyde in PBS (pH 7.4) for 2h at room temperature. Then, the cells were performed as described [69]. The specimens were cold sputter coated with gold and observed in a JSM-T300 scanning electron microscope (JEOL, Japan).

Statistical Analysis

All tests were done using PRISM version 5.0 (GraphPad Software Inc., San Diego, CA, U.S.A.) statistical software. All experiments were carried out for three times independently. Data were mean±standard deviation except special indication. Spearman's rank correlation test was used for association analysis between miR-124 and ROCK1 level data and pathological grading. Statistical significance between two groups was analyzed by Student's t test. One-way ANOVA was used to compare

multiple groups, with P<0.05 considered to be significant (*P<0.05; **P<0.01).

Supporting Information

Figure S1 Wound-healing assay of U251 glioma cells transfected with either control or the miR-124 expression vector, respectively.

(TIF)

Figure S2 qPCR assays of ROCK1 expression levels in glioma tissue samples. The expression level of ROCK1 was downregulated significantly in low grade human glioma tissues (three grade I and five grade II) than that in high grade human glioma tissues (five grade III and three grade IV) determined using qRT-PCR.

(TIF)

Acknowledgments

We thank Dr Xiaofei Zheng (Beijing Institute of Radiation Medicine, Beijing, PR China) for providing the pGL3M plasmid and Prof. Shuh Narumiya (Department of Pharmacology, Kyoto University Faculty of Medicine, Kyoto, Japan) and Yuhua Chen (Department of Developmental Biology, Key Laboratory of Cell Biology, Ministry of Public Health of China, China Medical University, Shenyang, PR China) for providing the pCAG-myc-p160^{ROCK}Δ3 and pCAG-myc plasmids.

Author Contributions

Conceived and designed the experiments: LA YL YG. Performed the experiments: LA YL. Analyzed the data: LA YL AW YG. Contributed reagents/materials/analysis tools: AW YG. Wrote the paper: LA YL YG.

References

- American Cancer Society.: Cancer Facts and Figures (2012). Atlanta GACS, 2012. Available: <http://www.cancer.org/research/cancerfactsfigures/cancerfactsfigures/cancer-facts-figures-2012>. Accessed July 31, 2012.
- Louis DN et al (2007) WHO classification of tumours of the central nervous system. Lyon: World Health Organization Press. 312 p.
- CBTRUS (2012). CBTRUS Statistical Report: Primary Brain and Central Nervous System Tumors Diagnosed in the United States in 2004–2008 (March 23, 2012 Revision).
- Louis DN, Ohgaki H, Wiestler OD, Cavenee WK, Burger PC, et al. (2007) The 2007 WHO classification of tumours of the central nervous system. *Acta Neuropathol* 114: 97–109.
- Eulalio A, Huntzinger E, Izaurralde E (2008) Getting to the Root of miRNA-Mediated Gene Silencing. *Cell* 132: 9–14.
- Li Q, Bian S, Hong J, Kawase Koga Y, Zhu E, et al. (2011) Timing specific requirement of microRNA function is essential for embryonic and postnatal hippocampal development. *PLoS One* 6: e26000.
- Chen CZ, Li L, Lodish HF, Bartel DP (2004) MicroRNAs modulate hematopoietic lineage differentiation. *Science* 303: 83–86.
- Brennecke J, Hipfner DR, Stark A, Russell RB, Cohen SM (2003) bantam encodes a developmentally regulated microRNA that controls cell proliferation and regulates the proapoptotic gene hid in *Drosophila*. *Cell* 113: 25–36.
- Chen CH, Guo M, Hay BA (2006) Identifying microRNA regulators of cell death in *Drosophila*. *Methods Mol Biol* 342: 229–240.
- Fu LL, Wen X, Bao JK, Liu B (2012) MicroRNA-modulated autophagic signaling networks in cancer. *Int J Biochem Cell Biol* 44: 733–736.
- Calin GA, Sevignani C, Dumitru CD, Hyslop T, Noch E, et al. (2004) Human microRNA genes are frequently located at fragile sites and genomic regions involved in cancers. *Proc Natl Acad Sci U S A* 101: 2999–3004.
- Cummins JM, He Y, Leary RJ, Pagliarini R, Diaz LA, et al. (2006) The colorectal microRNAome. *Proc Natl Acad Sci U S A* 103: 3687–3692.
- Hwang HW, Mendell JT (2006) MicroRNAs in cell proliferation, cell death, and tumorigenesis. *Br J Cancer* 94: 776–780.
- Zhang BG, Li JF, Yu BQ, Zhu ZG, Liu BY, et al. (2012) microRNA-21 promotes tumor proliferation and invasion in gastric cancer by targeting PTEN. *Oncol Rep* 27: 1019–1026.
- Manikandan J, Aarthi JJ, Kumar SD, Pushparaj PN (2008) Oncomirs: the potential role of non-coding microRNAs in understanding cancer. *Bioinformatics* 2: 330–334.
- Kong YW, Ferland McCollough D, Jackson TJ, Bushell M (2012) microRNAs in cancer management. *Lancet Oncol* 13: e249–258.
- Ciafre SA, Galardi S, Mangiola A, Ferracin M, Liu CG, et al. (2005) Extensive modulation of a set of microRNAs in primary glioblastoma. *Biochem Biophys Res Commun* 334: 1351–1358.
- Godlewski J, Newton HB, Chioocca EA, Lawler SE (2010) MicroRNAs and glioblastoma; the stem cell connection. *Cell Death Differ* 17: 221–228.
- Wurdinger T, Tannous BA (2009) Glioma angiogenesis: Towards novel RNA therapeutics. *Cell Adh Migr* 3: 230–235.
- Turner JD, Williamson R, Almeyty KK, Nakaji P, Porter R, et al. (2010) The many roles of microRNAs in brain tumor biology. *Neurosurg Focus* 28: E3.
- Pang JC, Kwok WK, Chen Z, Ng HK (2009) Oncogenic role of microRNAs in brain tumors. *Acta Neuropathol* 117: 599–611.
- Silber J, James CD, Hodgson JG (2009) microRNAs in gliomas: small regulators of a big problem. *Neuromolecular Med* 11: 208–222.
- Novakova J, Slaby O, Vyzula R, Michalek J (2009) MicroRNA involvement in glioblastoma pathogenesis. *Biochem Biophys Res Commun* 386: 1–5.
- Wong JW (2010) MicroRNA-induced silencing of glioma progression. *J Neurosci* 30: 3868–3869.
- Lagos-Quintana M, Rauhut R, Yalcin A, Meyer J, Lendeckel W, et al. (2002) Identification of tissue-specific microRNAs from mouse. *Curr Biol* 12: 735–739.
- Cao X, Pfaff SL, Gage FH (2007) A functional study of miR-124 in the developing neural tube. *Genes Dev* 21: 531–536.
- Pierson J, Hostager B, Fan R, Vibhakkar R (2008) Regulation of cyclin dependent kinase 6 by microRNA 124 in medulloblastoma. *J Neurooncol* 90: 1–7.
- Silber J, Lim DA, Petritsch C, Persson AI, Maunakea AK, et al. (2008) miR-124 and miR-137 inhibit proliferation of glioblastoma multiforme cells and induce differentiation of brain tumor stem cells. *BMC Med* 6: 14. Available: <http://www.biomedcentral.com/1741-7015/6/14>. Accessed 24 Jun 2008.
- Huang TC, Chang HY, Chen CY, Wu PY, Lee H, et al. (2011) Silencing of miR-124 induces neuroblastoma SK-N-SH cell differentiation, cell cycle arrest and apoptosis through promoting AHR. *FEBS Letters* 585: 3582–3586.
- Xia J, Wu Z, Yu C, He W, Zheng H, et al. (2012) MiR-124 Inhibits Cell Proliferation in Gastric Cancer through Downregulation of SPHK1. *The Journal of Pathology* 227: 470–478.
- Shi XB, Xue L, Ma AH, Tepper CG, Gandour Edwards R, et al. (2012) Tumor suppressive miR-124 targets androgen receptor and inhibits proliferation of prostate cancer cells. *Oncogene*. In press.
- Li KK, Pang JC, Ching AK, Wong CK, Kong X, et al. (2009) miR-124 is frequently down-regulated in medulloblastoma and is a negative regulator of SLC16A1. *Hum Pathol* 40: 1234–1243.

33. Fowler A, Thomson D, Giles K, Maleki S, Mreich E, et al. (2011) miR-124a is frequently down-regulated in glioblastoma and is involved in migration and invasion. *Eur J Cancer* 47: 953–963.
34. Hunt S, Jones AV, Hinsley EE, Whawell SA, Lambert DW (2011) MicroRNA-124 suppresses oral squamous cell carcinoma motility by targeting ITGB1. *FEBS Letters* 585: 187–192.
35. Zheng F, Liao YJ, Cai MY, Liu YH, Liu TH, et al. (2012) The putative tumour suppressor microRNA-124 modulates hepatocellular carcinoma cell aggressiveness by repressing ROCK2 and EZH2. *Gut* 61: 278–289.
36. Barnholtz Sloan JS, Williams VL, Maldonado JL, Shahani D, Stockwell HG, et al. (2008) Patterns of care and outcomes among elderly individuals with primary malignant astrocytoma. *J Neurosurg* 108: 642–648.
37. Hartmann C, Hentschel B, Wick W, Capper D, Felsberg J, et al. (2010) Patients with IDH1 wild type anaplastic astrocytomas exhibit worse prognosis than IDH1-mutated glioblastomas, and IDH1 mutation status accounts for the unfavorable prognostic effect of higher age: implications for classification of gliomas. *Acta Neuropathol* 120: 707–718.
38. Li D, Chen P, Li XY, Zhang LY, Xiong W, et al. (2011) Grade-Specific Expression Profiles of miRNAs/mRNAs and Docking Study in Human Grade I–III Astrocytomas. *OMICs: A Journal of Integrative Biology* 15: 673–682.
39. Xia H, Cheung WK, Ng SS, Jiang X, Jiang S, et al. (2012) Loss of brain-enriched miR-124 microRNA enhances stem-like traits and invasiveness of glioma cells. *The Journal of biological chemistry* 287: 9962–9971.
40. Katoh K (2001) Stress fiber organization regulated by MLCK and Rho-kinase in cultured human fibroblasts. *Am J Physiol Cell Physiol* 280: C1669–C1679.
41. Shum MSY, Pasquier E, Pouha ST, O'Neill GM, Chaponnier C, et al. (2011) γ -Actin regulates cell migration and modulates the ROCK signaling pathway. *The FASEB Journal* 25: 4423–4433.
42. Salhia B, Rutten F, Nakada M, Beaudry C, Berens M, et al. (2005) Inhibition of Rho-kinase affects astrocytoma morphology, motility, and invasion through activation of Rac1. *Cancer Res* 65: 8792–8800.
43. Ishizaki T, Naito M, Fujisawa K, Maekawa M, Watanabe N, et al. (1997) p16ROCK, a Rho-associated coiled-coil forming protein kinase, works downstream of Rho and induces focal adhesions. *FEBS Letters* 404: 118–124.
44. Gaur A, Jewell DA, Liang Y, Ridzon D, Moore JH, et al. (2007) Characterization of microRNA expression levels and their biological correlates in human cancer cell lines. *Cancer Res* 67: 2456–2468.
45. Malzkorn B, Wolter M, Liesenberg F, Grzendowski M, Stuhler K, et al. (2010) Identification and functional characterization of microRNAs involved in the malignant progression of gliomas. *Brain Pathol* 20: 539–550.
46. Hanahan D, Weinberg RA (2000) The hallmarks of cancer. *Cell* 100: 57–70.
47. Hanahan D, Weinberg RA (2011) Hallmarks of cancer: the next generation. *Cell* 144: 646–674.
48. Alves CC, Carneiro F, Hoefler H, Becker KF (2009) Role of the epithelial-mesenchymal transition regulator Slug in primary human cancers. *Frontiers in bioscience : a journal and virtual library* 14: 3035–3050.
49. Liang YJ, Wang QY, Zhou CX, Yin QQ, He M, et al. (2013) MiR-124 targets Slug to regulate epithelial-mesenchymal transition and metastasis of breast cancer. *Carcinogenesis* 34: 713–722.
50. Pellegrin S, Mellor H (2007) Actin stress fibres. *Journal of Cell Science* 120: 3491–3499.
51. Lee SH, Dominguez R (2010) Regulation of actin cytoskeleton dynamics in cells. *Mol Cells* 29: 311–325.
52. Tang AT, Campbell WB, Nithipatikom K (2012) ROCK1 feedback regulation of the upstream small GTPase RhoA. *Cellular Signalling* 24: 1375–1380.
53. Riento K, Ridley AJ (2003) Rocks: multifunctional kinases in cell behaviour. *Nature Reviews Molecular Cell Biology* 4: 446–456.
54. Nakagawa O, Fujisawa K, Ishizaki T, Saito Y, Nakao K, et al. (1996) ROCK-I and ROCK-II, two isoforms of Rho-associated coiled-coil forming protein serine/threonine kinase in mice. *FEBS Lett* 392: 189–193.
55. Kamai T, Tsujii T, Arai K, Takagi K, Asami H, et al. (2003) Significant association of Rho/ROCK pathway with invasion and metastasis of bladder cancer. *Clin Cancer Res* 9: 2632–2641.
56. Kaneko K, Satoh K, Masamune A, Satoh A, Shimosegawa T (2002) Expression of ROCK-1 in human pancreatic cancer: its down-regulation by morpholino oligo antisense can reduce the migration of pancreatic cancer cells in vitro. *Pancreas* 24: 251–257.
57. Spencer C, Montalvo J, McLaughlin SR, Bryan BA (2011) Small molecule inhibition of cytoskeletal dynamics in melanoma tumors results in altered transcriptional expression patterns of key genes involved in tumor initiation and progression. *Cancer Genomics Proteomics* 8: 77–85.
58. Somlyo AV, Bradshaw D, Ramos S, Murphy C, Myers CE, et al. (2000) Rho-kinase inhibitor retards migration and in vivo dissemination of human prostate cancer cells. *Biochem Biophys Res Commun* 269: 652–659.
59. Rattan R, Giri S, Singh AK, Singh I (2006) Rho/ROCK pathway as a target of tumor therapy. *J Neurosci Res* 83: 243–255.
60. Keller C, Kroening S, Zuehlke J, Kunath F, Krueger B, et al. (2012) Distinct mesenchymal alterations in N-cadherin and e-cadherin positive primary renal epithelial cells. *PLoS One* 7: e43584.
61. Deng L, Li G, Li R, Liu Q, He Q, et al. (2010) Rho-kinase inhibitor, fasudil, suppresses glioblastoma cell line progression in vitro and in vivo. *Cancer Biol Ther* 9: 875–884.
62. Lin CC, Chen JT, Yang JS, Lu HF, Hsu SC, et al. (2009) Danthron inhibits the migration and invasion of human brain glioblastoma multiforme cells through the inhibition of mRNA expression of focal adhesion kinase, Rho kinases-1 and metalloproteinase-9. *Oncol Rep* 22: 1033–1037.
63. Shields MA, Krantz SB, Bentrem DJ, Dangi Garimella S, Munshi HG (2012) Interplay between beta1-integrin and Rho signaling regulates differential scattering and motility of pancreatic cancer cells by snail and Slug proteins. *The Journal of biological chemistry* 287: 6218–6229.
64. Oellers P, Schroer U, Senner V, Paulus W, Thanos S (2009) ROCKs are expressed in brain tumors and are required for glioma-cell migration on myelinated axons. *Glia* 57: 499–509.
65. Liao JK, Seto M, Noma K (2007) Rho kinase (ROCK) inhibitors. *Journal of cardiovascular pharmacology* 50: 17–24.
66. Chen C, Ridzon DA, Broomer AJ, Zhou Z, Lee DH, et al. (2005) Real-time quantification of microRNAs by stem-loop RT-PCR. *Nucleic Acids Research* 33: e179. Available: <http://nar.oxfordjournals.org/content/33/20/e179.long>. Accessed 7 Nov 2005.
67. Tang F, Hajkova P, Barton SC, Lao K, Surani MA. (2006) MicroRNA expression profiling of single whole embryonic stem cells. *Nucleic Acids Research* 34: e9. Available: <http://nar.oxfordjournals.org/content/34/2/e9.long>. Accessed 24 Jan 2006.
68. Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-Delta Delta C(T)} Method. *Methods* 25: 402–408.
69. Machado CM, Schenka A, Vassallo J, Tamashiro WM, Goncalves EM, et al. (2005) Morphological characterization of a human glioma cell line. *Cancer cell international* 5: 13. Available: <http://www.cancerci.com/content/5/1/13>. Accessed 10 May 2005.