

Niche distribution and influence of environmental parameters in marine microbial communities: a systematic review

Felipe H. Coutinho^{1,2}, Pedro M. Meirelles¹, Ana Paula B. Moreira¹, Rodolfo P. Paranhos¹, Bas E. Dutilh^{1,2,4} and Fabiano L. Thompson^{1,3}

¹ Universidade Federal do Rio de Janeiro (UFRJ)/Instituto de Biologia (IB), Rio de Janeiro, Brazil

² Radboud University Medical Centre, Radboud Institute for Molecular Life Sciences, Centre for Molecular and Biomolecular Informatics (CMBI), Nijmegen, The Netherlands

³ Universidade Federal do Rio de Janeiro (UFRJ)/COPPE, SAGE, Rio de Janeiro, Brazil

⁴ University of Utrecht (UU), Theoretical Biology and Bioinformatics, Utrecht, The Netherlands

ABSTRACT

Associations between microorganisms occur extensively throughout Earth's oceans. Understanding how microbial communities are assembled and how the presence or absence of species is related to that of others are central goals of microbial ecology. Here, we investigate co-occurrence associations between marine prokaryotes by combining 180 new and publicly available metagenomic datasets from different oceans in a large-scale meta-analysis. A co-occurrence network was created by calculating correlation scores between the abundances of microorganisms in metagenomes. A total of 1,906 correlations amongst 297 organisms were detected, segregating them into 11 major groups that occupy distinct ecological niches. Additionally, by analyzing the oceanographic parameters measured for a selected number of sampling sites, we characterized the influence of environmental variables over each of these 11 groups. Clustering organisms into groups of taxa that have similar ecology, allowed the detection of several significant correlations that could not be observed for the taxa individually.

Subjects Computational Biology, Ecology, Genomics, Marine Biology, Microbiology

Keywords Metagenomics, Community ecology, Species interactions, Microbial ecology, Global ocean

INTRODUCTION

Assembly of microbial communities is believed to be simultaneously regulated by stochastic and deterministic processes (*Langenheder & Szekely, 2011; Jeraldo et al., 2012; Stegen et al., 2012*). Neutral theory postulates that the composition of biological communities is determined by stochastic processes only. In an extreme version of this theory, all species are considered ecologically equivalent, and their abundances between environments are influenced exclusively by random events of birth, death and dispersion (*Jeraldo et al., 2012*). In contrast, niche theory is based on the assumption that the species composition of an ecosystem is entirely determined by environmental conditions, a process known as habitat filtering (*Dumbrell et al., 2010; Pontarp et al., 2012*). This process results in communities

Submitted 6 December 2014

Accepted 19 May 2015

Published 16 June 2015

Corresponding author

Fabiano L. Thompson,
fabianothompson1@gmail.com

Academic editor

Ahmed Moustafa

Additional Information and
Declarations can be found on
page 13

DOI 10.7717/peerj.1008

© Copyright
2015 Coutinho et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

of co-existing organisms with largely overlapping ecological niches, meaning that they respond similarly to environmental conditions of their habitats and possibly compete for resources (Ulrich *et al.*, 2009; Maire *et al.*, 2012). In contrast, niche partitioning, allows co-occurring microorganisms to avoid competition by using different strategies to exploit the diversity of resources available at their environment (Macalady *et al.*, 2008).

Co-occurrence patterns between organisms can reveal ecological associations that take place between the members of a community. For example, if two organisms are frequently present together, and absent together, across multiple environments or samples, this can be interpreted as evidence that they occupy similar ecological niches (Horner-Devine *et al.*, 2007; Faust & Raes, 2012). Observing ecological associations among microbes *in situ* represents a much less trivial task than doing so for animals and plants. Therefore, analysis of co-occurrence networks represents an alternative to infer possible associations between microorganisms (Barberan *et al.*, 2012; Eiler, Heinrich & Bertilsson, 2012; Faust & Raes, 2012), and between microorganisms and environmental parameters (Ruan *et al.*, 2006; Gilbert *et al.*, 2012).

Here, we performed a meta-analysis of marine metagenomes from pelagic regions of the oceans around the globe, which includes previously published and new metagenomes from the South Atlantic Ocean, a poorly characterized marine realm. We identified patterns of co-variation between members of the marine microbiome. Our analysis identified hundreds of significant correlations that were used to build a co-occurrence network that sheds light into ecological processes taking place in the global ocean. Clustering the taxa of the network revealed groups of co-occurring prokaryotes that share a similar ecological niche. Next, we describe relationships between these groups and environmental parameters. Our results contribute to a better understanding of the processes that govern community assembly and inter-species co-occurrence patterns in the pelagic oceans, and provide important general insights for the understanding of microbial ecology.

METHODS

Samples

A collection of 180 metagenomes were retrieved from MG-RAST (Meyer *et al.*, 2008) (Table S1). Samples covered four major global oceans (Atlantic, Pacific, Indian, and Antarctic) and a broad depth range (0–4,800 m). Sampling sites of each metagenome are illustrated in Fig. 1. Among these samples, 71 metagenomes were obtained from the South Atlantic Ocean. These metagenomes were sampled, processed and analyzed as previously described (Bruce *et al.*, 2012; Alves Jr *et al.*, 2014). The remaining 109 metagenomes were obtained from distinct sites throughout the planet and were publicly available at the MG-RAST server. We chose our dataset aiming to cover a broad range of environmental conditions, allowing for enough variation in microbial abundance to occur between samples so that significant correlations can be detected. Our methodology has been shown to be appropriate to detect associations between microorganisms that can provide insightful information on their ecology (Fuhrman & Steele, 2008; Beman, Steele & Fuhrman, 2011; Steele *et al.*, 2011; Barberan *et al.*, 2012; Eiler, Heinrich &




Figure 1 Sample locations. Map of metagenome sampling sites: Blue circles represent metagenomes sampled at the South Atlantic Ocean. Yellow circles represent publicly available metagenomes from other regions of the planet.

Bertilsson, 2012; Faust & Raes, 2012). The differences in the environmental characteristics of the samples (e.g., location, season, depth) are required so that enough variation exists between samples so that relevant co-occurrence patterns can be detected. If we were to work with samples that were too homogeneous, very little variation would be observed concerning taxon abundances and environmental parameters, impairing the detection of relevant correlations. This broad range of environments provides the variation among samples that is necessary for non-spurious correlations to be detected, among taxa and also between taxa and environmental parameters (*Barberan et al., 2012*).

Prior to analysis, sequences from the MG-RAST metagenomes were de-replicated and filtered according to Phred score (≥ 20) and length (≥ 75 bp). No assembly was performed as to preserve the quantitative information within the metagenomes and to avoid the formation of chimeric sequences. All metagenomes were subjected to the same analysis pipeline. Taxonomic annotation was performed through the MG-RAST server best hit classification. Raw reads were translated in all 6 frames and aligned against Genbank as the reference database through BLAT (*Meyer et al., 2008*). This database was chosen due to the richness of complete genomes of marine microbes within it, such as those sequenced by the The Gordon and Betty Moore Foundation Marine Microbial Genome Sequencing Project. The cut-off parameters for annotation of a sequencing read were: e-value $\leq 10^{-5}$ and sequence identity $\geq 60\%$. Raw taxonomic counts were converted to relative abundances by dividing the count of each taxon by the total of annotated reads in each metagenome. A complete list of all 180 metagenomes including their MG-RAST identifiers, total number of reads, average read length, total bases, geographical coordinates, depth and original publication are available as [Table S1](#).

Physical and chemical parameters

The South Atlantic sampling sites were characterized regarding their water quality conditions, at the time of sampling, by the following methods: Chlorophyll-a analysis

was performed following positive pressure filtration of 2 L of seawater. Filters (glass fiber Whatman GF/F) were kept overnight under a solution of 90% acetone at 4 °C for extraction, and analyzed by spectrophotometry or fluorimetry. One liter of water from each sampling site was frozen and stored for further analysis of inorganic nutrients through the following methods: (1) ammonia by indophenol, (2) nitrite by diazotization, (3) nitrate by reduction in Cd–Cu column followed by diazotization, (4) total nitrogen by digestion with potassium persulfate following nitrate determination, (5) orthophosphate by reaction with ascorbic acid, (6) total phosphorous by acid digestion to phosphate, and (7) silicate by reaction with molybdate. All analyses were carried out as previously described ([Grasshoff, Kremling & Ehrhardt, 2009](#); [Alves Jr et al., 2014](#)).

Correlations network

Taxonomic annotations at the genus level were used as we considered any classification at deeper levels (i.e., species or strain) to be unreliable when dealing with the short reads from second generation sequencing technologies, which represent a significant fraction of our samples. Spearman rank correlation scores (R) were calculated between the relative abundances of all possible pairwise combinations of taxa. All taxa detected in less than 40% ($n = 72$) of the 180 samples, were excluded from this analysis to prevent sparsely distributed taxa with abundant zero values to yield spuriously high correlation scores. Multiple test correction was performed according to the False Discovery rate (FDR) procedure ([Benjamini & Hochberg, 1995](#)). Correlations for which the p -value and q -value were ≤ 0.001 and Spearman- R score ≤ -0.6 or $\geq +0.6$ were considered significant and plotted as a network through Cytoscape ([Saito et al., 2012](#)).

Node clustering

CFinder ([Palla, Barabasi & Vicsek, 2007](#)) was applied to identify clusters of highly connected taxa within the network through the Clique Percolation Method. “Cliques” are defined as groups of nodes (in these case the microbial taxa), which tend to have more connections with each other than with other members of a network. A k -step of 3 was chosen for this analysis so that cliques formed by three organisms or more could be identified. Our goal was to identify groups formed by taxa connected by significant positive correlations, therefore negative correlations were not considered by the Clique Percolation Method. Cliques were numbered according to the abundance of nodes assigned to each of them. Henceforth, these cliques will be referred to as “groups”.

Network consistency

The consistency of the correlations in the network was assessed through a sub-sampling strategy. One hundred new networks were calculated, each from a random sub-sample of 162 out of the 180 metagenomes. Next, the original and new networks were compared and we measured how often each one of the correlations from the original network were also detected in the new networks. The consistency of the groups identified through CPM was assessed by applying the algorithm over the 100 new networks and measuring how often

pairs of taxa that were assigned to the same group in the original network also clustered together by CPM in the new networks.

To account for occurrences of spurious correlations between genera due to the compositional (i.e., percentages) nature of our data, correlation scores of the original network were compared against those obtained through SparCC (Friedman & Alm, 2012). This tool was developed to calculate correlations between microbial abundances while eliminating errors that may emerge due to the use of compositional data. SparCC was run using default parameters.

Correlations between groups and environmental parameters

We addressed the influence of habitat variables over the groups identified by CFinder. For that end, Spearman rank correlation scores were calculated between the relative abundances of the groups and environmental parameters measured for the South Atlantic Ocean samples ($n = 71$, Table S2). Group abundances were calculated as the sum of the relative abundances of all of its members. The environmental parameters used for this analysis were: total nitrogen, nitrite, nitrate, ammonia, total phosphorus, orthophosphate, chlorophyll a, silicate, temperature, depth, latitude and longitude. To identify associations between groups, Spearman correlation scores were calculated for all the possible pairwise combinations of groups. This step of the analysis encompassed all metagenomes ($n = 180$). In both cases, only correlations which yielded a p -value ≤ 0.01 and q -value < 0.05 were considered significant.

RESULTS

The taxonomic composition of 180 marine metagenomes was used to build a network of correlations between genera of microorganisms. Next, the members of this network were clustered in groups based on the significant correlations between them. Correlation scores were measured between pairwise combinations of these groups and between the groups and environmental parameters of the South Atlantic sampling sites.

Network parameters

The resulting network was composed of 297 taxa (nodes) and 1,906 correlations (edges) (Fig. 2), of which 1,863 were positive correlations (mean $R = 0.68 \pm 0.06$) and 43 were negative correlations (mean $R = -0.62 \pm 0.02$). The network had a clustering coefficient (tendency of nodes to cluster together) of 0.54 and density (the number of edges in the network divided by the possible maximum) of 0.04. The average node degree (number of connections of a node) was 5.3 ± 8.04 . Using the Clique Percolation Method, we identified eleven groups of co-occurring microbial taxa, containing between three and eighty members. Of the 297 microbial taxa, 229 were assigned to a single group, while 65 were assigned to none. Three genera were assigned to more than a single group: *Marinomonas* (present in two groups), *Polynucleobacter* (two groups) and *Haliangium* (three groups). The composition of these groups is described in Table S3.

Out of the 1,906 correlations in the original network 1,843 (98%) were consistent in at least half of the sub-sampled networks. Groups assignments also yielded high consistency,




Figure 2 Correlations network: the 1,906 edges linking 297 nodes represent significant correlations between the relative abundances of the connected taxa. Positive correlations are showcased in green while negative ones are in blue. The width of the lines is proportional to the module of Spearman's R of each correlation. Node size represents the average abundance of the taxa across the 180 metagenomes. Nodes are color-coded according to the group to which they were assigned through the Clique Percolation Method. Nodes not assigned to any group are colored in white and nodes assigned to more than a single group (Polynucleobacter, Marinomonas and Haliangium) are colored in gray. For clarity, members of Groups 1 and 2 that are connected by negative correlations are displayed separately from the remaining taxa of their respective groups. *Pelagibacter*, *Prochlorococcus* and *Synechococcus* showed the highest average abundance. Positive correlations dominate the network. The majority of negative correlations were observed between members of groups 1 and 2, between classical examples of oligotrophs and copiotrophs (e.g., *Pelagibacter*/*Yersinia*). Groups 8 and 9 are isolated, while the remaining groups have at least one edge linking them to other nodes in the network. Strong positive correlations were observed between the members of groups 8 and 9 (e.g., *Synechococcus*/*Prochlorococcus* and *Cenaracheum*/*Nitrosopumilus*).

99% of the pairwise group assignments of the original network were consistent across at least 50% of the sub-sampled networks. Comparison between networks calculated through Spearman correlations and SparCC, revealed that 98,7% of all correlations detected in the first were also present in the latter (module of $R > 0.3$), indicating that correlations of the original network are not spurious due to the compositional nature of the data. The difference in correlation scores between the networks are likely the effect of non-linear associations between variables, that cannot be captured by SparCC.

Composition of groups identified through the Clique Percolation Method

Group 1 harbored 80 genera distributed among seven bacterial phyla (*Firmicutes*, *Chloroflexi*, *Fusobacteria*, *Planctomycetes*, *Proteobacteria*, *Bacteroidetes*, and *Actinobacteria*), this group had more members than any other and also harbored the highest number of phyla. Many of the members of Group 1 are typical genera of heterotrophic aquatic bacteria (e.g., *Vibrio* and *Shewanella*). Group 2, formed by 78 members, was dominated by

genera of *Alphaproteobacteria* (e.g., *Puniceispirillum* and *Rhodobacter*) and *Bacteroidetes* (e.g., *Chryseobacterium* and *Cytophaga*), this group also harbored the ubiquitous *Pelagibacter* genus. Most of the members of Group 3 were *Actinobacteria* or *Chloroflexi*, but other phyla (e.g., *Acidobacteria*, *Firmicutes*, *Proteobacteria* and *Chlorobi*) were also represented in this group. Group 4 was formed by 13 genera of *Gammaproteobacteria*, from the orders *Pseudomonadales*, *Alteromonadales*, *Chromatiales* and *Oceanospirillales*. The majority of genera assigned to Group 5 were either *Verrucomicrobia* or *Planctomycetes*, with only three exceptions: *Haliangiium*, *Stigmatella* and *Lentisphaera*. Group 6 was formed by genera of *Euryarchaeota*, which included several methanogenic organisms (e.g., *Methanothermobacter*). Group 7 was formed by genera of photosynthetic organisms of the phylum *Cyanobacteria* (e.g., *Anabaena*, *Cyanothece* and *Lyngbya*). Group 8 was composed entirely of *Archaea*, including the genera *Cenarchaeum* and *Nitrosopumilus* as well as unclassified organisms from the phyla *Thaumarchaeota* and *Crenarchaeota*. Group 9 was also composed entirely of *Cyanobacteria*. The photosynthesizers *Prochlorococcus* and *Synechococcus*, both among the most abundant organisms in the analyzed metagenomes, were assigned to this group, along with *Cyanobium*. Group 10 was formed by three genera of *Myxobacteria*: *Myxococcus*, *Plesiocystis* and *Haliangiium*, the latter was also a member of groups 4 and 5. Only three taxa are part of Group 11, all *Proteobacteria*: *Polynucleobacter*, *Methylococcus* and *Sideroxydans*.

Correlations between groups and environmental parameters

Correlations calculated between environmental parameters, of the South Atlantic Ocean and the relative abundance of the groups of microorganisms produced unique patterns for each group (Fig. 3A). With the exception of ammonia, all variables produced at least one significant correlation with at least one group. Total phosphorus produced significant correlations with eight groups, more than any of the other variable. The pattern of correlations detected between silicate and seven of the groups was similar to that observed for total phosphorus, but silicate showed no significant correlation with Group 4. The third variable with most significant correlations was depth (5 groups). The variables that produced least significant correlations were Chlorophyll a (1), latitude (1) and Ammonia (0).

The abundance of Group 1, which harbored many heterotrophic and pathogenic organisms, showed positive correlations with depth and several nutrients (i.e., orthophosphate, total phosphorus, silicate, nitrate and total nitrogen), also, a negative correlation was detected between salinity and the abundance of this group. The abundance of Group 2 (mainly *Alphaproteobacteria* and *Bacteroidetes*) was negatively correlated with three nutrients (nitrate, total phosphorus and silicate). Additionally, a positive correlation with nitrite was observed for this group. Significant positive correlations were obtained between the abundance of Group 3, composed of many extremophilic organisms, and total nitrogen, total phosphorus, orthophosphate and silicate. This same group had negative correlations with salinity, temperature, latitude and longitude. Positive correlations were detected between Group 4 (all *Gammaproteobacteria*) and both nitrate and total phosphorus. The abundance of group 5 showed positive correlations with orthophosphate and




Figure 3 Correlations between groups and environmental parameters. Heatmap of correlation scores: (A) Correlations calculated between group abundances and environmental parameters. (B) Correlations calculated between group abundances. Positive correlations are showcased in green while negative ones are in blue. Non-significant correlations ($p > 0.01$ or $q > 0.05$) are shown as white squares.

depth, and a negative one with temperature. Group 6, composed mainly of methanogenic *Archaea*, had a positive correlation with depth, no other significant correlations were detected between the abundance of these organisms and the other environmental variables. Group 7, composed of *Cyanobacteria*, showed negative correlations with total nitrogen, orthophosphate, total phosphorus and silicate. Positive correlations between salinity and longitude were also observed for Group 7. Positive correlations were detected between Group 8 and nitrate, total phosphorus, silicate and depth, and a negative correlation with temperature. Group 9, also formed by *Cyanobacteria*, showed negative correlations with total nitrogen, orthophosphate, total phosphorus, silicate and depth, positive correlations with salinity, temperature and longitude were also observed. Group 10 showed positive correlations with total phosphorus and silicate. A positive correlation between Chlorophyll and Group 11 was observed, no other significant correlations were observed for this group or this environmental parameter.

Several relationships emerged from correlating the relative abundances of the eleven groups between each other (Fig. 3B). Both positive and negative, significant correlations were detected and each group presented a unique pattern of correlations with the others.

DISCUSSION

Assessing technical heterogeneity biases

We opted for using samples derived from a broad spatial range (Fig. 1) so that we could identify microbial occurrence patterns that are applicable to the entirety of the oceans. Ideally, all of our samples would have been processed with the exact same protocols (e.g., for water filtering, DNA extraction and sequencing). Yet, restricting our dataset to one that fits these criteria would result in a very small number of samples, drastically impairing both the power of our approach and the relevance of our results, which emerge from the use of a large number of samples from a broad range of environments. The use of different sequencing technologies can indeed yield slightly different results for taxonomic composition. Nevertheless, these differences were shown to be of very little impact for the overall patterns of community composition (Danhorn, Young & Delong, 2012; Luo et al., 2012;

Solonenko et al., 2013). This is in agreement with previous studies which have shown that despite the potential biases that may be introduced by these methodologies, metagenomes carry a strong taxonomic and functional signal which is not surpassed by sample preparation biases (*Dinsdale et al., 2008; Willner, Thurber & Rohwer, 2009; Faust et al., 2012*).

We addressed the potential biases in metagenomes that could emerge from different sample preparation strategies in different ways. First, we assessed to what extent samples are grouped according to the laboratory by which they were processed. To do this, Euclidean distances were calculated between metagenomes based on their genera composition. These distances were used as input for Principal Coordinates Analysis (PCoA). In a scenario where the taxonomic composition of a metagenome is strongly determined by sample processing, metagenomes are expected to cluster tightly by laboratory. This pattern is not present among our samples ([Fig. S1](#)).

Second, we evaluated which of the 297 taxa of the network are over or under-represented in the samples from one laboratory, when compared to the samples from the remaining laboratories, by using the Mann–Whitney test, with multiple testing correction through the FDR. We assume that potentially biased taxa are those that are significantly over or under-represented on samples of a single laboratory. However, if a taxon is over or under-represented across multiple laboratories, it is more likely that this is a true biological signal of the samples analyzed by those projects, rather than a bias emerging from different sample processing methodologies. According to these criteria, only 14 taxa could be potentially biased (q value < 0.05 , see [Table S4](#) for the full list of taxa, the laboratories in which they are enriched and the groups to which they were assigned within the network according to CFinder). It is not possible to determine if this pattern emerges from a true biological signal within the samples, or if they are the result of sample preparation methodologies. Nevertheless, if the case is the latter this is likely to be over very little influence in the overall results, considering that only a very small fraction of all the taxa in the network fall within this category ($< 5\%$).

Third, to further explore issues that might arise from sample preparation, we recalculated a correlations network using only the 71 samples from the South Atlantic ocean sampled by Thompson et al. (see [Table S1](#)). These represent the largest consistent group regarding sample preparation strategies. We then compared the correlation scores from the global network (180 metagenomes) to those obtained using only the south Atlantic samples (71 metagenomes, see [Table S5](#)). The average absolute Spearman- R values of these two networks are respectively 0.67 ± 0.06 and 0.37 ± 0.21 and a significant correlation exists between these values (Pearson R 0.37, p -value $< 2.2^{-16}$), providing yet another evidence that the 1,906 correlations from the global network are consistent within a homogeneous dataset regarding sample preparation methods and do not result from sample preparation bias.

Groups are formed by organisms with shared ecological niches

Several genera assigned to Group 1 are copiotrophic bacteria (i.e., thrive in nutrient rich conditions) such as *Vibrio*, *Pseudomonas*, *Escherichia* and *Clostridium*. As a consequence

of their nutritional demands, these genera are frequently abundant in eutrophic waters (Gilbert *et al.*, 2010; Gregoracci *et al.*, 2012), in which nutrient concentrations are high. In contrast, many of the genera affiliated to Group 2 are oligotrophic bacteria, characteristic of aquatic environments where nutrients are scarce. The extremely abundant genus *Pelagibacter*, which possesses several adaptations to thrive in nutritionally poor waters (Carini *et al.*, 2013; Tripp, 2013) was assigned to this group, along with other genera adapted to live at nutrient deprived environments, such as the chemolithoautotrophic *Oligotropha* (Paul *et al.*, 2008) and the heterotrophic *Maricaulis* (Abraham *et al.*, 1999). Therefore, trophic strategies appear to be the unifying trait of the members of these two groups.

Many of the members of Group 3 are capable of surviving in extreme habitats, such as *Thermomicrobium*, capable of growing in elevated temperatures, and *Acidothermus*, capable of tolerating both low pH and high temperatures (Mohagheghi *et al.*, 1986; Wu *et al.*, 2009). These organisms were detected in metagenomes from samples retrieved at non-extreme environments, and encompassed several phyla. Despite their phylogenetic distance and differences regarding their adaptations to thrive in extreme environments, these genera of extremotolerants may share similar habitat preferences at mesophilic waters. Shared niche can also explain the correlations occurring within Group 6, which is dominated by genera of methanogenic *Archaea*. Co-variation in the abundance of these organisms is expected since methanogenesis usually takes place in very specific environments: rich in organic matter and poorly oxygenated (Peng *et al.*, 2008; Angel, Matthies & Conrad, 2011). Even though we applied a highly conservative e-value cutoff ($\leq 1 \times 10^{-5}$) it is also possible that some of the sequences received incorrect taxonomic assignments, which could explain the presence of extremophilic organisms at mesophilic environments.

The highly abundant genera *Prochlorococcus* and *Synechococcus* were assigned to Group 9, along with *Cyanobium*. Meanwhile the much less abundant genera of *Cyanobacteria* (e.g., *Anabaena* and *Cyanothece*) were all assigned to Group 7. No significant correlations were detected between any of the members of these two groups. Differences in how these two groups make use of the resources available at the marine ecosystem could be responsible for the increased ubiquity and abundance of *Prochlorococcus* and *Synechococcus* among Earth's oceans, traits which are not shared with the *Cyanobacteria* of Group 7. The success of *Prochlorococcus* sp., especially in oligotrophic waters, has been attributed to reduced genome and cell sizes, as well as high rates of nutrient uptake (Zubkov *et al.*, 2003; Partensky & Garczarek, 2010). Distinctive traits of bacteria from Group 7 include the formation of filamentous colonies in the water column (Rice, Mazur & Haselkorn, 1982; Sanudo-Wilhelmy *et al.*, 2001), blooms (Albert *et al.*, 2005), and also the diazotrophic metabolism present in several of its members (Reddy *et al.*, 1993; Omoregie *et al.*, 2004; Bergman *et al.*, 2013).

Based on these observations, we may conclude that the groups identified in the network are formed by organisms which occupy similar ecological niches (Chaffron *et al.*, 2010; Freilich *et al.*, 2010; Faust *et al.*, 2012). Positive correlations could also be the result of

collaborative associations (i.e., mutualism). Our data does not allow us to differentiate positive correlations emerging from sharing of an ecological niche from mutualistic associations (Chaffron *et al.*, 2010; Faust *et al.*, 2012). Although these associations may occur, previous studies have also concluded that positive correlations detected at microbial correlation networks are representative of niche overlap rather than cooperative associations and consider this to be a more parsimonious explanation for the occurrence of these correlations (Foster & Bell, 2012; Levy & Borenstein, 2013).

Previous analysis of microbial networks have reported that phylogenetically related organisms co-occur with each other more than expected by chance, as consequence of sharing a niche (Chaffron *et al.*, 2010; Barberan *et al.*, 2012). Some of the groups identified consisted of phylogenetically related taxa (e.g., groups 4, 6, and 10). However we also observed groups composed of distantly related organisms (e.g., groups 1, 2 and 3). Moreover, closely related organisms sometimes occurred in different groups (e.g., *Cyanobacteria* divided between groups 7 and 9). These patterns suggest that group composition was determined by niche distribution, rather than by phylogenetic relatedness.

It is important to take some matters into consideration when interpreting the results of the network. Organisms that were assigned to the same group do not necessarily occupy identical ecological niches. Instead, organisms of the same group have a higher degree of niche similarity between themselves when compared to other taxa of the network. *Synechococcus* and *Prochlorococcus* can be taken as an example of this pattern. These genera were grouped together by the Clique Percolation Method, the two taxa have well defined differences in their niche preferences, with regard to their geographic and temporal distributions, light harvesting apparatus and nutrient acquisition machinery (Partensky, Blanchot & Vaulot, 1999; Scanlan *et al.*, 2009). Despite those differences, the two groups of picocyanobacteria have many similarities in central aspects of their physiology, such as the photosynthetic metabolism, reduced cell and genome sizes, and carbon concentration mechanisms. These similarities may be the more important aspects that regulate their response to environmental conditions, thus giving rise to the strong positive correlations observed between their abundances. In addition, the taxonomy of *Synechococcus* and *Prochlorococcus* has not been fully elucidated, thus erroneous taxonomic assignments may have contributed for the correlations between the groups.

Also, it is likely that some of the organisms detected in the metagenomes are dormant or inactive. At this state, these organisms are not responding to the fluctuations of the environmental parameters taking place at their habitat. Therefore, no covariation should be expected to occur between these organisms and the active ones. Therefore, no false-positives are expected to arise due to the occurrence of inactive organisms.

Niche segregation in marine microbial communities

Each of the groups identified by the Clique Percolation Method represent genera that share a similar ecological niche. This means that our analysis detected 11 groups of prokaryotes with distinct ecologies. This is a conservative number since the heterogeneity and extension of the oceans contribute to a much wider diversity of niches to be occupied.

Additionally, the number of groups detected will depend on methodological parameters including the k -step chosen for the Clique Percolation Method ($k = 3$), the cutoff established for the Spearman correlations scores (≥ 0.6 or ≤ -0.6) and the minimum ubiquity of the taxa among the 180 samples (40%). Nevertheless, the detected correlations and group compositions were consistent in the sub-sampled networks. Finally, since the very rare organisms (i.e., detected in less than 40% of all metagenomes) were excluded from the network it is possible that the groups formed by them were disregarded as well.

Studies based on genomic analysis have shown that small variations in protein encoding genes may lead organisms that belong to the same genus, or even the same species to occupy different niches (Rocap *et al.*, 2003; Kashtan *et al.*, 2014). Species, and strains can have unique traits (e.g., strategies for assimilation of organic compounds and spatial distribution) that set them apart from the other members of their genus (Johnson *et al.*, 2006; Del Carmen Muñoz-Martín *et al.*, 2013; Thompson *et al.*, 2013). These differences could further segregate them into sub-divisions of the 11 groups identified through our network. Unfortunately, due to their length, metagenomic reads generated by second generation sequencing technologies cannot produce reliable annotations at taxonomic levels deeper than genus.

Influence of environmental parameters

Many associations were detected between the groups and environmental parameters. Yet, none of them yielded perfect correlation scores with the abundance the microbial groups. This is expected considering that the abundance of these groups, in the environment, is regulated by many variables simultaneously. Therefore, it is unlikely that a single measured physical or chemical parameter can adequately explain the abundance of a group across all the samples. Nevertheless, the pattern of significant correlations detected between group abundances and environmental parameters provides insights into how the members of these groups are influenced by environmental variables of their habitat and what niche is occupied by them.

Positive correlations were detected between several nutrients (i.e., nitrate, total nitrogen, orthophosphate, total phosphorus and silicate) and the abundance of Group 1 (Copiotrophs). Meanwhile, negative correlations were detected between Group 2 (Oligotrophs) and nitrate, silicate and total phosphorus (Fig. 3A). This pattern shows that members of Group 1 are more abundant in waters that are rich in these nutrients, while Group 2 is more abundant in regions of the ocean deprived of them. This is corroborated by our observation that an increase in the abundance of one group is accompanied by a reduction in the abundance of the other, leading to significant anti-correlations between the abundances of these organisms, which were observed when comparing the abundances of the genera individually (Fig. 2) and of the whole groups (Fig. 3B).

Groups 7 and 9, composed of *Cyanobacteria*, both showed negative correlations with total nitrogen, orthophosphate, total phosphorus, and silicate. These results are in accordance with genomic analyses of *Synechococcus* sp. and *Prochlorococcus* sp., members of

Group 9, which possess several features that contribute to the success of these organisms in oligotrophic conditions (Lauro *et al.*, 2009).

Clustering unveils the ecology of marine microorganisms

The associations between group abundances and the physicochemical environmental parameters (Fig. 3A) provide insights regarding the ecological roles and niche preferences of the identified groups. Interestingly, such correlations were not significant when these same habitat variables were compared with the abundance of the individual genera that are part of these groups. We suggest that clustering organisms based on co-occurrence is a useful and necessary tool to reveal elusive associations between microorganisms and habitat variables.

Considering the evidence that the organisms from the same group occupy similar niches, it is possible that they are ecologically redundant (i.e., they contribute with the same, or at least similar, roles for ecosystem stability). We propose that each of the groups identified in the network contributes to ecosystem stability by performing a distinct ecological role at the marine environment. Members of a microbial community eventually decline as a result of processes such as phage predation, grazing or a drastic change in environmental conditions. Withering organisms can be replaced by their ecological equivalent, which possesses the necessary features to fill the niche left unoccupied (Gifford *et al.*, 2012). This cycling of species would preserve the ecological roles that are necessary to sustain an ecosystem, as described by the insurance hypothesis (Yachi & Loreau, 1999; Allison & Martiny, 2008). In that case, the influence of environmental parameters would act upon all the members of a niche and not over the individual taxa (as these are redundant and interchangeable), which could explain why these correlations could only be detected at the group level.

CONCLUSIONS

The correlations network proved to be a valuable tool to disclose shared ecological niches in the global ocean microbiome. Habitat filtering and niche segregation may be considered important factors controlling the taxonomic composition of microbial communities from different locations of the global ocean. The data presented here provides insights into the ecological processes that structure marine microbial communities and show that clustering organisms into ecologically cohesive groups may reveal elusive associations between microbes and habitat variables. The advancement of technologies that allow microbial communities to be studied, *in situ* and at higher spatial resolution (i.e., at the micro, rather than the macro-scale) will help broaden the scope of the analysis presented here, allowing for deeper understanding of the ecology of marine microbial communities.

ACKNOWLEDGEMENTS

Special thanks to the members of the Centre of Molecular and Biomolecular Informatics and of the Laboratory of Microbiology at UFRJ for insightful discussions. The authors thank Chrisine Hörnlein for critical comments on the manuscript.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was funded by grants from CNPq, CAPES and FAPERJ. BED was supported by the Dutch Virgo Consortium (FES0908, NGI 050-060-452) and CAPES/BRASIL. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

CNPq.

CAPES.

FAPERJ.

Dutch Virgo Consortium: FES0908, NGI 050-060-452.

CAPES/BRASIL.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Felipe H. Coutinho conceived and designed the experiments, performed the experiments, analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Pedro M. Meirelles conceived and designed the experiments, performed the experiments, analyzed the data, wrote the paper, reviewed drafts of the paper.
- Ana Paula B. Moreira performed the experiments.
- Rodolfo P. Paranhos performed the experiments, contributed reagents/materials/analysis tools.
- Bas E. Dutilh and Fabiano L. Thompson conceived and designed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, reviewed drafts of the paper.

DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

MG-RAST

4464153.3

4464162.3

4464167.3

4464168.3

4464169.3

4464170.3

4464171.3

4464173.3
4465773.3
4465774.3
4465775.3
4484652.3
4484653.3
4484654.3
4484655.3
4484656.3
4484657.3
4487641.3
4487642.3
4487643.3
4487644.3
4487645.3
4487646.3.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.1008#supplemental-information>.

REFERENCES

- Abraham WR, Strompl C, Meyer H, Lindholst S, Moore ER, Christ R, Vancanneyt M, Tindall BJ, Bennasar A, Smit J, Tesar M. 1999. Phylogeny and polyphasic taxonomy of *Caulobacter* species. Proposal of *Maricaulis* gen. nov. with *Maricaulis maris* (Poindexter) comb. nov. as the type species, and emended description of the genera *Brevundimonas* and *Caulobacter*. *International Journal of Systematic Bacteriology* **49**(Pt 3):1053–1073 DOI [10.1099/00207713-49-3-1053](https://doi.org/10.1099/00207713-49-3-1053).
- Albert S, O’Neil JM, Udy JW, Ahern KS, O’Sullivan CM, Dennison WC. 2005. Blooms of the cyanobacterium *Lyngbya majuscula* in coastal Queensland, Australia: disparate sites, common factors. *Marine Pollution Bulletin* **51**:428–437 DOI [10.1016/j.marpolbul.2004.10.016](https://doi.org/10.1016/j.marpolbul.2004.10.016).
- Allison SD, Martiny JB. 2008. Colloquium paper: resistance, resilience, and redundancy in microbial communities. *Proceedings of the National Academy of Sciences of the United States of America* **105**(Suppl):11512–11519 DOI [10.1073/pnas.0801925105](https://doi.org/10.1073/pnas.0801925105).
- Alves Jr N, Meirelles PM, de Oliveira Santos E, Dutilh B, Silva GG, Paranhos R, Cabral AS, Rezende C, Iida T, de Moura RL, Kruger RH, Pereira RC, Valle R, Sawabe T, Thompson C, Thompson F. 2014. Microbial community diversity and physical–chemical features of the Southwestern Atlantic Ocean. *Archives of Microbiology* **197**(2):165–179 DOI [10.1007/s00203-014-1035-6](https://doi.org/10.1007/s00203-014-1035-6).
- Angel R, Matthies D, Conrad R. 2011. Activation of methanogenesis in arid biological soil crusts despite the presence of oxygen. *PLoS ONE* **6**:e20453 DOI [10.1371/journal.pone.0020453](https://doi.org/10.1371/journal.pone.0020453).
- Barberan A, Bates ST, Casamayor EO, Fierer N. 2012. Using network analysis to explore co-occurrence patterns in soil microbial communities. *The ISME Journal* **6**:343–351 DOI [10.1038/ismej.2011.119](https://doi.org/10.1038/ismej.2011.119).

- Beman JM, Steele JA, Fuhrman JA. 2011.** Co-occurrence patterns for abundant marine archaeal and bacterial lineages in the deep chlorophyll maximum of coastal California. *The ISME Journal* 5:1077–1085 DOI 10.1038/ismej.2010.204.
- Benjamini Y, Hochberg Y. 1995.** Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)* 289–300.
- Bergman B, Sandh G, Lin S, Larsson J, Carpenter EJ. 2013.** Trichodesmium—a widespread marine cyanobacterium with unusual nitrogen fixation properties. *FEMS Microbiology Reviews* 37:286–302 DOI 10.1111/j.1574-6976.2012.00352.x.
- Bruce T, Meirelles PM, Garcia G, Paranhos R, Rezende CE, de Moura RL, Filho RF, Coni EO, Vasconcelos AT, Amado Filho G, Hatay M, Schmieder R, Edwards R, Dinsdale E, Thompson FL. 2012.** Abrolhos bank reef health evaluated by means of water quality, microbial diversity, benthic cover, and fish biomass data. *PLoS ONE* 7:e36687 DOI 10.1371/journal.pone.0036687.
- Carini P, Steindler L, Beszteri S, Giovannoni SJ. 2013.** Nutrient requirements for growth of the extreme oligotroph “Candidatus *Pelagibacter ubique*” HTCC1062 on a defined medium. *The ISME Journal* 7:592–602 DOI 10.1038/ismej.2012.122.
- Chaffron S, Rehrauer H, Pernthaler J, von Mering C. 2010.** A global network of coexisting microbes from environmental and whole-genome sequence data. *Genome Research* 20:947–959 DOI 10.1101/gr.104521.109.
- Danhorn T, Young CR, Delong EF. 2012.** Comparison of large-insert, small-insert and pyrosequencing libraries for metagenomic analysis. *The ISME Journal* 6:2056–2066 DOI 10.1038/ismej.2012.35.
- Del Carmen Muñoz-Marín M, Luque I, Zubkov MV, Hill PG, Diez J, García-Fernández JM. 2013.** *Prochlorococcus* can use the Pro1404 transporter to take up glucose at nanomolar concentrations in the Atlantic Ocean. *Proceedings of the National Academy of Sciences of the United States of America* 110:8597–8602 DOI 10.1073/pnas.1221775110.
- Dinsdale EA, Edwards RA, Hall D, Angly F, Breitbart M, Brulc JM, Furlan M, Desnues C, Haynes M, Li L, McDaniel L, Moran MA, Nelson KE, Nilsson C, Olson R, Paul J, Brito BR, Ruan Y, Swan BK, Stevens R, Valentine DL, Thurber RV, Wegley L, White BA, Rohwer F. 2008.** Functional metagenomic profiling of nine biomes. *Nature* 452:629–632 DOI 10.1038/nature06810.
- Dumbrell AJ, Nelson M, Helgason T, Dytham C, Fitter AH. 2010.** Relative roles of niche and neutral processes in structuring a soil microbial community. *The ISME Journal* 4:337–345 DOI 10.1038/ismej.2009.122.
- Eiler A, Heinrich F, Bertilsson S. 2012.** Coherent dynamics and association networks among lake bacterioplankton taxa. *The ISME Journal* 6:330–342 DOI 10.1038/ismej.2011.113.
- Faust K, Raes J. 2012.** Microbial interactions: from networks to models. *Nature Reviews Microbiology* 10:538–550 DOI 10.1038/nrmicro2832.
- Faust K, Sathirapongsasuti JF, Izard J, Segata N, Gevers D, Raes J, Huttenhower C. 2012.** Microbial co-occurrence relationships in the human microbiome. *PLoS Computational Biology* 8:e1002606 DOI 10.1371/journal.pcbi.1002606.
- Foster KR, Bell T. 2012.** Competition, not cooperation, dominates interactions among culturable microbial species. *Current Biology* 22:1845–1850 DOI 10.1016/j.cub.2012.08.005.
- Freilich S, Kreimer A, Meilijson I, Gophna U, Sharan R, Ruppin E. 2010.** The large-scale organization of the bacterial network of ecological co-occurrence interactions. *Nucleic Acids Research* 38:3857–3868 DOI 10.1093/nar/gkq118.

- Friedman J, Alm EJ. 2012. Inferring correlation networks from genomic survey data. *PLoS Computational Biology* 8:e1002687 DOI 10.1371/journal.pcbi.1002687.
- Fuhrman J, Steele J. 2008. Community structure of marine bacterioplankton: patterns, networks, and relationships to function. *Aquatic Microbial Ecology* 53:69–81 DOI 10.3354/ame01222.
- Gifford SM, Sharma S, Booth M, Moran MA. 2012. Expression patterns reveal niche diversification in a marine microbial assemblage. *The ISME Journal* 7:281–298 DOI 10.1038/ismej.2012.96.
- Gilbert JA, Field D, Swift P, Thomas S, Cummings D, Temperton B, Weynberg K, Huse S, Hughes M, Joint I, Somerfield PJ, Muhling M. 2010. The taxonomic and functional diversity of microbes at a temperate coastal site: a “multi-omic” study of seasonal and diel temporal variation. *PLoS ONE* 5:e15545 DOI 10.1371/journal.pone.0015545.
- Gilbert JA, Steele JA, Caporaso JG, Steinbrück L, Reeder J, Temperton B, Huse S, McHardy AC, Knight R, Joint I, Somerfield P, Fuhrman JA, Field D. 2012. Defining seasonal marine microbial community dynamics. *The ISME Journal* 6:298–308 DOI 10.1038/ismej.2011.107.
- Grasshoff K, Kremling K, Ehrhardt M (eds.) 2009. *Methods of seawater analysis*. John Wiley & Sons.
- Gregoracci GB, Nascimento JR, Cabral AS, Paranhos R, Valentin JL, Thompson CC, Thompson FL. 2012. Structuring of bacterioplankton diversity in a large tropical bay. *PLoS ONE* 7:e31408 DOI 10.1371/journal.pone.0031408.
- Horner-Devine MC, Silver JM, Leibold MA, Bohannan BJM, Colwell RK, Fuhrman JA, Green JL, Kuske CR, Martiny JBH, Muyzer G, Øvreås L, Reysenbach AL, Smith VH. 2007. A comparison of taxon co-occurrence patterns for macro- and microorganisms. *Ecology* 88:1345–1353 DOI 10.1890/06-0286.
- Jeraldo P, Sipos M, Chia N, Brulc JM, Dhillon AS, Konkel ME, Larson CL, Nelson KE, Qu A, Schook LB, Yang F, White BA, Goldenfeld N. 2012. Quantification of the relative roles of niche and neutral processes in structuring gastrointestinal microbiomes. *Proceedings of the National Academy of Sciences of the United States of America* 109:9692–9698 DOI 10.1073/pnas.1206721109.
- Johnson ZI, Zinser ER, Coe A, McNulty NP, Woodward EM, Chisholm SW. 2006. Niche partitioning among *Prochlorococcus* ecotypes along ocean-scale environmental gradients. *Science* 311:1737–1740 DOI 10.1126/science.1118052.
- Kashtan N, Roggensack SE, Rodrigue S, Thompson JW, Biller SJ, Coe A, Ding H, Marttinen P, Malmstrom RR, Stocker R, Follows MJ, Stepanauskas R, Chisholm SW. 2014. Single-cell genomics reveals hundreds of coexisting subpopulations in wild *Prochlorococcus*. *Science* 344:416–420 DOI 10.1126/science.1248575.
- Langenheder S, Szekely AJ. 2011. Species sorting and neutral processes are both important during the initial assembly of bacterial communities. *The ISME Journal* 5:1086–1094 DOI 10.1038/ismej.2010.207.
- Lauro FM, McDougald D, Thomas T, Williams TJ, Egan S, Rice S, Demaree MZ, Ting L, Ertan H, Johnson J, Ferreira S, Lapidus A, Anderson I, Kyrpides N, Munk AC, Dettler C, Han CS, Brown MV, Robb FT, Kjelleberg S, Cavicchioli R. 2009. The genomic basis of trophic strategy in marine bacteria. *Proceedings of the National Academy of Sciences of the United States of America* 106:15527–15533 DOI 10.1073/pnas.0903507106.
- Levy R, Borenstein E. 2013. Metabolic modeling of species interaction in the human microbiome elucidates community-level assembly rules. *Proceedings of the National Academy of Sciences of the United States of America* 110:12804–12809 DOI 10.1073/pnas.1300926110.

- Luo C, Tsementzi D, Kyrpides N, Read T, Konstantinidis KT. 2012. Direct comparisons of Illumina vs. Roche 454 sequencing technologies on the same microbial community DNA sample. *PLoS ONE* 7:e30087 DOI 10.1371/journal.pone.0030087.
- Macalady JL, Dattagupta S, Schaperdoth I, Jones DS, Druschel GK, Eastman D. 2008. Niche differentiation among sulfur-oxidizing bacterial populations in cave waters. *The ISME Journal* 2:590–601 DOI 10.1038/ismej.2008.25.
- Maire V, Gross N, Börger L, Proulx R, Wirth C, Pontes LDS, Soussana JF, Louault F. 2012. Habitat filtering and niche differentiation jointly explain species relative abundance within grassland communities along fertility and disturbance gradients. *New Phytologist* 196:497–509 DOI 10.1111/j.1469-8137.2012.04287.x.
- Meyer F, Paarmann D, D'Souza M, Olson R, Glass EM, Kubal M, Paczian T, Rodriguez A, Stevens R, Wilke A, Wilkening J, Edwards RA. 2008. The metagenomics RAST server—a public resource for the automatic phylogenetic and functional analysis of metagenomes. *BMC Bioinformatics* 9:386 DOI 10.1186/1471-2105-9-386.
- Mohagheghi A, Grohmann K, Himmel M, Leighton L, Updegraff DM. 1986. Isolation and characterization of acidothermus-cellulolyticus Gen-Nov, Sp-Nov, a new genus of thermophilic, acidophilic, cellulolytic bacteria. *International Journal of Systematic Bacteriology* 36:435–443 DOI 10.1099/00207713-36-3-435.
- Omeregio EO, Crumbliss LL, Bebout BM, Zehr JP. 2004. Determination of nitrogen-fixing phylotypes in Lyngbya sp. and Microcoleus chthonoplastes cyanobacterial mats from Guerrero Negro, Baja California, Mexico. *Applied and Environmental Microbiology* 70:2119–2128 DOI 10.1128/AEM.70.4.2119-2128.2004.
- Palla G, Barabasi AL, Vicsek T. 2007. Quantifying social group evolution. *Nature* 446:664–667 DOI 10.1038/nature05670.
- Partensky F, Blanchot J, Vaulot D. 1999. Differential distribution and ecology of *Prochlorococcus* and *Synechococcus* in oceanic waters: a review. *Bulletin-Institut Oceanographique Monaco-Numero Special* 19:457–476.
- Partensky F, Garczarek L. 2010. *Prochlorococcus*: advantages and limits of minimalism. *Annual Review of Marine Science* 2:305–331 DOI 10.1146/annurev-marine-120308-081034.
- Paul D, Bridges S, Burgess SC, Dandass Y, Lawrence ML. 2008. Genome sequence of the chemolithoautotrophic bacterium *Oligotropha carboxidovorans* OM5T. *Journal of Bacteriology* 190:5531–5532 DOI 10.1128/JB.00614-08.
- Peng J, Lu Z, Rui J, Lu Y. 2008. Dynamics of the methanogenic archaeal community during plant residue decomposition in an anoxic rice field soil. *Applied and Environmental Microbiology* 74:2894–2901 DOI 10.1128/AEM.00070-08.
- Pontarp M, Canback B, Tunlid A, Lundberg P. 2012. Phylogenetic analysis suggests that habitat filtering is structuring marine bacterial communities across the globe. *Microbial Ecology* 64:8–17 DOI 10.1007/s00248-011-0005-7.
- Reddy KJ, Haskell JB, Sherman DM, Sherman LA. 1993. Unicellular, aerobic nitrogen-fixing cyanobacteria of the genus *Cyanothece*. *Journal of Bacteriology* 175:1284–1292.
- Rice D, Mazur BJ, Haselkorn R. 1982. Isolation and physical mapping of nitrogen fixation genes from the cyanobacterium *Anabaena* 7120. *Journal of Biological Chemistry* 257:13157–13163.
- Rocap G, Larimer FW, Lamerdin J, Malfatti S, Chain P, Ahlgren NA, Arellano A, Coleman M, Hauser L, Hess WR, Johnson ZI, Land M, Lindell D, Post AF, Regala W, Shah M, Shaw SL, Steglich C, Sullivan MB, Ting CS, Tolonen A, Webb EA, Zinser ER, Chisholm SW. 2003. Genome divergence in two *Prochlorococcus* ecotypes reflects oceanic niche differentiation. *Nature* 424:1042–1047 DOI 10.1038/nature01947.

- Ruan Q, Dutta D, Schwalbach MS, Steele JA, Fuhrman JA, Sun F. 2006. Local similarity analysis reveals unique associations among marine bacterioplankton species and environmental factors. *Bioinformatics* 22:2532–2538 DOI 10.1093/bioinformatics/btl417.
- Saito R, Smoot ME, Ono K, Ruscheinski J, Wang PL, Lotia S, Pico AR, Bader GD, Ideker T. 2012. A travel guide to Cytoscape plugins. *Nature Methods* 9:1069–1076 DOI 10.1038/nmeth.2212.
- Sanudo-Wilhelmy SA, Kustka AB, Gobler CJ, Hutchins DA, Yang M, Lwiza K, Burns J, Capone DG, Raven JA, Carpenter EJ. 2001. Phosphorus limitation of nitrogen fixation by *Trichodesmium* in the central Atlantic Ocean. *Nature* 411:66–69 DOI 10.1038/35075041.
- Scanlan DJ, Ostrowski M, Mazard S, Dufresne A, Garczarek L, Hess WR, Post AF, Hagemann M, Paulsen I, Partensky F. 2009. Ecological genomics of marine picocyanobacteria. *Microbiology and Molecular Biology Reviews* 73:249–299 DOI 10.1128/MMBR.00035-08.
- Solonenko SA, Ignacio-Espinoza JC, Alberti A, Cruaud C, Hallam S, Konstantinidis K, Tyson G, Wincker P, Sullivan MB. 2013. Sequencing platform and library preparation choices impact viral metagenomes. *BMC Genomics* 14:320 DOI 10.1186/1471-2164-14-320.
- Steele JA, Countway PD, Xia L, Vigil PD, Beman JM, Kim DY, Chow C-EET, Sachdeva R, Jones AC, Schwalbach MS, Rose JM, Hewson I, Patel A, Sun FZ, Caron DA, Fuhrman JA. 2011. Marine bacterial, archaeal and protistan association networks reveal ecological linkages. *The ISME Journal* 5:1414–1425 DOI 10.1038/ismej.2011.24.
- Stegen JC, Lin X, Konopka AE, Fredrickson JK. 2012. Stochastic and deterministic assembly processes in subsurface microbial communities. *The ISME Journal* 6:1653–1664 DOI 10.1038/ismej.2012.22.
- Thompson CC, Silva GG, Vieira NM, Edwards R, Vicente AC, Thompson FL. 2013. Genomic taxonomy of the genus *Prochlorococcus*. *Microbial Ecology* 66:752–762 DOI 10.1007/s00248-013-0270-8.
- Tripp HJ. 2013. The unique metabolism of SAR11 aquatic bacteria. *Journal of Microbiology* 51:147–153 DOI 10.1007/s12275-013-2671-2.
- Ulrich W, Hajdamowicz I, Zalewski M, Stańska M, Ciurzycki W, Tykarski P. 2009. Species assortment or habitat filtering: a case study of spider communities on lake islands. *Ecological Research* 25:375–381 DOI 10.1007/s11284-009-0661-y.
- Willner D, Thurber RV, Rohwer F. 2009. Metagenomic signatures of 86 microbial and viral metagenomes. *Environmental Microbiology* 11(7):1752–1766 DOI 10.1111/j.1462-2920.2009.01901.x.
- Wu D, Raymond J, Wu M, Chatterji S, Ren Q, Graham JE, Bryant DA, Robb F, Colman A, Tallon LJ, Badger JH, Madupu R, Ward NL, Eisen JA. 2009. Complete genome sequence of the aerobic CO-oxidizing thermophile *Thermomicrobium roseum*. *PLoS ONE* 4:e4207 DOI 10.1371/journal.pone.0004207.
- Yachi S, Loreau M. 1999. Biodiversity and ecosystem productivity in a fluctuating environment: the insurance hypothesis. *Proceedings of the National Academy of Sciences of the United States of America* 96:1463–1468 DOI 10.1073/pnas.96.4.1463.
- Zubkov MV, Fuchs BM, Tarran GA, Burkill PH, Amann R. 2003. High rate of uptake of organic nitrogen compounds by *Prochlorococcus* cyanobacteria as a key to their dominance in oligotrophic oceanic waters. *Applied and Environmental Microbiology* 69:1299–1304 DOI 10.1128/AEM.69.2.1299-1304.2003.