




Genome Sequence of Human Papillomavirus Type 20, Strain HPV-20/Lancaster/2015

Kate V. Atkinson,^{a,*} Lisa A. Bishop,^{a,b} Glenn Rhodes,^c Nicolas Salez,^d
Neil R. McEwan,^{e,*} Matthew J. Hegarty,^e Julie Robey,^f Nicola Harding,^f
Simon Wetherell,^f Robert M. Lauder,^a Roger W. Pickup,^a Mark Wilkinson,^b
 Derek Gatherer^a

Division of Biomedical & Life Sciences, Faculty of Health & Medicine, Lancaster University, Lancaster, United Kingdom^a; Royal Lancaster Infirmary, Lancaster, United Kingdom^b; Centre for Ecology & Hydrology, Lake Ecosystems Group, Lancaster Environment Centre, Lancaster University, Lancaster, United Kingdom^c; UMR_D 190, Emergence des Pathologies Virales, Aix-Marseille University, Marseille, France^d; Institute of Biological, Environmental & Rural Sciences, Aberystwyth University, Aberystwyth, United Kingdom^e; Queen Square Medical Practice, Lancaster, United Kingdom^f

ABSTRACT The genome sequence of human papillomavirus type 20 (HPV-20; family *Papillomaviridae*, genus *Betapapillomavirus*, species *Betapapillomavirus 1*, type 20) was assembled by deep sequencing from nasopharyngeal swabs. The assembled genome is 0.37% divergent over its full length from the single complete genome of HPV-20 in GenBank (U31778). We named the strain HPV-20/Lancaster/2015.

The family *Papillomaviridae* consists of more than 120 viral types, divided into 49 genera (1). Within the genus *Betapapillomavirus*, type 20 of the species *Betapapillomavirus 1* (HPV-20) is represented in GenBank by a single complete genome (2), a circular double-stranded DNA of 7.8 kb (U31778), and three shorter fragments (3–5). HPV-20 is a clinically significant papillomavirus, having been implicated in epidermodysplasia verruciformis (2), skin cancer (6), ocular syringoma (7), nongenital seborrheic keratosis (8), conjunctival papillomas (9), verrucous carcinoma of the lip (10), condylomatous lesions of the mamillae (11), squamous cell carcinoma of the esophagus (12), and toenail onycholysis (13).

Volunteers were recruited from a general practice surgery and a general hospital in Lancaster, UK (54.05°N, 2.80°W). Nasopharyngeal swabs were taken between 16 December 2014 and 25 February 2015. Ethical approval was granted by the UK National Research Ethics Service, reference 14/LO/1634, NIHR Clinical Research Network (UKCRN) portfolio, ID 17799. All methods were carried out in accordance with the relevant guidelines and regulations.

Pooled RNA from 51 swabs was deep sequenced using an Illumina Nextera XT library and HiSeq 2500 system (SRA accession no. SRP092324). An HPV-20 genome was assembled using BWA 0.7.12-r1039 (14), with U31778 as the template. The assembled genome is 7,742 bases in length, differing from U31778 by 29 substitutions (0.37%). A 15-base deletion in the new genome starts at position 110, within an AT repeat that has a total length of 52 residues in U31778. Three regions cannot be resolved: (i) 3351 to 3353, (ii) 3704 to 3736; and (iii) 3798 to 3839. The second and third of these are in repetitive regions. We cannot exclude the possibility that the lengths of these repetitive regions differ between U31778 and the new genome, as we have shown for the region around the deletion at position 110. The predicted protein sequences are derived by reference to U31778 and differ at 14 amino acid residues (0.53%), without nonsense substitutions.

de Villiers et al. (15) recommend that a nucleotide divergence of 15% be used as the threshold for designation of a new type of human papillomavirus. The new strain is

Received 12 June 2017 Accepted 13 June 2017 Published 3 August 2017

Citation Atkinson KV, Bishop LA, Rhodes G, Salez N, McEwan NR, Hegarty MJ, Robey J, Harding N, Wetherell S, Lauder RM, Pickup RW, Wilkinson M, Gatherer D. 2017. Genome sequence of human papillomavirus type 20, strain HPV-20/Lancaster/2015. *Genome Announc* 5:e00712-17. <https://doi.org/10.1128/genomeA.00712-17>.

Copyright © 2017 Atkinson et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Derek Gatherer, d.gatherer@lancaster.ac.uk.

* Present address: Kate V. Atkinson, University College London Hospitals NHS Foundation Trust, London, United Kingdom; Neil R. McEwan, Robert Gordon University, Aberdeen, United Kingdom.

therefore well within the range of diversity expected within type 20 and has been designated HPV-20/Lancaster/2015. It is only the second full-length genome of HPV-20 to be described.

Forslund et al. (16) found HPV types of the *Betapapillomavirus 1* species in 17% of nasal swabs in a study of 312 Danish health care staff members, but HPV-20 was detected in only 6 individuals (1.9%). Of 9 deep-sequenced nonoverlapping subsets of individuals from our 51 swabs, we detected HPV-20 in all but 2. Our frequency is therefore at least 7/51 (>13%) and possibly much higher. The significance of this clinically important papillomavirus at such prevalence in our sample remains a matter for speculation.

BAM files are available from <https://doi.org/10.17635/lancaster/researchdata/145>.

Accession number(s). The genome sequence of HPV-20/Lancaster/2015 has been deposited in GenBank under the accession number [KY969593](https://doi.org/10.17635/lancaster/researchdata/145).

ACKNOWLEDGMENTS

K.V.A. received a Service Increment from Teaching (SIFT) studentship from University Hospitals of Morecambe Bay (UHMB) National Health Service (NHS) Foundation Trust, UK, and performed this work as part of the requirements for the degree of Master of Science (M.Sc.). Rosetrees Trust, UK, provided additional funding for deep sequencing (grant M395).

REFERENCES

- Harari A, Chen Z, Burk RD. 2014. Human papillomavirus genomics: past, present and future. *Curr Probl Dermatol* 45:1–18. <https://doi.org/10.1159/000355952>.
- Kremsdorf D, Favre M, Jablonska S, Obalek S, Rueda LA, Lutzner MA, Blanchet-Bardon C, Van Voorst Vader PC, Orth G. 1984. Molecular cloning and characterization of the genomes of nine newly recognized human papillomavirus types associated with epidermodysplasia verruciformis. *J Virol* 52:1013–1018.
- Chan SY, Delius H, Halpern AL, Bernard HU. 1995. Analysis of genomic sequences of 95 papillomavirus types: uniting typing, phylogeny, and taxonomy. *J Virol* 69:3074–3083.
- Berkhout RJ, Tieben LM, Smits HL, Bavinck JN, Vermeer BJ, ter Schegget J. 1995. Nested PCR approach for detection and typing of epidermodysplasia verruciformis-associated human papillomavirus types in cutaneous cancers from renal transplant recipients. *J Clin Microbiol* 33:690–695.
- Adachi A, Kiyono T, Hayashi Y, Ohashi M, Ishibashi M. 1996. Detection of human papillomavirus (HPV) type 47 DNA in malignant lesions from epidermodysplasia verruciformis by protocols for precise typing of related HPV DNAs. *J Clin Microbiol* 34:369–375.
- de Villiers EM. 1998. Human papillomavirus infections in skin cancers. *Biomed Pharmacother* 52:26–33.
- Assadoullina A, Bialasiewicz AA, de Villiers EM, Richard G. 2000. Detection of HPV-20, HPV-23, and HPV-DL332 in a solitary eyelid syringoma. *Am J Ophthalmol* 129:99–101. [https://doi.org/10.1016/S0002-9394\(99\)00292-5](https://doi.org/10.1016/S0002-9394(99)00292-5).
- Li YH, Chen G, Dong XP, Chen HD. 2004. Detection of epidermodysplasia verruciformis-associated human papillomavirus DNA in non-genital seborrheic keratosis. *Br J Dermatol* 151:1060–1065. <https://doi.org/10.1111/j.1365-2133.2004.06244.x>.
- Mlakar J, Kocjan BJ, Hošnjak L, Pižem J, Beltram M, Gale N, Drnovšek-Olup B, Poljak M. 2015. Morphological characteristics of conjunctival squamous papillomas in relation to human papillomavirus infection. *Br J Ophthalmol* 99:431–436. <https://doi.org/10.1136/bjophthalmol-2014-306087>.
- Mitsuishi T, Ohara K, Kawashima M, Kobayashi S, Kawana S. 2005. Prevalence of human papillomavirus DNA sequences in verrucous carcinoma of the lip: genomic and therapeutic approaches. *Cancer Lett* 222:139–143. <https://doi.org/10.1016/j.canlet.2004.09.019>.
- de Villiers EM, Sandstrom RE, zur Hausen H, Buck CE. 2005. Presence of papillomavirus sequences in condylomatous lesions of the mamillae and in invasive carcinoma of the breast. *Breast Cancer Res* 7:R1–R11. <https://doi.org/10.1186/bcr940>.
- Lavergne D, de Villiers EM. 1999. Papillomavirus in esophageal papillomas and carcinomas. *Int J Cancer* 80:681–684.
- Umanoff N, Werner B, Rady PL, Tyring S, Carlson JA. 2015. Persistent toenail onycholysis associated with Beta-papillomavirus infection of the nail bed. *Am J Dermatopathol* 37:329–333. <https://doi.org/10.1097/DAD.000000000000110>.
- Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* 26:589–595. <https://doi.org/10.1093/bioinformatics/btp698>.
- de Villiers EM, Fauquet C, Broker TR, Bernard HU, zur Hausen H. 2004. Classification of papillomaviruses. *Virology* 324:17–27. <https://doi.org/10.1016/j.virol.2004.03.033>.
- Forslund O, Johansson H, Madsen KG, Kofoed K. 2013. The nasal mucosa contains a large spectrum of human papillomavirus types from the *Betapapillomavirus* and *Gammapapillomavirus* genera. *J Infect Dis* 208:1335–1341. <https://doi.org/10.1093/infdis/jit326>.