

RESEARCH ARTICLE

Defects in the GINS complex increase the instability of repetitive sequences *via* a recombination-dependent mechanism

Malgorzata Jedrychowska, Milena Denkiewicz-Kruk, Malgorzata Alabrudzinska, Adrianna Skoneczna , Piotr Jonczyk, Michal Dmowski *, Iwona J. Fijalkowska *

Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Warsaw, Poland

* mdmowski@ibb.waw.pl (MD); iwonaf@ibb.waw.pl (IJF)



 OPEN ACCESS

Citation: Jedrychowska M, Denkiewicz-Kruk M, Alabrudzinska M, Skoneczna A, Jonczyk P, Dmowski M, et al. (2019) Defects in the GINS complex increase the instability of repetitive sequences *via* a recombination-dependent mechanism. *PLoS Genet* 15(12): e1008494. <https://doi.org/10.1371/journal.pgen.1008494>

Editor: Julian E. Sale, MRC Laboratory of Molecular Biology, UNITED KINGDOM

Received: August 2, 2019

Accepted: October 25, 2019

Published: December 9, 2019

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pgen.1008494>

Copyright: © 2019 Jedrychowska et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Abstract

Faithful replication and repair of DNA lesions ensure genome maintenance. During replication in eukaryotic cells, DNA is unwound by the CMG helicase complex, which is composed of three major components: the Cdc45 protein, Mcm2-7, and the GINS complex. The CMG in complex with DNA polymerase epsilon (CMG-E) participates in the establishment and progression of the replisome. Impaired functioning of the CMG-E was shown to induce genomic instability and promote the development of various diseases. Therefore, CMG-E components play important roles as caretakers of the genome. In *Saccharomyces cerevisiae*, the GINS complex is composed of the Psf1, Psf2, Psf3, and Sld5 essential subunits. The Psf1-1 mutant form fails to interact with Psf3, resulting in impaired replisome assembly and chromosome replication. Here, we show increased instability of repeat tracts (mononucleotide, dinucleotide, trinucleotide and longer) in yeast *psf1-1* mutants. To identify the mechanisms underlying this effect, we analyzed repeated sequence instability using derivatives of *psf1-1* strains lacking genes involved in translesion synthesis, recombination, or mismatch repair. Among these derivatives, deletion of *RAD52*, *RAD51*, *MMS2*, *POL32*, or *PIF1* significantly decreased DNA repeat instability. These results, together with the observed increased amounts of single-stranded DNA regions and Rfa1 foci suggest that recombinational mechanisms make important contributions to repeat tract instability in *psf1-1* cells. We propose that defective functioning of the CMG-E complex in *psf1-1* cells impairs the progression of DNA replication what increases the contribution of repair mechanisms such as template switch and break-induced replication. These processes require sequence homology search which in case of a repeated DNA tract may result in misalignment leading to its expansion or contraction.

Author summary

Processes that ensure genome stability are crucial for all organisms to avoid mutations and decrease the risk of diseases. The coordinated activity of mechanisms underlying the maintenance of high-fidelity DNA duplication and repair is critical to deal with the

Funding: This work was supported by the National Science Centre, Poland (www.ncn.gov.pl) grant no. 2017/26/M/NZ3/01044 to MD and grant no. TEAM/2011-8/1 from the Foundation for Polish Science (www.fnp.org.pl), co financed from European Union - Regional Development Fund „New players involved in the maintenance of genomic stability” to IJF. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

malfunction of replication forks or DNA damage. Repeated sequences in DNA are particularly prone to instability; these sequences undergo expansions or contractions, leading in humans to various neurological, neurodegenerative, and neuromuscular disorders. A mutant form of one of the noncatalytic subunits of active DNA helicase complex impairs DNA replication. Here, we show that this form also significantly increases the instability of mononucleotide, dinucleotide, trinucleotide and longer repeat tracts. Our results suggest that in cells that harbor a mutated variant of the helicase complex, continuation of DNA replication is facilitated by recombination processes, and this mechanism can be highly mutagenic during repair synthesis through repetitive regions, especially regions that form secondary structures. Our results indicate that proper functioning of the DNA helicase complex is crucial for maintenance of the stability of repeated DNA sequences, especially in the context of recently described disorders in which mutations or deregulation of the human homologs of genes encoding DNA helicase subunits were observed.

Introduction

Mechanisms by which organisms efficiently and faithfully control DNA stability are subjects of primary scientific interest. Mutagenesis produces genetic variations that drive the evolution of all species but at the same time may affect the lives of individual organisms, resulting in enhanced risk of carcinogenesis and other disorders [1–3]. The instability of repeated DNA sequences, also called satellite sequences, causes more than 30 disorders. Microsatellites and minisatellites are DNA motifs consisting of 1–9 or 10–100 base pairs, respectively, that are repeated from five times up to hundreds of times [4,5]. Such sequences are frequently found in genomes and are characterized by high variability. Dinucleotide repeats are the most abundant DNA repeats (48–67%) identified in many species [6,7], but in primates, mononucleotide repeats were identified as the most numerous class of simple DNA repeats [4,8]. DNA repeats influence chromatin organization, gene activity, and regulation of DNA metabolic processes. Alleles of genes carrying altered minisatellites have been correlated with a number of severe diseases, such as progressive myoclonus epilepsy [9], insulin-dependent diabetes mellitus [10], attention-deficit hyperactivity disorder [11], asthma [12], ulcerative colitis [13] and several cancer subtypes [14–16]. Expansions in trinucleotide repeats in humans can cause Huntington’s disease, myotonic dystrophy, spinocerebellar ataxia, and many other neurodegenerative disorders [17–19].

Mutation rates in DNA repeats are very high (10^{-2} – 10^{-6} events per locus per generation) compared with the rates of point mutations at average gene loci (10^{-9} – 10^{-10}) [2,20]. Molecular mechanisms of DNA repeat instability have been studied in many experimental systems, including bacteria, yeast, fruit flies, mice, and human cells [21]. Various mechanisms were shown to be involved in DNA repeat instability, i.e., formation of unusual DNA structures during DNA replication or slipped-strand mispairing [22–24], DNA recombination [25–27], DNA repair [28–34], and transcription [35,36]. Moreover, these mechanisms may interact with each other [37–41]. For example, DNA regions that are processed by DNA repair mechanisms and contain repeat tracts are subject to expansion/contraction by slip-strand mispairing errors upon strand invasion and formation of secondary structures during repair synthesis [42]. Other examples of factors increasing the instability of repetitive sequences are mutations in the yeast DNA polymerases or their decreased levels [43–48], mutations in genes encoding the Rad27 nuclease involved in Okazaki fragments processing, DNA ligase I, the PCNA polymerase processivity clamp or the Rfc1 subunit of the clamp loader [30,49].

In all organisms, DNA replication is carried out by the replisome, a multiprotein complex [50]. In eukaryotic cells, the highly efficient unwinding of double-stranded DNA is catalyzed by a helicase known as the CMG (Cdc45-Mcm2-7-GINS) complex. The CMG complex is a macromolecular assembly of 11 essential replication factors: the Cdc45 protein, four subunits of the GINS complex (Psf1, Psf2, Psf3, and Sld5) and the heterohexameric Mcm2-7 complex, which functions as the helicase motor. This complex translocates along the leading strand to separate the duplex DNA [51]. CMG participates in the establishment and progression of the replisome serving not only as a helicase but also as a platform for coordination of the activity of different components of the replisome [52–55]. Moreover, CMG has been shown to interact with the Pol ϵ complex, forming the CMG-E complex [56,57] and stimulating polymerase activity [58]. Experiments in fission yeast suggest not only structural but also functional interplay between the CMG helicase and Pol ϵ [59]. Recently, it was shown that functional impairment of the CMG-E complex induces genomic instability and promotes the development of genetic diseases [60].

Among many interactions within the CMG-E complex, it was shown that the Psf1 subunit of the GINS complex interacts not only with Psf3 and Sld5 but also with the essential subunit of Pol ϵ - Dpb2 [61,62]. It was also demonstrated that Cdc45 interacts with the GINS complex via the Psf1 and Psf2 subunits [63,64]. The *psf1-1* mutant, isolated in Hiroyuki Araki's laboratory, is temperature-sensitive and does not grow at 37°C. The Psf1-1 mutant subunit (R84G) exhibits an unchanged interaction with Dpb2, but the interaction of this subunit with Psf3 is severely impaired [54,62], which compromises the integrity of the GINS complex. However, there are no data regarding the mechanism by which this mutant form of Psf1 influences interactions with other components of the replisome and influences the physiology of the cell. We previously showed that in a *psf1-1* strain, destabilization of the GINS complex influences the level of mutagenesis, increasing the levels of both base substitutions and insertions/deletions (indels). As observed with many other replisome defects, a significant fraction of spontaneous mutagenesis in this mutant is due to increased participation of the error-prone DNA polymerase zeta (Pol ζ) [62].

In this work, we analyzed the effects of impaired interactions within GINS on the instability of DNA repeat tracts (mononucleotide, dinucleotide, trinucleotide, and others). All the DNA repetition types analyzed were much more unstable in the *psf1-1* mutant than in the wild-type strain. Our results demonstrate that in the *psf1-1* mutant, Pol ζ does not contribute to the instability of the tested repeat tracts; instead, homologous recombination (HR) is the major mechanism responsible for the observed instability. We also show increased formation of single-stranded DNA regions in the *psf1-1* cells, which is likely the result of impaired replication. To deal with this problem, recombination-associated repair synthesis may proceed, increasing the risk of instability of repetitive regions, especially those forming secondary structures. These findings highlight the importance of proper interactions within the GINS complex for the stability of repetitive sequences.

Results

As the replisome components are highly conserved from yeasts to humans in terms of structure, chemistry, and functionality, we used *Saccharomyces cerevisiae* as a model organism to investigate the influence of the defective functioning of the essential GINS complex on the stability of DNA repeat tracts. In our experiments, we used the Psf1-1 mutant form of a subunit of this complex, which exhibits impaired interaction with the Psf3 subunit. We examined the effects of the *psf1-1* allele on the stability of a variety of repeat tracts by employing two widely

used assays, namely, the chromosomal trinucleotide repeat expansion assay [65] and the plasmid-based frameshift assay [66,67].

The *psf1-1* allele causes increased instability of DNA repeat tracts

Trinucleotide repeats are a class of microsatellite sequences, expansions of which are responsible for more than 30 human developmental and neurological disorders [18,19,68]. Previously, it was shown that impaired functioning of the CMG-E complex induces genomic instability and promotes the development of various diseases [60,69,70], demonstrating that CMG-E components have important roles as caretakers of the genome [71]. Therefore, we sought to investigate the impact of the defective functioning of the GINS complex on the stability of trinucleotide repeats using the chromosomal assay developed in the laboratory of R. Lahue [65]. In this assay, the promoter of the *URA3* reporter gene contains 25 repeats of tested trinucleotides located between the TATA box and the transcription start site. These repeats do not interfere with the expression of *URA3* and therefore yield sensitivity to the toxic drug 5-fluoroorotic acid (5-FOA). Expansion of at least five trinucleotides (>29 repeats) results in the production of a long transcript encompassing an out-of-frame ATG and, as a consequence, in translational incompetence which can be selected on medium containing 5-FOA which is toxic to cells producing Ura3 protein [72]. Appropriate sequences containing trinucleotide repeats (CTG, GAA or TTC) or the control sequence (75 random nucleotides) in the promoter of *URA3* were introduced into the *psf1-1* mutant strain and control wild-type cells. Wild-type cells with the “scrambled” (C, A, T, G) control sequence yield 5-FOA^R colonies at a rate of $<0.54 \times 10^{-8}$, while the *psf1-1* mutation results in increased rates of forward mutagenesis in *URA3* (19×10^{-8}) (Fig 1). Wild-type cells containing (GAA)₂₅, (TTC)₂₅ and (CTG)₂₅ tracts yield 5-FOA^R colonies at a rate of $<0.44 \times 10^{-8}$, $<0.47 \times 10^{-8}$ and 37×10^{-8} , respectively, which confirms that (CTG)₂₅ repeats are more unstable than (GAA)₂₅ and (TTC)₂₅ tracts (Fig 1) [73]. When (GAA)₂₅, (TTC)₂₅ and (CTG)₂₅ expansions were measured in the *psf1-1* mutant, the mutation rates were two orders of magnitude higher than those observed for the wild-type strain (114×10^{-8} , 132×10^{-8} , and 5795×10^{-8} , respectively) (Fig 1). Frequent expansions of tested trinucleotide tracts in the *psf1-1* strain highlight the importance of proper interactions within the GINS complex for the stability of repeated trinucleotides.

To further investigate the level and mechanisms of DNA tract instability in the *psf1-1* strains, we employed the widely used and well-documented plasmid system developed in T. Petes' laboratory based on the reporter gene *URA3* with in-frame insertions of various minor microsatellite sequences [66,67]. The stability of the repeat tracts depends on their length and nucleotide composition. Therefore, we tested (G)₁₈ (pMD28), (GT)₄₉ (p99GT), (AACGC AATGCG)₄ (pMD41) and (CAACGCAATGCGTTGGATCT)₃ (pEAS20) tracts in the *psf1-1* strain and, as a control, in *PSF1* cells. As an additional control, we designed the pKK2 plasmid carrying a random sequence that was also inserted in-frame and devoid of any repeats (see Materials and Methods). The alterations within the tracts leading to out-of-frame insertions or deletions can be selected on medium containing 5-FOA [72]. All the repeat tracts, i.e., (G)₁₈, (GT)₄₉, (AACGCAATGCG)₄ and (CAACGCAATGCGTTGGATCT)₃, were roughly two- to four-fold less stable in the *psf1-1* strain than in the wild-type strain (Fig 2).

To confirm that the 5-FOA resistance of wild-type and *psf1-1* cells with the tested plasmids resulted from altered repeat tracts, we characterized independent isolates derived from each strain. Using capillary electrophoresis, we analyzed the lengths of the PCR products encompassing the repeat tracts from 5-FOA-resistant mutants (Table 1).

As expected, in both the wild-type and *psf1-1* mutant strains harboring the control pKK2 plasmid, we did not observe changes in the length of the random sequence. In these mutants,




Fig 1. The *psf1-1* allele enhances chromosomal trinucleotide repeat instability in yeast. The 5-FOA^R mutation rates were determined at 23°C for yeast cells carrying the analyzed sequences; the values are medians with 95% confidence intervals calculated from data for at least ten cultures of each strain; the *p*-values for *psf1-1* mutants versus wild-type strains were calculated using the nonparametric Mann-Whitney U statistical test (**** *p* ≤ 0.0001; *** *p* ≤ 0.001). All associated *p*-values are presented in the S1 Table.

<https://doi.org/10.1371/journal.pgen.1008494.g001>




Fig 2. The *psf1-1* allele increases the instability of various repeat tracts. The 5-FOA^R mutation rates were determined at 23°C for yeast cells with the indicated genotypes carrying plasmids with analyzed repetitive sequences; the values are medians with 95% confidence intervals calculated from data for at least ten cultures of each strain; the *p*-values for *psf1-1* mutants versus wild-type strains were calculated using the nonparametric Mann-Whitney U statistical test (*** *p* ≤ 0.001). All associated *p*-values are presented in the S1 Table.

<https://doi.org/10.1371/journal.pgen.1008494.g002>

Table 1. Types of alterations within different repeat tracts in the *psf1-1* mutant and wild-type strains carrying plasmids with repeat sequences.

Tract sequence	Relevant genotype	Classes of tract alterations [%] ^a						Number of analyzed clones
		Large deletions	-2	-1	0	+1	+2	
random sequence	<i>PSF1</i>				100			66
	<i>psf1-1</i>				100			96
(GT) ₄₉	<i>PSF1</i>	28		2		66	4	85
	<i>psf1-1</i>	47		1	1	44	7	75
(AACGCAATGCG) ₄	<i>PSF1</i>		81	8	9		2	52
	<i>psf1-1</i>		85	2	11		2	47
(CAACGCAATGCGTTGGATCT) ₃	<i>PSF1</i>		46	46		8		63
	<i>psf1-1</i>		11	81	6	2		54

a. The numbers in the column headings are the number of repeat units added (+) or deleted (-).

<https://doi.org/10.1371/journal.pgen.1008494.t001>

we observed only changes in the *URA3* coding sequence, confirming that pKK2 is an appropriate control for our experiments (Table 1). Capillary analysis indicated that the alterations in the lengths of the poly(GT) tracts mainly involved insertions of one repeat in both the wild-type (66%) and *psf1-1* (44%) strains. The second important class of alterations was the deletion of more than two repeats in both the wild-type (28%) and *psf1-1* (47%) strains. In the plasmids containing (AACGCAATGCG)₄ repeats, the most common alterations were two-tract deletions (81% in the wild-type strain and 85% in the *psf1-1* strain). For minisatellites with long repeat units (CAACGCAATGCGTTGGATCT)₃, the most frequent changes were those involving one or two-repeat deletion (92% for both strains). These types of alterations are consistent with previously published results from T. Petes' laboratory [66,67] and the predicted structures that can be formed on the mother or daughter strand to produce contractions or expansions, respectively (S1 Fig). Alterations in mononucleotide microsatellites cannot be analyzed using capillary electrophoresis because these alterations are mainly deletions or additions of one repeat, which are beyond the sensitivity range of this method. In summary, the *psf1-1* allele causes an increase in repeat tract instability compared to that seen in the wild-type strain. To further analyze the mechanism underlying repeat tract instability in the *psf1-1* strains, we decided to use a plasmid-based system, which provides the opportunity to analyze the stability of various repeat tracts.

MMR repairs DNA loops which are at the origin of microsatellite sequences instability in *psf1-1* cells

What mechanisms are responsible for the instability observed in the *psf1-1* strains? The most common mechanism that explains tract instability is DNA/polymerase slippage [24,27,74]. Changes in microsatellite sequences resulting from rearrangement of the template and primer can be recognized and repaired by the mismatch repair (MMR) system [75]. In yeast, there are two MMR complexes composed of Msh2, Msh3, and Msh6, which are homologs of the prokaryotic MutS protein [76,77]. The Msh2-Msh6 and Msh2-Msh3 complexes are also called MutS α and MutS β , respectively. Inactivation of Msh6 decreases the stability of repeat tracts with repeat units of 1 or 2 bp, while inactivation of Msh2 or Msh3 decreases the stability of repeat tracts with repeat units ranging in length from 1 to 14 bp [67,78,79]. None of the complexes are capable of efficiently repairing 20-base loops [67,80,81]. Mutations in genes affecting MMR dramatically reduce microsatellite stability [82–87].

To determine whether the MMR mechanism corrects the alterations in repeated sequences in the *psf1-1* mutant, we analyzed the rate of mutagenesis in derivatives carrying the *psf1-1*




Fig 3. Effect of mismatch repair on repeat tract stability in *psf1-1* cells (A-D). The 5-FOA^R mutation rates were determined at 23°C for yeast cells with the indicated genotypes carrying plasmids with the analyzed sequences; the values are medians with 95% confidence intervals calculated from data for at least ten cultures of each strain; the *p*-values for *psf1-1 msh2Δ* versus *psf1-1* mutant strains were calculated using the nonparametric Mann-Whitney U statistical test (**** *p* ≤ 0.0001; ** *p* ≤ 0.01; ns-*p* > 0.5). All associated *p*-values are presented in the S1 Table.

<https://doi.org/10.1371/journal.pgen.1008494.g003>

allele and deletion of the *MSH2* gene (Fig 3). The *psf1-1* and *msh2Δ* single mutants carrying the random sequence on the control plasmid pKK2 show 4-fold and 15-fold mutator effects on the *URA3* coding sequence, respectively. Synergy in the mutator effect (54-fold increase) is observed in the double mutant *psf1-1 msh2Δ*/pKK2, which indicates that *psf1-1* significantly increases the number of replication errors recognized and repaired by the MMR system (Fig 3A). The *psf1-1* strain carrying a plasmid with a (GT)₄₉ tract shows a 4-fold elevated rate of microsatellite instability compared to that in the wild-type strain. The *msh2Δ* strain harboring the same plasmid elevates the level of microsatellite instability by 38-fold. The double mutant *psf1-1 msh2Δ* shows a 66-fold increase in instability. This destabilizing effect in the *psf1-1 msh2Δ* double mutant is stronger than an additive effect would have been (42-fold), suggesting a synergistic interaction of the two mutations (Fig 3B). Absence of the function of the MMR system has a moderate effect on (AACGCAATGCG)₄ tracts and no effect on relatively long tracts (CAACGCAATGCGTTGGATCT)₃ (Fig 3C and 3D), as expected from the specificity of the MMR mechanism. In the wild-type background, deletion of *MSH2* does not significantly affect the stability of these repeat tracts (Fig 3C and 3D). Together, these results indicate that Msh2 has an impact on microsatellite repair but is not involved in the repair of the tested minisatellite tracts in both wild-type and *psf1-1* strains.

The elevated rate of repeat tract instability in the *psf1-1* strain is not dependent on the error-prone DNA polymerase ζ

We previously showed that approximately 50% of spontaneous mutagenesis in *psf1-1* cells is associated with the participation of Pol ζ in DNA synthesis [62]. Pol ζ is composed of four subunits, namely, Rev3 (the catalytic subunit), Rev7, Pol31, and Pol32 [88,89]. Therefore, to test




Fig 4. Pol ζ has no effect on repeat tract stability in *psf1-1* (A-D). The 5-FOA^R mutation rates were determined at 23°C for yeast cells with the indicated genotypes carrying plasmids with the analyzed sequences; the values are medians with 95% confidence intervals calculated from data for at least ten cultures of each strain; the *p* values for *psf1-1 rev3Δ* versus *psf1-1* mutant strains were calculated using the nonparametric Mann-Whitney U statistical test (***) $p < 0.001$; ns- $p > 0.5$). All associated *p*-values are presented in the S1 Table.

<https://doi.org/10.1371/journal.pgen.1008494.g004>

whether Pol ζ is also involved in the instability of repeated sequences in the *psf1-1* strain, we performed measurements in the *rev3Δ* background. The level of forward *URA3* mutations in strains carrying a plasmid with the random sequence decreased by 45% in the *psf1-1 rev3Δ* strain (Fig 4A). This suggests that destabilized interactions within the GINS complex in the *psf1-1* mutant lead to increased participation of Pol ζ in DNA synthesis. In both *PSF1* and *psf1-1* strains possessing plasmids with repeat tracts, we observed no effect of Pol ζ deficiency on the rates of appearance of 5-FOA^R colonies (Fig 4B, 4C and 4D). This result suggests that Pol ζ does not participate in the mechanisms underlying the destabilization of repetitive sequences in both wild-type and *psf1-1* cells.

HR is engaged in repeat tract instability in the *psf1-1* strain

Many DNA repair mechanisms are based on HR. This is a highly conserved mechanism for the repair of DNA double-strand breaks (DSBs) and recovery of stalled or collapsed replication forks [90–92]. However, although recombination is thought to be error free, this mechanism is also potentially mutagenic, e.g., recombination can promote the instability of repeated DNA sequences [27] or cause genome rearrangements in the cells under constant replication stress [93].

The recombinases Rad52 and Rad51 are two major players in HR [94,95]. The most important function of Rad52 is the promotion of Rad51 filament formation on RPA-coated ssDNA. To investigate the role of Rad52 in the instability of repetitive sequences in the *psf1-1* strain, we combined *rad52Δ* with the *psf1-1* allele. Both alleles exhibit a mutator effect on the overall mutation rate at *URA3*. Strain *psf1-1* with the random control sequence shows a 3.8-fold

increase in *URA3* mutagenesis, while the *rad52Δ* allele exhibits a 9.6-fold mutator effect (Fig 5A). In the double mutant (*psf1-1 rad52Δ*), forward *URA3* mutagenesis increased 14.2-fold showing an additive effect (Fig 5A).




Fig 5. The instability of repeat tracts in *psf1-1* cells depends on the recombinases Rad2 (A-D) and Rad51 (E-H). The 5-FOA^R mutation rates were determined at 23°C for yeast cells with the indicated genotypes carrying plasmids with the analyzed sequences; the values are medians with 95% confidence intervals calculated from data for at least ten cultures of each strain; the *p* values for *psf1-1 rad52Δ* or *psf1-1 rad51Δ* versus *psf1-1* mutant strains were calculated using the nonparametric Mann-Whitney U statistical test (**** *p* < 0.0001; *** *p* < 0.001). All associated *p*-values are presented in the S1 Table.

<https://doi.org/10.1371/journal.pgen.1008494.g005>

Deletion of *RAD52* reduces the instability of (GT)₄₉ in the wild-type background by 70% (Fig 5B). The absence of *RAD52* also strongly reduces the increased instability of the (GT)₄₉ tract by 75%, in the *psf1-1* mutant (Fig 5B). For the (AACGCAATGCG)₄ and (CAACGCAATGCGTTGATCT)₃ tracts, deletion of *RAD52* in the wild-type background reduces the instability by 60% and 35%, respectively (Fig 5C and 5D). In the double *psf1-1 rad52Δ* mutant strain, the instability of these tracts is reduced by 82% and 66%, respectively, compared to that in the *psf1-1* strain (Fig 5C and 5D). These data indicate that Rad52-dependent recombination events are closely associated with the instability of the tested repeat tracts in the *psf1-1* background.

The central and highly conserved step in HR is the formation of Rad51 nucleoprotein filaments on the single-stranded DNA ends. Among its many functions in DNA replication and repair, Rad51 is crucial for the homology search and strand invasion steps [96–99]. Therefore, loss of this important player in most of the pathways of HR could be expected to affect DNA repeat tract instability in the *psf1-1* strain. In the wild-type strain with the random control sequence, deletion of *RAD51* results in an almost 8-fold increase in *URA3* mutagenesis (Fig 5E). The *psf1-1* and *rad51Δ* mutations together lead to a further additive increase in forward mutagenesis compared to the single mutants (Fig 5E). Based on the observed additive effect of *psf1-1* mutation with *RAD52* or *RAD51* deletion we conclude that recombination has no significant contribution to forward mutagenesis observed in *psf1-1* cells. In the wild-type strain, deletion of *RAD51* reduces the instability of the repeat tracts (GT)₄₉, (AACGCAATGCG)₄ and (CAACGCAATGCGTTGGATCT)₃ by 79%, 47% and 33%, respectively (Fig 5F–5H). Inactivation of *RAD51* in the *psf1-1* background reduces the instability of these tracts by 77%, 67%, and 55%, respectively, compared to that in the *psf1-1* strain (Fig 5F–5H). These results confirm that the instability of repeated sequences in *psf1-1* cells is highly dependent on recombination.

When HR is activated, the requirement for Rad51 nucleoprotein filaments increases, so the Rad51 level is elevated. Since in *psf1-1* cells, the stability of the repetitive sequences depends on HR, we asked whether the Rad51 level is elevated in this mutant. Indeed, as shown in Fig 6A and 6B, the Rad51 protein level is 1.5-fold higher in the *psf1-1* mutant than in the wild-type strain.

Accumulation of ssDNA in *psf1-1* cells

HR substrates are DSBs or ssDNA stretches that can be formed during impaired DNA replication, especially in cells with defective replisomes. The observed recombination-dependent increased instability of DNA repeat tracts in *psf1-1* cells suggested that ssDNA regions accumulate in this mutant. To verify this hypothesis, we embedded yeast chromosomes in agarose plugs and treated the plugs with S1 nuclease, which cleaves dsDNA at single-stranded regions such as nicks, gaps, or loops. Next, we visualized the integrity of the chromosomes by separating them using pulsed-field gel electrophoresis (PFGE). In addition to the wild-type and *psf1-1* strains, we also used the *pol32Δ* strain, which accumulates ssDNA gaps [100], as a control. In the *psf1-1* and *pol32Δ* strains, we observed strong degradation of chromosomes after S1 nuclease treatment compared to that in the wild-type strain (Fig 6C). This result is consistent with the idea that ssDNA stretches accumulate in *psf1-1* cells. Single-stranded DNA is coated by Replication Protein A (RPA). Therefore, to estimate the degree of ssDNA formation in *psf1-1* cells, we analyzed the formation of RPA-bound ssDNA through visualization of its subunit Rfa1 fused with YFP. We observed an increased number and higher intensity of Rfa1-YFP foci in *psf1-1* cells compared to that in wild-type cells (Fig 6D and 6E).

Mms2 influences the stability of repeat tracts in *psf1-1* cells

The recombination proteins Rad52 and Rad51 are also involved in template switching (TS), a recombination-related pathway that enables survival of fork stalling, DNA lesions, or




Fig 6. Accumulation of single-stranded DNA and activation of the HR pathway in *psf1-1* cells. (A) Rad51 protein levels in the wild-type and *psf1-1* strains were analyzed by western blotting with an anti-Rad51 antibody. (B) Quantification of the western blotting results showing the mean \pm s.d. of 8 to 12 repetitions of the assay. Strain BY4741 *rad51Δ* served as a negative control. Statistical significance was determined with Student's t-test (p -value $^* \leq 0.05$) (C) Pulsed-field gel electrophoresis (PFGE) analysis of chromosome integrity after S1 nuclease treatment of DNA from yeast cells cultured at the permissive temperature 23°C. Agarose plugs with DNA were treated with two units (2 U) or five units (5 U) of S1 nuclease for 30 min. Untreated plugs are shown as a control (NT). Strain *pol32Δ* served as a positive control. (D) Rfa1-YFP foci detected in wild-type and *psf1-1* strains. (E) Rfa1 foci frequency quantification. Three biological replicates were performed, each with at least 200 cells counted. The results represent the number of cells with indicated number of foci with SD. For statistical analysis contingency table and the chi-square test were used (S2 Table). The chi-square statistic is 223.4038, which corresponds to the p -value $^{****} < 0.0001$.

<https://doi.org/10.1371/journal.pgen.1008494.g006>




Fig 7. Template switch mechanism contributes to the instability of repeat tracts in *psf1-1* cells (A-D). The 5-FOA^R mutation rates were determined at 23°C for yeast cells with the indicated genotypes carrying plasmids with the analyzed sequences; the values are medians with 95% confidence intervals calculated from data for at least ten cultures of each strain; the *p* values for *psf1-1 mms2Δ* versus *psf1-1* strains were calculated using the nonparametric Mann-Whitney U statistical test (* *p* ≤ 0.05; *** *p* ≤ 0.0001). All associated *p*-values are presented in the S1 Table.

<https://doi.org/10.1371/journal.pgen.1008494.g007>

impediments [93,101,102]. This process uses the newly synthesized homologous daughter strand as a template to enable the continuation of DNA replication. TS requires further ubiquitylation of the monoubiquitinated proliferating cell nuclear antigen (PCNA) in a process driven by the Ubc13–Mms2–Rad5 E2–E3 polyubiquitylating complex [103,104]. Inactivation of Mms2 has an effect on forward *URA3* mutagenesis in both wild type and *psf1-1* cells (2.5-fold and 1.6-fold increase, respectively) (Fig 7A). The additive effect of *psf1-1* mutation and *MMS2* deletion demonstrate that TS does not contribute to the enhanced mutagenesis in *psf1-1* cells. *MMS2* deletion has no effect on the instability of repeat tracts in the wild-type strain (Fig 7B–7D). However, in *psf1-1* cells, inactivation of *MMS2* significantly reduces the instability of (GT)₄₉, (AACGCAATGCG)₄ and (CAACGCAATGCGTTGGATCT)₃ tracts by 47%, 39%, and 45%, respectively. This finding indicates that TS may be involved in the instability of repetitive sequences in *psf1-1* cells.

Break-induced replication is involved in the instability of repeated sequences in *psf1-1* cells

Another subpathway of HR is break-induced replication (BIR) [105]. This process, when involved in the synthesis of repeated DNA sequences leads to an elevated risk of genetic instability [106]. Therefore, we decided to test whether BIR may be involved in the instability of repeated sequences in the *psf1-1* mutant by deleting genes encoding proteins involved in this process. Pol32, is involved in BIR not only as a nonessential subunit of Pol δ, but also contributes to the establishment of a repair replication fork, i.e., strand displacement during bubble migration [107–110]. Therefore it is specifically involved in BIR but not in other HR-related mechanisms. The Pol32 subunit is not essential for normal DNA replication; however, deletion

of *POL32* causes cold sensitivity at 13°C [111]. The *psf1-1* strains also exhibit temperature sensitivity and do not grow at 37°C. We attempted to construct *psf1-1 pol32Δ* strains and obtained double mutants viable at 28°C that exhibit growth impairment at 15°C and 37°C. Deletion of *POL32* has no significant effect on forward mutagenesis in *URA3* in the wild-type strain (Fig 8A). Pol32 is one of two subunits shared by two DNA polymerases: Pol ζ and Pol δ. Because Pol ζ participates in mutagenesis in both wild-type and *psf1-1* strains, one can expect that deletion of *POL32* in these strains will reduce forward mutagenesis. However, similar to previously published data, our results show that this is not the case [112,113]. The lack of expected mutagenesis decrease is probably associated with the fact that Pol32 not only plays a role as a Pol ζ subunit but also participates in Pol δ activities. Therefore, in the *pol32Δ* strain, the antimutator effect on Pol ζ-dependent mutagenesis is compensated by a moderate mutator effect associated with the involvement of Pol32 in the reactions of Pol δ. For this reason, the effect of *pol32Δ* is moderate in the *psf1-1* background (Fig 8A).

Deletion of *POL32* reduces the instability of (GT)₄₉ tracts both in wild-type and *psf1-1* cells by 84% and 92%, respectively (Fig 8B). For the (AACGCAATGCG)₄ and (CAACGCAATGCGTTGGATCT)₃ tracts, inactivation of *POL32* had no effect on tract instability in wild-type cells (Fig 8C and 8D) but reduces the instability in the *psf1-1* background by 29% and 62%, respectively (Fig 8C and 8D). These results suggest that the mechanisms responsible for the instability are not the same for the microsatellite (GT)₄₉ tracts and minisatellite (AACGCAATGCG)₄ and (CAACGCAATGCGTTGGATCT)₃ tracts.

To extend our investigation of the involvement of BIR in repeated sequence instability, we constructed *pif1Δ* strain derivatives. *PIF1* encodes a DNA helicase involved in various DNA transactions, including BIR, or maturation of Okazaki fragments [108,114–116]. Deletion of *PIF1* has a significant effect on forward mutagenesis in wild-type cells (6.4-fold increase) probably due to inactivation of mitochondrial activity (loss of mtDNA, rho⁰/rho⁻) [117–119] leading to increased Pol ζ-dependent nuclear mutagenesis [120]. The observed similar mutagenesis levels in *pif1Δ* and *pif1Δ psf1-1* cells (Fig 8E) can be explained by the observation that participation of Pol ζ in DNA replication was already increased in the GINS mutant (Fig 4A). We also observed that inactivation of *PIF1* has no effect on the stability of the tested repeat tracts compared to that in the wild-type strain (Fig 8F–8H). However, analysis of the genetic interactions between *psf1-1* and *pif1Δ* shows that a lack of functional Pif1 helicase decreases the instability of the (GT)₄₉, (AACGCAATGCG)₄ and (CAACGCAATGCGTTGGATCT)₃ repeated sequences by 58%, 29%, and 53%, respectively. Together, the results obtained for the *pol32* or *pif1* deletion mutants indicate that BIR contributes to DNA repeat instability in the *psf1-1* strain.

Discussion

In this work, we investigated the impact of the proper functioning of GINS on the stability of repeated DNA sequences. GINS is a component of one of the crucial leading-strand replication elements of the replisome, i.e., the CMG-E complex of DNA helicase (Cdc45, Mcm2-7, GINS) with Pol ε [57]. Although the structural dynamics of this large complex have been extensively analyzed [121,122], the physiological role of the individual subunits is poorly understood, especially their role in the maintenance of genomic stability. In addition to its essential role in DNA unwinding, CMG is a platform that functionally and structurally coordinates the participation of various replisome elements in DNA replication. Previously, we showed that the *Psf1-1* mutant form of the GINS subunit increases the levels of both base substitutions and indels [62]. Here, we show that the *psf1-1* allele increases the instability of repeat tracts located on plasmids and various chromosomally encoded trinucleotide repeats. Of particular interest is




Fig 8. Break-induced replication contributes to the instability of repeat tracts in *psf1-1* cells. (A-D) Impact of *POL32* disruption. (E-H) Impact of *PIF1* disruption. The 5-FOA^R mutation rates were determined at 28°C for yeast cells with the indicated genotypes carrying plasmids with the analyzed sequences; the values are medians with 95% confidence intervals calculated from data for at least ten cultures of each strain; the *p* values for *psf1-1 pol32Δ* or *psf1-1 pif1Δ* versus *psf1-1* mutants versus wild-type strains were calculated using the nonparametric Mann-Whitney U statistical test (**** *p* ≤ 0.0001; *** *p* ≤ 0.001; ** *p* ≤ 0.01; * *p* ≤ 0.05). All associated *p*-values are presented in the S1 Table.

<https://doi.org/10.1371/journal.pgen.1008494.g008>

the observed Psf1-1-dependent instability of GAA and TTC repeats, expansions of which are involved in Friedreich ataxia in humans [123]. In the wild-type background, expansions of GAA and TTC are not frequent unless the number of repeats exceeds a length threshold (78 triplets) [124]. Additionally, the repeated CTG sequences, the instability of which is the cause of human disorders such as Huntington disease, myotonic dystrophy and spinocerebellar ataxia, also exhibit significantly increased instability in *psf1-1* cells (Fig 1). It was demonstrated that trinucleotide repeats can form structures that result in stalling of DNA synthesis *in vitro* [125–127] and/or impede replication fork progression *in vivo* [128–131]. Based on the results for *psf1-1*, we conclude that functional impairment of CMG-E can enhance these destabilizing effects.

Why is proper functioning of a noncatalytic subunit of the GINS complex so significant for the stability of repetitive sequences? Answering this important basic research question will provide a better understanding of the molecular processes that lead to the DNA repeat instability that has been observed in many human diseases. As the Psf1-1 protein exhibits a greatly impaired interaction with the Psf3 subunit of the GINS complex [54,62], this mutant may influence the integrity of the GINS complex, affecting communication with other components of CMG, Pol ϵ and/or DNA.

What might be the mechanisms involved in DNA repeat instability in *psf1-1* cells? Although deletion of the *REV3* gene, which inactivates Pol ζ , reduces by 40–50% the rate of mutagenesis in the *CANI* [62] and *URA3* reporter gene (Fig 4A), inactivation of Pol ζ does not affect the stability of repeat tract sequences (Fig 4B–4D). This result is consistent with previous studies showing that defects in trans-lesion synthesis polymerases do not influence repeat instability in wild-type budding yeast [44,132,133]. However, in cells with impeded replicative polymerases, some repeat expansions do occur *via* a Pol ζ -dependent mechanism [44], as do short duplications initiated by small hairpins [134]. Nevertheless, our results indicate that mutation in a subunit of GINS, a replisome component, results in Pol ζ -independent repeat instability.

Previous studies on mechanisms of microsatellite instability as well as analyses of mutations generated *via in vitro* replication assays demonstrated that changes in the number of repeat tracts often result from several mechanisms, e.g., formation of non-B DNA structures, strand slippage, reduced processivity or dissociation of DNA polymerase [67,74,135–138]. Some errors, such as base substitutions or small DNA loops, that arise during replication may be the target for the MMR system [127,139]. We showed that MMR corrects a majority of the spontaneous errors produced in *psf1-1* strains (Fig 3A), [62]. The statistically significant increase in polyGT instability in the *psf1-1 msh2 Δ* mutant compared to that in the single mutants indicates that the *psf1-1* allele enhances the frequency of MMR-corrected microsatellite instability. However, inactivation of *MSH2* has no effect on the instability of long minisatellite sequences in *psf1-1* cells, which is consistent with the specificity of loop-size recognition by the MMR system.

Another mechanism that is frequently involved in the instability of repeated DNA sequences is DNA recombination—for a review, see [27]. This process is involved in the reactivation of stalled or collapsed replication forks and in the repair of DNA gaps left behind the replication fork which might arise in the *psf1-1* cells as suggested by accumulation of increased number of single-stranded DNA regions and Rfa1 foci (Fig 6C–6E) shown in this work. HR, which is generally considered to be an error-free process, may be an important source of genomic instability due to rearrangements between the invading and homologous strands with repetitive regions, formation of hairpins during D-loop extension, and difficulties associated with synthesis across repetitive regions, especially those that form DNA secondary structures [27,140–142]. The most important recombination proteins are Rad52 and Rad51 [96,98,143]. Deletion of *RAD52* or *RAD51* reduced the instability of repeat tracts compared to the *psf1-1*

mutant (Fig 5B–5D) what supports the conclusion that HR mechanisms are involved in this process (Fig 5F–5H). An additional fact reinforcing this conclusion is the elevated level of Rad51 in *psf1-1* cells (Fig 6A and 6B). It seems that an increased level of Rad51 is the supportive mechanism for cells during replication slow-down or blockage. A similar effect was documented for cells lacking the transcription factor Swi6, which controls the expression of G1/S transition genes, resulting in limitation of proteins involved in DNA replication and repair and a prolonged cell cycle. Their survival is enhanced by Rad51-dependent illegitimate recombination [93]. Similar phenotypes were also described for human cells, in which mutation in the KRAS proto-oncogene caused replication fork stalling, DNA lesion accumulation and increased abundance of various proteins, including Rad51, which is obligatory for the survival of these mutant cells [144].

To further analyze the recombination-dependent mechanisms involved in *psf1-1*-mediated instability of repeat tracts, we inactivated the template switch mechanism, which depends on the activity of the Ubc13-Mms2-Rad5 complex [102–104,145,146]. The observed decrease in instability for all the tested repeated sequences in the *psf1-1 mms2Δ* strains (Fig 7B–7D) indicates that the Mms2-dependent template switch mechanism may be involved in the instability of repeat tracts in the *psf1-1* mutant.

Another possible mechanism that enables the continuation of DNA synthesis after replication fork stalling or collapse is BIR, in which the main proteins involved are Pol δ with its Pol32 subunit as well as the Pif1 helicase [108,114,116]. The involvement of BIR in the generation of large-scale expansions was shown previously [106,147]. Here, we show that deletion of *POL32* or *PIF1* significantly decreases the high instability of the tested tracts in the *psf1-1* background (Fig 8B–8D and 8F–8H) suggesting that BIR may be one of the mechanisms underlying the formation of repeated sequence instability in the *psf1-1* strains.

van Pel and colleagues [148] showed that the *psf1-1* mutant exhibits a dramatic increase in Rad52 foci frequency and a two-fold increase in the number of G2/M arrest large-budded cells. These data together with previously showed retarded S-phase progression [54], dumb-bell-cell formation and impaired interaction within the GINS complex and genomic instability [62], allow us to conclude that *Psf1-1* significantly impairs the replication process. It should be stressed here that the observed *psf1-1* phenotypes may result from a combination of processes activated in response to defective DNA replication and these mechanisms may be envisioned as sources of increased instability of repetitive DNA tracts in the *psf1-1* strains.

First, impaired replication and accumulation of ssDNA gaps may promote the formation of non-B DNA secondary structures by repeated sequences. Such structures formed on the nascent strand will promote expansions, whereas contractions may result from polymerase stalling before a hard-to-replicate hairpin structure followed by 3' end displacement to a new position within the repeated sequence template—for a review, see [27].

Second, impaired replication due to the low-stability CMG-E complex may promote a wide range of events, such as enhanced DNA / polymerase slippage, disruption of the helicase-polymerase interaction, uncoupling of leading and lagging DNA strand syntheses, fork restart and/or enhanced repriming. The mechanism allowing repriming on the leading DNA strand and recycling of stalled leading strand polymerase for downstream synthesis remains under extensive investigation. This mechanism was shown to be primase-coupled in bacteria [149,150] and helicase-coupled in yeast [151]. Interestingly, Hashimoto and colleagues showed that upon fork collapse, the active Cdc45–Mcm2-7–GINS (CMG) helicase complex loses its GINS subunit and interaction with Pol ϵ [152]. Then, Rad51 and Mre11 are required for functional replisome reassembly and reloading *via* a recombination-mediated process. In *psf1-1*, the need for recombination-dependent re-establishment of the replisome can be enhanced if the defects in GINS increase the uncoupling of GINS from the replisome.




Fig 9. Schematic model for the DNA repeats instability mechanisms in the *psf1-1* mutant. Defective interactions within the GINS complex of CMG-E may lead to replication perturbations and/or polymerase-helicase uncoupling and, as a consequence, formation of ssDNA gaps. To ensure the continuity of DNA duplication cells can employ various mechanisms including template switch (TS) and break-induced replication (BIR). These mechanisms require homology search which in case of a repeated DNA tract may result in misalignment leading to its expansion or contraction. Moreover, such DNA sequences may be prone to replication slippage enhancing their instability.

<https://doi.org/10.1371/journal.pgen.1008494.g009>

Finally, the increased amounts of ssDNA and Rfa1 foci (Fig 6C–6E) suggest that in the *psf1-1* cells, Pol ε and the CMG helicase may be uncoupled. Nicks or small gaps that arise during impaired DNA replication can be enlarged. Misalignments occurring during homology search between the invading and the template DNA strand may promote expansions and/or contractions. Nicks and gaps can also become DSBs if replicated or if they occur as a result of breakage due to the intrinsic fragility of ssDNA [153–155] serving as substrates for Rad51 and Rad52 to initiate recombination and enhance DNA repeat instability during D-loop extension (Fig 9). Importantly, as mentioned previously, the *psf1-1* mutant exhibits a dramatic increase in Rad52 foci [148]. These observations are consistent with the reported strong association between faulty replication-induced tract instability (i.e., slippage during synthesis, replication fork

pausing) and recombination [27,68,156]. The question of how different recombination pathways are preferentially activated over others and how the transition between different modes of action occurs requires further investigation. Nonetheless, our present results provide new insights into the source of repetitive sequence instability.

These findings shed new light on the impact of the GINS complex on genome stability in the context of recently described disorders in which mutations or deregulation of the human homolog of the *PSF1* gene was observed [60,157]. Importantly, our results expand the list of genes that, when mutated, affect the structures and functions of other genes and structural elements of the chromosome that contain repetitive DNA sequences.

Materials and methods

Strains, media, and growth conditions

Escherichia coli DH5 α (F⁻, *gyrA96*, *recA1*, *relA1*, *endA1*, *thi1*, *hsdR17*, *supE44*, *deoR*, Δ (*lacZYA-argF*)U169, [ϕ 80 Δ (*lacZ*)M15]) was used for cloning and plasmid amplification. Bacterial strains were grown at 37°C in LB medium (liquid or solidified with 1.5% agar), supplemented when needed with ampicillin 100 μ g/ml (Polfa Tarchomin S.A., Warsaw, Poland). The *Saccharomyces cerevisiae* strains were grown at 23 or 28°C in standard media [158,159]. YPD complete medium (1% Bacto-yeast extract, 2% Bacto-peptone, 2% glucose) was used when nutrient selection was not required. Transformants were selected on YPD with appropriate antibiotics: hygromycin B 300 μ g/ml (Bioshop, Burlington, Canada) or nourseothricin 100 μ g/ml (Werner BioAgents, Jena, Germany). Yeast strains were selected for prototrophy on synthetic SD minimal medium (0.67% yeast nitrogen base without amino acids, 2% glucose) supplemented with appropriate amino acids and nitrogenous bases. Solid media were obtained by the addition of 2% agar. SD medium with 5-FOA 1 mg/ml (USBiologicals, Salem, MA, USA) was used for the selection of *URA3* mutant cells [72].

General methods

Yeast strains were transformed using the LiAc/ssDNA/PEG method [160]. Total DNA was isolated from yeast cultures using the Genomic Mini AX Yeast Spin Kit (A&A Biotechnology, Gdansk, Poland). *E. coli* cells were transformed as described by Sambrook and Russell [161]. Plasmids were isolated from bacteria using the Plasmid Mini Kit (A&A Biotechnology). DNA was extracted from the agarose gel using the Gel-Out Kit (A&A Biotechnology) and purified after enzymatic reactions using the Clean-Up Kit (A&A Biotechnology). Restriction enzymes and DNA ligase (Thermo Scientific, Waltham, MA, USA) were used as recommended by the supplier.

Construction of yeast strains

Yeast strains used in this work are the derivatives of Δ I(-2)I-7B-YUNI300 [162], detailed in S3 Table. Strains carrying deletion of the *REV3*, *RAD51*, *RAD52*, *MSH2*, *MMS2*, *PIF1* or *POL32* gene were constructed based on the SC801 strain [62] by replacing the coding region of the appropriate gene with a DNA cassette containing the *HPH* or *NAT1* gene, which was PCR-amplified with the primers listed in S4 Table using pAG32 or pAG25 [163] as a template. Strains carrying the same deletions and the *psf1-1* allele were constructed by tetrad dissection from diploid strains constructed by crossing strain SC802 with appropriate single gene deletion MAT α strains listed in the S3 Table. Gene replacement was confirmed by PCR using primers listed in S4 Table. The presence of *PSF1* or the *psf1-1* allele was verified by a temperature sensitivity test (*psf1-1* strain does not grow at 37°C) and PCR (primers: InProm, dwPSF1, listed in S4 Table) followed by DNA sequencing.

CTTTCTCGAG3', into the Sall-XhoI-digested plasmid pSH44 [66]. The presence of the control sequence in the constructed plasmid was verified by PCR (with the primers GT_FOR and GT_REV, listed in S4 Table) followed by DNA sequencing. These plasmids were used for the transformation of yeast strains SC801, SC802 and their derivatives carrying deletions of the appropriate genes (S3 Table).

To determine mutation rates, 8–20 cultures of 2 or 3 independent isolates of each strain were inoculated in 2 ml of liquid SD medium supplemented with the required amino acids and nitrogenous bases, lacking tryptophan and leucine, and grown at 23°C. In the experiment with the *psf1-1 pol32Δ* mutant, all strains were grown at 28°C. When cultures reached the stationary phase, appropriate dilutions were plated on selective (containing 5-FOA for selection of *URA3* mutants) and nonselective media. Colonies were counted after 4–7 days of incubation at 23 or 28°C. The spontaneous mutation rates were determined as described below.

To define the spectrum of changes within the repeat tracts or the control sequences, total DNA from 47–85 5-FOA-resistant colonies from independent cultures of each strain was isolated and used for PCR amplification of the analyzed region with fluorescently labeled primers: REP1_FAM/ REP2_ROX/ REP3_HEX/ REP4_TAMRA and GT_FOR (listed in S4 Table). Changes in sequence length were identified by capillary electrophoresis and analyzed using Peak Scanner software v1.0.

Determination of spontaneous mutation rates

The mutation rates were calculated using the equation $\mu = f/\ln(N\mu)$, where μ is the mutation rate per round of DNA replication; f is the mutant frequency (cell count from selective media divided by the cell count from nonselective media), and N is the total population size [167]. The median values of the mutation rates and 95% confidence intervals were evaluated with STATISTICA 6.0 software. Statistical significance of differences in the mutation rates between the respective strains (p -values) was measured using the nonparametric Mann-Whitney U test while Kruskal-Wallis test with the posthoc Mann-Whitney U -test with Bonferroni correction for multiple comparisons was applied to analyze rates in three or more individual cultures.

Visualization of ssDNA in yeast chromosomes by S1 nuclease treatment and separation by pulsed-field gel electrophoresis

Yeast chromosome integrity was analyzed as described previously [168] with certain modifications. Yeast cells grown at 23°C were embedded in 20- μ l plugs of low-melting-point SeaKem Gold agarose (Lonza, Basel, Switzerland). The plugs were digested with Zymolyase 100T (BioShop) overnight at 37°C with gentle rotation and then with proteinase K (A&A Biotechnology) and RNase A (Sigma-Aldrich, St. Louis, MO, USA) overnight at 37°C with gentle rotation. Then, the plugs were treated with 2 U or 5 U of S1 nuclease (Thermo Scientific) for 30 min in S1 buffer provided by the manufacturer. Next, plugs were placed in the wells of a 1% D5 agarose gel (BioMaxima, Lublin, Poland) in 1x TAE and sealed with the same agarose. Electrophoresis was performed on a CHEF Mapper XA pulsed-field electrophoresis system (Bio-Rad, Hercules, CA, USA) for 18 h in 1x TAE buffer at 6 V/cm and 12°C, angle of 120°, and switch time of 60–85 s, with a ramp-up of 0.8. After electrophoresis, the DNA was stained with 0.5 μ g/ml ethidium bromide (Sigma-Aldrich) and visualized using UV light.

Determination of Rad51 protein levels by western blot

For western blotting, 1×10^8 cells from exponentially growing liquid culture (density 5×10^6 ml⁻¹) were collected by centrifugation. Protein extracts prepared using the TCA method were suspended in Laemmli sample buffer supplemented with 1 mM PMSF and cOmplete Protease

Inhibitor Cocktail (Roche, Base, Switzerland), and boiled for 5 min. After centrifugation (19,300 g for 2 min), equal volumes of the protein extracts were separated by SDS-PAGE (8% polyacrylamide gel), and the proteins were transferred onto PVDF membrane (GE Healthcare, Pittsburgh, PA, USA). Blots were blocked for 2 h in 5% (w/v) nonfat dried milk before anti-Rad51 detection or in 3% (w/v) BSA before anti-Act1 detection; both were dissolved in TBST [25 mM Tris-HCl pH 7.5, 137 mM NaCl, 27 mM KCl, 0.1% (v/v) Tween-20]. Rad51 protein was detected with the rabbit polyclonal antibody anti-Rad51 (1:2000, Thermo Fisher, PA5-34905) and goat anti-rabbit IgG conjugated to horseradish peroxidase (HRP) (1:2000, DAKO, P0448). Actin was detected by using a mouse anti-actin monoclonal antibody (1:3000, C4, CHEMICON, MAB-1501) and goat anti-mouse IgG (H+L) alkaline phosphatase (AP)-conjugated (1:3000, Bio-Rad, 1706520) antibody. Immunoreactive proteins on the blots were visualized using chemiluminescent substrates: SuperSignal WestPico, PIERCE for HRP and CDP-Star, ready-to-use, Roche for AP and documented with a charge-coupled device camera (FluorChem Q Multi Image III, Alpha Innotech, San Leandro, CA). The resulting bands were quantified by using Image Quant 5.2 (Molecular Dynamics, Inc., Sunnyvale, CA). The protein levels were normalized to those of Act1. The results of at least eight biological repetitions were averaged to determine the relative protein levels. Statistical significance was determined with Student's t-test.

Fluorescence microscopy

The yeast strains containing the chromosomal fusion of *RFA1-YFP* were grown at 23°C to exponential phase in SD medium supplemented with the required amino acids and nitrogenous bases. The Rfa1-YFP foci were examined using a Nikon E800 fluorescence microscope with a FITC filter (EX465-495, BA 515–555). Images were processed with ImageJ software [169]. At least 200 cells were screened for each of three biological repeats. To analyze the possible differences in Rfa1-Yfp foci in wild-type and *psf1-1* strains we used the contingency tables and further applied the chi-square test (S2 Table).

Supporting information

S1 Fig. Predicted structures formed by repeat tracts. (A) (AACGCAATGCG)₄ and (B) (CAACGCAATGCGTTGGATCT)₃. Predictions were made using the RNAstructure web server for nucleic acid secondary structure prediction (<https://rna.urmc.rochester.edu/RNAstructureWeb/Servers/Predict1/Predict1.html>).

(TIF)

S1 Table. *p*-values associated with data presented in Figs 1, 2, 3, 4, 5, 7 and 8.

(PDF)

S2 Table. Statistical analysis of Rfa1 foci in Wild-type and *psf1-1* cells presented in Fig 6E.

Contingency table and chi-square test.

(PDF)

S3 Table. Yeast strains used in this work.

(PDF)

S4 Table. Primers used in this work.

(PDF)

S5 Table. Random sequences used in control experiments.

(PDF)

Acknowledgments

We thank Hiroyuki Araki (National Institute of Genetics, Mishima, Japan), Robert S. Lahue (National University of Ireland, Galway, Ireland), and Thomas D. Petes (Duke University School of Medicine, Durham, NC, USA) for providing strains and plasmids. We are grateful to our colleagues from the Laboratory of Mutagenesis and DNA Repair for fruitful discussions. We thank Malgorzata Lichočka from the Laboratory of Confocal and Fluorescence Microscopy for assistance with fluorescence microscopy and Jaroslaw Poznanski for help with statistical analyses.

Author Contributions

Conceptualization: Malgorzata Alabrudzinska, Michal Dmowski, Iwona J. Fijalkowska.

Data curation: Malgorzata Jedrychowska, Milena Denkiewicz-Kruk, Malgorzata Alabrudzinska, Adrianna Skoneczna, Michal Dmowski.

Formal analysis: Michal Dmowski, Iwona J. Fijalkowska.

Funding acquisition: Michal Dmowski, Iwona J. Fijalkowska.

Investigation: Malgorzata Jedrychowska, Milena Denkiewicz-Kruk, Malgorzata Alabrudzinska, Adrianna Skoneczna, Piotr Jonczyk, Michal Dmowski.

Supervision: Michal Dmowski, Iwona J. Fijalkowska.

Visualization: Michal Dmowski.

Writing – original draft: Michal Dmowski, Iwona J. Fijalkowska.

Writing – review & editing: Michal Dmowski, Iwona J. Fijalkowska.

References

1. Loeb LA. Mutator phenotype may be required for multistage carcinogenesis. *Cancer Res. American Association for Cancer Research*; 1991; 51: 3075–9. Available: <http://www.ncbi.nlm.nih.gov/pubmed/2039987>
2. Drake JW, Charlesworth B, Charlesworth D, Crow JF. Rates of spontaneous mutation. *Genetics*. 1998; 148: 1667–86. Available: <http://www.ncbi.nlm.nih.gov/pubmed/9560386> PMID: 9560386
3. De Bont R, van Larebeke N. Endogenous DNA damage in humans: a review of quantitative data. *Mutagenesis*. 2004; 19: 169–85. Available: <http://www.ncbi.nlm.nih.gov/pubmed/15123782> <https://doi.org/10.1093/mutage/geh025> PMID: 15123782
4. Tóth G, Gáspári Z, Jurka J. Microsatellites in different eukaryotic genomes: survey and analysis. *Genome Res. Cold Spring Harbor Laboratory Press*; 2000; 10: 967–81. Available: <http://www.ncbi.nlm.nih.gov/pubmed/10899146> <https://doi.org/10.1101/gr.10.7.967> PMID: 10899146
5. Ramel C. Mini- and microsatellites. *Environ Health Perspect*. 1997; 105: 781–789. <https://doi.org/10.1289/ehp.97105s4781> PMID: 9255562
6. Wang Z, Weber JL, Zhong G, Tanksley SD. Survey of plant short tandem DNA repeats. *Theor Appl Genet*. 1994; 88: 1–6. <https://doi.org/10.1007/BF00222386> PMID: 24185874
7. Schug MD, Wetterstrand KA, Gaudette MS, Lim RH, Hutter CM, Aquadro CF. The distribution and frequency of microsatellite loci in *Drosophila melanogaster*. *Mol Ecol. John Wiley & Sons, Ltd* (10.1111); 1998; 7: 57–70. <https://doi.org/10.1046/j.1365-294x.1998.00304.x> PMID: 9465417
8. Wren JD, Forgacs E, Fondon JW, Pertsemidlis A, Cheng SY, Gallardo T, et al. Repeat Polymorphisms within Gene Regions: Phenotypic and Evolutionary Implications. *Am J Hum Genet. Cell Press*; 2000; 67: 345–356. <https://doi.org/10.1086/303013> PMID: 10889045
9. Lafrenière RG, Rochefort DL, Chrétien N, Rommens JM, Cochiuș JI, Kälviäinen R, et al. Unstable insertion in the 5' flanking region of the cystatin B gene is the most common mutation in progressive myoclonus epilepsy type 1, EPM1. *Nat Genet. Nature Publishing Group*; 1997; 15: 298–302. <https://doi.org/10.1038/ng0397-298> PMID: 9054946

10. Kennedy GC, German MS, Rutter WJ. The minisatellite in the diabetes susceptibility locus IDDM2 regulates insulin transcription. *Nat Genet.* Nature Publishing Group; 1995; 9: 293–298. <https://doi.org/10.1038/ng0395-293> PMID: 7773292
11. Yang B, Chan RCK, Jing J, Li T, Sham P, Chen RYL. A meta-analysis of association studies between the 10-repeat allele of a VNTR polymorphism in the 3'-UTR of dopamine transporter gene and attention deficit hyperactivity disorder. *Am J Med Genet Part B Neuropsychiatr Genet.* John Wiley & Sons, Ltd; 2007; 144B: 541–550. <https://doi.org/10.1002/ajmg.b.30453> PMID: 17440978
12. Kirkbride HJ, Bolscher JG, Nazmi K, Vinal LE, Nash MW, Moss FM, et al. Genetic polymorphism of MUC7: Allele frequencies and association with asthma. *Eur J Hum Genet.* Nature Publishing Group; 2001; 9: 347–354. <https://doi.org/10.1038/sj.ejhg.5200642> PMID: 11378823
13. Kyo K, Parkes M, Takei Y, Nishimori H, Vyas P, Satsangi J, et al. Association of ulcerative colitis with rare VNTR alleles of the human intestinal mucin gene, MUC3. *Hum Mol Genet.* 1999; 8: 307–11. Available: <http://www.ncbi.nlm.nih.gov/pubmed/9931338> <https://doi.org/10.1093/hmg/8.2.307> PMID: 9931338
14. Jeong YH, Kim MC, Ahn E-K, Seol S-Y, Do E-J, Choi H-J, et al. Rare Exonic Minisatellite Alleles in MUC2 Influence Susceptibility to Gastric Carcinoma. von Elm E, editor. *PLoS One.* Public Library of Science; 2007; 2: e1163. <https://doi.org/10.1371/journal.pone.0001163> PMID: 18000536
15. Krontiris TG, Devlin B, Karp DD, Robert NJ, Risch N. An Association between the Risk of Cancer and Mutations in the HRAS1 Minisatellite Locus. *N Engl J Med.* Massachusetts Medical Society; 1993; 329: 517–523. <https://doi.org/10.1056/NEJM1993081933290801> PMID: 8336750
16. Wang L, Ogawa S, Hangaishi A, Qiao Y, Hosoya N, Nanya Y, et al. Molecular characterization of the recurrent unbalanced translocation der(1;7)(q10;p10). *Blood.* American Society of Hematology; 2003; 102: 2597–604. <https://doi.org/10.1182/blood-2003-01-0031> PMID: 12816870
17. Mirkin SM. Expandable DNA repeats and human disease. *Nature.* 2007; 447: 932–940. <https://doi.org/10.1038/nature05977> PMID: 17581576
18. Pearson CE, Edamura KN, Cleary JD. Repeat instability: Mechanisms of dynamic mutations. *Nat Rev Genet.* 2005; 6: 729–742. <https://doi.org/10.1038/nrg1689> PMID: 16205713
19. Zhao XN, Usdin K. The Repeat Expansion Diseases: The dark side of DNA repair. *DNA Repair (Amst).* Elsevier B.V.; 2015; 32: 96–105. <https://doi.org/10.1016/j.dnarep.2015.04.019> PMID: 26002199
20. Fan H, Chu JY. A Brief Review of Short Tandem Repeat Mutation. *Genomics, Proteomics Bioinforma.* Beijing Institute of Genomics; 2007; 5: 7–14. [https://doi.org/10.1016/S1672-0229\(07\)60009-6](https://doi.org/10.1016/S1672-0229(07)60009-6) PMID: 17572359
21. Garrido-Ramos MA. Satellite DNA: An evolving topic. *Genes (Basel).* 2017; 8. <https://doi.org/10.3390/genes8090230> PMID: 28926993
22. Tachida H, Iizuka M. Persistence of repeated sequences that evolve by replication slippage. *Genetics.* 1992; 131.
23. Cleary JD, Pearson CE. Replication fork dynamics and dynamic mutations: the fork-shift model of repeat instability. *Trends Genet.* Elsevier Current Trends; 2005; 21: 272–280. <https://doi.org/10.1016/j.tig.2005.03.008> PMID: 15851063
24. Mirkin SM. DNA structures, repeat expansions and human hereditary disorders. *Curr Opin Struct Biol.* Elsevier Current Trends; 2006; 16: 351–358. <https://doi.org/10.1016/j.sbi.2006.05.004> PMID: 16713248
25. Harding RM, Boyce AJ, Clegg JB. The evolution of tandemly repetitive DNA: recombination rules. *Genetics.* Genetics Society of America; 1992; 132: 847–59. Available: <http://www.ncbi.nlm.nih.gov/pubmed/1468634> PMID: 1468634
26. Wells RD, Dere R, Hebert ML, Napierala M, Son LS. Advances in mechanisms of genetic instability related to hereditary neurological diseases. *Nucleic Acids Res.* Narnia; 2005; 33: 3785–3798. <https://doi.org/10.1093/nar/gki697> PMID: 16006624
27. Polleys EJ, House NCM, Freudenreich CH. Role of recombination and replication fork restart in repeat instability. *DNA Repair (Amst).* Elsevier; 2017; 56: 156–165. <https://doi.org/10.1016/j.dnarep.2017.06.018> PMID: 28641941
28. Jaworski A, Rosche WA, Gellibolian R, Kang S, Shimizu M, Bowater RP, et al. Mismatch repair in *Escherichia coli* enhances instability of (CTG)_n triplet repeats from human hereditary diseases. *Proc Natl Acad Sci U S A.* National Academy of Sciences; 1995; 92: 11019–23. <https://doi.org/10.1073/pnas.92.24.11019> PMID: 7479928
29. Panigrahi GB, Lau R, Montgomery SE, Leonard MR, Pearson CE. Slipped (CTG)_n(CAG) repeats can be correctly repaired, escape repair or undergo error-prone repair. *Nat Struct Mol Biol.* Nature Publishing Group; 2005; 12: 654–662. <https://doi.org/10.1038/nsmb959> PMID: 16025129

30. Daele DL, Mertz T, Lahue RS. Postreplication Repair Inhibits CAG · CTG Repeat Expansions in *Saccharomyces cerevisiae*. *Mol Cell Biol*. 2007; 27: 102–110. <https://doi.org/10.1128/MCB.01167-06> PMID: 17060452
31. Kovtun I V., Liu Y, Bjoras M, Klungland A, Wilson SH, McMurray CT. OGG1 initiates age-dependent CAG trinucleotide expansion in somatic cells. *Nature*. Nature Publishing Group; 2007; 447: 447–452. <https://doi.org/10.1038/nature05778> PMID: 17450122
32. McMurray CT. Hijacking of the mismatch repair system to cause CAG expansion and cell death in neurodegenerative disease. *DNA Repair (Amst)*. NIH Public Access; 2008; 7: 1121–34. <https://doi.org/10.1016/j.dnarep.2008.03.013> PMID: 18472310
33. McMurray CT. Mechanisms of trinucleotide repeat instability during human development. *Nat Rev Genet*. NIH Public Access; 2010; 11: 786–99. <https://doi.org/10.1038/nrg2828> PMID: 20953213
34. Usdin K, House NCM, Freudenreich CH. Repeat instability during DNA repair: Insights from model systems. *Crit Rev Biochem Mol Biol*. Informa Healthcare; 2015; 50: 142–167. <https://doi.org/10.3109/10409238.2014.999192> PMID: 25608779
35. Dion V, Wilson JH. Instability and chromatin structure of expanded trinucleotide repeats. *Trends Genet*. Elsevier Current Trends; 2009; 25: 288–297. <https://doi.org/10.1016/j.tig.2009.04.007> PMID: 19540013
36. Wierdl M, Greene CN, Datta A, Jinks-Robertson S, Petes TD. Destabilization of Simple Repetitive DNA Sequences by Transcription in Yeast. *Genetics*. 1996; 143.
37. Brohede J, Ellegren H. Microsatellite evolution: polarity of substitutions within repeats and neutrality of flanking sequences. *Proc R Soc London Ser B Biol Sci*. 1999; 266: 825–833. <https://doi.org/10.1098/rspb.1999.0712> PMID: 10343406
38. Gendrel CG, Boulet A, Dutreix M. (CA/GT)(n) microsatellites affect homologous recombination during yeast meiosis. *Genes Dev*. Cold Spring Harbor Laboratory Press; 2000; 14: 1261–8. Available: <http://www.ncbi.nlm.nih.gov/pubmed/10817760>
39. Li Y, Fahima T, Korol AB, Peng J, R MS, Kirzhner V, et al. Microsatellite Diversity Correlated with Ecological-Edaphic and Genetic Factors in Three Microsites of Wild Emmer Wheat in North Israel. *Mol Biol Evol*. 2000; 17: 851–862. <https://doi.org/10.1093/oxfordjournals.molbev.a026365> PMID: 10833191
40. Li Y-C, Fahima T, Röder MS, Kirzhner VM, Beiles A, Korol AB, et al. Genetic effects on microsatellite diversity in wild emmer wheat (*Triticum dicoccoides*) at the Yehudiyya microsite, Israel. *Heredity (Edinb)*. Nature Publishing Group; 2003; 90: 150–156. <https://doi.org/10.1038/sj.hdy.6800190> PMID: 12634821
41. Lenzmeier BA, Freudenreich CH. Trinucleotide repeat instability: a hairpin curve at the crossroads of replication, recombination, and repair. *Cytogenet Genome Res*. Karger Publishers; 2003; 100: 7–24. <https://doi.org/10.1159/000072836> PMID: 14526162
42. Iraqui I, Chekkal Y, Jmari N, Pietrobon V, Fréon K, Costes A, et al. Recovery of Arrested Replication Forks by Homologous Recombination Is Error-Prone. *PLoS Genet*. 2012; 8. <https://doi.org/10.1371/journal.pgen.1002976> PMID: 23093942
43. Kokoska RJ, Stefanovic L, DeMai J, Petes TD. Increased rates of genomic deletions generated by mutations in the yeast gene encoding DNA polymerase delta or by decreases in the cellular levels of DNA polymerase delta. *Mol Cell Biol*. 2000; 20: 7490–504. <https://doi.org/10.1128/mcb.20.20.7490-7504.2000> PMID: 11003646
44. Shah KA, Shishkin AA, Voineagu I, Pavlov YI, Shcherbakova P V., Mirkin SM. Role of DNA Polymerases in Repeat-Mediated Genome Instability. *Cell Rep*. The Authors; 2012; 2: 1088–1095. <https://doi.org/10.1016/j.celrep.2012.10.006> PMID: 23142667
45. Tran HT, Gordenin DA, Resnick MA. The 3'→5' exonucleases of DNA polymerases delta and epsilon and the 5'→3' exonuclease Exo1 have major roles in postreplication mutation avoidance in *Saccharomyces cerevisiae*. *Mol Cell Biol*. 1999; 19: 2000–7. <https://doi.org/10.1128/mcb.19.3.2000> PMID: 10022887
46. Schweitzer JK, Livingston DM. The effect of DNA replication mutations on CAG tract stability in yeast. *Genetics*. 1999; 152: 953–963. <https://doi.org/10.1534/genetics.112.541.test> PMID: 10388815
47. Zheng DQ, Petes TD. Genome instability induced by low levels of replicative DNA polymerases in yeast. *Genes (Basel)*. 2018; 9. <https://doi.org/10.3390/genes9110539> PMID: 30405078
48. Kokoska RJ, Stefanovic L, Tran HT, Resnick M a, Gordenin D a, Petes TD. Destabilization of yeast micro- and minisatellite DNA sequences by mutations affecting a nuclease involved in Okazaki fragment processing (rad27) and DNA polymerase delta (pol3-t). *Mol Cell Biol*. 1998; 18: 2779–2788. <https://doi.org/10.1128/mcb.18.5.2779> PMID: 9566897

49. Kokoska RJ, Stefanovic L, Buermeyer AB, Liskay RM, Petes TD. A Mutation of the Yeast Gene Encoding PCNA Destabilizes Both Microsatellite and Minisatellite DNA Sequences. *Genetics*. 1999;151.
50. Kurth I, O'Donnell M. New insights into replisome fluidity during chromosome replication. *Trends Biochem Sci*. Elsevier Ltd; 2013; 38: 195–203. <https://doi.org/10.1016/j.tibs.2012.10.003> PMID: 23153958
51. Onesti S, MacNeill SA. Structure and evolutionary origins of the CMG complex. *Chromosoma*. Springer-Verlag; 2013; 122: 47–53. <https://doi.org/10.1007/s00412-013-0397-x> PMID: 23412083
52. Kanemaki M, Sanchez-Diaz A, Gambus A, Labib K. Functional proteomic identification of DNA replication proteins by induced proteolysis in vivo. *Nature*. 2003; 423: 720–725. <https://doi.org/10.1038/nature01692> PMID: 12768207
53. Kubota Y, Takase Y, Komori Y, Hashimoto Y, Arata T, Kamimura Y, et al. A novel ring-like complex of *Xenopus* proteins essential for the initiation of DNA replication. *Genes Dev*. 2003; 17: 1141–1152. <https://doi.org/10.1101/gad.1070003> PMID: 12730133
54. Takayama Y, Kamimura Y, Okawa M, Muramatsu S, Sugino A, Araki H. GINS, a novel multiprotein complex required for chromosomal DNA replication in budding yeast. *Genes Dev*. 2003; 17: 1153–1165. <https://doi.org/10.1101/gad.1065903> PMID: 12730134
55. Gambus A, Jones RC, Sanchez-Diaz A, Kanemaki M, van Deursen F, Edmondson RD, et al. GINS maintains association of Cdc45 with MCM in replisome progression complexes at eukaryotic DNA replication forks. *Nat Cell Biol*. 2006; 8: 358–366. <https://doi.org/10.1038/ncb1382> PMID: 16531994
56. Langston LD, Mayle R, Schauer GD, Yurieva O, Zhang D, Yao NY, et al. Mcm10 promotes rapid isomerization of CMG-DNA for replisome bypass of lagging strand DNA blocks. *Elife*. 2017; 6. <https://doi.org/10.7554/elife.29118> PMID: 28869037
57. Langston LD, Zhang D, Yurieva O, Georgescu RE, Finkelstein J, Yao NY, et al. CMG helicase and DNA polymerase ϵ form a functional 15-subunit holoenzyme for eukaryotic leading-strand DNA replication. *Proc Natl Acad Sci*. 2014; 111: 15390–15395. <https://doi.org/10.1073/pnas.1418334111> PMID: 25313033
58. Bermudez VP, Farina A, Raghavan V, Tappin I, Hurwitz J. Studies on human DNA polymerase epsilon and GINS complex and their role in DNA replication. *J Biol Chem*. 2011; 286: 28963–77. <https://doi.org/10.1074/jbc.M111.256289> PMID: 21705323
59. Handa T, Kanke M, Takahashi TS, Nakagawa T, Masukata H. DNA polymerization-independent functions of DNA polymerase epsilon in assembly and progression of the replisome in fission yeast. *Mol Biol Cell*. 2012; 23: 3240–3253. <https://doi.org/10.1091/mbc.E12-05-0339> PMID: 22718908
60. Cottineau J, Kottemann MC, Lach FP, Kang YH, Vély F, Deenick EK, et al. Inherited GINS1 deficiency underlies growth retardation along with neutropenia and NK cell deficiency. *J Clin Invest*. 2017; 127: 1991–2006. <https://doi.org/10.1172/JCI90727> PMID: 28414293
61. Sengupta S, van Deursen F, de Piccoli G, Labib K. Dpb2 integrates the leading-strand DNA polymerase into the eukaryotic replisome. *Curr Biol*. Elsevier Ltd; 2013; 23: 543–552. <https://doi.org/10.1016/j.cub.2013.02.011> PMID: 23499531
62. Grabowska E, Wronska U, Denkiewicz M, Jaszczur M, Respondek A, Alabrudzinska M, et al. Proper functioning of the GINS complex is important for the fidelity of DNA replication in yeast. *Mol Microbiol*. 2014; 92: 659–680. <https://doi.org/10.1111/mmi.12580> PMID: 24628792
63. Costa A, Renault L, Swuec P, Petojevic T, Pesavento JJ, Ilves I, et al. DNA binding polarity, dimerization, and ATPase ring remodeling in the CMG helicase of the eukaryotic replisome. *Elife*. 2014; 3: e03273. <https://doi.org/10.7554/eLife.03273> PMID: 25117490
64. Yuan Z, Bai L, Sun J, Georgescu R, Liu J, O'Donnell ME, et al. Structure of the eukaryotic replicative CMG helicase suggests a pumpjack motion for translocation. *Nat Struct Mol Biol*. Nature Publishing Group; 2016; 23: 217–24. <https://doi.org/10.1038/nsmb.3170> PMID: 26854665
65. Miret JJ, Pessoa-Brandão L, Lahue RS. Orientation-dependent and sequence-specific expansions of CTG/CAG trinucleotide repeats in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci U S A*. 1998; 95: 12438–43. <https://doi.org/10.1073/pnas.95.21.12438> PMID: 9770504
66. Henderson ST, Petes TD. Instability of simple sequence DNA in *Saccharomyces cerevisiae*. *Mol Cell Biol*. 1992; 12: 2749–2757. <https://doi.org/10.1128/mcb.12.6.2749> PMID: 1588966
67. Sia EA, Kokoska RJ, Dominska M, Greenwell P, Petes TD. Microsatellite instability in yeast: dependence on repeat unit size and DNA mismatch repair genes. *Mol Cell Biol*. 1997; 17: 2851–8. <https://doi.org/10.1128/mcb.17.5.2851> PMID: 9111357
68. Mosbach V, Poggi L, Richard G-F. Trinucleotide repeat instability during double-strand break repair: from mechanisms to gene therapy. *Curr Genet*. Springer Berlin Heidelberg; 2018; 0: 0. <https://doi.org/10.1007/s00294-018-0865-1> PMID: 29974202

69. Gineau L, Cognet C, Kara N, Lach FP, Dunne J, Veturi U, et al. Partial MCM4 deficiency in patients with growth retardation, adrenal insufficiency, and natural killer cell deficiency. *J Clin Invest*. 2012; 122: 821–832. <https://doi.org/10.1172/JCI61014> PMID: 22354167
70. Hughes CR, Guasti L, Meimaridou E, Chuang C, Schimenti JC, King PJ, et al. MCM4 mutation causes adrenal failure, Short Stature, and Natural Killer Cell Deficiency in Humans. *J Clin Invest*. 2012; 122: 814–820. <https://doi.org/10.1172/JCI60224> PMID: 22354170
71. Seo Y-S, Kang Y-H. The Human Replicative Helicase, the CMG Complex, as a Target for Anti-cancer Therapy. *Front Mol Biosci*. *Frontiers*; 2018; 5: 26. <https://doi.org/10.3389/fmolb.2018.00026> PMID: 29651420
72. Boeke JD, LaCroute F, Fink GR. A positive selection for mutants lacking 5' phosphate decarboxylase activity in yeast: 5 fluoro-orotic acid resistance. *Mol Gen Genet*. 1984; 197: 345–346. <https://doi.org/10.1007/bf00330984> PMID: 6394957
73. Rolfsmeier ML, Dixon MJ, Pessoa-Brandão L, Pelletier R, Miret JJ, Lahue RS. Cis-elements governing trinucleotide repeat instability in *Saccharomyces cerevisiae*. *Genetics*. 2001; 157: 1569–1579. PMID: 11290713
74. Richards RI, Sutherland GR. Simple repeat DNA is not replicated simply. *Nat Genet*. 1994; 6: 114–116. <https://doi.org/10.1038/ng0294-114> PMID: 8162063
75. Eckert KA, Hile SE. Every microsatellite is different: Intrinsic DNA features dictate mutagenesis of common microsatellites present in the human genome. *Mol Carcinog*. John Wiley & Sons, Ltd; 2009; 48: 379–388. <https://doi.org/10.1002/mc.20499> PMID: 19306292
76. Kolodner RD, Marsischky GT. Eukaryotic DNA mismatch repair. *Curr Opin Genet Dev*. Elsevier Current Trends; 1999; 9: 89–96. [https://doi.org/10.1016/s0959-437x\(99\)80013-6](https://doi.org/10.1016/s0959-437x(99)80013-6) PMID: 10072354
77. Liu D, Keijzers G, Rasmussen LJ. DNA mismatch repair and its many roles in eukaryotic cells. *Mutat Res Mutat Res*. Elsevier; 2017; 773: 174–187. <https://doi.org/10.1016/j.MRREV.2017.07.001> PMID: 28927527
78. Johnson RE, Kovvali GK, Prakash L, Prakash S. Requirement of the yeast MSH3 and MSH6 genes for MSH2-dependent genomic stability. *J Biol Chem*. American Society for Biochemistry and Molecular Biology; 1996; 271: 7285–8. <https://doi.org/10.1074/jbc.271.13.7285> PMID: 8631743
79. Marsischky GT, Filosi N, Kane MF, Kolodner R. Redundancy of *Saccharomyces cerevisiae* MSH3 and MSH6 in MSH2-dependent mismatch repair. *Genes Dev*. 1996; 10: 407–20. Available: <http://www.ncbi.nlm.nih.gov/pubmed/8600025> <https://doi.org/10.1101/gad.10.4.407> PMID: 8600025
80. Bishop DK, Williamson MS, Fogel S, Kolodner RD. The role of heteroduplex correction in gene conversion in *Saccharomyces cerevisiae*. *Nature*. Nature Publishing Group; 1987; 328: 362–364. <https://doi.org/10.1038/328362a0> PMID: 3299108
81. Tran HT, Gordenin DA, Resnick MA. The prevention of repeat-associated deletions in *Saccharomyces cerevisiae* by mismatch repair depends on size and origin of deletions. *Genetics*. 1996. pp. 1579–1587. PMID: 8844147
82. Sia EA, Jinks-Robertson S, Petes TD. Genetic control of microsatellite stability. *Mutat Res Repair*. Elsevier; 1997; 383: 61–70. [https://doi.org/10.1016/S0921-8777\(96\)00046-8](https://doi.org/10.1016/S0921-8777(96)00046-8)
83. Greene CN, Jinks-Robertson S. Frameshift intermediates in homopolymer runs are removed efficiently by yeast mismatch repair proteins. *Mol Cell Biol*. American Society for Microbiology Journals; 1997; 17: 2844–50. <https://doi.org/10.1128/mcb.17.5.2844> PMID: 9111356
84. Greene CN, Jinks-Robertson S. Spontaneous frameshift mutations in *saccharomyces cerevisiae*: Accumulation during dna replication and removal by proofreading and mismatch repair activities. *Genetics*. 2001; 159: 65–75. PMID: 11560887
85. Heale SM, Petes TD. The stabilization of repetitive tracts of DNA by variant repeats requires a functional DNA mismatch repair system. *Cell*. Cell Press; 1995; 83: 539–545. [https://doi.org/10.1016/0092-8674\(95\)90093-4](https://doi.org/10.1016/0092-8674(95)90093-4) PMID: 7585956
86. Strand M, Earley MC, Crouse GF, Petes TD. Mutations in the MSH3 gene preferentially lead to deletions within tracts of simple repetitive DNA in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci U S A*. National Academy of Sciences; 1995; 92: 10418–21. <https://doi.org/10.1073/pnas.92.22.10418> PMID: 7479796
87. Sia EA, Dominska M, Stefanovic L, Petes TD. Isolation and Characterization of Point Mutations in Mismatch Repair Genes That Destabilize Microsatellites in Yeast. *Mol Cell Biol*. American Society for Microbiology Journals; 2001; 21: 8157–8167. <https://doi.org/10.1128/MCB.21.23.8157-8167.2001> PMID: 11689704
88. Makarova A V, Stodola JL, Burgers PM. A four-subunit DNA polymerase ζ complex containing Pol δ accessory subunits is essential for PCNA-mediated mutagenesis. *Nucleic Acids Res*. 2012; 40: 11618–26. <https://doi.org/10.1093/nar/gks948> PMID: 23066099

89. Johnson RE, Prakash L, Prakash S. Pol31 and Pol32 subunits of yeast DNA polymerase δ are also essential subunits of DNA polymerase ζ . *Proc Natl Acad Sci U S A*. 2012; 109: 12455–60. <https://doi.org/10.1073/pnas.1206052109> PMID: 22711820
90. Pâques F, Haber JE. Multiple pathways of recombination induced by double-strand breaks in *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev*. 1999; 63: 349–404. Available: <http://www.ncbi.nlm.nih.gov/pubmed/10357855> PMID: 10357855
91. Symington LS, Rothstein R, Lisby M. Mechanisms and regulation of mitotic recombination in *saccharomyces cerevisiae*. *Genetics*. 2014; 198: 795–835. <https://doi.org/10.1534/genetics.114.166140> PMID: 25381364
92. Hum YF, Jinks-Robertson S. DNA strand-exchange patterns associated with double-strand break-induced and spontaneous mitotic crossovers in *Saccharomyces cerevisiae*. Copenhaver GP, editor. *PLOS Genet*. Public Library of Science; 2018; 14: e1007302. <https://doi.org/10.1371/journal.pgen.1007302> PMID: 29579095
93. Krol K, Antoniuk-Majchrzak J, Skoneczny M, Sienko M, Jendrysek J, Rumieniczuk I, et al. Lack of G1/S control destabilizes the yeast genome via replication stress-induced DSBs and illegitimate recombination. *J Cell Sci*. The Company of Biologists Ltd; 2018; 131: jcs226480. <https://doi.org/10.1242/jcs.226480> PMID: 30463853
94. Skoneczna A, Krol K, Skoneczny M. How Do Yeast and Other Fungi Recognize and Respond to Genome Perturbations? *Stress Response Mechanisms in Fungi*. 2018. https://doi.org/10.1007/978-3-030-00683-9_3
95. Piazza A, Heyer WD. Homologous Recombination and the Formation of Complex Genomic Rearrangements. *Trends Cell Biol*. Elsevier Ltd; 2019; 29: 135–149. <https://doi.org/10.1016/j.tcb.2018.10.006> PMID: 30497856
96. Godin SK, Sullivan MR, Bernstein KA. Novel insights into RAD51 activity and regulation during homologous recombination and DNA replication. *Biochem Cell Biol*. NIH Public Access; 2016; 94: 407–418. <https://doi.org/10.1139/bcb-2016-0012> PMID: 27224545
97. Bhat KP, Cortez D. RPA and RAD51: fork reversal, fork protection, and genome stability. *Nat Struct Mol Biol*. Nature Publishing Group; 2018; 25: 446–453. <https://doi.org/10.1038/s41594-018-0075-z> PMID: 29807999
98. Andriuskevicius T, Kotenko O, Makovets S. Putting together and taking apart: assembly and disassembly of the Rad51 nucleoprotein filament in DNA repair and genome stability. *Cell Stress*. Shared Science Publishers; 2018; 2: 96–112. <https://doi.org/10.15698/cst2018.05.134> PMID: 31225474
99. Jenkins SS, Gore S, Guo X, Liu J, Ede C, Veaute X, et al. Role of the Srs2-Rad51 Interaction Domain in Crossover Control in *Saccharomyces cerevisiae*. *Genetics*. Genetics; 2019; genetics.302337.2019. <https://doi.org/10.1534/genetics.119.302337> PMID: 31142613
100. Karras GI, Jentsch S. The RAD6 DNA damage tolerance pathway operates uncoupled from the replication fork and is functional beyond S phase. *Cell*. Elsevier Ltd; 2010; 141: 255–67. <https://doi.org/10.1016/j.cell.2010.02.028> PMID: 20403322
101. Vanoli F, Fumasoni M, Szakal B, Maloisel L, Branzei D. Replication and Recombination Factors Contributing to Recombination-Dependent Bypass of DNA Lesions by Template Switch. Haber JE, editor. *PLoS Genet*. Public Library of Science; 2010; 6: e1001205. <https://doi.org/10.1371/journal.pgen.1001205> PMID: 21085632
102. Branzei D, Szakal B. Building up and breaking down: mechanisms controlling recombination during replication. *Crit Rev Biochem Mol Biol*. 2017; 52: 381–394. <https://doi.org/10.1080/10409238.2017.1304355> PMID: 28325102
103. Boiteux S, Jinks-Robertson S, Hartwell LH, Crouse GF. DNA repair mechanisms and the bypass of DNA damage in *Saccharomyces cerevisiae*. *Genetics*. Genetics; 2013; 193: 1025–64. <https://doi.org/10.1534/genetics.112.145219> PMID: 23547164
104. Saugar I, Ortiz-Bazán MÁ, Tercero JA. Tolerating DNA damage during eukaryotic chromosome replication. *Exp Cell Res*. Academic Press; 2014; 329: 170–177. <https://doi.org/10.1016/j.yexcr.2014.07.009> PMID: 25038291
105. Kramara J, Osia B, Malkova A. Break-Induced Replication: The Where, The Why, and The How. *Trends Genet*. Elsevier Current Trends; 2018; 34: 518–531. <https://doi.org/10.1016/j.tig.2018.04.002> PMID: 29735283
106. Leffak M. Break-induced replication links microsatellite expansion to complex genome rearrangements. *BioEssays*. 2017; 39: 1–8. <https://doi.org/10.1002/bies.201700025> PMID: 28621832
107. Lydeard JR, Lipkin-Moore Z, Sheu YJ, Stillman B, Burgers PM, Haber JE. Break-induced replication requires all essential DNA replication factors except those specific for pre-RC assembly. *Genes Dev*. 2010; 24: 1133–1144. <https://doi.org/10.1101/gad.1922610> PMID: 20516198

108. Wilson MA, Kwon Y, Xu Y, Chung WH, Chi P, Niu H, et al. Pif1 helicase and Pol δ promote recombination-coupled DNA synthesis via bubble migration. *Nature*. Nature Publishing Group; 2013; 502: 393–396. <https://doi.org/10.1038/nature12585> PMID: 24025768
109. Donnianni RA, Symington LS. Break-induced replication occurs by conservative DNA synthesis. *Proc Natl Acad Sci*. 2013; 110: 13475–13480. <https://doi.org/10.1073/pnas.1309800110> PMID: 23898170
110. Lydeard JR, Jain S, Yamaguchi M, Haber JE. Break-induced replication and telomerase-independent telomere maintenance require Pol32. *Nature*. 2007; 448: 820–823. <https://doi.org/10.1038/nature06047> PMID: 17671506
111. Gerik KJ, Gerik KJ, Pautz A, Pautz A. Characterization of the Two Small Subunits of *Saccharomyces cerevisiae* DNA Polymerase δ . *Mol Biol*. 1998; 273: 19747–19755. <https://doi.org/10.1074/jbc.273.31.19747> PMID: 9677405
112. Szwajczak E, Fijalkowska IJ, Suski C. The CysB motif of Rev3p involved in the formation of the four-subunit DNA polymerase ζ is required for defective-replisome-induced mutagenesis. *Mol Microbiol*. 2017; 106: 659–672. <https://doi.org/10.1111/mmi.13846> PMID: 28941243
113. Huang ME, de Calignon a, Nicolas a, Galibert F. POL32, a subunit of the *Saccharomyces cerevisiae* DNA polymerase delta, defines a link between DNA replication and the mutagenic bypass repair pathway. *Curr Genet*. 2000; 38: 178–187. <https://doi.org/10.1007/s002940000149> PMID: 11126776
114. Buzovetsky O, Kwon Y, Pham NT, Kim C, Ira G, Sung P, et al. Role of the Pif1-PCNA Complex in Pol δ -Dependent Strand Displacement DNA Synthesis and Break-Induced Replication. *Cell Rep*. Elsevier-Company.; 2017; 21: 1707–1714. <https://doi.org/10.1016/j.celrep.2017.10.079> PMID: 29141206
115. Boulé JB, Zakian VA. Roles of Pif1-like helicases in the maintenance of genomic stability. *Nucleic Acids Res*. 2006; 34: 4147–4153. <https://doi.org/10.1093/nar/gkl561> PMID: 16935874
116. Saini N, Ramakrishnan S, Elango R, Ayyar S, Zhang Y, Deem A, et al. Migrating bubble during break-induced replication drives conservative DNA synthesis. *Nature*. 2013; 502: 389–392. <https://doi.org/10.1038/nature12584> PMID: 24025772
117. Van Dyck E, Foury F, Stillman B, Brill SJ. A single-stranded DNA binding protein required for mitochondrial DNA replication in *S. cerevisiae* is homologous to *E. coli* SSB. *EMBO J*. 1992; 11: 3421–3430. <https://doi.org/10.1002/j.1460-2075.1992.tb05421.x> PMID: 1324172
118. Foury F, Van Dyck E. A PIF-dependent recombinogenic signal in the mitochondrial DNA of yeast. *EMBO J*. 1985; 4: 3525–3530. <https://doi.org/10.1002/j.1460-2075.1985.tb04112.x> PMID: 16453651
119. Serero A, Jubin C, Loeillet S, Legoix-Né P, Nicolas AG. Mutational landscape of yeast mutator strains. *Proc Natl Acad Sci U S A*. 2014; 111: 1897–902. <https://doi.org/10.1073/pnas.1314423111> PMID: 24449905
120. Rasmussen AK, Chatterjee A, Rasmussen LJ, Singh KK. Mitochondria-mediated nuclear mutator phenotype in *Saccharomyces cerevisiae*. *Nucleic Acids Res*. 2003; 31: 3909–3917. <https://doi.org/10.1093/nar/gkg446> PMID: 12853606
121. Zhou JC, Janska A, Goswami P, Renault L, Abid Ali F, Kotecha A, et al. CMG–Pol epsilon dynamics suggests a mechanism for the establishment of leading-strand synthesis in the eukaryotic replisome. *Proc Natl Acad Sci*. 2017; 114: 201700530. <https://doi.org/10.1073/pnas.1700530114> PMID: 28373564
122. Douglas ME, Ali FA, Costa A, Diffley JFX. The mechanism of eukaryotic CMG helicase activation. *Nature*. Nature Publishing Group; 2018; 555: 265–268. <https://doi.org/10.1038/nature25787> PMID: 29489749
123. Campuzano V, Montermini L, Moltò MD, Pianese L, Cossée M, Cavalcanti F, et al. Friedreich's Ataxia: Autosomal Recessive Disease Caused by an Intronic GAA Triplet Repeat Expansion. *Science* (80-). 1996; 271: 1423 LP– 1427. <https://doi.org/10.1126/science.271.5254.1423> PMID: 8596916
124. Shishkin AA, Voineagu I, Matera R, Cherng N, Chernet BT, Krasilnikova MM, et al. Large-Scale Expansions of Friedreich's Ataxia GAA Repeats in Yeast. *Mol Cell*. Elsevier Inc.; 2009; 35: 82–92. <https://doi.org/10.1016/j.molcel.2009.06.017> PMID: 19595718
125. Ohshima K, Kang S, Larson JE, Wells RD. Cloning, characterization, and properties of seven triplet repeat DNA sequences. *J Biol Chem*. American Society for Biochemistry and Molecular Biology; 1996; 271: 16773–83. <https://doi.org/10.1074/jbc.271.28.16773> PMID: 8663377
126. Gacy AM, Goellner GM, Spiro C, Chen X, Gupta G, Bradbury EM, et al. GAA Instability in Friedreich's Ataxia Shares a Common, DNA-Directed and Intraallelic Mechanism with Other Trinucleotide Diseases. *Mol Cell*. Cell Press; 1998; 1: 583–593. [https://doi.org/10.1016/s1097-2765\(00\)80058-1](https://doi.org/10.1016/s1097-2765(00)80058-1) PMID: 9660942
127. Baptiste BA, Ananda G, Strubczewski N, Lutzkanin A, Khoo SJ, Srikanth A, et al. Mature Microsatellites: Mechanisms Underlying Dinucleotide Microsatellite Mutational Biases in Human Cells. G3

- Genes, Genomes, Genet. G3: Genes, Genomes, Genetics; 2013; 3: 451–463. <https://doi.org/10.1534/G3.112.005173> PMID: 23450065
128. Ohshima K, Montermini L, Wells RD, Pandolfo M. Inhibitory effects of expanded GAA.TTC triplet repeats from intron I of the Friedreich ataxia gene on transcription and replication in vivo. *J Biol Chem. American Society for Biochemistry and Molecular Biology*; 1998; 273: 14588–95. <https://doi.org/10.1074/jbc.273.23.14588> PMID: 9603975
 129. Krasilnikova MM, Mirkin SM. Replication stalling at Friedreich's ataxia (GAA)_n repeats in vivo. *Mol Cell Biol. American Society for Microbiology (ASM)*; 2004; 24: 2286–95. <https://doi.org/10.1128/MCB.24.6.2286-2295.2004> PMID: 14993268
 130. Pollard LM, Sharma R, Gómez M, Shah S, Delatycki MB, Pianese L, et al. Replication-mediated instability of the GAA triplet repeat mutation in Friedreich ataxia. *Nucleic Acids Res. Oxford University Press*; 2004; 32: 5962. <https://doi.org/10.1093/nar/gkh933> PMID: 15534367
 131. Gadgil R, Barthelemy J, Lewis T, Leffak M. Replication stalling and DNA microsatellite instability. *Bio-phys Chem. Elsevier B.V.*; 2017; 225: 38–48. <https://doi.org/10.1016/j.bpc.2016.11.007> PMID: 27914716
 132. Freudenreich CH, Kantrow SM, Zakian VA. Expansion and length-dependent fragility of CTG repeats in yeast. *Science (80-)*. 1998; 279: 853–856. <https://doi.org/10.1126/science.279.5352.853> PMID: 9452383
 133. Dixon MJ, Lahue RS. Examining the potential role of DNA polymerases eta and zeta in triplet repeat instability in yeast. *DNA Repair (Amst)*. 2002; 1: 763–70. Available: <http://www.ncbi.nlm.nih.gov/pubmed/12509280>
 134. Northam MR, Moore EA, Mertz TM, Binz SK, Stith CM, Stepchenkova EI, et al. DNA polymerases ζ and Rev1 mediate error-prone bypass of non-B DNA structures. *Nucleic Acids Res*. 2014; 42: 290–306. <https://doi.org/10.1093/nar/gkt830> PMID: 24049079
 135. Kunkel TA. Frameshift mutagenesis by eucaryotic DNA polymerases in vitro. *J Biol Chem*. 1986; 261: 13581–13587. PMID: 3759982
 136. Kunkel T a, Bebenek K. DNA replication fidelity. *Annu Rev Biochem*. 2000; 69: 497–529. <https://doi.org/10.1146/annurev.biochem.69.1.497> PMID: 10966467
 137. Streisinger G, Okada Y, Emrich J, Newton J, Tsugita A, Terzaghi E, et al. Frameshift mutations and the genetic code. This paper is dedicated to Professor Theodosius Dobzhansky on the occasion of his 66th birthday. *Cold Spring Harb Symp Quant Biol*. 1966; 31: 77–84. Available: <http://www.ncbi.nlm.nih.gov/pubmed/5237214> <https://doi.org/10.1101/sqb.1966.031.01.014> PMID: 5237214
 138. Polyzos AA, McMurray CT. Close encounters: Moving along bumps, breaks, and bubbles on expanded trinucleotide tracts. *DNA Repair (Amst)*. Elsevier; 2017; 56: 144–155. <https://doi.org/10.1016/j.dnarep.2017.06.017> PMID: 28690053
 139. Kunkel TA, Erie DA. Eukaryotic Mismatch Repair in Relation to DNA Replication. *Annu Rev Genet*. 2015; 49: 291–313. <https://doi.org/10.1146/annurev-genet-112414-054722> PMID: 26436461
 140. Strathern JN, Shafer BK, McGill CB. DNA synthesis errors associated with double-strand-break repair. *Genetics*. Genetics Society of America; 1995; 140: 965–72. Available: <http://www.ncbi.nlm.nih.gov/pubmed/7672595> PMID: 7672595
 141. Yang Y, Sterling J, Storici F, Resnick MA, Gordenin DA. Hypermutability of Damaged Single-Strand DNA Formed at Double-Strand Breaks and Uncapped Telomeres in Yeast *Saccharomyces cerevisiae*. Cohen-Fix O, editor. *PLoS Genet. Public Library of Science*; 2008; 4: e1000264. <https://doi.org/10.1371/journal.pgen.1000264> PMID: 19023402
 142. Malkova A, Haber JE. Mutations Arising During Repair of Chromosome Breaks. *Annu Rev Genet*. 2012; 46: 455–473. <https://doi.org/10.1146/annurev-genet-110711-155547> PMID: 23146099
 143. Gao Y, Mutter-Rottmayer E, Zlatanou A, Vaziri C, Yang Y. Mechanisms of post-replication DNA repair. *Genes (Basel)*. 2017; 8. <https://doi.org/10.3390/genes8020064> PMID: 28208741
 144. Kalimutho M, Bain AL, Mukherjee B, Nag P, Nanayakkara DM, Harten SK, et al. Enhanced dependency of KRAS-mutant colorectal cancer cells on RAD51-dependent homologous recombination repair identified from genetic interactions in *Saccharomyces cerevisiae*. *Mol Oncol. John Wiley & Sons, Ltd*; 2017; 11: 470–490. <https://doi.org/10.1002/1878-0261.12040> PMID: 28173629
 145. Anand RP, Tsaponina O, Greenwell PW, Lee C-S, Du W, Petes TD, et al. Chromosome rearrangements via template switching between diverged repeated sequences. *Genes Dev. Cold Spring Harbor Laboratory Press*; 2014; 28: 2394–406. <https://doi.org/10.1101/gad.250258.114> PMID: 25367035
 146. Halas A, Krawczyk M, Sledziewska-Gojska E. PCNA SUMOylation protects against PCNA polyubiquitination-mediated, Rad59-dependent, spontaneous, intrachromosomal gene conversion. *Mutat Res Mol Mech Mutagen. Elsevier*; 2016; 791–792: 10–18. <https://doi.org/10.1016/J.MRFMMM.2016.08.001> PMID: 27505077

147. Kim JC, Harris ST, Dinter T, Shah KA, Mirkin SM. The role of break-induced replication in large-scale expansions of (CAG)_n/(CTG)_n repeats. *Nat Struct Mol Biol.* Nature Publishing Group; 2017; 24: 55–60. <https://doi.org/10.1038/nsmb.3334> PMID: 27918542
148. Pel DM van, Stirling PC, Minaker SW, Sipahimalani P, Hieter P. *Saccharomyces cerevisiae* Genetics Predicts Candidate Therapeutic Genetic Interactions at the Mammalian Replication Fork. *G3 Genes, Genomes, Genet.* G3: Genes, Genomes, Genetics; 2013; 3: 273–282. <https://doi.org/10.1534/G3.112.004754> PMID: 23390603
149. Yeeles JTP, Marians KJ. The *Escherichia coli* replisome is inherently DNA damage tolerant. *Science* (80-). 2011; 334: 235–238. <https://doi.org/10.1126/science.1209111> PMID: 21998391
150. Heller RC, Marians KJ. Replisome assembly and the direct restart of stalled replication forks. *Nat Rev Mol Cell Biol.* Nature Publishing Group; 2006; 7: 932–943. <https://doi.org/10.1038/nrm2058> PMID: 17139333
151. Fumasoni M, Zwicky K, Vanoli F, Lopes M, Branzei D. Error-Free DNA Damage Tolerance and Sister Chromatid Proximity during DNA Replication Rely on the Polα/Primase/Ctf4 Complex. *Mol Cell.* Cell Press; 2015; 57: 812–823. <https://doi.org/10.1016/j.molcel.2014.12.038> PMID: 25661486
152. Hashimoto Y, Puddu F, Costanzo V. RAD51- and MRE11-dependent reassembly of uncoupled CMG helicase complex at collapsed replication forks. *Nat Struct Mol Biol.* Nature Publishing Group; 2012; 19: 17–24. <https://doi.org/10.1038/nsmb.2177> PMID: 22139015
153. Byun TS, Pacek M, Yee MC, Walter JC, Cimprich K a. Functional uncoupling of MCM helicase and DNA polymerase activities activates the ATR-dependent checkpoint. *Genes Dev.* 2005; 19: 1040–1052. <https://doi.org/10.1101/gad.1301205> PMID: 15833913
154. Hashimoto Y, Ray Chaudhuri A, Lopes M, Costanzo V. Rad51 protects nascent DNA from Mre11-dependent degradation and promotes continuous DNA synthesis. *Nat Struct Mol Biol.* Europe PMC Funders; 2010; 17: 1305–11. <https://doi.org/10.1038/nsmb.1927> PMID: 20935632
155. Branzei D, Psakhye I. DNA damage tolerance. *Curr Opin Cell Biol.* Elsevier Current Trends; 2016; 40: 137–144. <https://doi.org/10.1016/j.ceb.2016.03.015> PMID: 27060551
156. Neil AJ, Kim JC, Mirkin SM. Precarious maintenance of simple DNA repeats in eukaryotes. *BioEssays.* 2017; 39: 1–10. <https://doi.org/10.1002/bies.201700077> PMID: 28703879
157. Barkley LR, Song IY, Zou Y, Vaziri C. Reduced expression of GINS complex members induces hallmarks of pre-malignancy in primary untransformed human cells. *Cell Cycle.* 2009; 8: 1577–1588. <https://doi.org/10.4161/cc.8.10.8535> PMID: 19377277
158. Adams A, Gottschling DE, Kaiser CA, Stearns T. *Methods in Yeast Genetics: A Cold Spring Harbor Laboratory Course Manual.* Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press; 1997.
159. Amberg DC, Burke DJ, Strathern JN. *Methods in Yeast Genetics. A Cold Spring Harbor Laboratory Course Manual.* Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press; 2005.
160. Gietz RD, Woods RA. Transformation of Yeast by the Lithium Acetate/Single-Stranded Carrier DNA/PEG Method [Internet]. *Methods in Microbiology.* Elsevier; 1998. [https://doi.org/10.1016/S0580-9517\(08\)70325-8](https://doi.org/10.1016/S0580-9517(08)70325-8)
161. Sambrook J, Russell DW. *Molecular cloning: a laboratory manual.* 3rd ed. Cold Spring Harbor, N.Y.: Cold Spring Harbor Laboratory Press, c2001.; 2001.
162. Pavlov YI, Newlon CS, Kunkel T a. Yeast origins establish a strand bias for replicational mutagenesis. *Mol Cell.* 2002; 10: 207–213. [https://doi.org/10.1016/s1097-2765\(02\)00567-1](https://doi.org/10.1016/s1097-2765(02)00567-1) PMID: 12150920
163. Goldstein AL, McCusker JH. Three new dominant drug resistance cassettes for gene disruption in *Saccharomyces cerevisiae*. *Yeast.* 1999; 15: 1541–1553. [https://doi.org/10.1002/\(SICI\)1097-0061\(199910\)15:14<1541::AID-YEA476>3.0.CO;2-K](https://doi.org/10.1002/(SICI)1097-0061(199910)15:14<1541::AID-YEA476>3.0.CO;2-K) PMID: 10514571
164. Dmowski M, Gołębiewski M, Kern-Zdanowicz I. Characteristics of the Conjugative Transfer System of the IncM Plasmid pCTX-M3 and Identification of Its Putative Regulators. *J Bacteriol.* American Society for Microbiology Journals; 2018; 200: e00234–18. <https://doi.org/10.1128/JB.00234-18> PMID: 29986941
165. Dixon MJ, Bhattacharyya S, Lahue RS. Genetic Assays for Triplet Repeat Instability in Yeast. *Trinucleotide Repeat Protoc.* 2004; 277: 029–046. <https://doi.org/10.1385/1-59259-804-8:029>
166. Wierdl M, Dominska M, Petes TD. Microsatellite Instability in Yeast: Dependence on the Length of the Microsatellite. *Genetics.* 1997; 779: 769–779.
167. Drake JW. A constant rate of spontaneous mutation in DNA-based microbes. *Proc Natl Acad Sci.* 1991; 88: 7160–7164. <https://doi.org/10.1073/pnas.88.16.7160> PMID: 1831267
168. Krol K, Jendrysek J, Debski J, Skoneczny M, Kurlandzka A, Kaminska J, et al. Ribosomal DNA status inferred from DNA cloud assays and mass spectrometry identification of agarose-squeezed proteins interacting with chromatin (ASPIC-MS). *Oncotarget.* 2017; 5: 24988–25004. <https://doi.org/10.18632/oncotarget.15332> PMID: 28212567

169. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods*. NIH Public Access; 2012; 9: 671–5. <https://doi.org/10.1038/nmeth.2089> PMID: 22930834