

Weight management communications in idiopathic intracranial hypertension: challenges and recommendations from the patients' perspective

Sally Abbott,^{1,2} Amanda Denton,³ Sui H Wong,^{4,5,6,7} Susan P Mollan,^{8,9} Kim CM Bul ^{10,11}

To cite: Abbott S, Denton A, Wong SH, *et al*. Weight management communications in idiopathic intracranial hypertension: challenges and recommendations from the patients' perspective. *BMJ Neurology Open* 2023;5:e000527. doi:10.1136/bmjno-2023-000527

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjno-2023-000527>).

Received 15 September 2023
Accepted 05 November 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Kim CM Bul;
ac2658@coventry.ac.uk

ABSTRACT

Background Idiopathic intracranial hypertension (IIH) is a neurometabolic condition severely impacting the quality of life of people living with IIH (PwIIH). Most PwIIH are overweight or live with obesity, and weight loss is recommended by healthcare professionals (HCPs) as it is central to disease management. There is currently no research evaluating patient–clinician interactions when discussing weight management in IIH. The aim of this study was to evaluate the patient experience of communication with HCPs regarding weight management from the perspective of PwIIH.

Methods A cross-sectional online survey was developed and distributed by the IIH UK charity via their mailing list and social media network. Eligible participants were adults with IIH who have been recommended to lose weight by their HCP. Descriptive statistics were used to summarise quantitative responses and content analysis was used to inductively draw out themes from open-ended free-text responses.

Results There were 625 respondents. One-fifth of PwIIH (n=127/603, 21%) felt that HCPs were supportive and empathetic about weight management. Five themes were identified on how experiences regarding weight management for IIH can be improved, with PwIIH recommending for HCPs to: (1) detail the relationship between IIH and weight, (2) individualise care, (3) give advice, (4) provide support and (5) adapt communication.

Conclusion The majority of PwIIH recalled a poor experience and negative emotions when engaged in discussions regarding weight management with their HCPs. Further research should explore the HCPs perspective and evaluate interventions aiming to improve the quality of patient–HCPs communication in IIH.

INTRODUCTION

Idiopathic intracranial hypertension (IIH) is characterised by increased intracranial pressure leading to reported signs and symptoms such as headache, pulsatile tinnitus and papilloedema, with the potential risk of permanent visual loss.^{1,2} Although the exact causes of this condition remain unknown it occurs more often in women of childbearing

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Discussions about weight management are a sensitive topic for patients and evidence shows that healthcare professionals (HCPs) lack confidence in discussing body weight with patients. Priority setting research by the James Lind Alliance highlighted that weight management in the context of idiopathic intracranial hypertension (IIH) is a top research priority for patients and HCPs. To date, there is no published research evaluating the experiences of weight management interactions from the perspective of people living with IIH (PwIIH).

WHAT THIS STUDY ADDS

⇒ This is the first study evaluating the experiences of weight management interactions with HCPs from the perspective of PwIIH. The results of this study demonstrate that PwIIH feel unsupported and experience negative emotions when weight management discussions are initiated by their HCPs. This study also provides unique insights into how HCPs could improve their approach to weight management; by explaining the relationship between IIH and weight, providing personalised care, giving tailored advice, providing support and adapting communication style and the setting of communication.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ The results of this study provide for the first time the experience of PwIIH and their recollection of weight management discussions with HCPs and they offer suggestions that would make the interactions more effective. This research highlights the need for specific training for HCPs in the context of weight management for IIH to support improved healthcare accessibility, promote non-stigmatising practices and improve both patient and clinician satisfaction in this context.

age compared with men and children as well as in people who are overweight (body mass index, BMI, 25–30 kg/m²) or who live with obesity (BMI≥30 kg/m²).³ The global annual

IIH incidence rates range from 0.03 to 2.36 per 100 000 and are positively associated with country-specific obesity rates.⁴ In the UK, there has been a stepwise increase in the incidence and prevalence of IIH with increasing numbers of people being admitted to hospital care.^{15–7}

The use of specialised healthcare by people living with IIH (PwIIH) causes a significant economic burden on the National Health Services (NHS) with estimated direct healthcare resource use costs of £462 million per annum by 2030.⁵ This has also been demonstrated in the USA.⁸ Additionally, research states that PwIIH experience low quality of life mainly due to their headache symptoms⁹; however, research into other factors contributing to this is limited. This stresses the importance of more research into effective and acceptable weight management interventions for PwIIH.

There is a striking association with increased BMI and moderate weight gain.^{1 4 6 10} The disease is modified by weight loss and a recent randomised controlled trial evaluating bariatric surgery and a multicomponent lifestyle intervention found that weight loss mirrored reduction in intracranial pressure.^{11 12} A weight loss in the region of between 3% and 24% has been shown to reduce symptoms and lead to remission of the disease whereas regaining 6% of body weight has been shown to be associated with recurrence of IIH among some PwIIH as demonstrated by one study.^{2 12 13} The condition is managed by neurologists, ophthalmologists, neurosurgeons and more recently interventional radiologists, none of whom have formal training in weight management.¹⁴

A recent qualitative systematic literature review highlighted that weight stigma within patient–healthcare professionals (HCPs) interactions has a negative impact on healthcare access and quality of healthcare provision from the perspective of people living with obesity.¹⁵ People who have obesity perceive there to be negativity around their weight status based on the language used by HCPs and often feel ashamed, humiliated and blame themselves for having obesity. At the same time, due to a lack of knowledge base, skill set and guidelines, HCPs experience many challenges when communicating about weight with people who have obesity. There is a clear unmet clinical need for education and specialised training among HCPs on how to communicate about body weight change and weight status. This is important as HCP interactions about weight management can negatively affect future weight trajectory and the dialogue with people who have obesity.¹⁶ Although HCPs specialising in neurology are medically trained to diagnose PwIIH, they are not likely to have received training in obesity management. This could potentially lead to ineffective consultations where the importance of sustained weight reduction is not addressed or managed effectively.

Patient and public involvement and engagement in research and healthcare practices is becoming increasingly important to generate meaningful healthcare impact.¹⁷ This has been reflected in a recent research study in which a priority setting exercise was performed

with PwIIH and HCPs in accordance with the James Lind Alliance Priority Setting Partnership.¹⁸ There is limited evidence in IIH where patients are central to the conduct of the research, and a previous study has shown the benefit of this close working relationship.¹⁹ Research into the role of weight management in PwIIH was highlighted as a research priority, including approaches that provide sensitivity in discussing weight and address the stigma associated with obesity.^{2 18} Despite this, there are currently no published research studies on the experiences of PwIIH when HCPs communicate about the role of body weight and weight management in IIH. Hence, the aim of this study was to evaluate the experiences of PwIIH in their interactions with HCPs regarding weight management.

METHODS

Study design and participants

A cross-sectional study was undertaken and collected data using the SurveyMonkey platform. The survey was open to responses between 26 April 2021 and 14 May 2021. Eligible participants were people with a diagnosis of IIH, aged over the age of 18 years old, and who had been advised to lose weight by any HCP.

Survey development

The questionnaire was developed in response from patients citing concerns around the patient–HCPs weight management interactions as expressed by PwIIH across diverse social media groups. A steering committee of Trustees of the charity IIH UK, which included two PwIIH, led on the questionnaire development. The questions within the survey were formulated based on patient experiences voiced to the charity IIH UK by PwIIH, primarily via Facebook and Twitter social media channels. The survey was piloted with three PwIIH which led to minor modifications to the questionnaire wording to aid the use of plain English language readability (see online supplemental file 1).

Data collection

The survey was distributed to 395 members of the charity IIH UK via their membership mailing list. The survey was also distributed via social media channels, including the ‘IIH UK National Charity’ Facebook page and the @IIHUK Twitter handle. All responses were anonymous, and questions permitted both closed and open responses from participants, giving participants the opportunity to expand on their given answer in their own words. Participants were able to answer all the questions however no single question was mandated to be filled out. Additionally, the participants could select one or more options at certain questions. IP addresses for each response were cross-checked to ensure that responses were submitted only by PwIIH residing in the UK.

Data analysis

Responses to closed questions were analysed using frequency descriptive statistics (n, %). Open free-text

Table 1 Quotation excerpts from unique participants recalling weight loss targets set by the HCP and whether they did or did not feel this was realistic

Realistic weight loss goals	Unrealistic weight loss goals
'Was asked to lose 5% and achieved 7%.'	'It sounded like a big number. 13% of your body weight sounds huge.'
'The doctors told me to create my own goals and they should not be super goals or unachievable in a short period. So I am managing smaller goals one at a time.'	'It was the lowest end of healthy BMI and it just seemed impossible that my body would ever be that small.'
'Approx 1 kg per week of steady loss. Calorie reduction of approx 100–200 per day.'	'I was told to lose at least 10% of my body weight within a few months. No support or help was given to help me achieve that.'
'Between 5% and 10% of my body weight in a year.'	'10st is just so huge, it feels like a mountain I can't climb.'
'I was told 10% of my original weight was often enough to see improvement. In reality, I had to lose a lot more than that but 10% was quite easy to achieve.'	'Yes my initial goal was 10% which I achieved but it made no difference to symptoms so I was told to keep going until it works. Now no goal and no end in sight.'
	'They kept moving it... 5%, 6%–10% 12%, then I didn't lose it fast enough.'
	'Got told to lose 15 stone I am no good with massive goals like that.'
	'He just looked at me and said I needed to lose a lot of weight. Wasn't even weighed.'
	'The goal I was given was not realistic as it is what I weighed 20 years ago when I was a 15 year old adolescent.'
	'I was told 10% of my body weight. I have lost quite a bit more than that, but yet my doctor's still insist if I lost more weight my symptoms would improve.'

BMI, body mass index; HCP, healthcare professional.

(figure 2). Table 1 shows participant quotation excerpts illustrating the weight loss targets advised by their HCP and whether they felt the target was realistic or not.

Improving the weight management patient experience

Online supplemental file 2 outlines the 5 categories and 18 subcategories that were derived from free-text responses to questions about how the experience of treating IiH via weight management could be improved from the perspective of PwIiH, alongside illustrative quotations.

The relationship between IiH and weight

PwIiH wished that HCPs explore the relationship between IiH and excess weight and to 'provide an explanation' of the biological mechanisms of excess weight that may cause or exacerbate symptoms of IiH; but to also 'acknowledge weight as a risk factor, not a cause' of IiH. PwIiH referred to the condition itself being named idiopathic (meaning of 'unknown cause'), thus perceiving that IiH cannot be caused by excess weight alone. And they reasoned that not all PwIiH have excess weight and that comparatively very few women of childbearing age with obesity in the population have IiH, either. There was a sense of scepticism among PwIiH around the relationship between IiH and excess weight and, therefore, wanted HCPs to 'Provide evidence' and 'actual scientific proof' to them.

Individualising care

PwIiH felt that weight management discussions should be more individualised. HCPs could 'consider the impact of IiH symptoms' when providing any lifestyle advice for weight management; citing that symptoms of pain and fatigue were a challenge to adopting helpful physical activity and eating behaviours. Some PwIiH expressed that HCPs should 'investigate comorbidities' aside from IiH, as barriers to their weight management. PwIiH also felt that HCPs should 'explore mental health' since they felt that the symptoms of IiH contributed to mental illness and vice versa that mental illness presented challenges for their weight management. To provide individualised care, PwIiH expressed that HCPs should 'get to know the 'person' without making assumptions and expressed that they did not want to be 'treated like a number'.

Giving advice

When being given advice on weight management, PwIiH felt that HCPs should acknowledge that IiH is 'complex' to self-manage and that HCPs should 'avoid oversimplistic advice of 'eat less, move more'. PwIiH wanted HCPs to 'provide realistic weight loss targets' which should be quantified both in terms of amount of weight loss and specified time frame.

Providing support

PwIiH wished for improved weight management support to be provided by HCPs. PwIiH wanted HCPs to 'discuss

the available options' with them, which would contribute to PwIIH feeling more 'in control' of their care and solutions were being offered. PwIIH did not expect that the HCP managing their IIH should be expected to provide isolated weight management support and wanted their HCP to 'organise referrals to other HCPs', in particular to weight management services or HCPs trained in lifestyle management (eg, dietitians, exercise physiologists). PwIIH expressed that HCPs could also 'provide resources for self-management' in the form of leaflets which offered 'practical' support.

Communication

PwIIH felt that HCPs could adapt their communication style and approach when discussing weight management. PwIIH felt that HCPs should 'ask for permission' before discussing or measuring weight. PwIIH felt HCPs should 'Show empathy' and alter their choice of language when discussing weight management; including 'use of people-first language' not using terms such as 'obese' and 'avoid using shaming language' which could result in PwIIH feeling 'blamed' for their illness. PwIIH wished for HCPs to 'engage in a two-way dialogue' about their excess weight as part of shared decision-making, rather than being 'talked at'. The setting in which weight management conversations took place was important to PwIIH. They expressed that HCPs should 'ensure privacy' and not have weight management discussions where privacy could not be maintained, such as in inpatient settings, or vocalising in waiting rooms what investigation they were being called in for that is, to be weighed.

DISCUSSION

This study provides novel insights into the experiences of PwIIH regarding their current, mostly negative, interactions with HCPs about body weight and weight management. It offers suggestions from PwIIH on how the communication of weight management could be improved in the context of IIH disease management. Despite significant evidence demonstrating that weight loss is an effective treatment for IIH¹⁰⁻¹² we found that only a fifth of PwIIH recalled being offered support for weight management and only 16% of PwIIH found the support to be 'helpful and appropriate'. This evidence demonstrates an unmet clinical need to provide training and support in regard to approaching weight management conversations and onward referrals to weight management service, where appropriate, for HCPs who care for PwIIH.

PwIIH expressed that it was important to them that the HCP should first ask their permission to discuss weight. However, less than 1 in 10 (7%) of PwIIH in our study recalled being asked for permission first. Asking questions, rather than issuing statements, is a core principle of motivational interviewing which has been shown to facilitate patient-driven behaviour change³² and is recommended

in fields such as pain neuroscience to enhance a patient's receptiveness to advice and education.²³

The way in which HCPs communicated about weight management invoked a negative emotive response among PwIIH, with most reporting feelings of self-blame and 'fault' for having IIH due to their weight. Research has shown that providing a biological explanation for disease processes is effective at reducing disease-related stigma²⁴ and shifting the focus from personal blame to physiological mechanisms can be an effective means to avoid using stigmatising language and inciting blame.²⁵

Setting targets and goals is considered an important behavioural change technique in weight management. In our study, PwIIH expressed that they found weight targets to be helpful for their weight management; however, most PwIIH recounted that the targets set were unrealistic to attain. Providing targets for weight loss in IIH are yet to be determined, as while most studies recommend a level at which the disease remits, none have yet to assess this as their primary aim. Unrealistic weight loss targets are cited in the literature as a barrier to weight management in the general obesity population,²⁶ with the 'false hope' hypothesis suggesting that very ambitious targets are less likely to be met and leads to disappointment, dissatisfaction and decreased effort.²⁷ On the contrary, systematic review data have found that realistic and specific goal setting is effective for weight management among people with obesity.²⁸

Our findings also provide suggestions to improve clinical communication from the perspective of PwIIH. More effective weight management encounters as part of treatment for IIH could be enabled by HCPs explaining the evidence of the relationship between IIH and weight, individualising care, giving advice, providing support and adapting communication. This could potentially contribute to better treatment adherence and improve self-management practices as seen in chronic conditions, including obesity.^{29 30}

Our study has several strengths. Embedding active patient and public involvement ensured the research was relevant to service users and user-friendly survey question.^{19 31} Although it is unknown how many PwIIH have been exposed to the online survey, the method of survey distribution via IIH UK charity platforms meant that a large UK sample of PwIIH completed the survey. However, this study also has some limitations. Due to the nature of the survey distribution, it was not possible to calculate a response rate. For future occurrences, the online survey could be posted on IIH UK charity webpage so the number of views and responses can be tracked to calculate a response rate. Although this was an anonymous survey which may have contributed to a higher response rate, there were no data on participant characteristics and therefore it is not certain how representative the study sample is of the target population. As the survey was distributed online, people without access to the internet and people with lower levels of digital literacy are likely not have been included.³² As the responses were collected

retrospectively of healthcare encounters, results are likely to be susceptible to a recall bias, and no question inquired about the time between the consultation and completion of the questionnaire. The richness of the qualitative data was limited by responses being provided only in written form and therefore semistructured interviews would be recommended for future research.³³

Further research could focus on examining experiences from both the perspective of PwIIH and the HCP and how this can affect the patient–HCPs relationship, accessibility and ultimately the health outcomes. Increasing the knowledge base for both PwIIH and HCPs regarding the role of weight in IIH is important and seen as a priority.¹⁸ Understanding of the actual pathways and pathogenic role of body weight and weight gain in IIH will help HCPs discuss this aspect of the disease confidently in IIH and reduce HCP weight stigmatisation.¹⁰ Effective communication on this challenging topic will help make PwIIH feeling more ‘heard’ and ‘accepted’.

CONCLUSION

This study showed many PwIIH experience negative emotions when HCPs communicate with them about their body weight, its role in IIH and how to manage weight reduction. They call for improvements to be made to the weight management experience in the context of IIH. Further research is needed to understand the HCP perspective on body weight and weight management communication in IIH. Targeted training and ready access to weight management services are required to address weight stigma and facilitate effective communication between HCPs and PwIIH.

Author affiliations

¹Institute for Health and Wellbeing, Centre for Healthcare and Communities, Coventry University, Coventry, UK

²Department of Dietetics, University Hospitals Coventry and Warwickshire NHS Trust, Coventry, UK

³IIH UK Registered Charity, Washington, UK

⁴Department of Neuro-Ophthalmology, Moorfields Eye Hospital NHS Foundation Trust, London, UK

⁵Medical Eye Unit, Guy's and St Thomas' Hospitals NHS Trust, London, UK

⁶Faculty of Life Sciences and Medicine, King's College London, London, UK

⁷Institute of Neurology, Department of Clinical and Movement Neuroscience, University College London, London, UK

⁸Birmingham Neuro-Ophthalmology, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK

⁹Institute of Metabolism and Systems Research, Translational Brain Science, University of Birmingham, Birmingham, UK

¹⁰Institute for Health and Wellbeing, Centre for Intelligent Healthcare, Coventry University, Coventry, UK

¹¹Renal Services, University Hospitals Coventry and Warwickshire NHS Trust, Coventry, UK

Acknowledgements The authors would like to thank the charity IIH UK for initiating and performing this study from the perspective of PwIIH and for supporting this study financially. The authors would also like to thank PwIIH for their willingness to complete the online survey.

Contributors All authors contributed equally to the manuscript with the following specific contributions: SA—writing methodology and result sections, data analysis, review final manuscript and provide intellectual contribution; AD—

conceptualisation and methodology, review final manuscript and provide intellectual contribution; SHW—writing, full manuscript review and intellectual contribution; SPM—writing, full manuscript review and intellectual contribution; KCMB—writing introduction and discussion sections, review final manuscript and provide intellectual contribution. SA acts as guarantor and takes full responsibility for the work and conduct of the study, had access to the data and controlled the decision to publish. All authors read and approved the final manuscript.

Funding This study was funded by the IIH UK charity (grant/award number N/A).

Competing interests All authors have completed the ICMJE uniform disclosure form and declared not having any competing interests.

Patient and public involvement Two PwIIH have been involved in the development of the online survey and questions have been formulated based on the experiences of PwIIH as collected by charity IIH UK. The charity IIH UK represents the voice of PwIIH and distributed the online survey through their mailing list and social media channels. Three PwIIH piloted the online survey before it was finalised. The coauthor AD is a patient with the lived experience of IIH and reviewed and contributed to the current manuscript.

Patient consent for publication Not applicable.

Ethics approval Approval was not required from the Health Research Authority (HRA) nor National Health Service (NHS) Research Ethics Committee (REC) as participants were not recruited through NHS sites. Participants of the survey were made aware of the intended use of their data before filling out the survey, and therefore, gave their consent by filling the survey out. All data collected was anonymised.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available. The data are not shareable due to the qualitative nature of data collection, in order to maintain participant anonymity.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Kim CM Bul <http://orcid.org/0000-0001-8984-9921>

REFERENCES

- Miah L, Strafford H, Fonferko-Shadrach B, *et al*. Incidence, prevalence and healthcare outcomes in idiopathic intracranial hypertension: a population study. *Neurology* 2021;96:e1251–61.
- Mollan SP, Grech O, Alimajstorovic Z, *et al*. New horizons for idiopathic intracranial hypertension: advances and challenges. *Br Med Bull* 2020;136:118–26.
- Mollan SP, Ali F, Hassan-Smith G, *et al*. Evolving evidence in adult idiopathic intracranial hypertension: pathophysiology and management. *J Neurol Neurosurg Psychiatry* 2016;87:982–92.
- McCluskey G, Doherty-Allan R, McCarron P, *et al*. Meta-analysis and systematic review of population-based epidemiological studies in idiopathic intracranial hypertension. *Eur J Neurol* 2018;25:1218–27.
- Mollan SP, Aguiar M, Evison F, *et al*. The expanding burden of idiopathic intracranial hypertension. *Eye (Lond)* 2019;33:478–85.
- Adderley NJ, Subramanian A, Nirantharakumar K, *et al*. Association between idiopathic intracranial hypertension and risk of cardiovascular diseases in women in the United Kingdom. *JAMA Neurol* 2019;76:1088–98.
- Mollan SP, Mytton J, Tsermoulas G, *et al*. Idiopathic intracranial hypertension: evaluation of admissions and emergency readmissions

- through the hospital episode statistic dataset between 2002–2020. *Life (Basel)* 2021;11:417.
- 8 Friesner D, Rosenman R, Lobb BM, *et al.* Idiopathic intracranial hypertension in the USA: the role of obesity in establishing prevalence and healthcare costs: prevalence and healthcare costs in IIH. *Obes Rev* 2011;12:e372–80.
 - 9 Mulla Y, Markey KA, Woolley RL, *et al.* Headache determines quality of life in idiopathic intracranial hypertension. *J Headache Pain* 2015;16:45:521..
 - 10 Mollan SP, Tahrani AA, Sinclair AJ. The potentially modifiable risk factor in idiopathic intracranial hypertension: body weight. *Neurol Clin Pract* 2021;11:e504–7.
 - 11 Mollan SP, Mitchell JL, Ottridge RS, *et al.* Effectiveness of bariatric surgery vs community weight management intervention for the treatment of idiopathic intracranial hypertension: a randomized clinical trial. *JAMA Neurol* 2021;78:678–86.
 - 12 Mollan SP, Mitchell JL, Yiangou A, *et al.* Association of amount of weight lost after bariatric surgery with intracranial pressure in women with idiopathic intracranial hypertension. *Neurology* 2022;99:e1090–9.
 - 13 Ko MW, Chang SC, Ridha MA, *et al.* Weight gain and recurrence in idiopathic intracranial hypertension a case-control study. *Neurology* 2011;76:1564–7.
 - 14 Mollan SP, Sinclair AJ, Tsermoulas G. Letter: cerebrospinal fluid shunting for idiopathic intracranial hypertension: a systematic review, meta-analysis, and implications for a modern management protocol. *Neurosurgery* 2023;92:e59–60.
 - 15 Ryan L, Coyne R, Heary C, *et al.* Weight stigma experienced by patients with obesity in healthcare settings: a qualitative evidence synthesis. *Obes Rev* 2023;24:e13606.
 - 16 Auckburally S, Davies E, Logue J. The use of effective language and communication in the management of obesity: the challenge for healthcare professionals. *Curr Obes Rep* 2021;10:274–81.
 - 17 Crocker JC, Boylan A-M, Bostock J, *et al.* Is it worth it? Patient and public views on the impact of their involvement in health research and its assessment: a UK-based qualitative interview study. *Health Expect* 2017;20:519–28.
 - 18 Mollan S, Hemmings K, Herd CP, *et al.* What are the research priorities for idiopathic intracranial hypertension? A priority setting partnership between patients and healthcare professionals. *BMJ Open* 2019;9:e026573.
 - 19 Scotton WJ, Mollan SP, Walters T, *et al.* Characterising the patient experience of diagnostic lumbar puncture in idiopathic intracranial hypertension: a cross-sectional online survey. *BMJ Open* 2018;8:e020445.
 - 20 Hsieh H-F, Shannon SE. Three approaches to qualitative content analysis. *Qual Health Res* 2005;15:1277–88.
 - 21 Kondracki NL, Wellman NS, Amundson DR. Content analysis: review of methods and their applications in nutrition education. *J Nutr Educ Behav* 2002;34:224–30.
 - 22 M V, H P, AM S, *et al.* n.d. Minimal intervention for obesity counseling in primary care.
 - 23 Nijs J, Wijma AJ, Willaert W, *et al.* Integrating motivational interviewing in pain neuroscience education for people with chronic pain: a practical guide for clinicians. *Phys Ther* 2020;100:846–59.
 - 24 Doley JR, Hart LM, Stukas AA, *et al.* Interventions to reduce the stigma of eating disorders: a systematic review and meta-analysis: doley *et al.* *Int J Eat Disord* 2017;50:210–30.
 - 25 Fearon N, Sudlow A, le Roux CW, *et al.* Say what you mean, mean what you say: the importance of language in the treatment of obesity. *Obesity (Silver Spring)* 2022;30:1189–96.
 - 26 Wamsteker EW, Geenen R, Zelissen PMJ, *et al.* Unrealistic weight-loss goals among obese patients are associated with age and causal attributions. *J Am Diet Assoc* 2009;109:1903–8.
 - 27 Polivy J, Herman CP. The false-hope syndrome: unfulfilled expectations of self-change. *Curr Dir Psychol Sci* 2000;9:128–31.
 - 28 Pearson ES. Goal setting as a health behavior change strategy in overweight and obese adults: a systematic literature review examining intervention components. *Patient Educ Couns* 2012;87:32–42.
 - 29 Iacobucci G. Treat obesity with the same effort used to reduce smoking, says lead psychologist. *BMJ* 2019;l5713.
 - 30 Fernandes JB, Teixeira F, Godinho C. Personalized care and treatment compliance in chronic conditions. *J Pers Med* 2022;12:737.
 - 31 Brett J, Staniszewska S, Mockford C, *et al.* Mapping the impact of patient and public involvement on health and social care research: a systematic review. *Health Expect* 2014;17:637–50.
 - 32 Datta A, Bhatia V, Noll J, *et al.* Bridging the digital divide: challenges in opening the digital world to the elderly, poor, and digitally illiterate. *IEEE Consumer Electron Mag* 2019;8:78–81.
 - 33 Reja U, Manfreda KL, Hlebec V, *et al.* Open-ended vs. close-ended questions in web questionnaires; 2003.