



Lung Hyperinflation as Treatable Trait in Chronic Obstructive Pulmonary Disease: A Narrative Review

Maud Koopman ¹⁻³, Rein Posthuma ¹⁻³, Lowie EGW Vanfleteren ⁴, Sami O Simons ^{2,3}, Frits ME Franssen ¹⁻³

¹Research and Development, Ciro+, Horn, the Netherlands; ²NUTRIM, Institute of Nutrition and Translational Research in Metabolism, University Maastricht, Maastricht, the Netherlands; ³Department of Respiratory Medicine, Maastricht University Medical Center (MUMC+), Maastricht, the Netherlands; ⁴COPD Center, Institute of Medicine, Sahlgrenska University Hospital, University of Gothenburg, Gothenburg, Sweden

Correspondence: Frits ME Franssen, Ciro+, Hornerheide 1, Horn, 6085, NM, the Netherlands, Email fritsfranssen@ciro-horn.nl

Abstract: Lung hyperinflation (LH) is a common clinical feature in patients with chronic obstructive pulmonary disease (COPD). It results from a combination of reduced elastic lung recoil as a consequence of irreversible destruction of lung parenchyma and expiratory airflow limitation. LH is an important determinant of morbidity and mortality in COPD, partially independent of the degree of airflow limitation. Therefore, reducing LH has become a major target in the treatment of COPD over the last decades. Advances were made in the diagnostics of LH and several effective interventions became available. Moreover, there is increasing evidence suggesting that LH is not only an isolated feature in COPD but rather part of a distinct clinical phenotype that may require a more integrated management. This narrative review focuses on the pathophysiology and adverse consequences of LH, the assessment of LH with lung function measurements and imaging techniques and highlights LH as a treatable trait in COPD. Finally, several suggestions regarding future studies in this field are made.

Keywords: COPD, hyperinflation, treatable trait, emphysema, phenotype

Introduction

Chronic obstructive pulmonary disease (COPD) has in recent decades been recognized as a complex condition consisting of more than airflow limitation alone. Both additional pulmonary as well as extra-pulmonary features contribute to the burden of the disease.¹⁻³ The concept of identifying treatable traits in individual patients was a step forward in addressing the complexity and heterogeneity of COPD.⁴ Lung hyperinflation (LH) is considered a pulmonary trait for which diagnostic criteria and specific interventions are available.⁴ Indeed, LH, resulting from destruction of lung parenchyma and loss of lung elastic recoil, is one of the hallmarks of the disease.⁵ Its partial independence from the degree of airflow limitation^{1,6} and its strong link with morbidity⁷ and mortality,⁸ combined with the availability of targeted treatments, justify its status as a treatable trait. However, rather than an isolated trait, LH may be part of a broader phenotype of the disease, with typical pulmonary features, symptoms, disease trajectory, extra-pulmonary features and comorbidities.⁹

This narrative review focusses on the pathophysiology and adverse consequences of LH, its assessment and position as a treatable trait in COPD. Identified knowledge gaps are summarized as suggestions for future studies.

Definition, Mechanisms and Pathophysiology

Hyperinflation is defined by an increase in the amount of gas in the lungs and airways at the end of a spontaneous expiration, amounting to a functional residual capacity (FRC) above the upper 95th percentile of the predicted values.^{10,11} LH can most accurately be defined as an increase in total lung capacity (TLC) above the upper limit of normal, with an FRC/TLC or residual volume (RV)/TLC above the upper limit of normal.¹² An increase in TLC >120%

of the predicted value or an increase in RV above the upper limit of normal are also often used as definitions of LH in literature.¹³

Static Hyperinflation

Static LH, defined by increased resting lung volumes, is determined by a modification of the elastic properties of the respiratory system due to a decrease of the inward elastic recoil of the lungs without changes of the natural outward elastic properties of the chest wall.¹⁴ Pulmonary emphysema, characterized by destruction of lung parenchyma and loss of elastic fibers in the lungs, is the most important cause of LH.¹⁵ This results in a displacement of the volume–pressure curve of the lungs leftward and upward (Figure 1).^{13,16} At the end of a spontaneous expiration, the lung compensates for a reduced recoil pressure by acquiring a larger volume than normal to balance the outward recoil of the chest wall in patients with COPD. The static/elastic equilibrium point of the total respiratory system is moved towards a higher-than-normal lung volume, and the respiratory system's relaxation volume or FRC is permanently increased.

RV in turn is increased in COPD, not only due to reduced elastic recoil but also due to premature closure of the small airways and expiratory airflow limitation at low lung volume, also known as gas trapping.¹⁰ This is already observed in COPD patients with mild airflow limitations.^{17,18} Moreover, a reduced duration for expiration due to the development of unendurable respiratory discomfort contributes to the increase in RV.¹³ TLC also increases due to loss of lung recoil. With increasing severity of the disease, the rise in RV becomes greater than the increase in TLC, because TLC is limited by the thoracic cage. This, together with a fall in vital capacity (VC) due to premature airway closure, will lead to a higher RV/TLC ratio, which also impacts on a further reduction in forced expiratory volume in the first second (FEV₁). In these severe cases, spirometry in COPD can show not only an obstructive but also a restrictive pattern.¹⁹

While the degree of airflow limitation and static hyperinflation are directly correlated in most patients,^{1,19} some patients have distinct hyperinflation without severe airflow limitation,¹ as is also shown in Figure 2. The figure shows a self-organizing map of 467 patients with stable COPD that participated in the Chance Study: an observational,




Figure 1 Change in end-expiratory lung volume (EELV) with COPD. EELV is set at the point at which the elastic recoil pressures of the lung and chest wall are equal and opposite in direction. In COPD patients with hyperinflation, emphysema decreases lung elastic recoil pressure and causes a reset of functional residual capacity, or EELV, at a higher absolute lung volume. The difference between expected (long-dashed horizontal lines) and observed EELV (short-dashed horizontal lines) represent static hyperinflation. Reprinted from Dubé B, Guerdier, (A), Morelot-Panzini, (C) et al. The clinical relevance of the emphysema-hyperinflated phenotype in COPD. *COPD Research and Practice*. 2015;2(1). This source is Open Access, the figure was copied under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>).¹⁶




Figure 2 Heterogeneity of lung function impairment in patients with COPD.

Notes: The panels were generated using Viscosity (Viscosity Software GmbH, Vienna, Austria). The Viscosity program placed all patients in a specific position on the map based on their profile of a comprehensive lung function assessment. When looking at the different lung function attributes, patients appear red on the chart when the attribute is relatively high within this sample, appear green if the attribute is moderate and appear blue when the attribute is relatively low within this sample. The more patients show resemblance in terms of their overall lung function, the closer they are on the map. By drawing lines on the map, the Viscosity program could identify eight different clusters of patients with COPD with significantly different lung function profile. AL+: airflow limitation ($FEV_1 < 60\%$ predicted); EMF+: emphysema ($KCO < 60\%$ predicted); LH+: lung hyperinflation ($RV > 120\%$ predicted); -: not fulfilling any of these criteria for severity of lung function impairment (neither airflow limitation, emphysema or lung hyperinflation).

prospective, single-center study about COPD, health status and cardiovascular co-morbidities (METC 11-3-070).²⁰ Patients were ordered based on the absence or presence of airflow limitation (AL): ($FEV_1 < 60\%$ predicted); emphysema (EMF): carbon monoxide transfer coefficient (KCO) $< 60\%$ predicted; and lung hyperinflation (LH): $RV > 120\%$ predicted. Eight clusters could be identified with a significantly different lung function profile. In this cohort, 14% of

the patients had LH without significant expiratory airflow limitation. Moreover, 8% of the patients had emphysema without significant lung hyperinflation. Hence, although most COPD patients have a combination of AL, LH and EMF, these data implicated that the presence of LH does not always mean coexistence of airflow limitation or emphysema in patients with COPD.

Dynamic Hyperinflation

Dynamic hyperinflation (DH) is defined as a temporary increase of end expiratory volume (EELV) above its resting value when ventilation increases in patients with expiratory airflow limitation.¹³ It results from a discrepancy between the time required for the lungs to empty during exhalation and the time available between two consecutive inspiratory efforts. Reduced expiratory airflow limits the exhaled volume over time, while increased breathing frequency shortens the time available to expire. Several factors can further worsen DH: high inspired volume combined with insufficient expiration, ie during exercise; high frequency of breathing, eg during spontaneous breathing, exercise, anxiety or panic attacks; decreased driving pressure for expiration; increased airflow resistance, during exacerbation; expiratory airflow limitation, when small airways become excessively compressed due to destruction of supporting alveolar attachments.²¹

Although most COPD patients have DH during exercise, Guenette et al confirmed that approximately 15–20% of patients with moderate-to-severe COPD do not, even with only slightly less resting hyperinflation.²² It was suggested they counteract DH by contracting the expiratory muscles, although DH can occur during exercise despite increased expiratory muscle activity.²³

Airway Remodeling

It has been well established that small conducting airways less than 2 mm in diameter, which account for less than 10% of the total airflow resistance in normal lungs,²⁴ become the major site of airflow obstruction in COPD.²⁵ The terminal bronchioles are also known to disappear in patients with very severe COPD, and it is suggested that this destruction might precede emphysematous tissue destruction in COPD.²⁶ Koo et al performed micro-CT's and histological assessments of the matching lung tissue and showed that the lungs of patients with mild and moderate COPD have significantly fewer conducting terminal bronchioles and respiratory transitional bronchioles compared with age-matched smokers with normal lung function, and that this loss is also present in apparently non-emphysematous lungs.²⁷ However, since Koo et al performed cross-sectional analyses it is unclear if this is due to abnormal development of the lungs or due to destruction of airways. Imaging studies using parametric response mapping to assess gas trapping resulting from small airways disease showed that this is also the dominant cause of hyperinflation in mild to moderate COPD.²⁸ Other studies demonstrate that loss of elastic recoil due to emphysematous destruction potentially causes further airway obstruction,²⁹ which in turn can lead to LH. Acknowledgement of these findings is important in light of the development of effective treatments and interpreting outcomes of clinical trials in COPD. By inclusion of advanced COPD populations, most previous trials studied the effects of treatments on lungs with substantially fewer terminal and transitional bronchioles. To develop disease-modifying treatments for COPD, an understanding of disease pathogenesis within these small airways and in patients without clinically apparent (or with very early disease) is essential.

Diagnosing Static and Dynamic Hyperinflation

The most commonly used methods for lung volume measurements are body plethysmography, gas dilution techniques (helium dilution method and nitrogen washout method) and imaging (computed tomography).³⁰ Magnetic resonance imaging (MRI) can also be used but appears to be more suitable for functional imaging.³¹

Measuring Static Hyperinflation by Lung Function Techniques

Body plethysmography is the reference method for the measurement of lung volume and measures intrathoracic gas volume (ITGV), FRC, and allows calculation of RV and TLC.^{32,33} FRC measured by plethysmography can overestimate the true FRC in the presence of abdominal gas or by significant pressure losses between alveoli and airway opening in the presence of severely obstructed airways.³⁴ Moreover, in plethysmography mouth pressure and alveolar pressure are assumed to be equal, which is not the case in severe COPD. This can lead to overestimation of lung volumes.¹² Both RV/

TLC ratio and RV have recently been demonstrated to be relevant indicators of hyperinflation and patient-related outcomes in patients with severe emphysema.³⁵

The gas dilution techniques can be used as an alternative in patients who cannot perform body plethysmography, including patients with claustrophobia or severe obesity.³⁶ However, typically in patients with severe emphysema, FRC can be largely underestimated as the measured volume corresponds with the well-ventilated areas only.³⁴

Spirometrically assessed inspiratory capacity (IC) can indirectly indicate the presence of LH. The IC is the maximal volume of air that can be inhaled after a spontaneous expiration to FRC and is the difference between TLC and FRC. Pharmacological studies in COPD have established that IC as a measure of hyperinflation correlates stronger with dyspnea and exercise performance than FEV₁.^{37,38} However, the use of IC for measuring LH ignores an isolated increase in RV and assumes that TLC does not change during the course of the disease as a result of an exacerbation or bronchodilation therapy.²¹ A recent study showed that gas trapping or LH in terms of RV/TLC ratio could be estimated using only FEV₁ and FVC.³⁹ Das et al showed that severe hyperinflation in COPD patients can potentially be indicated by measuring the area under the forced expiratory flow-volume loop.⁴⁰ However, as the authors emphasized, a reference value for this method does not yet exist, and the study focused only on severe static hyperinflation.

Measuring Static Hyperinflation by Imaging Techniques

Conventional chest radiographs can show signs of increased lung volume but are not a standardized method for measuring LH. The most supportive radiographic signs of LH are a flattened diaphragm,⁴¹ an increase in the retrosternal air space of more than 2.5 cm and an increase in lung height.²¹

A more important method for assessment of LH is computer tomography (CT). Lung volumes measured by CT are less prone to inaccuracies related to chest wall or image distortion compared to chest x-rays. Usually, TLC measured by plethysmography and CT have a very high correlation ($r > 0.9$).⁴² The difference between TLC through plethysmography and CT is greatest among patients with FEV₁ < 30% of predicted.⁴³ A limitation of measurement of LH with CT is the difficulty with maximum inspiratory maneuvers and breath-holding techniques during scanning. Also, because CT measures TLC in supine position and plethysmographic techniques are performed in the sitting position, TLC is greater in plethysmography.²¹ This discrepancy can be even larger in obese patients.

CT imaging has made an important contribution to the diagnosis and management of COPD for many years,⁴⁴ especially, it allows the identification of emphysema.⁴⁵ The extent of emphysema on CT has been shown to correlate significantly with hyperinflation determined by lung function testing.⁴⁶ Moreover, by combining LH and emphysema additional information that is useful for predicting the course of disease can be obtained.⁴⁷ Volumetric acquired high-resolution CT (HRCT) is excellent for measuring the presence, pattern, distribution, and extent of emphysema.⁴⁸ The NETT study demonstrated that the regional distribution and heterogeneity of emphysema on HRCT are clinically important features to help select patients who better respond to lung volume reduction surgery.⁴⁹ Also, the extent of emphysema on CT predicts (absence of) response to pharmacotherapies in COPD.⁵⁰ Additionally, CT has widespread availability and rapid image acquisition and may aid in the early detection of COPD, as the comparison between inspiratory and expiratory images can show air trapping and help to distinguish COPD phenotypes.⁵¹ However, the presence of emphysema does not always imply the presence of hyperinflation (Figure 2).

Quantitative CT (QCT) uses software to quantify, either at lobar or whole lung level, the extent of emphysema, based on voxel-by-voxel attenuation; the degree of bronchial wall thickening, based on the ratio of average lumen diameter to wall area in the small airways and the extent of gas trapping, based on differences between inspiratory and expiratory phase images.^{28,52,53} Parametric response mapping, a voxel-based image analysis technique, enables differentiation between gas trapping from emphysema or small airways disease.²⁸ QCT indices correlate independently with clinical outcomes including disease progression, mortality and symptoms.^{28,53} Currently, QCT is mainly used in emphysematous patients with severe LH to select candidates for bronchoscopic lung volume reduction.⁵⁴

Imaging emphysema with magnetic resonance imaging (MRI) is a major challenge due to the loss of lung tissue and reduced blood volume, resulting in a marked reduction of lung parenchymal signal.⁵⁵ MRI might be used in COPD for assessment of functional parameters like perfusion and respiratory dynamics.⁵⁶ There is no literature describing imaging of LH with MRI.

Measuring Dynamic Hyperinflation

DH can be measured by performing IC maneuvers during a cycle ergometer test or 6WMT, with a decline in IC during exercise being indicative of the presence of DH. A decrease in IC from baseline >150 mL during exercise indicates DH.⁵⁷ Several studies consider a decline of 10% as clinically relevant.^{58,59} IC maneuvers during exercise provide valuable information on ventilatory constraints during exercise,^{60,61} but require logistics and repetitive IC maneuvers that can be difficult to perform for patients,⁶² especially for those with advanced stages of COPD. Metronome-paced tachypnea (MPT) is considered to be a feasible alternative to measure DH.⁶² By applying mandatory tachypnea for a short period while sitting at rest, and performing an IC maneuver before and after, it mimics the dynamic respiratory pattern that occurs during exertion.⁶³ It is suggested that a minimal decrease in IC of 11.1% is the optimal cut-off for MPT-induced DH.⁵⁸ It must be noted that the sensitivity of this method is 85%, but with limited specificity, which can lead to overdiagnosis of DH.⁶⁴

Esophageal pressure is a well-known surrogate of the pleural pressure and is considered the golden standard to measure dynamic hyperinflation during exercise.⁶⁵ A balloon catheter is placed nasally at approximately the lower third of the esophagus to minimize cardiogenic oscillations.⁶⁶ The use of esophageal catheters to measure hyperinflation is customary in research settings, while it requires a high technical effort and the insertion can be cumbersome and time-consuming in daily clinical practice.

Optoelectronic plethysmography is another established technique to visualize dynamic hyperinflation, by measuring tidal changes in the volume of the chest wall and its compartments.⁶⁷ A number of reflective markers are placed on selected anatomical reference sites of the thoraco-abdominal surface of the subject. A set of cameras placed nearby the subject and a system for human motion analysis measures the three-dimensional coordinates and motion of these markers.⁶⁸ Also this technique is mostly suited for laboratory research settings.

Hyperinflation as Part of the Systemic Disease COPD

Although hyperinflation may be approached as an isolated pulmonary trait in COPD that only partially correlates with other lung function measurements¹ (Figure 1), it may alternatively be seen as one of the distinct features of a specific disease phenotype.⁹ The emphysematous phenotype was traditionally described as the “pink puffer”, known for its characteristic barrel chest deformity as a consequence of emphysema and low body weight,¹⁰ which can also be found in patients with hyperinflation (Figure 3). In the past decades, additional features of the emphysematous phenotype but also the hyperinflated patient and its therapeutic implications have been studied extensively and these will be discussed here.

Dyspnea, Exercise Limitation, Exacerbations, Gas Exchange

LH importantly contributes to the degree of (exertional) dyspnea and exercise limitation,³⁸ impacting daily life activities.⁶⁹ DH also exists in mild COPD with no or only mild static hyperinflation.⁷⁰

The presence of a higher RV/TLC ratio or IC/TLC ratio, as a marker of hyperinflation, is independently associated with frequent exacerbations, hospitalizations and all-cause mortality.^{71–74} Zeng et al even showed that smoking subjects with a higher RV/TLC ratio but with preserved spirometry were more likely to have higher all-cause mortality.⁷⁵

It has been shown that in patients with stable COPD, severe hypoxemia is independently associated with higher RV, RV/TLC, FRC and TLC.^{76,77} Also, hypercapnic COPD patients have greater degrees of hyperinflation.⁷⁸ Hyperinflation places the diaphragm and other inspiratory muscles at severe mechanical disadvantage, producing alveolar hypoventilation and hypercapnia.⁷⁹ In the presence of DH, there is an increase in ventilation-perfusion mismatching, which in turn leads to hypoxemia.⁸⁰ Also, DH can lead to hypercapnia, due to increased fatigue of the inspiratory muscles, leading to rapid shallow breathing and diminished CO₂ removal.²¹

Cardiac Impairment and Cardiovascular Disease

The link between LH and cardiac dysfunction in COPD patients is increasingly recognized.⁸¹ Lung hyperinflation and the corresponding increased intrathoracic pressure directly affect the cardiovascular system. This, along with larger swings in intrapleural pressure, can cause diminished venous return and increase left ventricle afterload by compressing the large




Figure 3 Clinical and radiologic presentation of lung hyperinflation.

and small pulmonary arterial vessels and the heart chambers.^{82,83} Due to left ventricular failure and alveolar hypoxia, in association with high pulmonary capillary pressures, this effect increases the right ventricle afterload.⁸³ In patients with severe COPD and normal cardiac function at rest, the rise in EELV during tachypnea was associated with an increase in right atrial pressure, mean pulmonary artery pressure and wedge pressure.⁸⁴ Hyperinflated lungs in severe emphysematous patients have been shown to be associated with intrathoracic hypovolemia, decreased biventricular preload and reduced cardiac performance.⁸⁵ Next to systolic dysfunction, it has been established that airway obstruction and LH are associated with impaired left heart diastolic filling that can be detected by echocardiography.⁸⁶ Conversely, lung deflation by long-acting bronchodilators improves diastolic function and stroke volume in COPD. Reduction of hyperinflation using bronchoscopic lung volume reduction with endobronchial valves can improve cardiac preload and output.^{87–89}

Recently, coronary artery disease was found to be strongly associated with lung hyperinflation in current and former smokers.⁹⁰

Pulmonary hypertension can develop in COPD patients as a result of loss of alveolar capillary bed or as a consequence of chronic alveolar hypoxia with secondary vasoconstriction.⁹¹ LH was associated with raised pulmonary artery pressure during exercise and larger pulmonary artery pressure at rest.^{92,93} However, evidence regarding LH as the cause of pulmonary hypertension is lacking.

Muscle Dysfunction

Muscle dysfunction is a well-established systemic manifestation of COPD, affecting both respiratory and peripheral muscles in all COPD stages.^{30,94} LH increases the work of breathing as it flattens and shortens the diaphragm, which

results in a non-optimal strength-length position of the diaphragm. It also shifts the tidal volume towards a less favorable part of the compliance curve. This higher work of breathing might lead to progressively higher nutrient and oxygen demands of respiratory muscles.⁹⁵

A direct link between LH and peripheral muscle dysfunction has not been fully established. Peripheral muscle dysfunction is common in COPD, probably mainly driven by muscle deconditioning by disuse of the locomotor muscles.^{94,96} Disuse results from exertional dyspnea, DH and redistribution of blood flow during exercise from locomotor muscles to respiratory muscles.⁹⁷ However, other intrinsic and systemic factors are also involved. This is indicated by the only partial reversal of functional and structural limb muscle changes with exercise training.⁹⁵ Therefore, hyperinflation might also be indirectly related to peripheral muscle dysfunction.⁹⁸

Body Composition, Sarcopenia and Bone Density

Emphysema is often associated with low body mass index (BMI) and low fat-free mass index (FFMI).⁹⁹ COPD patients with advanced emphysema constitute a specific phenotype characterized by an enhanced loss of tissue mass over time in multiple organs.¹⁰⁰ This is likely to be in relation to abnormal tissue loss/repair capacity. Patients with severe emphysema not only showed lower BMI and FFMI at baseline but also accelerated loss of BMI and FFMI over time. Moreover, these patients had more exacerbations, hospitalizations, and higher mortality during three-year follow-up,¹⁰⁰ indicating that proper identification of patients with severe emphysema and excessive loss of tissue is clinically relevant. However, data describing BMI and FFMI in relation to LH are scarce. One study researched the impact of increasing BMI on static lung volumes and showed a reduction of RV, FRC and TLC with increasing BMI.⁶

Patients with severe emphysema and LH undergoing lung volume reduction surgery can show an increase in FFMI after six months, implying that the systemic effects of the disease on body composition can, at least partly, be reversed.¹⁰¹ Weight gain and skeletal muscle remodeling following bronchoscopic lung volume reduction have also been observed and were associated with functional improvements.¹⁰²

Osteopenia and osteoporosis are especially associated with the presence and severity of emphysema.¹⁰³ Studies evaluating the pathophysiologic mechanisms underlying this relationship are limited, although systemic inflammation^{104,105} and abnormal tissue loss/repair capacity¹⁰⁰ are commonly suggested as underlying mechanisms for multi-organ tissue loss. Again, these were studies describing relationships between emphysema and bone density, but not LH. One small retrospective study showed improvements in bone mineral density after lung volume reduction surgery; however, these results have not been confirmed in a randomized trial.¹⁰⁶

Lung Cancer

Several studies have reported an association between emphysema and lung cancer.^{107,108} Recently, LH has been shown to be an independent risk factor for the development of lung cancer in COPD patients.¹⁰⁹ However, emphysema as a confounding factor was not ruled out.

Pharmacologic Treatment

Bronchodilators

Long-acting bronchodilators (LABDs) are the cornerstone of pharmacological treatment of patients with COPD.¹¹⁰ In general, they increase FEV₁, improve dyspnea and exercise performance and reduce exacerbation frequency. Bronchodilators improve airway conductance,¹¹¹ relax airway smooth muscle, improve small airway patency and enhance lung deflation, which is reflected by a reduction in static and dynamic LH.¹¹² Breathing at lower operating lung volumes decreases work of breathing while placing the respiratory muscles in a more efficient arrangement for pressure generation,¹¹³ delaying time to respiratory muscle fatigue.¹¹⁴ Bronchodilators seem to be even more effective in patients with relatively limited airway obstruction but marked hyperinflation compared to patients without hyperinflation,¹¹⁵ and LH might serve as an objective marker for aggressive pharmacological intervention.¹¹⁶ Clinical trials demonstrated that a combination of two long-acting LABDs is superior to a single bronchodilator^{117,118} and to long-acting beta2-agonist (LABA) and inhaled corticosteroids (ICS) in improving spirometry measures of hyperinflation.¹¹⁹

Administration of bronchodilators in LH also reduces exacerbation frequency, probably by improving lung mechanics, reducing the work of breathing and degree of dyspnea and a potential reduction of mechanical stress in the lungs.¹²⁰ In addition, it was shown that LABDs improve diastolic function in COPD, in accordance with the finding that lung deflation can improve cardiac filling.⁸⁸ Moreover, lung deflation by tiotropium has been demonstrated to increase oxygen pulse as a non-invasive surrogate of stroke volume.⁸⁹ Thus, it can be speculated that LABDs antagonize the decreased left atrium size in COPD resulting from LH, improve early left ventricular filling and increase left heart preload and filling, and improve stroke volume.

Anti-Inflammatory Agents

Few studies investigated the add-on effects of ICS on bronchodilators with regard to LH in COPD. Significant improvements in inspiratory capacity, as a marker of LH, and exercise endurance time during ICS/LABA treatment were reported compared with LABA and with placebo.^{121,122} Another study, however, did not confirm benefits with ICS over LABA monotherapy.^{121,122} Thus, LH is not a specific indication for ICS therapy in current international guidelines. Roflumilast, an oral phosphodiesterase-4 (PDE4) inhibitor, is indicated to prevent exacerbations in COPD patients with severe airflow obstruction, chronic bronchitis, and frequent exacerbations. It was also shown to improve IC, but without effect on resting hyperinflation.¹²³

Opioids

Opioids can be useful in patients with COPD and symptoms of severe dyspnea and impaired exercise capacity, to relieve breathlessness,¹²⁴ improve exercise endurance,¹²⁵ and health status,¹²⁶ and that these effects might occur without actually decreasing DH.¹²⁷

Non-Pharmacologic Treatment

Management of Dyspnea

Pursed-Lips Breathing

Pursed-lips breathing (PLB) is a strategy frequently adopted spontaneously by patients with COPD to reduce dyspnea by prolonging expiration¹²⁸ and counterbalancing intrinsic positive end-expiratory pressure (PEEP). PLB is frequently taught during respiratory physical therapy and pulmonary rehabilitation programs to reduce dyspnea and improve exercise tolerance.¹²⁹ By breathing out through partly closed lips, expiratory airway pressure is increased, expiratory airway collapse is reduced and thus expiratory airflow rate is improved. A recent meta-analysis showed that PLB during exercise is effective in reducing minute ventilation and respiratory rate, however, it does not significantly improve walking distance or dyspnea.¹³⁰ Probably, lack of methodological quality and the restricted number of patients included in the trials contributed to this result.

Inspiratory Muscle Training

Inspiratory muscle training (IMT) is effective in increasing inspiratory muscle function and exercise performance in COPD when used as a stand-alone intervention,¹³¹ and has been recommended in patients with inspiratory muscle weakness.¹³² It is postulated that IMT decreases inspiratory time during loaded breathing tasks. This is likely the result of increased velocity of diaphragmatic contraction due to increase in the number of fast-twitch type 2 muscle fibers.¹³³ Consequently, expiratory time is prolonged, thereby decreasing DH and reducing dyspnea.¹³⁴ However, its additional benefit as an adjunct to whole body exercise training is questionable.¹³⁵

Exercise Training

Exercise training decreases ventilatory requirements, resulting in lower respiratory rate at a given level of exercise. The longer available time for expiration results in less DH and therefore less dyspnea.¹³⁶ There is a lot of evidence regarding leg training decreasing breathing frequency during exercise and increasing IC.²¹ Even in patients with (very) severe LH and accompanying low exercise capacity, pulmonary rehabilitation has been shown to be effective.¹³⁷

Oxygen Therapy

Oxygen supplementation during exercise can improve exercise endurance by reducing the respiratory rate and thereby DH and dyspnea in normoxemic COPD patients,¹³⁸ and those with resting hypoxemia or exercise-induced desaturation.¹³⁹ However, a recently published randomized controlled trial investigating the effect of oxygen supplementation during an 8-week supervised exercise training program in normoxemic COPD patients with exercise induced desaturation, who also had LH, showed no greater improvements in exercise capacity with oxygen supplementation than with medical air.¹⁴⁰ The current evidence does not support the routine use of oxygen supplementation during exercise in COPD patients with DH, except in patients who already receive long-term oxygen therapy.¹³²

Heliox (mixture of helium/oxygen 80:20) can improve exercise endurance, peak ventilation time and IC, reducing DH and dyspnea at isotime in COPD.¹⁴¹ The mixture decreases turbulent airflow in large airways.¹⁴¹ However, the results are variable and their application during exercise training remains to be established.¹³²

Relieving Inspiratory Threshold Load

PLB can relieve inspiratory threshold load, as described above. Another method to accomplish this is applying external PEEP.

Intrinsic PEEP can account for about 1/3 of the total work of breathing in patients with dynamic airway collapse, and small amounts of external PEEP or continuous positive airway pressure (CPAP) can be beneficial during exercise.¹⁴² It unloads inspiratory muscles from the inspiratory threshold load induced by intrinsic PEEP. The application of PEEP and inspiratory pressure support or proportional assisted ventilation can decrease the inspiratory workload in mechanically ventilated patients.^{143,144} Inspiratory pressure support applied during exercise can improve endurance time and reduce dyspnea.¹⁴⁵ Proportional assisted ventilation unloads respiratory muscles, and thereby reduces dyspnea and prevents exercise-induced diaphragmatic fatigue.¹⁴⁶ Expiratory positive airway pressure (EPAP) aims to increase resistance in the expiratory phase, which is thought to reduce physiological dead space and increase tidal volume. However, a recent systematic review showed that the use of EPAP in COPD patients did not modify DH and reduced exercise time.¹⁴⁷ Bilevel positive airway pressure (BiPAP) decreases inspiratory workload and FRC through a modification of the breathing pattern.¹⁴⁸ BiPAP can reduce or delay DH during exercise⁹⁸ or during an acute exacerbation of COPD¹⁴⁹ and can reduce the load of overburdened inspiratory muscles and the work of breathing.¹⁴⁵ However, patients without hypercapnia currently do not have an indication for non-invasive ventilation (NIV) during exercise or during an acute exacerbation of COPD.¹⁵⁰

Freitas et al showed a reduction in RV in 72 patients with hypercapnia after two years of treatment with long-term NIV.¹⁵¹ However, a two-year RCT in 122 stable COPD patients with chronic hypercapnia showed no difference in lung function after introducing long-term NIV.¹⁵² However, non-hypercapnic patients with hyperinflation currently do not have an indication for long-term NIV.^{150,153}

Lung Volume Reduction

Static hyperinflation can also be reduced by improving elastic recoil. There are several therapies that aim to eliminate the most emphysematous or hyperinflated portions of the lung, allowing the remaining pulmonary parenchyma and the respiratory muscles to function more effectively. A classic invasive technique to eliminate emphysematous lung tissue is lung-volume-reduction surgery (LVRS). The NETT was a large clinical trial that confirmed that patients with COPD with upper-lobe emphysema and low exercise capacity had improved survival when treated with LVRS compared to standard medical treatment.⁴⁹ Also, LVRS can restore body weight and body composition, improve right ventricular function, stroke volume and bone mineral density.^{106,154} However, patients with high post-pulmonary rehabilitation exercise capacity had no difference in survival after LVRS.⁴⁹ Moreover, LVRS results in higher mortality in patients with severe emphysema with an FEV₁ less than or equal to 20% predicted, and either homogeneous emphysema or diffusing capacity of the lung for carbon monoxide of less than or equal to 20% predicted.¹⁵⁵

As a less-invasive and safe alternative for surgical lung volume reduction,⁴⁹ endobronchial valve placement can improve lung volumes, reduce FRC, improve exercise endurance and quality of life in a highly selected group of COPD patients with severe emphysema and severe LH.¹⁵⁶ Recently, it has been shown that reduction of LH by endobronchial

valve placement can improve cardiac preload and output.⁸⁷ Moreover, the therapy might prevent further loss or even increase muscle mass.¹⁰² Long-term efficacy, such as impact on exacerbations, hospitalizations and survival remains to be established,¹⁵⁷ although preliminary data are promising in terms of exacerbation reduction and survival benefit.¹⁵⁸ Endobronchial coil placement is a different technique for patients without intact lobar fissures, inducing regional parenchymal volume reduction and restoring lung recoil with comparable effects.¹⁵⁹

Currently, health insurance companies worldwide require mandatory pulmonary rehabilitation prior to LVRS and endobronchial valve placement, and clinical trials evaluating LVRS and endobronchial valves required pulmonary rehabilitation prior to randomization.¹⁶⁰ The NETT trial showed that pulmonary rehabilitation prior to LVRS resulted in improvements in exercise capacity and quality of life.¹⁶¹ A post-hoc analysis of the VENT trial showed modest improvements in exercise capacity and dyspnea.¹⁶⁰ The SOLVE trial was designed to study the impact of optimal timing of pulmonary rehabilitation on exercise capacity and patient reported outcomes in patients found eligible for endobronchial valve placement. They found that the combination of pulmonary rehabilitation with endobronchial valve placement did not result in increased exercise capacity, improved patient-reported outcomes or increased daily step count compared to endobronchial valve placement alone.¹⁶² However, their study also included participants who had undergone rehabilitation in the past. Subsequent rehabilitation has been demonstrated to be less impactful than the initial treatment.¹⁶¹

Both the use of biological sealant agents and thermal vapor ablation induce atelectasis due to airway occlusion and subsequent remodeling. The remodeling will cause contraction of lung parenchyma and loss of hyperinflated areas.^{163,164} Thermal vapor ablation has been acknowledged as a possible lung volume reduction treatment; however, the availability for clinical practice is limited to a small number of centers.¹⁶⁵ Biological sealant agents have been found to lead to relatively high incidence of adverse events.^{166,167}

How to Approach the COPD Patient with Hyperinflation

Although LH is common in patients with COPD and plays a central role in the burden and outcomes of the disease, its presence and severity cannot reliably be derived from the degree of airflow limitation (Figure 2) and thus requires specific assessment. As patients with mild or no airflow limitation can develop LH and/or DH, there is a rationale for assessment of LH in all patients with dyspnea during their daily life activities. Spirometrically assessed IC can indirectly indicate the presence of LH. Subsequently, body plethysmography is used to indicate the presence and severity of hyperinflation. When DH is suspected, with or without the presence of severe airflow limitation and/or LH, IC maneuvers during exercise testing can be used.

Bronchodilators can reduce static and dynamic LH, improve dyspnea, exercise performance and reduce exacerbation frequency. There is no specific indication for ICS therapy in current guidelines for LH, but additional therapy might improve exercise endurance.

PLB can be easily taught during physical therapy to reduce dyspnea and improve exercise tolerance. Exercise training in patients with LH decreases respiratory rate at a given level of exercise. Also, pulmonary rehabilitation has been shown to be effective in patients with very severe LH. Current evidence does not support the routine use of oxygen supplementation in normoxemic patients with DH. Moreover, patients without hypercapnia currently do not have an indication for NIV during exercise or long-term nocturnal NIV.

In patients with severe hyperinflation, assessment of regional distribution and heterogeneity of emphysema by (quantitative) CT can be performed in consideration of bronchoscopic or surgical lung volume reduction. This can improve exercise endurance, body weight and composition and cardiac function. Efficacy on exacerbations, hospitalizations and survival remains to be established, although preliminary data are promising in terms of exacerbation reduction and survival benefit.

Cardiac impairment and muscle dysfunction might be present in patients with LH and can contribute to dyspnea and exercise limitation. The presence of low BMI, osteopenia and osteoporosis can be considered, although the presence of these treatable traits is mainly reported in patients with emphysema. Lastly, it should be noted that LH has been shown to be a risk factor for the development of lung cancer.

Box 1 Unanswered Questions Regarding Hyperinflation in COPD

- What are the trajectories of hyperinflation in COPD?
- Do patients with hyperinflation benefit to the same extent from exercise training as patients without hyperinflation?
- What is the optimal exercise modality in patients with hyperinflation?
- Is hyperinflation variable throughout the day?
- Which patients do or do not benefit from LABDs, are there actually non-responders?
- How do patients with hyperinflation respond to a combination of interventions?
- Is treatment of hyperinflation useful and effective in patients with hyperinflation but without obstruction?
- What is the association between reduction of hyperinflation and improvements in patient-related outcomes in individual patients?
- Does reduction of hyperinflation lower cardiovascular risk?
- Does reduction of hyperinflation improve mortality in COPD patients?

The interplay of factors contributing to the clinical presentation of LH in an individual patient is complex and may alter over time. Thus, an integrated approach is warranted to provide personalized management. Although LH is recognized as a treatable trait of COPD, breaking down the disease into a list of traits for which specific treatments are available may deny its true complexity and multi-level relationships between disease characteristics. Also, underlying pathophysiologic mechanisms will remain unrevealed and clinical phenotypes might not be recognized. At the same time, it should be acknowledged that COPD phenotypes are not mutually exclusive, emphasizing the need for comprehensive assessment and personalized treatments in COPD. In **Box 1**, a few potentially interesting future research questions regarding LH in COPD are listed.

Conclusion

Lung hyperinflation represents a treatable feature of patients with COPD that requires specific lung function and/or radiological assessment. Effective pharmacological and non-pharmacological treatments are available for this trait but must be applied in the broader context of the health condition of the individual patient.

Acknowledgments

The person in **Figure 3** provided written informed consent for publication of his image. This manuscript has not been submitted to another journal and has not been published in whole or in part elsewhere previously.

Disclosure

Dr Lowie Vanfleteren reports grants and/or personal fees from AstraZeneca, GSK, Novartis, Boehringer, Pulmonx and Chiesi outside the submitted work. Dr Sami Simons reports grants and personal fees from AstraZeneca, Chiesi, Boehringer Ingelheim, GlaxoSmithKline and Roche outside the submitted work. Prof. Dr. Frits Franssen reports personal fees from AstraZeneca, Chiesi, GlaxoSmithKline, MSD and Verona Pharma outside the submitted work. The authors report no other conflicts of interest in this work.

References

1. Augustin IML, Spruit MA, Houben-Wilke S, et al. The respiratory physiome: clustering based on a comprehensive lung function assessment in patients with COPD. *PLoS One*. 2018;13(9):e0201593. doi:10.1371/journal.pone.0201593
2. Han MK, Agusti A, Calverley PM, et al. Chronic obstructive pulmonary disease phenotypes: the future of COPD. *Am J Respir Crit Care Med*. 2010;182(5):598–604. doi:10.1164/rccm.200912-1843CC
3. Agusti A, Calverley PM, Celli B, et al. Characterisation of COPD heterogeneity in the ECLIPSE cohort. *Respir Res*. 2010;11:122. doi:10.1186/1465-9921-11-122
4. Agusti A, Bel E, Thomas M, et al. Treatable traits: toward precision medicine of chronic airway diseases. *Eur Respir J*. 2016;47(2):410–419. doi:10.1183/13993003.01359-2015
5. Corbridge T, Irvin C. Pathophysiology of chronic obstructive pulmonary disease with emphasis on physiologic and pathologic correlations. In: *Casaburi R and Petty T, Principles and Practice of Pulmonary Rehabilitation*. Philadelphia, PA: Saunders; 1993:18–32.
6. O'Donnell DE, Deesomchok A, Lam YM, et al. Effects of BMI on static lung volumes in patients with airway obstruction. *Chest*. 2011;140(2):461–468. doi:10.1378/chest.10-2582

7. Ofir D, Laveneziana P, Webb KA, Lam YM, O'Donnell DE. Mechanisms of dyspnea during cycle exercise in symptomatic patients with GOLD stage I chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 2008;177(6):622–629. doi:10.1164/rccm.200707-1064OC
8. Casanova C, Cote C, de Torres JP, et al. Inspiratory-to-total lung capacity ratio predicts mortality in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 2005;171(6):591–597. doi:10.1164/rccm.200407-867OC
9. Vanfleteren LE, Spruijt MA, Wouters EF, Franssen FM. Management of chronic obstructive pulmonary disease beyond the lungs. *Lancet Respir Med.* 2016;4:911–924. doi:10.1016/S2213-2600(16)00097-7
10. Gibson GJ. Pulmonary hyperinflation a clinical overview. *Eur Respir J.* 1996;9(12):2640–2649. doi:10.1183/09031936.96.09122640
11. Hall GL, Filipow N, Ruppel G, et al. Official ERS technical standard: global Lung Function Initiative reference values for static lung volumes in individuals of European ancestry. *Eur Respir J.* 2021;57(3):2000289. doi:10.1183/13993003.00289-2020
12. Stanojevic S, Kaminsky DA, Miller MR, et al. ERS/ATS technical standard on interpretive strategies for routine lung function tests. *Eur Respir J.* 2022;60(1):1.
13. O'Donnell DE, Laveneziana P. Physiology and consequences of lung hyperinflation in COPD. *Eur Respir Rev.* 2006;15(100):61–67. doi:10.1183/09059180.00010002
14. Similowski T, Milic-Emili J, Derenne J-P. Respiratory mechanics during acute respiratory failure of chronic obstructive pulmonary disease. *Lung Bio Health Dis.* 1996;92:23–46.
15. Carbonara P, Eidelman DH. *Pulmonary Statics in Diseases. Physiologic Basis of Respiratory Diseases Hamilton.* Ont, Canada: Dekker Inc; 2005:69–76.
16. Dubé B, Guerder A, Morelot-Panzini C, et al. The clinical relevance of the emphysema-hyperinflated phenotype in COPD. *COPD Research and Practice.* 2015;2(1). doi:10.1186/s40749-015-0017-7
17. Bates DV, Woolf CR, Paul GI. A report on the first two stages of the co-ordinated study of chronic bronchitis in the Department of Veterans Affairs, Canada. *Med Serv J Can.* 1962;18:211–303.
18. Deesomchok A, Webb KA, Forkert L, et al. Lung hyperinflation and its reversibility in patients with airway obstruction of varying severity. *COPD.* 2010;7(6):428–437. doi:10.3109/15412555.2010.528087
19. Macklem PT. Therapeutic implications of the pathophysiology of COPD. *Eur Respir J.* 2010;35(3):676–680. doi:10.1183/09031936.00120609
20. Smid DE, Wilke S, Jones PW, et al. Impact of cardiovascular comorbidities on COPD Assessment Test (CAT) and its responsiveness to pulmonary rehabilitation in patients with moderate to very severe COPD: protocol of the Chance study. *BMJ Open.* 2015;5(7):e007536. doi:10.1136/bmjopen-2014-007536
21. Rossi A, Aisanov Z, Avdeev S, et al. Mechanisms, assessment and therapeutic implications of lung hyperinflation in COPD. *Respir Med.* 2015;109(7):785–802. doi:10.1016/j.rmed.2015.03.010
22. Guenette JA, Webb KA, O'Donnell DE. Does dynamic hyperinflation contribute to dyspnoea during exercise in patients with COPD? *Eur Respir J.* 2012;40(2):322–329. doi:10.1183/09031936.00157711
23. Laveneziana P, Webb KA, Wadell K, Neder JA, O'Donnell DE. Does expiratory muscle activity influence dynamic hyperinflation and exertional dyspnea in COPD? *Respir Physiol Neurobiol.* 2014;199:24–33. doi:10.1016/j.resp.2014.04.005
24. Macklem PT, Mead J. Resistance of central and peripheral airways measured by a retrograde catheter. *J Appl Physiol.* 1967;22(3):395–401. doi:10.1152/jappl.1967.22.3.395
25. Hogg JC, Macklem PT, Thurlbeck WM. Site and nature of airway obstruction in chronic obstructive lung disease. *N Engl J Med.* 1968;278(25):1355–1360. doi:10.1056/NEJM196806202782501
26. McDonough JE, Yuan R, Suzuki M, et al. Small-airway obstruction and emphysema in chronic obstructive pulmonary disease. *N Engl J Med.* 2011;365(17):1567–1575. doi:10.1056/NEJMoa1106955
27. Koo HK, Vasilescu DM, Booth S, et al. Small airways disease in mild and moderate chronic obstructive pulmonary disease: a cross-sectional study. *Lancet Respir Med.* 2018;6(8):591–602. doi:10.1016/S2213-2600(18)30196-6
28. Bhatt SP, Soler X, Wang X, et al. Association between functional small airway disease and FEV1 decline in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 2016;194(2):178–184. doi:10.1164/rccm.201511-2219OC
29. Hogg JC, Pare PD, Hackett T-L. The contribution of small airway obstruction to the pathogenesis of chronic obstructive pulmonary disease. *Physiol Rev.* 2017;97(2):529–552. doi:10.1152/physrev.00025.2015
30. Maltais F, Decramer M, Casaburi R, et al. An official American Thoracic Society/European Respiratory Society statement: update on limb muscle dysfunction in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 2014;189(9):e15–62. doi:10.1164/rccm.201402-0373ST
31. Wielputz M, Kauczor HU. MRI of the lung: state of the art. *Diagn Interv Radiol.* 2012;18(4):344–353. doi:10.4261/1305-3825.DIR.5365-11.0
32. Wanger J, Clausen JL, Coates A, et al. Standardisation of the measurement of lung volumes. *Eur Respir J.* 2005;26(3):511–522. doi:10.1183/09031936.05.00035005
33. Criece CP, Sorichter S, Smith HJ, et al. Body plethysmography--its principles and clinical use. *Respir Med.* 2011;105(7):959–971. doi:10.1016/j.rmed.2011.02.006
34. Rodenstein DO, Stanescu DC, Francis C. Demonstration of failure of body plethysmography in airway obstruction. *J Appl Physiol Respir Environ Exerc Physiol.* 1982;52(4):949–954. doi:10.1152/jappl.1982.52.4.949
35. de Weger WW, Klooster K, Ten Hacken NH, van Dijk M, Hartman JE, Slebos DJ. Determining static hyperinflation in patients with severe emphysema: relation between lung function parameters and patient-related outcomes. *Lung.* 2020;198(4):629–636. doi:10.1007/s00408-020-00368-9
36. Brown R, Leith DE, Enright PL. Multiple breath helium dilution measurement of lung volumes in adults. *Eur Respir J.* 1998;11(1):246–255. doi:10.1183/09031936.98.11010246
37. O'Donnell DE, Lam M, Webb KA. Spirometric correlates of improvement in exercise performance after anticholinergic therapy in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 1999;160(2):542–549. doi:10.1164/ajrccm.160.2.9901038
38. O'Donnell DE, Fluge T, Gerken F, et al. Effects of tiotropium on lung hyperinflation, dyspnoea and exercise tolerance in COPD. *Eur Respir J.* 2004;23(6):832–840. doi:10.1183/09031936.04.00116004
39. Alter P, Orszag J, Kellerer C, et al. Prediction of air trapping or pulmonary hyperinflation by forced spirometry in COPD patients: results from COSYCONET. *ERJ Open Res.* 2020;6(3):00092–2020. doi:10.1183/23120541.00092-2020

40. Das N, Topalovic M, Aerts JM, Janssens W. Area under the forced expiratory flow-volume loop in spirometry indicates severe hyperinflation in COPD patients. *Int J Chron Obstruct Pulmon Dis.* 2019;14:409–418. doi:10.2147/COPD.S185931
41. Webb WR. Radiology of obstructive pulmonary disease. *AJR Am J Roentgenol.* 1997;169(3):637–647. doi:10.2214/ajr.169.3.9275869
42. Garfield JL, Marchetti N, Gaughan JP, Steiner RM, Criner GJ. Total lung capacity by plethysmography and high-resolution computed tomography in COPD. *Int J Chron Obstruct Pulmon Dis.* 2012;7:119–126. doi:10.2147/COPD.S26419
43. O'Donnell CR, Bankier AA, Stiebellehner L, Reilly JJ, Brown R, Loring SH. Comparison of plethysmographic and helium dilution lung volumes: which is best for COPD? *Chest.* 2010;137(5):1108–1115. doi:10.1378/chest.09-1504
44. Petousi N, Talbot NP, Pavord I, Robbins PA. Measuring lung function in airways diseases: current and emerging techniques. *Thorax.* 2019;74:797–805. doi:10.1136/thoraxjnl-2018-212441
45. Bergin C, Muller N, Nichols DM, et al. The diagnosis of emphysema. A computed tomographic-pathologic correlation. *Am Rev Respir Dis.* 1986;133(4):541–546. doi:10.1164/arrd.1986.133.4.541
46. Gurney JW, Jones KK, Robbins RA, et al. Regional distribution of emphysema: correlation of high-resolution CT with pulmonary function tests in unselected smokers. *Radiology.* 1992;183(2):457–463. doi:10.1148/radiology.183.2.1561350
47. Kim YW, Lee CH, Hwang HG, et al. Resting hyperinflation and emphysema on the clinical course of COPD. *Sci Rep.* 2019;9(1):3764. doi:10.1038/s41598-019-40411-1
48. Goldin JG, Abtin F. Update on radiology of emphysema and therapeutic implications. *Thorac Surg Clin.* 2009;19(2):159–167. doi:10.1016/j.thorsurg.2009.04.006
49. Fishman A, Martinez F, Naunheim K, et al. A randomized trial comparing lung-volume-reduction surgery with medical therapy for severe emphysema. *N Engl J Med.* 2003;348(21):2059–2073.
50. Lee JH, Lee YK, Kim EK, et al. Responses to inhaled long-acting beta-agonist and corticosteroid according to COPD subtype. *Respir Med.* 2010;104(4):542–549. doi:10.1016/j.rmed.2009.10.024
51. Siddiqui S, Usmani OS. Small airways, big challenge: measuring the unseen? *Nat Med.* 2012;18(11):1619–1621. doi:10.1038/nm.2981
52. Matin TN, Rahman N, Nickol AH, et al. Chronic obstructive pulmonary disease: lobar analysis with hyperpolarized (129)Xe MR imaging. *Radiology.* 2017;282(3):857–868. doi:10.1148/radiol.2016152299
53. Lynch DA, Moore CM, Wilson C, et al. CT-based visual classification of emphysema: association with mortality in the COPDGen study. *Radiology.* 2018;288(3):859–866. doi:10.1148/radiol.2018172294
54. Lynch DA, Al-Qaisi MA. Quantitative computed tomography in chronic obstructive pulmonary disease. *J Thorac Imaging.* 2013;28(5):284–290. doi:10.1097/RTI.0b013e318298733c
55. Bankier AA, O'Donnell CR, Mai VM, et al. Impact of lung volume on MR signal intensity changes of the lung parenchyma. *J Magn Reson Imaging.* 2004;20(6):961–966. doi:10.1002/jmri.20198
56. Ley-Zaporozhan J, Ley S, Kauczor HU. Morphological and functional imaging in COPD with CT and MRI: present and future. *Eur Radiol.* 2008;18(3):510–521. doi:10.1007/s00330-007-0772-1
57. Radtke T, Crook S, Kaltsakas G, et al. ERS statement on standardisation of cardiopulmonary exercise testing in chronic lung diseases. *Eur Respir Rev.* 2019;28(154):180101. doi:10.1183/16000617.0101-2018
58. Lahaije AJ, Willems LM, van Hees HW, Dekhuijzen PN, van Helvoort HA, Heijdra YF. Diagnostic accuracy of metronome-paced tachypnea to detect dynamic hyperinflation. *Clin Physiol Funct Imaging.* 2013;33(1):62–69. doi:10.1111/j.1475-097X.2012.01164.x
59. O'Donnell DE, Revill SM, Webb KA, et al. Dynamic hyperinflation and exercise intolerance in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 2001;164(5):770–777. doi:10.1164/ajrccm.164.5.2012122
60. Guenette JA, Chin RC, Cory JM, Webb KA, O'Donnell DE. Inspiratory capacity during exercise: measurement, analysis, and interpretation. *Pulm Med.* 2013;2013:956081. doi:10.1155/2013/956081
61. James MD, Phillips DB, Vincent SG, et al. Exertional dyspnoea in patients with mild-to-severe chronic obstructive pulmonary disease: neuromechanical mechanisms. *J Physiol.* 2022;600(18):4227–4245. doi:10.1113/JP283252
62. Calligaro GL, Raine RI, Bateman ME, Cooper CB. Comparing dynamic hyperinflation and associated dyspnea induced by metronome-paced tachypnea versus incremental exercise. *COPD.* 2014;11(1):105–112. doi:10.3109/15412555.2013.841669
63. Weigt SS, Abrazado M, Kleerup EC, Tashkin DP, Cooper CB. Time course and degree of hyperinflation with metronome-paced tachypnea in COPD patients. *COPD.* 2008;5(5):298–304. doi:10.1080/15412550802363428
64. Mannee D, Vis E, Hoekstra-Kuik A, van der Maten J, van 't Hul AJ, van Helvoort H. Is the Metronome-Paced Tachypnea Test (MPT) ready for clinical use? Accuracy of the MPT in a prospective and clinical study. *Respiration.* 2019;97(6):569–575. doi:10.1159/000496290
65. Yoshida T, Amato MBP, Grieco DL, et al. Esophageal manometry and regional transpulmonary pressure in lung injury. *Am J Respir Crit Care Med.* 2018;197(8):1018–1026. doi:10.1164/rccm.201709-1806OC
66. Milic-Emili J, Mead J, Turner JM, Glauser EM. Improved technique for estimating pleural pressure from esophageal balloons. *J Appl Physiol.* 1964;19:207–211. doi:10.1152/jappl.1964.19.2.207
67. Aliverti A, Dellaca R, Pelosi P, Chiumello D, Gaihnoni L, Pedotti A. Compartmental analysis of breathing in the supine and prone positions by optoelectronic plethysmography. *Ann Biomed Eng.* 2001;29(1):60–70. doi:10.1114/1.1332084
68. Laveneziana P, Albuquerque A, Aliverti A, et al. ERS statement on respiratory muscle testing at rest and during exercise. *Eur Respir J.* 2019;53(6). doi:10.1183/13993003.01214-2018
69. Castro AA, Kumpel C, Rangueri RC, et al. Daily activities are sufficient to induce dynamic pulmonary hyperinflation and dyspnea in chronic obstructive pulmonary disease patients. *Clinics.* 2012;67(4):319–325. doi:10.6061/clinics/2012(04)04
70. Chin RC, Guenette JA, Cheng S, et al. Does the respiratory system limit exercise in mild chronic obstructive pulmonary disease? *Am J Respir Crit Care Med.* 2013;187(12):1315–1323. doi:10.1164/rccm.201211-1970OC
71. Budweiser S, Harlacher M, Pfeifer M, Jorres RA. Co-morbidities and hyperinflation are independent risk factors of all-cause mortality in very severe COPD. *COPD.* 2014;11(4):388–400. doi:10.3109/15412555.2013.836174
72. Shin TR, Oh YM, Park JH, et al. The prognostic value of residual volume/total lung capacity in patients with chronic obstructive pulmonary disease. *J Korean Med Sci.* 2015;30(10):1459–1465. doi:10.3346/jkms.2015.30.10.1459
73. Oh YM, Sheen SS, Park JH, et al. Emphysematous phenotype is an independent predictor for frequent exacerbation of COPD. *Int J Tuberc Lung Dis.* 2014;18(12):1407–1414. doi:10.5588/ijtld.14.0205

74. Trudzinski FC, Jorres RA, Alter P, et al. Sex-specific associations of comorbidity and pulmonary morbidity with mortality in chronic obstructive pulmonary disease: results from COSYCONET. *Sci Rep.* 2022;12(1):8790. doi:10.1038/s41598-022-12828-8
75. Zeng S, Tham A, Bos B, Jin J, Giang B, Arjomandi M. Lung volume indices predict morbidity in smokers with preserved spirometry. *Thorax.* 2019;74(2):114–124. doi:10.1136/thoraxjnl-2018-211881
76. Zysman M, Deslee G, Perez T, et al. Burden and Characteristics of Severe Chronic Hypoxemia in a Real-World Cohort of Subjects with COPD. *Int J Chron Obstruct Pulmon Dis.* 2021;16:1275–1284. doi:10.2147/COPD.S295381
77. Saure EW, Eagan TM, Jensen RL, et al. Predictors for PaO₂ and hypoxemic respiratory failure in COPD-A three-year follow-up. *COPD.* 2014;11(5):531–538. doi:10.3109/15412555.2014.898027
78. Begin P, Grassino A. Inspiratory muscle dysfunction and chronic hypercapnia in chronic obstructive pulmonary disease. *Am Rev Respir Dis.* 1991;143(5):905–912. doi:10.1164/ajrccm/143.5_Pt_1.905
79. Rochester DF, Braun NM. Determinants of maximal inspiratory pressure in chronic obstructive pulmonary disease. *Am Rev Respir Dis.* 1985;132(1):42–47. doi:10.1164/arrd.1985.132.1.42
80. Rodriguez-Roisin R, Drakulovic M, Rodriguez DA, Roca J, Barbera JA, Wagner PD. Ventilation-perfusion imbalance and chronic obstructive pulmonary disease staging severity. *J Appl Physiol.* 2009;106(6):1902–1908. doi:10.1152/jappphysiol.00085.2009
81. Smith JR, Johnson BD, Olson TP. Impaired central hemodynamics in chronic obstructive pulmonary disease during submaximal exercise. *J Appl Physiol.* 2019;127:691–697. doi:10.1152/jappphysiol.00877.2018
82. Aliverti A, Macklem PT. How and why exercise is impaired in COPD. *Respiration.* 2001;68(3):229–239. doi:10.1159/000050502
83. Neder JA, Rocha A, Berton DC, O'Donnell DE. Clinical and physiologic implications of negative cardiopulmonary interactions in coexisting chronic obstructive pulmonary disease-heart failure. *Clin Chest Med.* 2019;40(2):421–438. doi:10.1016/j.ccm.2019.02.006
84. Butler J, Schrijen F, Henriquez A, Polu JM, Albert RK. Cause of the raised wedge pressure on exercise in chronic obstructive pulmonary disease. *Am Rev Respir Dis.* 1988;138(2):350–354. doi:10.1164/ajrccm/138.2.350
85. Jorgensen K, Muller MF, Nel J, Upton RN, Houltz E, Ricksten SE. Reduced intrathoracic blood volume and left and right ventricular dimensions in patients with severe emphysema: an MRI study. *Chest.* 2007;131(4):1050–1057. doi:10.1378/chest.06-2245
86. Alter P, Watz H, Kahnert K, et al. Airway obstruction and lung hyperinflation in COPD are linked to an impaired left ventricular diastolic filling. *Respir Med.* 2018;137:14–22. doi:10.1016/j.rmed.2018.02.011
87. van der Molen MC, Hartman JE, Vanfleteren L, et al. Reduction of lung hyperinflation improves cardiac preload, contractility, and output in emphysema: a clinical trial in patients who received endobronchial valves. *Am J Respir Crit Care Med.* 2022;206(6):704–711. doi:10.1164/rccm.202201-0214OC
88. Hohlfeld JM, Vogel-Claussen J, Biller H, et al. Effect of lung deflation with indacaterol plus glycopyrronium on ventricular filling in patients with hyperinflation and COPD (CLAIM): a double-blind, randomised, crossover, placebo-controlled, single-centre trial. *Lancet Respir Med.* 2018;6(5):368–378. doi:10.1016/S2213-2600(18)30054-7
89. Travers J, Laveneziana P, Webb KA, Kesten S, O'Donnell DE. Effect of tiotropium bromide on the cardiovascular response to exercise in COPD. *Respir Med.* 2007;101(9):2017–2024. doi:10.1016/j.rmed.2007.03.008
90. Chandra D, Gupta A, Kinney GL, et al. The association between lung hyperinflation and coronary artery disease in smokers. *Chest.* 2021;160(3):858–871. doi:10.1016/j.chest.2021.04.066
91. Rennard SI. Clinical approach to patients with chronic obstructive pulmonary disease and cardiovascular disease. *Proc Am Thorac Soc.* 2005;2(1):94–100. doi:10.1513/pats.200410-051SF
92. Akcay M, Yeter E, Durmaz T, et al. Treatment of acute chronic obstructive pulmonary disease exacerbation improves right ventricle function. *Eur J Echocardiogr.* 2010;11(6):530–536. doi:10.1093/ejehocardiography/jeq013
93. Schrijen FV, Henriquez A, Carton D, Delorme N, Butler J. Pulmonary vascular resistance rises with lung volume on exercise in obstructed airflow disease. *Clin Physiol.* 1989;9(2):143–150. doi:10.1111/j.1475-097X.1989.tb00965.x
94. Donaldson AV, Maddocks M, Martolini D, Polkey MI, Man WD. Muscle function in COPD: a complex interplay. *Int J Chron Obstruct Pulmon Dis.* 2012;7:523–535. doi:10.2147/COPD.S28247
95. Gea J, Pascual S, Casadevall C, Orozco-Levi M, Barreiro E. Muscle dysfunction in chronic obstructive pulmonary disease: update on causes and biological findings. *J Thorac Dis.* 2015;7(10):E418–438. doi:10.3978/j.issn.2072-1439.2015.08.04
96. Gea J, Agustí A, Roca J. Pathophysiology of muscle dysfunction in COPD. *J Appl Physiol.* 2013;114(9):1222–1234. doi:10.1152/jappphysiol.00981.2012
97. St Croix CM, Morgan BJ, Wetter TJ, Dempsey JA. Fatiguing inspiratory muscle work causes reflex sympathetic activation in humans. *J Physiol.* 2000;529:493–504. doi:10.1111/j.1469-7793.2000.00493.x
98. Gagnon P, Guenette JA, Langer D, et al. Pathogenesis of hyperinflation in chronic obstructive pulmonary disease. *Int J Chron Obstruct Pulmon Dis.* 2014;9:187–201. doi:10.2147/COPD.S38934
99. Rutten EP, Grydeland TB, Pillai SG, et al. Quantitative CT: associations between emphysema, airway wall thickness and body composition in COPD. *Pulm Med.* 2011;2011:419328. doi:10.1155/2011/419328
100. Celli BR, Locantore N, Tal-Singer R, et al. Emphysema and extrapulmonary tissue loss in COPD: a multi-organ loss of tissue phenotype. *Eur Respir J.* 2018;51(2):1702146. doi:10.1183/13993003.02146-2017
101. Nezu K, Yoshikawa M, Yoneda T, et al. The change in body composition after bilateral lung volume reduction surgery for underweight patients with severe emphysema. *Lung.* 2000;178(6):381–389. doi:10.1007/s004080000041
102. Sanders KJC, Klooster K, Vanfleteren L, Slebos DJ, Schols A. CT-derived muscle remodelling after bronchoscopic lung volume reduction in advanced emphysema. *Thorax.* 2019;74(2):206–207. doi:10.1136/thoraxjnl-2018-211931
103. Ohara T, Hirai T, Muro S, et al. Relationship between pulmonary emphysema and osteoporosis assessed by CT in patients with COPD. *Chest.* 2008;134(6):1244–1249. doi:10.1378/chest.07-3054
104. Bai P, Sun Y, Jin J, et al. Disturbance of the OPG/RANK/RANKL pathway and systemic inflammation in COPD patients with emphysema and osteoporosis. *Respir Res.* 2011;12:157. doi:10.1186/1465-9921-12-157
105. Suh YJ, McDonald MN, Washko GR, et al. Lung, fat and bone: increased adiponectin associates with the combination of smoking-related lung disease and osteoporosis. *Chronic Obstr Pulm Dis.* 2018;5(2):134–143. doi:10.15326/jcopdf.5.2.2016.0174

106. Mineo TC, Ambrogi V, Mineo D, Fabbri A, Fabbrini E, Massoud R. Bone mineral density improvement after lung volume reduction surgery for severe emphysema. *Chest*. 2005;127(6):1960–1966. doi:10.1378/chest.127.6.1960
107. de Torres JP, Bastarrika G, Wisnivesky JP, et al. Assessing the relationship between lung cancer risk and emphysema detected on low-dose CT of the chest. *Chest*. 2007;132(6):1932–1938. doi:10.1378/chest.07-1490
108. Gonzalez J, Henschke CI, Yankelevitz DF, et al. Emphysema phenotypes and lung cancer risk. *PLoS One*. 2019;14(7):e0219187. doi:10.1371/journal.pone.0219187
109. Zamarron E, Prats E, Tejero E, et al. Static lung hyperinflation is an independent risk factor for lung cancer in patients with chronic obstructive pulmonary disease. *Lung Cancer*. 2019;128:40–46. doi:10.1016/j.lungcan.2018.12.012
110. Laura M, Leong W, Murray NM, et al. Chronic inflammatory demyelinating polyradiculoneuropathy: MRI study of brain and spinal cord. *Neurology*. 2005;64(5):914–916. doi:10.1212/01.WNL.0000152842.11864.D0
111. O'Donnell DE, Hamilton AL, Webb KA. Sensory-mechanical relationships during high-intensity, constant-work-rate exercise in COPD. *J Appl Physiol*. 2006;101(4):1025–1035. doi:10.1152/jappphysiol.01470.2005
112. Rossi A, Khirani S, Cazzola M. Long-acting beta2-agonists (LABA) in chronic obstructive pulmonary disease: efficacy and safety. *Int J Chron Obstruct Pulmon Dis*. 2008;3(4):521–529. doi:10.2147/COPD.S1353
113. Man WD, Mustfa N, Nikolettou D, et al. Effect of salmeterol on respiratory muscle activity during exercise in poorly reversible COPD. *Thorax*. 2004;59(6):471–476. doi:10.1136/thx.2003.019620
114. Beeh KM, Beier J. The short, the long and the "ultra-long": why duration of bronchodilator action matters in chronic obstructive pulmonary disease. *Adv Ther*. 2010;27(3):150–159. doi:10.1007/s12325-010-0017-6
115. O'Donnell DE, Forkert L, Webb KA. Evaluation of bronchodilator responses in patients with "irreversible" emphysema. *Eur Respir J*. 2001;18(6):914–920. doi:10.1183/09031936.01.00216501
116. Chen C, Jian W, Gao Y, Xie Y, Song Y, Zheng J. Early COPD patients with lung hyperinflation associated with poorer lung function but better bronchodilator responsiveness. *Int J Chron Obstruct Pulmon Dis*. 2016;11:2519–2526. doi:10.2147/COPD.S110021
117. Mahler DA, D'Urzo A, Bateman ED, et al. Concurrent use of indacaterol plus tiotropium in patients with COPD provides superior bronchodilation compared with tiotropium alone: a randomised, double-blind comparison. *Thorax*. 2012;67(9):781–788. doi:10.1136/thoraxjnl-2011-201140
118. Vincken W, Aumann J, Chen H, Henley M, McBryan D, Goyal P. Efficacy and safety of coadministration of once-daily indacaterol and glycopyrronium versus indacaterol alone in COPD patients: the GLOW6 study. *Int J Chron Obstruct Pulmon Dis*. 2014;9:215–228. doi:10.2147/COPD.S51592
119. Hoshino M, Ohtawa J, Akitsu K. Comparison of airway dimensions with once daily tiotropium plus indacaterol versus twice daily Advair(R) in chronic obstructive pulmonary disease. *Pulm Pharmacol Ther*. 2015;30:128–133. doi:10.1016/j.pupt.2014.08.002
120. Beeh KM, Burgel PR, Franssen FME, et al. How do dual long-acting bronchodilators prevent exacerbations of chronic obstructive pulmonary disease? *Am J Respir Crit Care Med*. 2017;196(2):139–149. doi:10.1164/rccm.201609-1794CI
121. O'Donnell DE, Sciruba F, Celli B, et al. Effect of fluticasone propionate/salmeterol on lung hyperinflation and exercise endurance in COPD. *Chest*. 2006;130(3):647–656. doi:10.1378/chest.130.3.647
122. Worth H, Forster K, Eriksson G, Nihlen U, Peterson S, Magnussen H. Budesonide added to formoterol contributes to improved exercise tolerance in patients with COPD. *Respir Med*. 2010;104(10):1450–1459. doi:10.1016/j.rmed.2010.07.006
123. O'Donnell DE, Bredenkoter D, Brose M, Webb KA. Physiological effects of roflumilast at rest and during exercise in COPD. *Eur Respir J*. 2012;39(5):1104–1112. doi:10.1183/09031936.00096511
124. Ekstrom M, Nilsson F, Abernethy AA, Currow DC. Effects of opioids on breathlessness and exercise capacity in chronic obstructive pulmonary disease. A systematic review. *Ann Am Thorac Soc*. 2015;12(7):1079–1092. doi:10.1513/AnnalsATS.201501-034OC
125. Abdallah SJ, Wilkinson-Maitland C, Saad N, et al. Effect of morphine on breathlessness and exercise endurance in advanced COPD: a randomised crossover trial. *Eur Respir J*. 2017;50(4):1701235. doi:10.1183/13993003.01235-2017
126. Verberkt CA, van den Beuken-van Everdingen MHJ, Schols J, Hameleers N, Wouters EFM, Janssen DJA. Effect of sustained-release morphine for refractory breathlessness in chronic obstructive pulmonary disease on health status: a randomized clinical trial. *JAMA Intern Med*. 2020;180(10):1306–1314. doi:10.1001/jamainternmed.2020.3134
127. Jensen D, Alshail A, Viola R, Dudgeon DJ, Webb KA, O'Donnell DE. Inhaled fentanyl citrate improves exercise endurance during high-intensity constant work rate cycle exercise in chronic obstructive pulmonary disease. *J Pain Symptom Manage*. 2012;43(4):706–719. doi:10.1016/j.jpainsymman.2011.05.007
128. Thoman RL, Stoker GL, Ross JC. The efficacy of pursed-lips breathing in patients with chronic obstructive pulmonary disease. *Am Rev Respir Dis*. 1966;93(1):100–106. doi:10.1164/arrd.1966.93.1.100
129. Nici L, Donner C, Wouters E, et al. American Thoracic Society/European Respiratory Society statement on pulmonary rehabilitation. *Am J Respir Crit Care Med*. 2006;173(12):1390–1413. doi:10.1164/rccm.200508-1211ST
130. Mayer AF, Karloh M, Dos Santos K, de Araujo CLP, Gulart AA. Effects of acute use of pursed-lips breathing during exercise in patients with COPD: a systematic review and meta-analysis. *Physiotherapy*. 2018;104(1):9–17. doi:10.1016/j.physio.2017.08.007
131. Gosselink R, De Vos J, van den Heuvel SP, Segers J, Decramer M, Kwakkel G. Impact of inspiratory muscle training in patients with COPD: what is the evidence? *Eur Respir J*. 2011;37(2):416–425. doi:10.1183/09031936.00031810
132. Spruit MA, Singh SJ, Garvey C, et al. An official American Thoracic Society/European Respiratory Society statement: key concepts and advances in pulmonary rehabilitation. *Am J Respir Crit Care Med*. 2013;188(8):e13–64. doi:10.1164/rccm.201309-1634ST
133. Ramirez-Sarmiento A, Orozco-Levi M, Guell R, et al. Inspiratory muscle training in patients with chronic obstructive pulmonary disease: structural adaptation and physiologic outcomes. *Am J Respir Crit Care Med*. 2002;166(11):1491–1497. doi:10.1164/rccm.200202-0750C
134. Marin JM, Carrizo SJ, Gascon M, Sanchez A, Gallego B, Celli BR. Inspiratory capacity, dynamic hyperinflation, breathlessness, and exercise performance during the 6-minute-walk test in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2001;163(6):1395–1399. doi:10.1164/ajrccm.163.6.2003172
135. Beaumont M, Mialon P, Le Ber C, et al. Effects of inspiratory muscle training on dyspnoea in severe COPD patients during pulmonary rehabilitation: controlled randomised trial. *Eur Respir J*. 2018;51(1):1701107. doi:10.1183/13993003.01107-2017

136. Porszasz J, Emtner M, Goto S, Somfay A, Whipp BJ, Casaburi R. Exercise training decreases ventilatory requirements and exercise-induced hyperinflation at submaximal intensities in patients with COPD. *Chest*. 2005;128(4):2025–2034. doi:10.1378/chest.128.4.2025
137. Vanfleteren MJ, Koopman M, Spruit MA, et al. Effectiveness of pulmonary rehabilitation in patients with chronic obstructive pulmonary disease with different degrees of static lung hyperinflation. *Arch Phys Med Rehabil*. 2018;99(11):2279–2286 e2273. doi:10.1016/j.apmr.2018.05.011
138. Somfay A, Porszasz J, Lee SM, Casaburi R. Dose-response effect of oxygen on hyperinflation and exercise endurance in nonhypoxaemic COPD patients. *Eur Respir J*. 2001;18(1):77–84. doi:10.1183/09031936.01.00082201
139. Jarosch I, Gloeckl R, Damm E, et al. Short-term effects of supplemental oxygen on 6-min walk test outcomes in patients with COPD: a Randomized, placebo-controlled, single-blind, crossover trial. *Chest*. 2017;151(4):795–803. doi:10.1016/j.chest.2016.11.044
140. Alison JA, McKeough ZJ, Leung RWM, et al. Oxygen compared to air during exercise training in COPD with exercise-induced desaturation. *Eur Respir J*. 2019;2019:1.
141. Palange P, Valli G, Onorati P, et al. Effect of heliox on lung dynamic hyperinflation, dyspnea, and exercise endurance capacity in COPD patients. *J Appl Physiol*. 2004;97(5):1637–1642. doi:10.1152/jappphysiol.01207.2003
142. Petrof BJ, Calderini E, Gottfried SB. Effect of CPAP on respiratory effort and dyspnea during exercise in severe COPD. *J Appl Physiol*. 1990;69(1):179–188. doi:10.1152/jappl.1990.69.1.179
143. Appendini L, Patessio A, Zanaboni S, et al. Physiologic effects of positive end-expiratory pressure and mask pressure support during exacerbations of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 1994;149(5):1069–1076. doi:10.1164/ajrccm.149.5.8173743
144. Appendini L, Purro A, Gudjonsdottir M, et al. Physiologic response of ventilator-dependent patients with chronic obstructive pulmonary disease to proportional assist ventilation and continuous positive airway pressure. *Am J Respir Crit Care Med*. 1999;159(5):1510–1517. doi:10.1164/ajrccm.159.5.9804130
145. van 't Hul A, Kwakkel G, Gosselink R. The acute effects of noninvasive ventilatory support during exercise on exercise endurance and dyspnea in patients with chronic obstructive pulmonary disease: a systematic review. *J Cardiopulm Rehabil*. 2002;22(4):290–297. doi:10.1097/00008483-200207000-00013
146. Bianchi L, Foglio K, Pagani M, Vitacca M, Rossi A, Ambrosino N. Effects of proportional assist ventilation on exercise tolerance in COPD patients with chronic hypercapnia. *Eur Respir J*. 1998;11(2):422–427. doi:10.1183/09031936.98.11020422
147. Cardoso DM, Gass R, Sbruzzi G, Berton DC, Knorst MM. Effect of the expiratory positive airway pressure on dynamic hyperinflation and exercise capacity in patients with COPD: a meta-analysis. *Sci Rep*. 2020;10(1):13292. doi:10.1038/s41598-020-70250-4
148. Diaz O, Begin P, Torrealba B, Jover E, Lisboa C. Effects of noninvasive ventilation on lung hyperinflation in stable hypercapnic COPD. *Eur Respir J*. 2002;20(6):1490–1498. doi:10.1183/09031936.02.00034402
149. Navalesi P, Costa R, Ceriana P, et al. Non-invasive ventilation in chronic obstructive pulmonary disease patients: helmet versus facial mask. *Intensive Care Med*. 2007;33(1):74–81. doi:10.1007/s00134-006-0391-3
150. Windisch W, Geiseler J, Simon K, Waltersbacher S, Dreher M; on behalf of the Guideline C. German national guideline for treating chronic respiratory failure with invasive and non-invasive ventilation - revised edition 2017: part 2. *Respiration*. 2018;96(2):171–203. doi:10.1159/000488667
151. Freitas C, Serino M, Araujo D, et al. Long-term home non-invasive ventilation in chronic hypercapnic chronic obstructive pulmonary disease patients: real-world impact on lung function, acute exacerbations and survival. *Clin Respir J*. 2021;15(10):1130–1138. doi:10.1111/crj.13419
152. Clini E, Sturani C, Rossi A, et al. The Italian multicentre study on noninvasive ventilation in chronic obstructive pulmonary disease patients. *Eur Respir J*. 2002;20(3):529–538. doi:10.1183/09031936.02.02162001
153. Lopez-Campos JL, Soler-Cataluna JJ, Miravittles M. Global strategy for the diagnosis, management, and prevention of chronic obstructive lung disease 2019 report: future challenges. *Arch Bronconeumol*. 2020;56(2):65–67. doi:10.1016/j.arbr.2019.06.014
154. Lammi MR, Ciccolella D, Marchetti N, Kohler M, Criner GJ. Increased oxygen pulse after lung volume reduction surgery is associated with reduced dynamic hyperinflation. *Eur Respir J*. 2012;40(4):837–843. doi:10.1183/09031936.00169311
155. Group NETTR. Patients at high risk of death after lung-volume-reduction surgery. *N Engl J Med*. 2001;345(15):1075–1083. doi:10.1056/NEJMoa11798
156. Hopkinson NS, Toma TP, Hansell DM, et al. Effect of bronchoscopic lung volume reduction on dynamic hyperinflation and exercise in emphysema. *Am J Respir Crit Care Med*. 2005;171(5):453–460. doi:10.1164/rccm.200407-961OC
157. Hartman JE, Vanfleteren L, van Rikxoort EM, Klooster K, Slebos DJ. Endobronchial valves for severe emphysema. *Eur Respir Rev*. 2019;28(152):180121. doi:10.1183/16000617.0121-2018
158. Hartman JE, Welling JBA, Klooster K, Carpaij OA, Augustijn SWS, Slebos DJ. Survival in COPD patients treated with bronchoscopic lung volume reduction. *Respir Med*. 2022;196:106825. doi:10.1016/j.rmed.2022.106825
159. Sciruba FC, Criner GJ, Strange C, et al. Effect of endobronchial coils vs usual care on exercise tolerance in patients with severe emphysema: the RENEW randomized clinical trial. *JAMA*. 2016;315(20):2178–2189. doi:10.1001/jama.2016.6261
160. Brock JM, Kontogianni K, Sciruba FC, Criner GJ, Herth F. Utility of rehabilitation prior to bronchoscopic lung volume reduction: post hoc analysis of the VENT trial. *ERJ Open Res*. 2024;10(1):00735–2023. doi:10.1183/23120541.00735-2023
161. Ries AL, Make BJ, Lee SM, et al. The effects of pulmonary rehabilitation in the national emphysema treatment trial. *Chest*. 2005;128(6):3799–3809. doi:10.1378/chest.128.6.3799
162. van der Molen MC, Posthuma R, Hartman JE, et al. Impact and timing of pulmonary rehabilitation in patients undergoing bronchoscopic lung volume reduction with endobronchial valves: a multicentre randomized controlled trial in patients with severe emphysema. *Respirology*. 2024. doi:10.1111/resp.14734
163. Herth FJ, Gompelmann D, Stanzel F, et al. Treatment of advanced emphysema with emphysematous lung sealant (AeriSeal(R)). *Respiration*. 2011;82(1):36–45. doi:10.1159/000322649
164. Gompelmann D, Eberhardt R, Ernst A, et al. The localized inflammatory response to bronchoscopic thermal vapor ablation. *Respiration*. 2013;86(4):324–331. doi:10.1159/000354175
165. Singh D, Agusti A, Anzueto A, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive lung disease: the GOLD science committee report 2019. *Eur Respir J*. 2019;53(5):1900164. doi:10.1183/13993003.00164-2019

166. Refaely Y, Dransfield M, Kramer MR, et al. Biologic lung volume reduction therapy for advanced homogeneous emphysema. *Eur Respir J*. 2010;36(1):20–27. doi:10.1183/09031936.00106009
167. Kramer MR, Refaely Y, Maimon N, Rosengarten D, Fruchter O. Bilateral endoscopic sealant lung volume reduction therapy for advanced emphysema. *Chest*. 2012;142(5):1111–1117. doi:10.1378/chest.12-0421

International Journal of Chronic Obstructive Pulmonary Disease

Dovepress

Publish your work in this journal

The International Journal of COPD is an international, peer-reviewed journal of therapeutics and pharmacology focusing on concise rapid reporting of clinical studies and reviews in COPD. Special focus is given to the pathophysiological processes underlying the disease, intervention programs, patient focused education, and self management protocols. This journal is indexed on PubMed Central, MedLine and CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-chronic-obstructive-pulmonary-disease-journal>