

Association between Anemia and COPD in Iranian Population

Roshanak Hasheminasab Zavarreh, MD¹. Mohammad-Mehdi Zahmatkesh, MD¹. Masood Vakili, MD². Ali Shahriari-Ahmadi, MD². Mohammad Ali zohal, MD³. Mohsen Arabi, MD². Alireza Mahmoudian, PhD⁴. Abbas gheisuri, MD². Abdolhamed Kian, MD². Ali Fahimi, MD².

¹Pulmonary Diseases Research Center, Rasoul-e-Akram hospital, Iran University of Medical Sciences, Tehran, Iran

²Hematology-Oncology Department Of Rasoul-e Akram Hospital, Iran University Of medical sciences, Tehran, Iran

³Associate professor of internal medicine, pulmonologist, Qazvin University of medical since, Qazvin, Iran

⁴Microbiology Department of Oroumieh University, Oroumieh, Iran

Corresponding author: Mohsen Arabi, MD

Hematology-Oncology Department Of Rasoul-e Akram Hospital, Tehran University Of medical sciences, Tehran, Iran

Email: arabimohsen@yahoo.com

Tel: 09127667297

Received: 20, Feb, 2013

Accepted: 1, Mar, 2013

ABSTRACT

Background and Aim: Chronic obstructive pulmonary disease (COPD) is one of the major causes of morbidity and mortality in adults. Anemia is known as comorbidity in many chronic diseases that can increase morbidity and mortality of COPD. Recent studies have shown that anemia may be more prevalent than expected in COPD patients and can increase disabilities of COPD. In this study we have evaluated the correlation between anemia and the severity of COPD in patients referred to teaching hospitals of the Tehran University of Medical Sciences (TUMS), Tehran, Iran.

Materials and Methods: In this cross-sectional study the severity of COPD in 760 patients with dyspnea who referred to teaching hospitals of Tehran University of Medical Sciences and 96 stable COPD patients were categorized using a GOLD criteria from mild to moderate, severe and very severe. Anemia was determined as hemoglobin <13 g/dL in men and <12 g/dL in women, respectively. Demographic characteristics, spirometry parameters and laboratory findings were compared between anemic and non-anemic groups using Student *t*-test and regression tests (SPSS v.18 software).

Results: The Mean age of patients was 65 ± 13.07 years (59.4% male). Overall prevalence of anemia was 27% and there was no correlation between severity of COPD and anemia. Anemic patients were significantly older than non-anemic patients (71.1 ± 8.5 years vs. 65.4 ± 12.8 years; $p=0.030$). RBC count of anemic patients were significantly lower than non-anemic group (4.3 ± 0.5 vs. $5.02 \pm 0.8 \times 10^6/\mu\text{L}$; $p<0.001$). Erythropoietin levels in anemic group was significantly higher than non-anemic group (16.33 ± 2.43 vs. 10.22 ± 2.67 mu/ml; $p<0.001$) and there was a significant inverse correlation of hemoglobin vs erythropoietin ($r= -0.8$).

Conclusion: there was a high prevalence of anemia in COPD patients. Anemia can increase disabilities of COPD. Thus, treatment of anemia may improve quality of life in these patients. Further comprehensive studies are needed for determination of exact prevalence of anemia and its physiologic effects in COPD.

KEY WORDS: COPD, Anemia.

INTRODUCTION

Chronic obstructive pulmonary disease (COPD) is one of the major causes of morbidity and mortality in adults. As a result of oxygen deficiency, polycythemia is common in patients with COPD, but prevalence rate of anemia in COPD patients is

reported about 13-33 %.¹⁻⁴ COPD can affect main extrapulmonary organs, including heart, brain, and musculoskeletal system. It has been shown that there is an increase in level of proinflammatory cytokines, including interleukin-1 (IL-1), interleukin-6 (IL-6) and tumor necrosis factor- α (TNF- α).⁵⁻⁶

Anemia is a known comorbidity in some disorders, including chronic renal failure, congestive heart failure, infections and cancer, and increases morbidity and mortality and impacts the quality of life.⁷⁻¹¹ Anemia can be a poor prognostic factor in COPD patients.¹⁴ Recently, some studies have been shown that anemia may be more prevalent than previous estimations in COPD patients. It has been shown that 10-15 percent of patients with severe forms of COPD concurrently suffer anemia.¹² In another cohort study, a direct association between hemoglobin levels and mortality of COPD patients was shown.¹³ It has also showed that anemia can be associated with limitation in physical activity and worsen dyspnea in chronic disorders.¹⁵ There are limited and controversial information about the relationship between the anemia and COPD, thus in the current study we tried to investigate the relation between anemia and severity of COPD.

MATERIALS AND METHODS

In this study, 760 COPD patients who were referred to pulmonary clinic of teaching hospitals of Tehran University of Medical Sciences, including 96 stable COPD were investigated during 2010 to 2012. Spirometry was used to confirm the COPD in the patients. All COPD patients were clinically stable and under therapy. 74 patients with mean age of 65±13.07 (23 to 87 years old) entered the study. Patients who had no history of infection or had not been admitted during past 3 months, had no phlebotomy history during last year, had no coexisting diseases, including cancer, thyroid disease, liver or kidney disease and sever heart diseases, and had no history of active bleeding or other hemorrhages and folic acid or vitamin B12 deficiency were included in this study. Demographic characteristics (age, gender and smoking status), laboratory variables (complete blood count and

differential), plasma ferritin level, reticulocyte index, plasma TSH level, blood urea nitrogen and plasma creatinine level, aspartate transaminase (AST) and alanine transaminase (ALT) levels, plasma erythropoietin (Epo), total iron binding capacity (TIBC), Serum iron, erythrocyte sedimentation rate (ESR) and C reactive protein (CRP) were measured. Lung function tests were done. Cigarette smoking was measured as pack-years. By using Global Initiative for Chronic Obstructive Lung Disease (GOLD) criteria to determine, the severity of COPD, patients were classified into 4 groups (mild, moderate, severe, and very severe). Anemia was confirmed by hemoglobin <13 g/dL in men and < 12 g/dL in women, respectively, as explained previously.¹⁴⁻¹⁷ This study was approved by institutional review board in Tehran University of Medical Sciences, Tehran, Iran.

Data Analysis

All statistical analysis, including prevalence rate, mean and standard deviation, were performed using SPSS version 18.0. Chi-square, T-test, regression, and correlation were used to determine the association between different variants.

RESULTS

From 760 patients with dyspnea, 96 patients had an stable COPD, of those 74 patient had no other diseases. Forty-five patients were men (61%) and 29 were women (39%). Forty-three patients (58%) had a positive history of smoking and 31 patients had a history of inhalation of indoor air pollution, including cooking stoves or exposure to passive cigarette smoke. The mean cigarette consumption in smokers (with age of 20-65 years) was 36.6 ± 13.2 pack/year. Demographic characters of patients are shown in Table 1.

Table1. Comparison between Anemic and non-Anemic patients' Features

	Anemic group		Non-anemic group		P value
Mean age	8.5 ± 71.1		12.8 ± 65.4		0.03
COPD Severity	Mild	0 (0.0%)	(4.1%) 3	> 0.05	
	Moderate	9 (45%)	21 (40.5%)		
	Severe	7 (35%)	18 (33.8%)		
	Very severe	4 (20%)	12 (21.6%)		
Gender	Male	14 (70%)	31(57.4%)	> 0.05	
	Female	6 (30%)	23 (42.6%)		
Smoking	Smoker	13 (65%)	30 (55.6%)	> 0.05	
	Nonsmoker	7 (35%)	24 (44.4%)		

Based on the severity of COPD, patients were classified into 4 groups including: mild (4.1%), moderate (40.5%), severe (33.8%), and very severe (21.6%). Overall, 20 patients (27%) were identified to suffer concurrent anemia. The mean age of COPD patients with and without anemia were 71.1 ± 8.5 vs. 65.4 ± 12.8 years ($p=0.03$), respectively. RBC count in anemic patients with COPD was significantly lower than non-anemic patients (4.3 ± 0.5 vs. 5.02 ± 0.8 million per micro-liter; $p<0.001$). ESR rate in anemic patients with COPD was significantly higher than non-anemic patients (29.15 ± 29.36 mm/hr vs. 17.42 ± 13.7 mm/hr; $p=0.02$). The erythropoietin level in anemic patients with COPD was significantly higher than non-anemic patients (16.33 ± 2.43 mu/ml vs. 10.22 ± 2.67 ; $p<0.001$) (Table 2).

Table2. Hematological Findings in Patients with COPD

Variable	Range	SD± Mean
Hb(g/dl)	9.2-19	1.8 ± 13.65
FEV1%	17.9-90	17 ± 48.2
FEV1/FVC %	27.9-90	13.09 ± 56.4
WBC ($10^3/\mu\text{L}$)	3.4-21	3.5 ± 8.6
RBC ($10^6/\mu\text{L}$)	3.2-7.6	0.8 ± 4.8
Hct (%)	30.4-58	5.5 ± 43.4
Plt ($10^3/\mu\text{L}$)	105-394	16.9 ± 63.06
ESR (mm/hr)	1-25	9.5 ± 20.65
BUN (mg/dl)	9-28	8.97 ± 19.2
Cr(mg/dl)	.6-1.6	0.6 ± 1.09
SGOT (IU/L)	10-42	8.3 ± 33.7
SGPT(IU/L)	7-38	5.2 ± 28.8
EPO (mu/ml)	3-21	3.7 ± 11.75
Ferritine (ng/dl)	77-285	8.6 ± 90.5
BMI	15.57-43.96	6.9 ± 26.07

There was a reverse association between hemoglobin and erythropoietin levels. ($r= -0.8$) (Figure1)

The severity of the COPD determined by spirometry was not significantly different between anemic and non-anemic groups ($p>0.05$) (Table 3).




Fig1. Relation between hemoglobin and erythropoietin levels in anemic and non-anemic

Table3. Comparison of Spirometric and Laboratory Results in Anemic and non-Anemic Patients with COPD

	Anemic group	Non anemic group	P value
Age	8.5 ± 71.1	12.8 ± 65.4	0.03
FEV1%	17.46 ± 49.37	17.37 ± 46.33	>0.05
FEV1/FVC %	13.92 ± 56.43	13.11 ± 56.09	>0.05
WBC ($10^3/\mu\text{L}$)	5.33 ± 9.87	2.67 ± 8.22	>0.05
RBC($10^6/\mu\text{L}$)	0.52 ± 4.22	0.72 ± 5.02	<0.001
Hct (%)	0.9 ± 2.45	1.1 ± 2.1	>0.05
Plt ($10^3/\mu\text{L}$)	66.51 ± 243.62	65.6 ± 237.4	>0.05
ESR (mm/hr)	29.36 ± 29.15	13.17 ± 17.42	0.02
BUN (mg/dl)	13.7 ± 2.94	6.53 ± 18.59	>0.05
Cr (mg/dl)	0.25 ± 1.02	0.24 ± 1.04	>0.05
SGOT (IU/L)	15.69 ± 26.3	43.9 ± 36.6	>0.05
SGPT(IU/L)	25.17 ± 24.1	38.64 ± 30.68	>0.05
EPO (mu/ml)	2.46 ± 16.33	2.67 ± 10.22	<0.001
Ferritin (ng/dl)	68.07 ± 80.56	53.23 ± 127.53	>0.05
BMI	6.1 ± 23.83	6.2 ± 25.58	>0.05

DISCUSSION

The current study showed that the prevalence of anemia in COPD patients according to the WHO criteria was 27%. Previous studies have shown that prevalence of anemia in COPD patients can vary 10-15 %. In our study the prevalence of anemia was 27%, which is higher than the findings from previous studies^{12, 19-21} that they mostly investigate greater sample size. However, most of those studies had been retrospective and were performed in several hospitals or different laboratories that failed to use the same clinical or laboratory methods to confirm the disease. In the current study, we examined all patients using a standard criteria for diagnosis of COPD and all laboratory procedures were performed at the same center by using the same diagnostic materials. Recently it has been shown that the high levels of erythropoietin in COPD patients indicates an erythropoietin resistance.¹⁸ In that study, a reverse relationship was found between hemoglobin and erythropoietin level that attributed to a strong stimulating effect of severe hypoxia at higher stages of disease on erythropoietin production.¹⁴ Several studies have been shown that regulation of blood oxygen level and EPO production is a complicated process during which, different factors such as cytokines²² or therapeutic agents, including the inhibitors of angiotensin converting enzyme or angiotensin receptor blockers,^{25, 26} can play an important role in inhibition of body responses to oxygen level changes. Chronic COPD has a greater role in initiation of anemia than a severe COPD. However, several studies stated that anemia can be induced by extensive inflammation following COPD or failure in response towards the erythropoietin.^{6,16,17,21,22} but this hypothesis needs to be backed by more evidences. Dysfunction in transportation of the Iron reticuloendothelial resources can cause anemia in COPD patients.^{23, 24}

Ageing has been also believed to be responsible for greater prevalence of anemia in COPD patients.^{22,10,15,18} Annual costs for health services in anemic COPD patients are as twice more than non-anemic patients.¹⁸ It has also been shown that anemic COPD patients had more comorbidities and mortalities.¹⁰ In addition to anemia, other health issues, including neurologic, respiratory and

musculoskeletal disorders are more common in COPD patients.¹⁰

CONCLUSION

Our study showed that the prevalence of anemia in COPD patients is higher than that was previously stated by other studies (27%) and there was no significant relation between anemia and the severity of COPD. further investigation are required to better understand the association between the COPD and anemia and to gain a broader insights towards the impacts of these diseases in Iranian population.

REFERENCES

1. Krishnan G, Grant BJ, Muti PC, Mishra A, Ochs-Balcom HM, Freudenheim JL. Association between anemia and quality of life in a population sample of individuals with chronic obstructive pulmonary disease. *BMC Pulm Med.* 2006 Sep 5; 6:23.
2. Almagro P, Calbo E, OchoadeEchaguen A, Mortality after hospitalization for COPD. *Chest* 2002; 121:1441–1448.
3. Gerardi DA, Lovett L, Benoit-ConnorsML, Reardon JZ, Zu Wallack RL. Variables related to increased mortality Following out-patient pulmonary rehabilitation. *Eur Respir J* 1996;9:431–435.
4. Hersh CP, DeMeo DL, Al-Ansari E, Predictors of survival in severe, early onset COPD. *Chest* 2004; 126:1443–1451.
5. Landbo C, Prescott E, LangeP, Vestbo J, Almdal TP. Prognostic value of nutritional status in chronic obstructive Pulmonary disease. *Am J Respir Crit Care Med* 1999;160: 1856–1861.
6. John M, Hoernig S, Doehner W, Okonko DD, Witt C, Anker SD. Anemia and inflammation in COPD. *Chest.* 2005 Mar; 127(3):825-9.
7. Mascitelli L, Pezzetta F. Anemia and COPD. *Chest.* 2005 Oct; 128(4):3084.
8. Standardization of spirometry 1994 update, American thoracic society *Amj Respi rit care med* 1995, 152 (3):7107-1136
9. Hurd s, pauwels R. Global initiative for chronic obstructive long diseases (GOLD) *pulm pharmacol ther* 2002,15(4):353-355.
10. Halpern MT, Zilberberg MD, Schmier JK, Lau EC, Shorr AF. Anemia, costs and mortality in chronic obstructive pulmonary disease. *Cost Eff Resour Alloc.* 2006 Oct 16; 4:17.
11. American thoracic society standards for the diagnosis & care of patients with chronic obstructive

- pulmonary disease. *Amj respircit care med.* 1995; 152: 77-121.
12. Penninx BW, Pahor M, Cesari M, et al. Anaemias Associated with disability and decreased physical performance and muscle strength in the elderly. *J Am Geriatr Soc* 2004; 52:719–724.
 13. Mansen TJ, McCance KL, Parker-Cohen PD. Alterations of erythrocyte function. In: McCance KL, Huether SE, eds. *Pathophysiology. The Biologic Basis for Disease in Adults and Children.* 2nd Edn. St.Louis, Mosby-Year Book Inc., 1994; pp.860–877.
 14. Reilly JJ, Edwin JR, Silverman K, et al. chronic obstructive pulmonary disease in: Braunwald, Fauci, Kasper et al (editors), *Harrisons principals of internal medicine*, 18th ed, New York. Mc Graw hill 2012. p.1635-1643
 15. Cote C, Zilberberg MD, Mody SH, Dordelly LJ, Celli B. Haemoglobin level and its clinical impact in a cohort of patients with COPD. *Eur Respir J.* 2007 May; 29(5):923-9.
 16. John M, Lange A, Hoernig S, Witt C, Anker SD. Prevalence of anemia in chronic obstructive pulmonary disease: comparison to other chronic diseases. *Int J Cardiol.* 2006 Aug 28; 111(3):365-70.
 17. Kilburn KH, Asmundsson T. Factors influencing the course of COPD. *Postgrad Med.* 1973 Sep; 54(3):135-41.
 18. Shorr AF, Doyle J, Stern L, Dolgitsers M, Zilberberg MD. Anemia in chronic obstructive pulmonary disease: epidemiology and economic implications. *Curr Med Res Opin.* 2008 Apr; 24(4):1123-30.
 19. Similowski T, Agustí A, MacNee W, Schönhofer B. The potential impact of anaemia of chronic disease in COPD. *Eur Respir J.* 2006 Feb; 27(2):390-6.
 20. Chambellan A, Chailleux E, Similowski T; ANTADIR Observatory Group. Prognostic value of the hematocrit in patients with severe COPD receiving long-term oxygen therapy. *Chest.* 2005 Sep; 128(3):1201-8.
 21. Singh A. Is 13 g.dL(-1) the threshold to correct anaemia in COPD *Eur Respir J.* 2007;30(5):1024-5.
 22. Pavlisa G, Vrbanic V, Kusec V, Jaksic B. Erythropoietin response after correction of severe hypoxaemia due to acute respiratory failure in chronic obstructive pulmonary disease patients. *Clin Sci (Lond).* 2004 Jan; 106(1):43-51.
 23. Schols AM. Pulmonary cachexia. *Int J Cardiol* 2002; 85:101–110.
 24. Means RT Jr. Advances in the anemia of chronic disease. *Int J Hematol* 1999; 70:7–12.
 25. Marathias KP, Agroyannis B, Mavromoustakos T, et al. Hematocrit-lowering effect following inactivation of rennin angiotensin system with angiotensin converting enzyme inhibitors and angiotensin receptor blockers. *Curr Top Med Chem* 2004;4:483–486
 26. Ishani A, Weinhandl E, Zhao Z, et al. Angiotensin converting enzyme inhibitor as a risk factor for the development of anemia, and the impact of incident anemia on mortality in patients with Left ventricular dysfunction. *Jam Coll Cardiol* 2005;45:391–399