

# Which foods are displaced in the diets of adults with type 2 diabetes with the inclusion of eggs in their diets? A randomized, controlled, crossover trial

Valentine Y Njike, Rachel Annam, Victoria Christina Costales, Niloufarsadat Yarandi, David L Katz

**To cite:** Njike VY, Annam R, Costales VC, *et al.* Which foods are displaced in the diets of adults with type 2 diabetes with the inclusion of eggs in their diets? A randomized, controlled, crossover trial. *BMJ Open Diab Res Care* 2017;**5**:e000411. doi:10.1136/bmjdr-2017-000411

Received 6 March 2017  
Revised 2 May 2017  
Accepted 5 June 2017

## ABSTRACT

**Background** The inclusion or exclusion of specific foods from the overall diet inevitably affects other food choices, and this matter is routinely neglected in dietary guidance and nutritional epidemiology. We examined how the inclusion of eggs in the diets of type 2 diabetics affected dietary pattern.

**Methods** Randomized, controlled, single-blind, crossover trial of 34 adults (mean age 64.5 years; 14 women, 20 men) with type 2 diabetes assigned to one of two possible sequence permutations of two different 12-week treatments (two eggs/day or egg exclusion), with 6-week washout periods. For the egg inclusion phase, participants received advice from a dietitian on how to preserve an isocaloric condition relative to the egg exclusion phase. To assess changes in dietary pattern in the diets of our study participants, we analyzed the 12 components of the 2010 Healthy Eating Index.

**Results** The inclusion of eggs was associated with reduced consumption of refined grains nearing statistical significance ( $-0.7 \pm 3.4$  vs  $0.7 \pm 2.2$ ;  $p=0.0530$ ). The consumption of total protein foods significantly increased from baseline ( $0.3 \pm 0.7$ ;  $p=0.0153$ ) with the inclusion of eggs for 12 weeks, while the consumption of dairy products significantly decreased with the exclusion of eggs from their diets ( $-1.3 \pm 2.9$ ;  $p=0.0188$ ).

**Conclusions** Eggs in the diets of type 2 diabetics may lead to increased consumption of some healthful foods and reduced consumption of some less healthful foods.

**Trial registration number** NCT02052037; Post-results.

## BACKGROUND

Diabetes is a public health problem of epidemic proportions. According to the 2014 National Diabetes Statistics Report, 29.1 million people, or 9% of the US population, are estimated to have diabetes, of which 21.0 million people are diagnosed.<sup>1</sup> About 86 million people, or one in three people in the USA, are estimated to have pre-diabetes,<sup>1</sup> yet 9 out of 10 people with pre-diabetes are unaware that they have this condition.<sup>1</sup> Of the people who have pre-diabetes, 15%–30% are likely to develop type 2 diabetes with 5 years.<sup>1</sup>

## Significance of this study

### What is already known about this subject?

- ▶ Foods displaced from ad libitum diets while accommodating for the inclusion of a tested food are routinely not investigated as part of clinical research studies. Few dietary intervention studies have included this type of investigation as part of their study protocol design.
- ▶ We examined the foods displaced from the diets with type 2 diabetes while including eggs as part of their otherwise ad libitum diets.

### What are the new findings?

- ▶ The inclusion of eggs in the diets of type 2 diabetics may improve protein foods consumption.
- ▶ The inclusion of eggs in the diets of type 2 diabetics may reduce consumption of refined grains.

### How might these results change the focus of research or clinical practice?

- ▶ The results of this study may serve as guidance to nutrition researchers and professionals who provide dietary guidance to people with type 2 diabetes on foods to displace when including eggs in their diets.

The risk of death in individuals with diabetes is more than 50% higher than for adults without diabetes. Type 2 diabetes accounts for about 90%–95% of all diagnosed cases of diabetes. Diabetes is the seventh leading cause of death in the USA. Diabetes complications include cardiovascular disease (CVD), stroke, hypertension, blindness, kidney disease, nervous system damage, limb amputations, and biochemical imbalances that can cause acute life-threatening events. Rates of cardiovascular mortality are 2–4 times higher among adults with diabetes than among those without diabetes.<sup>2</sup>

There is no cure for type 2 diabetes. The cornerstone of the management and prevention of type 2 diabetes is lifestyle intervention.<sup>2</sup>



CrossMark

Yale University, Prevention Research Center, Griffin Hospital, Derby, Connecticut, USA

### Correspondence to

Dr David L Katz; davkatz7@gmail.com

Previous studies have shown that a reduction of 5%–7% in body weight can lead to a significant improvement in insulin sensitivity, glycemic control, reduced medication use, and reduced risk of developing type 2 diabetes.<sup>3–6</sup> Low glycemic index foods are typically recommended for patients with type 2 diabetes or at risk for type 2 diabetes. Diets with a low glycemic load have been reported to improve serum lipid profiles, reduce C reactive protein (CRP) levels, and aid in weight control. In cross-sectional studies, they have been associated with higher levels of high-density lipoprotein cholesterol, with reduced CRP concentrations, and, in cohort studies, with decreased risk of developing diabetes and CVD.<sup>7</sup>

In a previous publication,<sup>8</sup> we demonstrated that the inclusion of eggs in the diets of adults with type 2 diabetes led to improved anthropometric measures, but did not describe the foods displaced with the inclusion of eggs in their diets. This report further elaborates the findings from that study by evaluating the foods displaced from the diets of these adults with type 2 diabetes with the inclusion of eggs.

## METHODS

### Design

This is a randomized, single-blind crossover trial designed with a 4-week run-in period and two treatment assignments to compare the effects of 12 weeks of daily inclusion or daily exclusion of eggs on dietary pattern in adults with type 2 diabetes. After a 4-week run-period of ad libitum diet, participants were randomized and then underwent repeated measures following inclusion of 10–14 eggs per week or egg exclusion for 3 months in their diets in random sequence (ie, one of two sequence permutations), with a 6-week washout between treatments. The investigators were blinded to the treatment assignments, while the study participants were not. Details of the study design—including inclusion/exclusion criteria, and randomization process—are presented in our previous report.<sup>8</sup> This study was approved by the Griffin Hospital Institutional Review Board.

### Egg inclusion phase

Participants met with a registered dietitian and received instructions for including two eggs per day (10–14 eggs/week) in their diets, while preserving an isocaloric condition relative to the egg exclusion phase. The study dietitian provided individualized guidance to participants on how to make room for eggs in their diets, while giving them latitude in determining how to adjust for the extra calories from the eggs, to better approximate real-world conditions.

### Egg exclusion phase

Participants also met with the dietitian and receive relevant meal planning guidance and instructions to avoid eggs and specific egg-containing products.

During both intervention phases, study participants were advised to eat to their usual state of fullness, and

dietary monitoring and weighing were conducted to ensure that an isocaloric condition was maintained.

### Outcome measure

We tracked variation in dietary patterns over the course of the study by asking study participants to provide information on the foods and beverages that they consumed by completing 24 hours recalls using a web-based Automated Self-Administered 24 hours Recall (ASA24) (<http://risk-factor.cancer.gov/tools/instruments/asa24/>). Three 24 hours recalls (ie, for two weekdays and one weekend day) were collected at each timepoint. The average of the 3-day recalls at each timepoint was used to assess diet quality.

The food groups were assessed using the subscale of the Healthy Eating Index (HEI) 2010. The HEI is a measure of the diet quality, independent of quantity, which is used to assess compliance with the U.S. Dietary Guidelines for Americans and monitor changes in dietary patterns. The HEI is a scoring metric that is used to determine the quality of a given dietary pattern, set of foods, or menu. There are 12 components in the HEI–2010. All of the key Dietary Guidelines food choice recommendations that relate to diet quality are reflected in the HEI–2010's 12 components. Nine of the components focus on adequacy (dietary components to increase), and three focus on moderation (dietary components to decrease). The performance of the HEI–2010 has been evaluated through an assessment of its content validity, construct validity, and reliability.<sup>9</sup> To assess the foods displaced with the inclusion of eggs in the diets, we analyzed the 12 components of the HEI–2010 of our study participants.

### Statistical analysis

Generalized linear models were used to assess difference between the egg inclusion versus egg exclusion diets. Paired Student's *t*-tests were used to assess difference from baseline to endpoints. Regression models were used to adjust for potential confounding factors (ie, age, gender, race, compliance, and treatment sequence). All analyses at end points were based on intention-to-treat principle. *p* Values of <0.05 were considered statistically significant. SAS software for Windows V.9.3 was used to carry out all statistical analyses. Data are presented as mean ± SD except otherwise stated.

## Results

### Demographic characteristics and baseline data

Our study participants were predominantly male (ie, 58.8%) and were mostly white (ie, 76.5%). The average age of the study participants was 64.5 years, and their average diet quality score was 52.9 out of a maximum possible score of 100. The average intakes of the different components of the subscales of the HEI are presented in table 1.

### Efficacy data

When compared with egg exclusion, the inclusion of eggs in the diets of type 2 diabetes for 12 weeks

**Table 1** Baseline characteristics

| Variable                                    | Mean ± SD |
|---|-----------|
| 2010 Healthy Eating Index score             | 52.9±11.0 |
| Total vegetables*                           | 3.5±1.2   |
| Greens and beans*                           | 1.9±1.6   |
| Total fruit <sup>†</sup>                    | 2.6±1.9   |
| Whole fruit <sup>‡</sup>                    | 2.6±1.9   |
| Whole grains                                | 3.3±2.8   |
| Dairy <sup>§</sup>                          | 5.7±2.7   |
| Total protein foods <sup>¶</sup>            | 4.3±0.8   |
| Seafood and plant proteins <sup>¶, **</sup> | 2.0±1.5   |
| Fatty acids <sup>††</sup>                   | 4.2±2.3   |
| Sodium                                      | 3.3±2.4   |
| Refined grains                              | 6.1±2.7   |
| Empty calories <sup>‡‡</sup>                | 13.8±3.9  |

\*Includes any beans and peas not counted as total protein foods.

<sup>†</sup>Includes 100% fruit juice.

<sup>‡</sup>Includes all forms except juice.

<sup>§</sup>Includes all milk products, such as fluid milk, yogurt, and cheese, and fortified soy beverages.

<sup>¶</sup>Beans and peas are included here (and not with vegetables) when the total protein foods standard is otherwise not met.

\*\*Includes seafood, nuts, seeds, soy products (other than beverages), as well as beans and peas counted as total protein foods.

<sup>††</sup>Ratio of polyunsaturated and monounsaturated fatty acids to saturated fatty acids.

<sup>‡‡</sup>Calories from solid fats, alcohol, and added sugars; threshold for counting alcohol is >13 g/1000 kcal.

non-significantly decreased the consumption of refined grains ( $-0.7\pm 3.4$  vs  $0.7\pm 2.2$ ;  $p=0.0530$ ) and non-significantly increased the consumption of total protein foods (ie, beans and peas are included here (and not with vegetables)) ( $0.3\pm 0.7$  vs  $-0.1\pm 1.0$ ;  $p=0.0789$ ). No significant ( $p>0.05$ ) changes were observed during the egg inclusion phase as compared with the egg exclusion phase in intake of total vegetables; greens and beans; total fruit (ie, 100% fruit juice); whole fruit (ie, all forms except juice); whole grains; seafood and plant proteins; fatty acids (ie, ratio of polyunsaturated and monounsaturated fatty acids to saturated fatty acids); sodium; empty calories (ie, calories from solid fats, alcohol, and added sugars); and overall diet quality.

The consumption of total protein foods significantly increased from baseline ( $0.3\pm 0.7$ ;  $p=0.0153$ ) with the inclusion of eggs for 12 weeks. The consumption of dairy products (ie, all milk products, such as fluid milk, yogurt, and cheese, and fortified soy beverages) significantly decreased from baseline with the exclusion of eggs in the diets ( $-1.3\pm 2.9$ ;  $p=0.0188$ ) (see table 2).

## DISCUSSION

The inclusion of eggs in the diets of adults with type 2 diabetes significantly increased the consumption of total

**Table 2** Change in outcome measures from baseline

| Variable                        | Eggs exclusion phase | Eggs inclusion phase | p Value |
|---------------------------------|----------------------|----------------------|---------|
| 2010 Healthy Eating Index score | $-3.5\pm 12.7$       | $-0.1\pm 11.4$       | 0.2572  |
| Total vegetables                | $-0.1\pm 1.5$        | $-0.1\pm 1.1$        | 0.9605  |
| Greens and beans                | $-0.4\pm 2.0$        | $-0.3\pm 1.9$        | 0.7050  |
| Total fruit                     | $0.2\pm 1.0$         | $0.3\pm 1.5$         | 0.8299  |
| Whole fruit                     | $0.4\pm 1.2$         | $0.1\pm 1.7$         | 0.3611  |
| Whole grains                    | $0.4\pm 3.5$         | $-0.6\pm 3.1$        | 0.2143  |
| Dairy                           | $-1.3\pm 2.9^*$      | $-0.5\pm 2.9$        | 0.7767  |
| Total Protein Foods             | $-0.1\pm 1.0$        | $0.3\pm 0.7^*$       | 0.0789  |
| Seafood and plant proteins      | $-0.1\pm 1.8$        | $0.0\pm 1.5$         | 0.7767  |
| Fatty acids                     | $-0.3\pm 3.6$        | $0.1\pm 3.2$         | 0.6103  |
| Sodium                          | $0.7\pm 2.7$         | $0.1\pm 3.0$         | 0.4340  |
| Refined grains                  | $0.7\pm 2.2$         | $-0.7\pm 3.4$        | 0.0530  |
| Empty calories                  | $1.0\pm 5.7$         | $-0.6\pm 3.7$        | 0.1786  |

Values are mean ± SD.

\*Significant ( $p$  value<0.05) change from baseline.

protein foods from baseline. While the exclusion of eggs in the diets of type 2 diabetes significantly decreased the consumption of dairy from baseline. Compared with the exclusion of eggs, the inclusion of eggs non-significantly decreased the intake of refined grains. No meaningful improvements were observed in intake of total vegetables; greens and beans; total fruit; whole fruit; whole grains; seafood and plant proteins; fatty acids; sodium; or empty calories with the inclusion of eggs.

In a meta-analysis by Vigiouliouk *et al*,<sup>10</sup> replacing animal proteins with plant proteins improved glycemic control in individuals with diabetes. In previous epidemiological studies,<sup>11-13</sup> consumption of vegetable proteins was associated with a reduction in type 2 diabetes, while the consumption of animal proteins was associated with an increased risk. In another epidemiological study, Pan *et al*<sup>14</sup> demonstrated an association between animal protein consumption and the risk of type 2 diabetes. A mediation analysis by Li *et al*<sup>12</sup> demonstrated that the effects of proteins on the risk of type 2 diabetes were mediated through the insulin sensitivity. Plant-based proteins such as beans, lentils, peas, and nuts have a low glycemic index, and they provide quality protein, healthy fatty acids, and fiber that help with weight loss and blood glucose control and that improve insulin sensitivity.

While we observed a meaningful reduction of refined grain consumption with the inclusion of eggs in the diets, we did not see a statistically significant reduction. The lack of statistical significance may be due to the small sample size, a placebo effect, and/or the Hawthorne effect. In a recent randomized trial by Kim *et al*,<sup>15</sup> short-term consumption of diets rich in red processed meat

and refined grains led to a decreased in insulin sensitivity compared with diets high in whole grains, nuts, dairy products, and legumes in insulin-resistant adults. In another randomized control trial, whole-grain cereal-based diets compared with refined cereal diets reduced postprandial insulin in individuals with metabolic syndrome.<sup>16</sup> Diets rich in red meat and refined grains have been shown to increase glucose and insulin response, which may increase pancreatic stress that could in turn increase the risk of type 2 diabetes.<sup>17</sup>

The inclusion of eggs in the diets of our study participants did not show any meaningful change in the consumption of dairy products, while the exclusion of eggs in the diets led to a significant reduction. The effects of dairy products on type 2 diabetes are controversial. In prospective epidemiological studies,<sup>18,19</sup> low-fat fermented dairy product consumption was associated with a lower risk of developing type 2 diabetes. In an epidemiological study by Chen *et al*,<sup>20</sup> higher consumption of yogurt was associated with lower risk of type 2 diabetes, while intake of other dairy foods or total dairy was not associated with the incidence of type 2 diabetes risk. A large European prospective study<sup>21</sup> showed no association between total dairy consumption and the risk of type 2 diabetes, but observed an inverse association with cheese consumption and combined fermented dairy. In another study,<sup>22</sup> the consumption of dairy foods was inversely associated with the prevalence of obesity. Wang *et al*<sup>23</sup> showed that higher total dairy and yogurt consumption were associated with body weight and waist circumference maintenance.

We did not see meaningful beneficial effects in consumption of total vegetables; greens and beans; total fruit; whole fruit; whole grains; seafood and plant proteins; fatty acids; sodium; and empty calories with the inclusion of eggs in the diets, possibly due to small sample size, placebo effect, and/or or the Hawthorne effect. Increased dietary consumption of total vegetables; greens and beans; total fruit; whole fruit; whole grains; seafood and plant proteins; and healthful fatty acids, and the reduced consumption of sodium and empty calories has been shown to improve cardiometabolic risk in type 2 diabetes.<sup>24–40</sup>

### Limitations

This study has several limitations. First, the sample size of the study was small, limiting the ability to detect significant findings in some variables. However, the statistical power of the study was improved by using a crossover study design. Second, the study population was predominantly male and white, which limits the ability to extrapolate the study findings to a much broader population demographic. Third, due to the nature of the study intervention, the study participants were not blinded, which could have led to placebo or Hawthorne effects. Fourth, the dietary intakes of the study participants were self-reported. It is possible that the study participants would have underestimated or overestimated their food and beverage consumption. However, their dietary

data were captured using a reliable, validated tool (ie, ASA-24) that provided specific directions to estimate their portion sizes and the percent of each portion size that they actually consumed, which minimized the likelihood of underestimation or overestimation of their dietary intake. In addition, the study dietitian provided guidance to the study participants on how to enter their dietary data. Fifth, the study participants were not supervised while consuming the foods and beverages during each study phase. Therefore, the veracity of these dietary data reported depends on the honesty of the study participants. However, this may be viewed also like strength of this study because it emulates a real-world scenario.

### Conclusions

These data suggest that the inclusion of eggs in the diets of type 2 diabetics facilitated the intake of protein foods and reduced the intake of refined grains. In general, this study highlights the impact of single-food inclusions or exclusions on other dietary choices, and potentially, overall diet quality. The small sample studied here limited the statistical power to identify relevant associations and argues for larger studies to examine the impact of food-specific recommendations on dietary pattern, particularly in at-risk groups such as those with type 2 diabetes.

**Acknowledgements** The authors wish to acknowledge the technical assistance of Susan Acheychek RDMS; Rockiy Ayettey MS; and Judith A. Treu, MS.

**Contributors** VYN: study design, project oversight, data analysis, data interpretation, developed manuscript draft, and approved the final manuscript. RA: assisted with development of manuscript draft. VCC: assisted in writing the manuscript and provided critical review. DLK: study design, project oversight, data interpretation, critical review of paper, and final approval.

**Funding** Funding for this study has been provided by the Egg Nutrition Center.

**Competing interests** None declared.

**Patient consent** Participants filled out study consent form approved by IRB prior to enrollment in the study.

**Ethics approval** Griffin Hospital Institutional Review Board.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** No additional data are available.

**Open Access** This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

### REFERENCES

1. National Diabetes Statistics Report. 2014. <http://www.cdc.gov/diabetes/pubs/statsreport14/national-diabetes-report-web.pdf>
2. Centers for Disease Control and Prevention. National diabetes fact sheet: national estimates and general information on diabetes and prediabetes in the United States. 2011. [http://www.cdc.gov/diabetes/pubs/pdf/ndfs\\_2011.pdf](http://www.cdc.gov/diabetes/pubs/pdf/ndfs_2011.pdf)
3. Knowler WC, Barrett-Connor E, Fowler SE, *et al*. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346:393–403.
4. Pronk NP. Structured diet and physical activity programmes provide strong evidence of effectiveness for type 2 diabetes prevention

- and improvement of cardiometabolic health. *Evid Based Med* 2016;21:18.
5. Ackermann RT. Diabetes Prevention at the Tipping Point: Aligning Clinical and Public Health Recommendations. *Ann Intern Med* 2015;163:475.
  6. Balk EM, Earley A, Raman G, et al. Combined Diet and Physical Activity Promotion Programs to Prevent Type 2 Diabetes Among Persons at Increased Risk: A Systematic Review for the Community Preventive Services Task Force. *Ann Intern Med* 2015;163:437.
  7. Esfahani A, Wong JM, Mirrahimi A, et al. The glycemic index: physiological significance. *J Am Coll Nutr* 2009;28 Suppl:439S–45.
  8. Njike VY, Ayettey RG, Rajebi H, et al. Egg ingestion in adults with type 2 diabetes: effects on glycemic control, anthropometry, and diet quality—a randomized, controlled, crossover trial. *BMJ Open Diabetes Res Care* 2016;4:e000281.
  9. Guenther PM, Kirkpatrick SI, Reedy J, et al. The Healthy Eating Index–2010 is a valid and reliable measure of diet quality according to the 2010 Dietary Guidelines for Americans. *J Nutr* 2014;144:399–407.
  10. Viguiiok E, Stewart SE, Jayalath VH, et al. Effect of Replacing Animal Protein with Plant Protein on Glycemic Control in Diabetes: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Nutrients* 2015;7:9804–24.
  11. Malik VS, Li Y, Tobias DK, et al. Dietary Protein Intake and Risk of Type 2 Diabetes in US Men and Women. *Am J Epidemiol* 2016;183:715–28.
  12. Li J, Sun C, Liu S, et al. Dietary Protein Intake and Type 2 Diabetes Among Women and Men in Northeast China. *Sci Rep* 2016;6:37604.
  13. Satija A, Bhupathiraju SN, Rimm EB, et al. Plant-Based Dietary Patterns and Incidence of Type 2 Diabetes in US Men and Women: Results from Three Prospective Cohort Studies. *PLoS Med* 2016;13:e1002039.
  14. Pan A, Sun Q, Bernstein AM, et al. Changes in red meat consumption and subsequent risk of type 2 diabetes mellitus: three cohorts of US men and women. *JAMA Intern Med* 2013;173:1328–35.
  15. Kim Y, Keogh JB, Clifton PM, et al. Consumption of red and processed meat and refined grains for 4 weeks decreases insulin sensitivity in insulin-resistant adults: a randomized crossover study. *Metabolism* 2017;68:173–83.
  16. Giacco R, Costabile G, Della Pepa G, et al. A whole-grain cereal-based diet lowers postprandial plasma insulin and triglyceride levels in individuals with metabolic syndrome. *Nutr Metab Cardiovasc Dis* 2014;24:837–44.
  17. Kim Y, Keogh J, Clifton P, et al. Differential Effects of Red Meat/ Refined Grain Diet and Dairy/Chicken/Nuts/Whole Grain Diet on Glucose, Insulin and Triglyceride in a Randomized Crossover Study. *Nutrients* 2016;8:11:687.
  18. Díaz-López A, Bulló M, Martínez-González MA, et al. Dairy product consumption and risk of type 2 diabetes in an elderly Spanish Mediterranean population at high cardiovascular risk. *Eur J Nutr* 2016;55:349–60.
  19. O'Connor LM, Lentjes MA, Luben RN, et al. Dietary dairy product intake and incident type 2 diabetes: a prospective study using dietary data from a 7-day food diary. *Diabetologia* 2014;57:909–17.
  20. Chen M, Sun Q, Giovannucci E, et al. Dairy consumption and risk of type 2 diabetes: 3 cohorts of US adults and an updated meta-analysis. *BMC Med* 2014;12:215.
  21. Sluijs I, Forouhi NG, Beulens JW, et al. The amount and type of dairy product intake and incident type 2 diabetes: results from the EPIC-InterAct Study. *Am J Clin Nutr* 2012;96:382–90.
  22. Crichton GE, Alkerwi A. Whole-fat dairy food intake is inversely associated with obesity prevalence: findings from the Observation of Cardiovascular Risk Factors in Luxembourg study. *Nutr Res* 2014;34:936–43.
  23. Wang H, Troy LM, Rogers GT, et al. Longitudinal association between dairy consumption and changes of body weight and waist circumference: the Framingham Heart Study. *Int J Obes* 2014;38:299–305.
  24. Li M, Fan Y, Zhang X, et al. Fruit and vegetable intake and risk of type 2 diabetes mellitus: meta-analysis of prospective cohort studies. *BMJ Open* 2014;4:e005497.
  25. Wu Y, Zhang D, Jiang X, et al. Fruit and vegetable consumption and risk of type 2 diabetes mellitus: a dose-response meta-analysis of prospective cohort studies. *Nutr Metab Cardiovasc Dis* 2015;25:140–7.
  26. Carter P, Gray LJ, Troughton J, et al. Fruit and vegetable intake and incidence of type 2 diabetes mellitus: systematic review and meta-analysis. *BMJ* 2010;341:c4229.
  27. Bazzano LA, Li TY, Joshipura KJ, et al. Intake of fruit, vegetables, and fruit juices and risk of diabetes in women. *Diabetes Care* 2008;31:1311–7.
  28. Horikawa C, Yoshimura Y, Kamada C, et al. Dietary sodium intake and incidence of diabetes complications in Japanese patients with type 2 diabetes: analysis of the Japan Diabetes Complications Study (JDCS). *J Clin Endocrinol Metab* 2014;99:3635–43.
  29. Poggio R, Gutierrez L, Matta MG, et al. Daily sodium consumption and CVD mortality in the general population: systematic review and meta-analysis of prospective studies. *Public Health Nutr* 2015;18:695–704.
  30. Kalogeropoulou AP, Georgiopoulou VV, Murphy RA, et al. Dietary sodium content, mortality, and risk for cardiovascular events in older adults: the Health, Aging, and Body Composition (Health ABC) Study. *JAMA Intern Med* 2015;175:410–9.
  31. Due A, Larsen TM, Mu H, et al. Comparison of 3 ad libitum diets for weight-loss maintenance, risk of cardiovascular disease, and diabetes: a 6-mo randomized, controlled trial. *Am J Clin Nutr* 2008;88:1232–41.
  32. Risérus U, Willett WC, Hu FB, et al. Dietary fats and prevention of type 2 diabetes. *Prog Lipid Res* 2009;48:44–51.
  33. Blanco Mejia S, Kendall CW, Viguiiok E, et al. Effect of tree nuts on metabolic syndrome criteria: a systematic review and meta-analysis of randomised controlled trials. *BMJ Open* 2014;4:e004660.
  34. Ye EQ, Chacko SA, Chou EL, et al. Greater whole-grain intake is associated with lower risk of type 2 diabetes, cardiovascular disease, and weight gain. *J Nutr* 2012;142:1304–13.
  35. Aune D, Norat T, Romundstad P, et al. Whole grain and refined grain consumption and the risk of type 2 diabetes: a systematic review and dose-response meta-analysis of cohort studies. *Eur J Epidemiol* 2013;28:845–58.
  36. Sun Q, Spiegelman D, van Dam RM, et al. White rice, brown rice, and risk of type 2 diabetes in US men and women. *Arch Intern Med* 2010;170:961–9.
  37. Pan A, Malik VS, Schulze MB, et al. Plain-water intake and risk of type 2 diabetes in young and middle-aged women. *Am J Clin Nutr* 2012;95:1454–60.
  38. Fagherazzi G, Vilier A, Saes Sartorelli D, et al. Consumption of artificially and sugar-sweetened beverages and incident type 2 diabetes in the Etude Epidemiologique aupres des femmes de la Mutuelle Generale de l'Education Nationale-European Prospective Investigation into Cancer and Nutrition cohort. *Am J Clin Nutr* 2013;97:517–23.
  39. Patel PS, Forouhi NG, Kuijsten A, et al. The prospective association between total and type of fish intake and type 2 diabetes in 8 European countries: EPIC-InterAct Study. *Am J Clin Nutr* 2012;95:1445–53.
  40. Wu JH, Micha R, Imamura F, et al. Omega-3 fatty acids and incident type 2 diabetes: a systematic review and meta-analysis. *Br J Nutr* 2012;107 Suppl 2:S214–27.