

Automated Analysis of Diffusion-Weighted Magnetic Resonance Imaging for the Differential Diagnosis of Multiple System Atrophy from Parkinson's Disease

Florian Krismer, PhD,^{1,2} Vincent Beliveau, PhD,^{1,2} Klaus Seppi, MD,^{1,2} Christoph Mueller, MD,^{1,2} Georg Goebel, PhD,³ Elke R. Gizewski, MD,^{2,4} Gregor K. Wenning, PhD,¹ Werner Poewe, MD,^{1,2} and Christoph Scherfler, MD^{1,2*}

¹Department of Neurology, Medical University of Innsbruck, Innsbruck, Austria ²Neuroimaging Research Core Facility, Medical University of Innsbruck, Innsbruck, Austria ³Department of Medical Statistics, Informatics and Health Economics, Medical University of Innsbruck, Innsbruck, Austria ⁴Department of Neuroradiology, Medical University of Innsbruck, Innsbruck, Austria

ABSTRACT: Background: Manual region-of-interest analysis of putaminal and middle cerebellar peduncle diffusivity distinguishes patients with multiple system atrophy (MSA) and Parkinson's disease (PD) with high diagnostic accuracy. However, a recent meta-analysis found substantial between-study heterogeneity of diagnostic accuracy due to the lack of harmonized imaging protocols and standardized analyses pipelines.

Objective: Evaluation of diagnostic accuracy of observer-independent analysis of microstructural integrity as measured by diffusion-tensor imaging in patients with MSA and PD.

Methods: A total of 29 patients with MSA and 19 patients with PD (matched for age, gender, and disease duration) with 3 years of follow-up were investigated with diffusion-tensor imaging and T1-weighted magnetic resonance imaging. Automated localization of relevant brain regions was obtained, and mean

diffusivity and fractional anisotropy values were averaged within the regions of interest. The classification was performed using a C5.0 hierarchical decision tree algorithm.

Results: Mean diffusivity of the middle cerebellar peduncle and cerebellar gray and white matter compartment as well as the putamen were significantly increased in patients with MSA and showed superior effect sizes compared to the volumetric analysis of these regions. A classifier model identified mean diffusivity of the middle cerebellar peduncle and putamen as the most predictive parameters. Cross-validation of the classification model yields a Cohen's κ and overall diagnostic accuracy of 0.823 and 0.914, respectively.

Conclusion: Analysis of microstructural integrity within the middle cerebellar peduncle and putamen yielded a superior effect size compared to the volumetric measures, resulting in excellent diagnostic accuracy to discriminate patients with MSA from PD in the early to moderate disease stages. © 2020 The Authors. *Movement Disorders* published by Wiley Periodicals LLC on behalf of International Parkinson and Movement Disorder Society

Key Words: multiple system atrophy; MRI; diffusion-weighted imaging; Freesurfer

Introduction

Multiple system atrophy (MSA) is an orphan and progressive neurodegenerative disorder featuring parkinsonian, cerebellar, and autonomic symptoms.¹ Symptoms of early-stage MSA frequently mimic Parkinson's disease (PD), hampering an early differential diagnosis.²⁻⁴ Accuracy rates of a clinical diagnosis of PD may range from 65% to 93%.^{2,5,6} However, a reliably early differential diagnosis is critical to patient counseling and for recruitment into interventional trials and other types of academic research.

Numerous magnetic resonance imaging (MRI) studies attempted to improve the diagnostic accuracy for the early differential diagnosis of parkinsonian disorders.⁷ Recently, advances in MRI postprocessing algorithms have provided an opportunity to localize and grade MSA-specific brain atrophy patterns.⁸⁻¹² However, prior to tissue loss becoming measurable as brain atrophy, the neurodegenerative cascade causes dysfunction at the cellular level, including membrane destruction. Alterations of tissue integrity can be quantified by diffusion-tensor MRI (DTI).^{13,14} DTI estimates the microstructural integrity and the degree of axonal pathology in brain tissue by quantifying the amount and direction of diffusing water molecules.^{13,15} DTI signal changes in the putamen and the middle cerebellar peduncle (MCP), assessed by manually labeled regions of interest (ROI), were shown to

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

*Correspondence to: Dr. Christoph Scherfler, Department of Neurology, Medical University Innsbruck, Anichstrasse 35, A-6020 Innsbruck, Austria; E-mail: christoph.scherfler@i-med.ac.at

Relevant conflicts of interest/financial disclosures: Nothing to report.

Funding agencies: This study was supported by grants from the Austrian Central Bank Anniversary Fund, project number: 14174, the Austrian Science Fund, project number: KLI82-B00 and the MSA Coalition.

Received: 24 June 2020; **Revised:** 14 August 2020; **Accepted:** 17 August 2020

Published online 16 September 2020 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mds.28281

differentiate the Parkinson variant of MSA from PD reliably.¹⁶⁻¹⁹ However, a meta-analysis reported substantial methodological differences, including variability of manual ROI placement, causing a considerable between-study heterogeneity.²⁰

In the present study, we sought to apply established automated subcortical volume segmentation to precisely delineate relevant brain regions and estimate averaged DTI parameters within these ROIs in a uniform fashion by observer-independent colocalization. We then evaluated the diagnostic accuracy of this multimodal approach for the differential diagnosis of MSA and PD.

Materials and Methods

Patients

Patients with MSA and PD from the MRI database of the Movement Disorders outpatient clinic at the Medical University of Innsbruck were identified based on the following inclusion criteria: (1) a clinical diagnosis of probable MSA (all patients showed some degree of parkinsonism) or PD at the last visit according to consensus operational criteria,^{21,22} (2) a clinical follow-up of ≥ 36 months, and (3) the availability of high-quality DTI registrations. Exclusion criteria were white-matter lesions grades 2 and 3, vascular or space-occupying lesions within the cerebrum, or motion artifacts on MRI. Presynaptic nigrostriatal dopaminergic dysfunction was confirmed by dopamine transporter single photon emission computed tomography in all patients with MSA. The present cohort represents a subset (those patients who have DTI registrations of sufficient quality) of a previously published cohort.⁹

The clinical evaluation included the Hoehn and Yahr staging scheme, the Unified Parkinson's Disease Rating Scale, and the Mini-Mental State Examination. All rating scales in patients with PD were performed in the *on* state.

Magnetic Resonance Sequences

MRI measurements were performed on a 3.0 Tesla magnetic resonance scanner (Magnetom Verio, Siemens, Erlangen, Germany) equipped with a 12-channel head coil. All participants underwent the same MRI protocol, including a coronal 3-dimensional T1-weighted magnetization prepared rapid gradient echo and a transversal diffusion-weighted echo-planar imaging with diffusion-sensitizing gradients in 20 directions with a b value of 1000 s/mm^2 and 1 reference image with $b = 0 \text{ s}/\text{mm}^2$.

Segmentation of the subcortical regions and estimation of structure volumes were performed using the FreeSurfer package 6.0 (<http://surfer.nmr.mgh.harvard.edu/>).²³⁻²⁶ Moreover, a published technique to integrate MCP segmentation into FreeSurfer was applied.⁹ The preprocessing steps were visually inspected to ensure that no misalignment of brain structures had occurred.

We then processed native DTI data and performed structural-to-diffusion data coregistration using FreeSurfer and FMRIB Software Library tools. This process included (1) eddy current and motion correction, (2) DTI general linear model fit and tensor construction, (3) registration of low b image to same-subject anatomical T1, and (4) fractional anisotropy (FA)/mean diffusivity (MD) mapping to Talairach space.

Data Analysis

Data analyses were performed using R 3.6.1 (R Foundation for Statistical Computing, Vienna, Austria). Gaussian distribution was confirmed by visual analysis of Q-Q plots and the Kolmogorov-Smirnov test. Group differences of normally distributed data were analyzed by parametric tests and non-Gaussian distributed variables by nonparametric tests. Distributional differences were determined by the Pearson chi-square test for independence. Bonferroni correction for multiple testing was applied where applicable. A C5.0 decision tree to classify patients with PD and MSA informed by MD, FA, and volume measurements from subcortical brain regions was developed (<http://www.rulequest.com/see5-info.html>). Leave-one-out cross-validation was employed to estimate the classification performance of the decision tree.

Results

Demographics

A total of 28 patients with MSA (19 Parkinson predominant, 9 cerebellar predominant) and 19 patients with PD as well as 25 age-matched and sex-matched healthy individuals were included in the present study. There was no significant difference between the study groups regarding gender distribution, age, and disease duration. Motor impairment was significantly greater in patients with MSA than in patients with PD. Detailed information on the participants' demographics is provided in Table 1.

Group Analysis of DTI Measures

Group differences of key brain regions and corresponding effect sizes are provided in Table 2. Significantly increased MD and significantly decreased FA values were observed in numerous subcortical brain regions of the MSA group compared with the PD group and healthy controls. There were no group differences between patients with PD and healthy controls. When comparing patients with MSA and PD, the effect sizes were largest within the MCP, the cerebellar cortex, the cerebellar white matter, and the putamen.

Classification Accuracy

Based on a C5.0 decision tree algorithm generated from DTI and volume measurements, increased MD values of the

TABLE 1. Demographics and basic clinical information

Variables	HC, N = 25	PD, N = 19	MSA, N = 28	Total, N = 72	P Value
Sex, n (%)					0.957 ^a
Female	11 (44.0)	8 (42.1)	13 (46.4)	32 (44.4)	
Male	14 (56.0)	11 (57.9)	15 (53.6)	40 (55.6)	
Age at MRI scan					0.048 ^b
Mean (SD)	60.0 (5.9)	64.9 (5.8)	63.4 (7.8)	62.6 (6.9)	
Disease duration at MRI					0.194 ^b
Mean (SD)	NA	3.1 (1.9)	2.4 (1.7)	2.7 (1.8)	
Predominant motor presentation, n (%)					
Parkinson predominant	NA	NA	19 (67.9)	NA	
Cerebellar predominant	NA	NA	9 (32.1)	NA	
Hoehn and Yahr staging					<0.001 ^c
Mean (SD)	NA	2.39 (0.54)	3.30 (0.69)	2.94 (0.77)	
Range	NA	1.50–3.00	2.00–4.00	1.50–4.00	
MMSE					0.124 ^b
Mean (SD)	NA	28.7 (1.5)	27.8 (2.2)	28.2 (2.0)	
UPDRS III					<0.001 ^b
Mean (SD)	NA	24.6 (6.5)	40.2 (12.9)	33.9 (13.2)	
UPDRS sum score					<0.001 ^b
Mean (SD)	NA	37.1 (9.9)	64.2 (19.1)	53.3 (20.8)	

^aPearson chi-squared test.

^bLinear model analysis of variance.

^cKruskal-Wallis rank-sum test.

Abbreviations: HC, healthy controls; PD, Parkinson’s disease; MSA, multiple system atrophy; MRI, magnetic resonance imaging; SD, standard deviation; MMSE, Mini-Mental State Examination; UPDRS III, Unified Parkinson’s Disease Rating Scale Part III.

TABLE 2. Mean diffusivity and fractional anisotropy values of relevant brain regions

DTI measure	HC, N = 25, Mean (SD)	PD, N = 19, Mean (SD)	MSA, N = 28, Mean (SD)	Total, N = 72, Mean (SD)	P Value	Effect Size
Mean diffusivity						
Cerebellum, white matter	0.719 (0.023)	0.715 (0.026)	0.862 (0.152)	0.773 (0.119)	<0.001	1.24
Cerebellar cortex	0.945 (0.048)	0.937 (0.040)	1.159 (0.192)	1.026 (0.163)	<0.001	1.47
Putamen	0.747 (0.035)	0.772 (0.034)	1.000 (0.222)	0.852 (0.184)	<0.001	1.32
Middle cerebellar peduncle	0.752 (0.033)	0.744 (0.020)	0.886 (0.124)	0.802 (0.104)	<0.001	1.47
Superior cerebellar peduncle	1.028 (0.179)	1.063 (0.183)	1.461 (0.460)	1.206 (0.377)	<0.001	0.68
Thalamus	0.866 (0.044)	0.892 (0.038)	0.914 (0.059)	0.892 (0.053)	0.003	0.43
Caudate nucleus	0.977 (0.103)	1.005 (0.144)	1.013 (0.153)	0.998 (0.134)	0.611	0.06
Pallidum	0.770 (0.109)	0.774 (0.096)	0.800 (0.106)	0.783 (0.104)	0.534	0.25
Midbrain	0.915 (0.058)	0.914 (0.061)	0.975 (0.079)	0.938 (0.073)	0.002	0.84
Pons	0.850 (0.064)	0.836 (0.050)	0.951 (0.136)	0.886 (0.109)	<0.001	1.05
Fractional anisotropy						
Cerebellum, white matter	0.396 (0.033)	0.389 (0.027)	0.327 (0.054)	0.367 (0.052)	<0.001	1.36
Cerebellar cortex	0.178 (0.014)	0.182 (0.016)	0.155 (0.028)	0.170 (0.024)	0.001	1.10
Putamen	0.248 (0.030)	0.248 (0.026)	0.260 (0.033)	0.253 (0.030)	0.319	–0.37
Middle cerebellar peduncle	0.477 (0.027)	0.483 (0.027)	0.424 (0.048)	0.458 (0.045)	<0.001	1.42
Superior cerebellar peduncle	0.380 (0.082)	0.407 (0.076)	0.390 (0.071)	0.391 (0.076)	0.509	0.04
Thalamus	0.323 (0.021)	0.322 (0.017)	0.317 (0.022)	0.320 (0.020)	0.541	0.26
Caudate nucleus	0.200 (0.028)	0.221 (0.034)	0.219 (0.039)	0.213 (0.035)	0.082	0.06
Pallidum	0.368 (0.063)	0.401 (0.048)	0.428 (0.053)	0.400 (0.061)	<0.001	0.53
Midbrain	0.443 (0.028)	0.461 (0.026)	0.438 (0.034)	0.446 (0.031)	0.032	0.75
Pons	0.514 (0.025)	0.523 (0.018)	0.473 (0.043)	0.500 (0.039)	<0.001	1.42

Group differences were calculated using linear model analysis of variance. Effect sizes are presented as Cohen’s d.

Abbreviations: HC, healthy controls; SD, standard deviation; PD, Parkinson’s disease; MSA, multiple system atrophy.

MCP and the putamen were the most discriminative metrics differentiating MSA from PD. Increased apparent diffusion coefficient (ADC) values (exploiting the corresponding cut-off values of the decision tree) in the MCP were present in

all patients with cerebellar-predominant MSA and in 68.4% of patients with Parkinson-predominant MSA, whereas putaminal ADC increases were found in 84.2% of patients with Parkinson-predominant MSA and 33.3% of

patients with cerebellar-predominant MSA. Cross-validation of the classification model yields a Cohen's κ of 0.823 and an overall diagnostic accuracy of 0.914. The final decision tree is presented in the Figure S1.

Discussion

In the present study, automated analysis of the MCP and putamen revealed significant differences of DTI metrics between patients with PD and MSA with larger effect sizes compared to volumetric measures of these structures. Notably, an attribute selection by a C5.0 decision tree, which was informed by volumetric MRI as well as DTI, identified increased MD values within the putamen and the MCP as the most predictive features for a diagnosis of MSA. The overall accuracy of image classification relative to the final clinical diagnosis based on cross-validation was 91.4%. This study extends previous research by demonstrating that automated analysis of DTI appears to be more sensitive to disease-specific tissue changes than volumetric measures and therefore more relevant for diagnostic purposes in the early stages of the disease. ■

Diagnostic Accuracy of Microstructural Changes

Detection of structural abnormalities within subcortical brain regions yields high diagnostic accuracy for a diagnosis of MSA. Automated segmentation of structural MRI with consecutive subcortical volume calculation and adjustment for total intracranial volume further increased the diagnostic accuracy for discriminating MSA and PD.^{8,9}

In addition, increased MD values were shown to be a useful measure of microstructural changes in neurodegenerative disorders.^{13,27,28} In patients with MSA, previous DTI studies reported increased MD and reduced FA values in brain regions that are affected by MSA pathology.²⁰ Interestingly, a case report of a patient with Parkinson-predominant MSA who had serial, multimodal MRIs during the course of 5 years showed that DTI changes preceded volume loss.²⁹ Furthermore, other MRI studies in patients with MSA supported this notion, suggesting that changes in diffusivity emerge prior to regional, disease-specific brain atrophy.²⁷⁻³⁰ Such findings were also reported in other neurodegenerative diseases, including Alzheimer's disease. A recent study in Alzheimer's disease mutation carriers suggested that regionally selective white matter degeneration as measured by DTI occurs years before the estimated onset of clinical symptoms.³¹ Another advantage of DTI over MRI volumetry is that DTI sequences do not require whole-brain volume correction, which facilitates data analysis. The delineation of subcortical structures on DTI sequences, however, requires expert knowledge and extensive training of the reader. Moreover, it is time consuming and prone to variability of metric parameters because of inconsistencies regarding the delineation of

boundaries and selection of adequate ROI size and shape. Such difficulties can be resolved by extending a well-established, automated MRI segmentation pipeline with DTI to T1-weighted image coregistration that transfers the anatomic information from high-resolution 3-dimensional T1 images to lower resolution DTI acquisitions.

In the present study, the MD of the MCP yielded the highest effect size in differentiating patients with PD and MSA. Accordingly, increased MCP diffusivity was selected as the most predictive measure in our diagnostic decision algorithm. This is not surprising given the results of previous research with manual ROI-based DTI measurements and voxel-wise group comparisons demonstrating that increased diffusivity in the MCP is common in MSA.^{11,17,28,32-36} The second most discriminative feature in the present diagnostic algorithm represented putaminal MD increases. This observation was also to be expected because previous studies that manually delineated the putamen on DTI images constantly showed increased diffusivity within the putamen in patients with MSA.^{17,19,20,28,34-37}

One of the particular strengths of the present study is the observer-independent ROI placement. Standardized, observer-independent placement of the ROIs could be helpful in harmonizing results and might provide better test-retest reliability.

Axial and radial diffusivity measures were considered less helpful for the present study because those are 2-dimensional measures within a single voxel and therefore inappropriate for averaged volume measurements.

Limitations

Some limitations need to be considered when interpreting the results of the present study. Despite the long follow-up period and the application of operational diagnostic criteria, the lack of postmortem verification remains a potential limitation because clinical misdiagnosis cannot be fully excluded. Although automated segmentation expedites the volume measurements of brain regions, visual inspection by an expert human reader is inevitable to ensure that no misalignment of brain structures had occurred. Moreover, automated volumetry and extraction of DTI data are labor-intensive techniques that require high-level technology and expertise that currently limits the availability of these imaging approaches to specialized centers. Thus, future efforts are required to move the innovative diagnostic procedure to general clinical practice.

Conclusions

Overall, the present study confirms the excellent diagnostic accuracy of DTI measures and demonstrates that observer-independent diffusion-imaging analysis is feasible and reliable in the differential diagnosis of MSA versus PD. Future studies are needed to validate diagnostic algorithms in independent data sets and describe the benefits of different approaches.

Acknowledgments: This study was supported by a grant of the MSA Coalition.

References

- Krismer F, Wenning GK. Multiple system atrophy: insights into a rare and debilitating movement disorder. *Nat Rev Neurol* 2017;13(4):232–243.
- Hughes AJ, Daniel SE, Ben-Shlomo Y, Lees AJ. The accuracy of diagnosis of parkinsonian syndromes in a specialist movement disorder service. *Brain* 2002;125(Pt. 4):861–870.
- Koga S, Aoki N, Uitti RJ, et al. When DLB, PD, and PSP masquerade as MSA: an autopsy study of 134 patients. *Neurology* 2015;85(5):404–412.
- Rizzo G, Copetti M, Arcuti S, Martino D, Fontana A, Logroscino G. Accuracy of clinical diagnosis of Parkinson disease: a systematic review and meta-analysis. *Neurology* 2016;86(6):566–576.
- Adler CH, Beach TG, Hentz JG, et al. Low clinical diagnostic accuracy of early vs advanced Parkinson disease: clinicopathologic study. *Neurology* 2014;83(5):406–412.
- Rajput AH, Rozdilsky B, Rajput A. Accuracy of clinical diagnosis in parkinsonism—a prospective study. *Can J Neurol Sci* 1991;18(3):275–278.
- Heim B, Krismer F, Seppi K. Structural imaging in atypical parkinsonism. *Int Rev Neurobiol* 2018;142:67–148.
- Scherfler C, Gobel G, Muller C, et al. Diagnostic potential of automated subcortical volume segmentation in atypical parkinsonism. *Neurology* 2016;86(13):1242–1249.
- Krismer F, Seppi K, Gobel G, et al. Morphometric MRI profiles of multiple system atrophy variants and implications for differential diagnosis. *Mov Disord* 2019;34(7):1041–1048.
- Huppertz HJ, Moller L, Sudmeyer M, et al. Differentiation of neurodegenerative parkinsonian syndromes by volumetric magnetic resonance imaging analysis and support vector machine classification. *Mov Disord* 2016;31(10):1506–1517.
- Peran P, Barbagallo G, Nemmi F, et al. MRI supervised and unsupervised classification of Parkinson's disease and multiple system atrophy. *Mov Disord* 2018;33(4):600–608.
- Moller L, Kassubek J, Sudmeyer M, et al. Manual MRI morphometry in Parkinsonian syndromes. *Mov Disord* 2017;32(5):778–782.
- Le Bihan D. Looking into the functional architecture of the brain with diffusion MRI. *Nat Rev Neurosci* 2003;4(6):469–480.
- Lehericy S, Vaillancourt DE, Seppi K, et al. The role of high-field magnetic resonance imaging in parkinsonian disorders: pushing the boundaries forward. *Mov Disord* 2017;32(4):510–525.
- Basser PJ, Pierpaoli C. Microstructural and physiological features of tissues elucidated by quantitative-diffusion-tensor MRI. *J Magn Reson B* 1996;111(3):209–219.
- Esterhammer R, Seppi K, Reiter E, et al. Potential of diffusion tensor imaging and Relaxometry for the detection of specific pathological alterations in Parkinson's disease (PD). *PLoS One* 2015;10(12):e0145493.
- Nicoletti G, Lodi R, Condino F, et al. Apparent diffusion coefficient measurements of the middle cerebellar peduncle differentiate the Parkinson variant of MSA from Parkinson's disease and progressive supranuclear palsy. *Brain* 2006;129(Pt. 10):2679–2687.
- Schocke MF, Seppi K, Esterhammer R, et al. Diffusion-weighted MRI differentiates the Parkinson variant of multiple system atrophy from PD. *Neurology* 2002;58(4):575–580.
- Schocke MF, Seppi K, Esterhammer R, et al. Trace of diffusion tensor differentiates the Parkinson variant of multiple system atrophy and Parkinson's disease. *Neuroimage* 2004;21(4):1443–1451.
- Bajaj S, Krismer F, Palma JA, et al. Diffusion-weighted MRI distinguishes Parkinson disease from the parkinsonian variant of multiple system atrophy: a systematic review and meta-analysis. *PLoS One* 2017;12(12):e0189897.
- Gilman S, Wenning GK, Low PA, et al. Second consensus statement on the diagnosis of multiple system atrophy. *Neurology* 2008;71(9):670–676.
- Fischl B, Salat DH, Busa E, et al. Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. *Neuron* 2002;33(3):341–355.
- Fischl B, van der Kouwe A, Destrieux C, et al. Automatically parcellating the human cerebral cortex. *Cereb Cortex* 2004;14(1):11–22.
- Segonne F, Dale AM, Busa E, et al. A hybrid approach to the skull stripping problem in MRI. *Neuroimage* 2004;22(3):1060–1075.
- Fischl B, Dale AM. Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proc Natl Acad Sci U S A* 2000;97(20):11050–11055.
- Ito M, Watanabe H, Kawai Y, et al. Usefulness of combined fractional anisotropy and apparent diffusion coefficient values for detection of involvement in multiple system atrophy. *J Neurol Neurosurg Psychiatry* 2007;78(7):722–728.
- Tha KK, Terae S, Yabe I, et al. Microstructural white matter abnormalities of multiple system atrophy: in vivo topographic illustration by using diffusion-tensor MR imaging. *Radiology* 2010;255(2):563–569.
- Park J, Joo JJ, Ahn BJ, Kwon KY. Serial MRI findings in a case of the Parkinson variant of multiple system atrophy: clinical usefulness of diffusion-weighted imaging at B1000 in early stages of the disease. *J Neurol Sci* 2016;362:136–138.
- Seppi K, Schocke MF. An update on conventional and advanced magnetic resonance imaging techniques in the differential diagnosis of neurodegenerative parkinsonism. *Curr Opin Neurol* 2005;18(4):370–375.
- Araque Caballero MA, Suarez-Calvet M, Duering M, et al. White matter diffusion alterations precede symptom onset in autosomal dominant Alzheimer's disease. *Brain* 2018;141(10):3065–3080.
- Barbagallo G, Sierra-Pena M, Nemmi F, et al. Multimodal MRI assessment of nigro-striatal pathway in multiple system atrophy and Parkinson disease. *Mov Disord* 2016;31(3):325–334.
- Nemmi F, Pavy-Le Traon A, Phillips OR, et al. A totally data-driven whole-brain multimodal pipeline for the discrimination of Parkinson's disease, multiple system atrophy and healthy control. *Neuroimage Clin* 2019;23:101858.
- Paviour DC, Thornton JS, Lees AJ, Jager HR. Diffusion-weighted magnetic resonance imaging differentiates Parkinsonian variant of multiple-system atrophy from progressive supranuclear palsy. *Mov Disord* 2007;22(1):68–74.
- Pellecchia MT, Barone P, Mollica C, et al. Diffusion-weighted imaging in multiple system atrophy: a comparison between clinical subtypes. *Mov Disord* 2009;24(5):689–696.
- Prodoehl J, Li H, Planetta PJ, et al. Diffusion tensor imaging of Parkinson's disease, atypical parkinsonism, and essential tremor. *Mov Disord* 2013;28(13):1816–1822.
- Seppi K, Schocke MF, Donnemiller E, et al. Comparison of diffusion-weighted imaging and [123I]IBZM-SPECT for the differentiation of patients with the Parkinson variant of multiple system atrophy from those with Parkinson's disease. *Mov Disord* 2004;19(12):1438–1445.
- Postuma RB, Berg D, Stern M, et al. Clinical diagnostic criteria for Parkinson's disease. *Mov Disord* 2015;30(12):1591–1601. <http://dx.doi.org/10.1002/mds.26424>

Supporting Data

Additional Supporting Information may be found in the online version of this article at the publisher's web-site.