



# Offering a putative neurobiological “dopamine homeostatic” solution to overcome the perils of the reward deficiency syndrome pandemic: emergence of “precision behavioral management”

Kenneth Blum<sup>1,2,3,4</sup>, Catherine A. Dennen<sup>5</sup>, David Baron<sup>1</sup>, Panayotis K. Thanos<sup>6</sup>, Rajendra D. Badgaiyan<sup>4</sup>

<sup>1</sup>Division Of Addiction Research & Education, Center for Sports, Exercise and Global Mental Health, Western University Health Sciences, Pomona, CA, USA; <sup>2</sup>Department of Psychiatry, Wright University Boonshoft School of Medicine, Dayton, OH, USA; <sup>3</sup>The Kenneth Blum Institute of Behavior & Neurogenetics, Austin, TX, USA; <sup>4</sup>Department of Psychiatry, South Texas Veteran Health Care System, Audie L. Murphy Memorial VA Hospital, Long School of Medicine, University of Texas Medical Center, San Antonio, TX, USA; <sup>5</sup>Department of Family Medicine, Jefferson Health Northeast, Philadelphia, PA, USA; <sup>6</sup>Department of Psychology, University of Buffalo, the State University of New York, Buffalo, NY, USA

Correspondence to: Kenneth Blum, PhD. Division Of Addiction Research & Education, Center for Sports, Exercise and Global Mental Health, Western University Health Sciences, Pomona, CA, USA. Email: drd2gene@gmail.com.

Submitted Sep 29, 2022. Accepted for publication Oct 19, 2022.

doi: 10.21037/atm-2022-67

View this article at: <https://dx.doi.org/10.21037/atm-2022-67>

In addition to the COVID-19 pandemic, an addiction crisis is currently sweeping the entire world (1). This is extremely troubling, especially given that illicit drug usage can increase the risks associated with the coronavirus. Specifically, the psychological and social risks of the pandemic, which can also promote and augment drug abuse, in a possibly toxic cycle. Our latest figures have revealed that approximately 26.8 million people globally suffered from opioid use disorder (OUD) in 2016, with more than 100,000 opioid overdose deaths yearly, including more than 100,000 in the United States (US) in 2021 (2). While drug overdose is most prevalent in the US, it is a global issue requiring “out of the box” thinking to prevent further devastation and death (3).

Currently, the standard treatment of OUD and opioid addiction involves the use of powerful opioids, which seems inane and traps people in an unwanted cycle of addiction (4,5). This standard treatment is known as opioid agonist therapy (OAT), which functions by interacting with opioid receptors to reduce cravings and harm. However, OAT does not address the root cause, and this form of therapy may actually worsen the addiction pandemic worldwide. Therefore, a more prudent and effective treatment is needed. One solution, as espoused by Lee *et al.* (4), includes targeting various portions of the dopamine-dependent addiction pathway, identifying vulnerable genes, and

modifying gene products such as, for example, precision Pro-dopamine Regulation (KKB220) or other gentle dopamine agonists (6).

Our group seeks to lessen harm while addressing the core underlying issue (7). Therefore, we are hereby suggesting that a hypodopaminergic state (epigenetic) and/or trait (genetic) is critical in terms of continued motivation to use/abuse licit and illicit substances and can lead to relapse. Although there are FDA-approved medications that can be utilized to treat substance addictions (e.g., alcohol, opiates, nicotine, etc.), these agents typically only offer a short-term advantage by blocking dopamine. Thus, it has been argued that instead of the utilization of long-term administration of these FDA-approved drugs, the goal should be to induce “dopamine homeostasis”, or in simpler terms, “normalcy”. Miller *et al.* (7), suggested that this could be accomplished through a variety of holistic modalities including, but not limited to, dopamine-boosting diets, exercise, yoga, meditation, hyper-oxygenation, heavy metal detoxification, and, most crucially, nutraceuticals like KB220 variants, which can help balance brain neurotransmitters.

Another strategy is to use a narcotic antagonist, such as naltrexone, to induce “psychological extinction” by blocking D2 receptors for not only opioids but methamphetamines as well (8). The latter approach appears to be more palatable, but compliance is a major problem. In addition, there is

poor compliance even with the extended-release injectable form of naltrexone. Another option is acamprosate, which is an NMDA receptor antagonist as well as a positive allosteric modulator of GABAA receptors that disrupts dopaminergic signaling (9). Our laboratory showed the novelty and profound enhancement of naltrexone compliance in patients who rapidly detoxed from long-term methadone utilization (10). In an open-label investigation in humans, Blum *et al.* (10) found that dopamine augmentation with the pro-dopamine regulator KB220 (262 days) improved naltrexone compliance and results when compared to naltrexone monotherapy (37 days). Compared to standard treatments, this well-studied complex consists of amino-acid neurotransmitter precursors, catabolism inhibition, and enkephalinase inhibitor therapy (11).

Understanding the above premise and the further emerging acceptance of the umbrella term reward deficiency syndrome (RDS), which was established by Blum in 1995, promotes the common mechanism hypothesis for drug and non-addictive behaviors and even anhedonia, an important endophenotype linked to relapse (12). Previously, we postulated that anhedonia is a by-product of hypodopaminergic traits and states resulting from the combination of hereditary traits and epigenetic neurobiological alterations as a result of environmental factors. Dopaminergic activity can cause a variety of pathophysiological conditions, including aberrant learning, incentive sensitization, and stress-like “anti-reward” phenomena (13). It seems prudent to suggest that neurogenetic antecedents to RDS and elucidation of reward gene polymorphisms could provide a framework for determining an individual’s genetic risk for developing anhedonia as a subset of RDS. Thus, our prevention strategy encompasses the restoration of homeostatic balance through physiological activation of dopaminergic receptors (D2/D3) and might have heuristic value for targeting not only anhedonia but also drug craving and relapse.

Our simple proposal to help restore brain neurotransmitter balance in the afflicted individual with a possible pharmacogenomic personalized approach involves the coupling of a genetic based addiction risk assessment, for example, the Genetic Addiction Risk Severity (GARS) test, and customized KB220 (14). Understanding the common neuromodulating aspects of neurotransmission and its disruption via chronic exposure to drugs and behavioral addictions requires a known approach involving “dopamine homeostasis”. While there is an emerging push for the utilization of “psychedelic medicine” (15) in the short-


term at low doses via patch delivery systems, we further propose that long-term treatment requires induction of “dopamine homeostasis” (16). However, along these lines of thinking, Bill Wilson’s psychedelic experience, which led to his becoming alcohol-free (but not nicotine-free, which eventually killed him) and the founding of Alcoholics Anonymous, appears to be consistent with the current re-excitement of psychedelic medicine. A 2012 meta-analysis of lysergic acid diethylamide (LSD) therapy was found to be at least as efficacious a treatment as anything we currently have today (17). The work of Mash and associates has paved the way to implicate the idea of psychedelics like Ibogaine to treat addiction (18).

### Snapshot review of evidence

Our “out of the box” novel approach requires the coupling of genetic risk polymorphic testing with the safe and well-studied complex KB220 or KB220Z customized to match the presence of resultant alleles and, as such, provides a precision nutraceutical known to have pro-dopamine regulatory pharmacological properties, including H-Wave therapy (19). There is a plethora of high-tier publications providing unequivocal evidence to establish a common overlap neuro-mechanism between substance and non-substance addiction (alcohol, opioids, food, etc.) (20). In addition, there are currently over 55 clinical studies involving this amino-acid based enkephalinase inhibitory Pro-Dopamine Regulator (PDR) with KB220 nutraceutical complex as the primary constituent ingredient (11,21). The basis of this complex is that it mimics the brain reward cascade (see *Figure 1*), an established model of reward processing.

KB220 has also been shown to have BOLD activation across the brain reward circuit, which has been observed in both animal and human heroin-dependent individuals (23,24). These include the nucleus accumbens, anterior thalamic nuclei, anterior cingulate gyrus, hippocampus, infralimbic, and prelimbic loci. The evidence for genetic vulnerability as an important antecedent to this unwanted behavior may be a determinant factor which must be identified early in life.

Based on a rather moderate published literature, the role of reward gene polymorphisms puts individuals at a higher risk for developing all RDS behavior subtypes, including anhedonia. With this in mind, our laboratory has created the novel patented GARS test to help identify one’s risk for these addictive-like behaviors. Specifically, published studies illustrate the coupling of GARS with KB220Z variants



**Figure 1** The brain reward cascade. The interaction of at least seven major neurotransmitter-pathways involved in the BRC. In the hypothalamus environmental stimulation results in the release of serotonin, which in turn via, for example, 5HT-2a receptors activate (green equal sign) the subsequent release of opioid peptides from opioid peptide neurons, also in the hypothalamus. Then, in turn, the opioid peptides have two distinct effects, possibly via two different opioid receptors. One that inhibits (red hash sign) through the mu-opioid receptor (possibly via enkephalin) and projects to the Substantia nigra to GABA<sub>A</sub> neurons. Another stimulates (green equal sign) cannabinoid neurons (the Anandamide and 2-archydonoglycerol, for example) through Beta-Endorphin linked delta receptors, which in turn inhibit GABA<sub>A</sub> neurons at the Substantia nigra. Also, when activated, cannabinoids primarily 2-archydonoglycerol, can indirectly disinhibit (red hash sign) GABA<sub>A</sub> neurons through activation of G1/0 coupled to CB1 receptors in the Substantia nigra. In the DRN, glutamate neurons can then indirectly disinhibit GABA<sub>A</sub> neurons in the Substantia nigra through activation of GLU M<sub>3</sub> receptors (red hash sign). GABA<sub>A</sub> neurons, when stimulated, will, in turn, powerfully (red hash signs) inhibit VTA glutaminergic drive via GABA<sub>B</sub> 3 neurons. It is also possible that stimulation of ACH neurons at the Nucleus Accumbens ACH can stimulate both muscarinic (red hash) or Nicotinic (green hash). Finally, glutamate neurons in the VTA will project to dopamine neurons through NMDA receptors (green equal sign) to preferentially release dopamine at the NAc shown as a bullseye indicates a euphoria, or “wanting” response. The result is that when dopamine release is low (endorphin deficiency), unhappiness is felt while general (healthy) happiness depends on the dopamine homeostatic tonic set point (22) (used with permission). BRC, brain reward cascade; DRN, dorsal raphe nuclei; NAc, nucleus accumbens.

utilizing a semi-customized precision matched to one’s GARS. This novel modality linked to a systems biological approach provides an increased efficacy as well as clinical outcomes in terms of treating RDS (25,26).

We present herein new evidence related to the utilization of precision behavioral management (PBM) to effect clinical outcomes of (I) alcohol use disorder (AUD) (27); (II) identification of high addiction genetic risk in pain clinics (26); (III) bariatric surgery outcomes associated with GARS (28); (IV) conversion of incarceration to rehabilitation in Drug Court (29). There is even growing evidence that epigenetic insult, for example, methylation and histone modification of the *DRD2* gene, has been shown in people with lifetime gambling use disorder (30). This argues for the need to induce dopamine homeostasis.

## Conclusions

It is generally accepted that balancing the brain reward circuit or achievement of “dopamine homeostasis” is a laudable goal instead of inhibiting natural dopamine or prescribing a potent opioid to treat opioid addiction (31). We are encouraging both the neuroscience and clinical science communities to potentially embrace this disruptive technology with a futuristic view of addressing the notion of what constitutes “standard of care” in the face of the ongoing addiction (alcohol, opioid, psychostimulant, food, etc.) pandemic (32).

While additional research is needed, it is pertinent to begin establishing guidelines that incorporate the knowledge of RDS as an umbrella term for all addictive behaviors. Comprehending the neurogenetics and using a

systems biological approach (PBM), as previously stated, appears to be the most prudent and marks a breakthrough in restoring joy to the billions suffering globally, particularly in terms of early detection (33).

### Acknowledgments

*Funding:* The study was supported by R41 MD012318/MD/NIMHD NIH HHS/United States (to KB) and 1101 CX002099-01 Merit review grant, Veterans Administration (to RDB).

### Footnote

*Provenance and Peer Review:* This article was commissioned by the editorial office, *Annals of Translational Medicine*. The article did not undergo external peer review.

*Conflicts of Interest:* All authors have completed the ICMJE uniform disclosure form (available at <https://atm.amegroups.com/article/view/10.21037/atm-2022-67/coif>). KB serves as an unpaid editorial board member of *Annals of Translational Medicine* from September 2022 to August 2024. KB receives grants from NIH and reports consulting fees from Electronic Waveform Lab and United Scientific Group, leadership role in Society of Brain Mapping & Therapeutics and Royal Society of Medicine. He is the member of 19 editorial boards across the world. He is the inventor of GARS and KB220 and holds both domestic and foreign patents issued and pending. His company Synaptamine Inc. licensed Victory Nutrition International (VNI) to market resultant products. PKT receives consulting fees from AMCA. RDB reports grants from Veterans Administration. The other authors have no conflicts of interest to declare.

*Ethical Statement:* The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links

to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

### References

- Ornell F, Moura HF, Scherer JN, et al. The COVID-19 pandemic and its impact on substance use: Implications for prevention and treatment. *Psychiatry Res.* 2020 Jul;289:113096.
- Strang J, Volkow ND, Degenhardt L, et al. Opioid use disorder. *Nat Rev Dis Primers* 2020;6:3.
- Darke S, Farrell M. Would legalizing illicit opioids reduce overdose fatalities? Implications from a natural experiment. *Addiction* 2014;109:1237-42.
- Lee YK, Gold MS, Fuehrlein BS. Looking beyond the opioid receptor: A desperate need for new treatments for opioid use disorder. *J Neurol Sci* 2022;432:120094.
- Blum K, Baron D. Opioid Substitution Therapy: Achieving Harm Reduction While Searching for a Prophylactic Solution. *Curr Pharm Biotechnol* 2019;20:180-2.
- Blum K, Kazmi S, Modestino EJ, et al. A Novel Precision Approach to Overcome the "Addiction Pandemic" by Incorporating Genetic Addiction Risk Severity (GARS) and Dopamine Homeostasis Restoration. *J Pers Med* 2021;11:212.
- Miller D, Miller M, Blum K, et al. Addiction Treatment in America: After Money or Aftercare? *J Reward Defic Syndr* 2015;1:87-94.
- Wang ZY, Guo LK, Han X, et al. Naltrexone attenuates methamphetamine-induced behavioral sensitization and conditioned place preference in mice. *Behav Brain Res* 2021;399:112971.
- Olive MF, Nannini MA, Ou CJ, et al. Effects of acute acamprosate and homotaurine on ethanol intake and ethanol-stimulated mesolimbic dopamine release. *Eur J Pharmacol* 2002;437:55-61.
- Blum K, Lott L, Baron D, et al. Improving naltrexone compliance and outcomes with putative pro- dopamine regulator KB220, compared to treatment as usual. *J Syst Integr Neurosci* 2020. doi:10.15761/JSIN.1000229.
- Kenneth B, Edward MJ, Marjorie GLC, et al. Pro-Dopamine Regulator (KB220) A Fifty Year Sojourn to Combat Reward Deficiency Syndrome (RDS): Evidence Based Bibliography (Annotated). *CPQ Neurol Psychol* 2018;1:<https://www>.



12. Gold MS, Blum K, Febo M, et al. Molecular role of dopamine in anhedonia linked to reward deficiency syndrome (RDS) and anti-reward systems. *Front Biosci (Schol Ed)* 2018;10:309-25.
13. Borsook D, Linnman C, Faria V, et al. Reward deficiency and anti-reward in pain chronification. *Neurosci Biobehav Rev* 2016;68:282-97.
14. Blum K, Madigan MA, Fried L, et al. Coupling Genetic Addiction Risk Score (GARS) and Pro Dopamine Regulation (KB220) to Combat Substance Use Disorder (SUD). *Glob J Addict Rehabil Med* 2017;1:555556.
15. Nutt D. Psychedelic drugs-a new era in psychiatry? *Dialogues Clin Neurosci* 2019;21:139-47.
16. Blum K, Febo M, Fried L, et al. Hypothesizing That Neuropharmacological and Neuroimaging Studies of Glutamatergic-Dopaminergic Optimization Complex (KB220Z) Are Associated With "Dopamine Homeostasis" in Reward Deficiency Syndrome (RDS). *Subst Use Misuse* 2017;52:535-47.
17. Krebs TS, Johansen PØ. Lysergic acid diethylamide (LSD) for alcoholism: meta-analysis of randomized controlled trials. *J Psychopharmacol* 2012;26:994-1002.
18. Mash DC, Duque L, Page B, et al. Ibogaine Detoxification Transitions Opioid and Cocaine Abusers Between Dependence and Abstinence: Clinical Observations and Treatment Outcomes. *Front Pharmacol* 2018;9:529.
19. Gupta A, Bowirrat A, Gomez LL, et al. Hypothesizing in the Face of the Opioid Crisis Coupling Genetic Addiction Risk Severity (GARS) Testing with Electrotherapeutic Nonopioid Modalities Such as H-Wave Could Attenuate Both Pain and Hedonic Addictive Behaviors. *Int J Environ Res Public Health* 2022;19:552.
20. Kotyuk E, Magi A, Eisinger A, et al. Co-occurrences of substance use and other potentially addictive behaviors: Epidemiological results from the Psychological and Genetic Factors of the Addictive Behaviors (PGA) Study. *J Behav Addict* 2020;9:272-88.
21. Blum K, Trachtenberg MC, Elliott CE, et al. Enkephalinase inhibition and precursor amino acid loading improves inpatient treatment of alcohol and polydrug abusers: double-blind placebo-controlled study of the nutritional adjunct SAAVE. *Alcohol* 1988;5:481-93.
22. Gold MS, Baron D, Bowirrat A, Blum K. Neurological correlates of brain reward circuitry linked to opioid use disorder (OUD): Do homo sapiens acquire or have a reward deficiency syndrome? *J Neurol Sci* 2020;418:117137.
23. Febo M, Blum K, Badgaiyan RD, et al. Enhanced functional connectivity and volume between cognitive and reward centers of naïve rodent brain produced by pro-dopaminergic agent KB220Z. *PLoS One* 2017;12:e0174774.
24. Blum K, Liu Y, Wang W, et al. rsfMRI effects of KB220Z<sup>TM</sup> on neural pathways in reward circuitry of abstinent genotyped heroin addicts. *Postgrad Med* 2015;127:232-41.
25. Blum K, Bowirrat A, Baron D, et al. Biotechnical development of genetic addiction risk score (GARS) and selective evidence for inclusion of polymorphic allelic risk in substance use disorder (SUD). *J Syst Integr Neurosci* 2020. doi: 10.15761/JSIN.1000221.
26. Moran M, Blum K, Ponce JV, et al. High Genetic Addiction Risk Score (GARS) in Chronically Prescribed Severe Chronic Opioid Proband Attending Multi-pain Clinics: an Open Clinical Pilot Trial. *Mol Neurobiol* 2021;58:3335-46.
27. Blum K, Steinberg B, Gondre-Lewis MC, et al. A Review of DNA Risk Alleles to Determine Epigenetic Repair of mRNA Expression to Prove Therapeutic Effectiveness in Reward Deficiency Syndrome (RDS): Embracing "Precision Behavioral Management". *Psychol Res Behav Manag* 2021;14:2115-34.
28. Blum K, Bailey J, Gonzalez AM, et al. Neuro-Genetics of Reward Deficiency Syndrome (RDS) as the Root Cause of "Addiction Transfer": A New Phenomenon Common after Bariatric Surgery. *J Genet Syndr Gene Ther* 2011;2012:S2-001.
29. Blum k, Green R, Mullen P et al. Determinism v. Free Will & Genetic Evidence of Addiction in Plea Bargaining and Sentence Mitigation: Conversion of Incarceration to Probation and Rehabilitation Based on Genetic Addiction Risk Severity (GARS) Test. *St Marys Law J* 2022. (in Press).
30. Hillemecher T, Frieling H, Buchholz V, et al. Alterations in DNA-methylation of the dopamine-receptor 2 gene are associated with abstinence and health care utilization in individuals with a lifetime history of pathologic gambling. *Prog Neuropsychopharmacol Biol Psychiatry* 2015;63:30-4.
31. Blum K, Chen AL, Chen TJ, et al. Activation instead of blocking mesolimbic dopaminergic reward circuitry is a preferred modality in the long term treatment of reward deficiency syndrome (RDS): a commentary. *Theor Biol Med Model* 2008;5:24.
32. Fischer B, O'Keefe-Markman C, Lee AM, et al. 'Resurgent', 'twin' or 'silent' epidemic? A select data

overview and observations on increasing psycho-stimulant use and harms in North America. *Subst Abuse Treat Prev Policy* 2021;16:17.

33. Gold MS, Blum K, Oscar-Berman M, et al. Low dopamine

function in attention deficit/hyperactivity disorder: should genotyping signify early diagnosis in children? *Postgrad Med* 2014;126:153-77.

**Cite this article as:** Blum K, Dennen CA, Baron D, Thanos PK, Badgaiyan RD. Offering a putative neurobiological “dopamine homeostatic” solution to overcome the perils of the reward deficiency syndrome pandemic: emergence of “precision behavioral management”. *Ann Transl Med* 2022;10(23):1291. doi: 10.21037/atm-2022-67