

Identifying promoters for gene expression in *Clostridium thermocellum*



Daniel G. Olson^{a,d,*}, Marybeth Maloney^{a,d}, Anthony A. Lanahan^{a,d}, Shuen Hon^{a,d},
Loren J. Hauser^{b,d}, Lee R. Lynd^{a,b,c,d,*}

^a Thayer School of Engineering at Dartmouth College, Hanover, NH 03755, United States

^b Oak Ridge National Laboratory, Oak Ridge, TN 37830, United States

^c Mascoma Corporation, NH 37830, United States

^d BioEnergy Science Center, TN, United States

ARTICLE INFO

Article history:

Received 31 October 2014

Received in revised form

26 February 2015

Accepted 21 March 2015

Available online 30 March 2015

Keywords:

Promoter

Copy number

Structural instability

Rolling circle replication

lacZ

adhB

ABSTRACT

A key tool for metabolic engineering is the ability to express heterologous genes. One obstacle to gene expression in non-model organisms, and especially in relatively uncharacterized bacteria, is the lack of well-characterized promoters. Here we test 17 promoter regions for their ability to drive expression of the reporter genes β -galactosidase (*lacZ*) and NADPH-alcohol dehydrogenase (*adhB*) in *Clostridium thermocellum*, an important bacterium for the production of cellulosic biofuels. Only three promoters have been commonly used for gene expression in *C. thermocellum*, *gapDH*, *cbp* and *eno*. Of the new promoters tested, 2638, 2926, 966 and 815 showed reliable expression. The 2638 promoter showed relatively higher activity when driving *adhB* (compared to *lacZ*), and the 815 promoter showed relatively higher activity when driving *lacZ* (compared to *adhB*).

© 2015 The Authors. Published by Elsevier B.V. International Metabolic Engineering Society. This is an open access article under the CC BY-NC-SA license (<http://creativecommons.org/licenses/by-nc-sa/4.0/>).

1. Introduction

Clostridium thermocellum is a promising candidate for the conversion of biomass to ethanol due to its native ability to use cellulose and produce ethanol (Olson et al., 2012). Recently there has been significant progress engineering *C. thermocellum* to improve ethanol production (Argyros et al., 2011; Deng et al., 2013; Van der Veen et al., 2013), however ethanol yield needs further improvement for commercial viability.

To date, most metabolic engineering of *C. thermocellum* has focused on gene deletion but many metabolic engineering strategies require increased or heterologous gene expression, in addition to gene deletion. Only a few genes have been successfully expressed in *C. thermocellum*, including the antibiotic resistance markers *cat* and *neo*; the counter-selectable markers *pyrF*, *hpt* and *tdk* (Olson and Lynd, 2012a); the cellulosome scaffoldin *cipA* (Olson et al., 2013) and the metabolic enzyme pyruvate kinase (Deng et al., 2013).

The most commonly used promoters are from the regions upstream of the *gapDH* (Clo1313_2095), *cbp* (Clo1313_1954) and

eno (Clo1313_2090) genes in *C. thermocellum*. The *gapDH* gene encodes the glyceraldehyde-3-phosphate dehydrogenase enzyme, which is one of the most highly expressed proteins in the *C. thermocellum* proteome (Olson et al., 2013, 2010; Rydzak et al., 2012). While none of these promoters have been extensively characterized, the *cbp* promoter has been reliably used to drive expression of the *pyrF* (Tripathi et al., 2010) and *tdk* (Argyros et al., 2011) selectable markers. The *eno* promoter has been used to drive expression of an exogenous pyruvate kinase from *Thermoanaerobacterium saccharolyticum* (Deng et al., 2013). The transcription start sites have not been determined nor is anything known about which sigma factor is responsible for promoter recognition.

In the absence of detailed analysis, it is still useful to search for suitable promoters for expression by scanning those available. It is important to appreciate that any such study based on the assay of a reporter gene activity will also depend on mRNA stability and enzyme efficiency. An ideal promoter for metabolic engineering would have the following characteristics:

1. Low expression of the gene of interest in cloning strains of *Escherichia coli*.
2. Consistently high expression in *C. thermocellum*, independent of genetic context.
3. Low homology to the chromosome (in the case of native promoters, shorter is better).

* Corresponding authors.

E-mail addresses: dan268@gmail.com (D.G. Olson),
Lee.R.Lynd@dartmouth.edu (L.R. Lynd).

The goal of this work is to identify new promoters for gene expression in *C. thermocellum*, and compare them with ones currently in use.

2. Materials and methods

2.1. Plasmid construction

Plasmids were constructed using standard molecular biology techniques (Sambrook and Russell, 2001) and isothermal DNA assembly (Gibson, 2011). Plasmid sequence was confirmed by Sanger sequencing of the promoter region and reporter gene. Plasmids were transformed into *C. thermocellum* DSM 1313 using standard techniques (Olson and Lynd, 2012a). Plasmids were also transformed in to *E. coli* C2566 (New England Biolabs) using standard techniques (Sambrook and Russell, 2001). The thermostable *lacZ* gene was a gift from James Liao (Lin et al., 2014). A list of the promoters and reporter genes in each plasmid is given in Table 1.

2.2. Choice of promoter regions

Promoter regions were chosen based on comparison of several sets of published gene expression data. These sets include Riederer et al. (2011), Raman et al. (2011), Gowen and Fong (2010) and Van der Veen et al. (2013). Note that for the last reference, strain characterization is described in the referred paper, but the transcription data is available from the GEO database, accession GSE27046. Promoters were selected on the basis of showing moderate to high expression of the genes they control across several of the data sets. For a given gene, the region upstream of the translation start site of the gene was selected. In general this region was 100–500 bp in length (Table 2). For promoters that are listed twice (i.e. *cbp* and *cbp_2*), the first version of the promoter caused problems with cloning, so a slightly different region was selected.

Table 1
Plasmids used in this study.

Plasmid	Promoter	Putative SigA/RpoD motif	Predicted conventional RBS Sequence	Predicted TIE strength (arbitrary units) ^a	Reporter gene
pDGO-66	gapDH	TTGA(A)A-N17-TA(A)AAT	AGGAGG	–	none
pDGO80	none			369	<i>adhB</i>
pDGO81	0544		GGAGG	5190	<i>adhB</i>
pDGO83	<i>cbp</i>	TTGA(A)(T)-N17-TATAAT	AGGAGG	244,912	<i>adhB</i>
pDGO84	<i>eno</i>	TTGA(A)A-N18-(C)AT(T)AT	GGAG	8763	<i>adhB</i>
pDGO86	1194	TTG(T)(T)(T)-N15-TATAAT	AGG(G)GG	19,141	<i>adhB</i>
pDGO87	0966	TTG(C)(A)(T)-N15-TGNTATAAT	AGGA	5587	<i>adhB</i>
pDGO88	2638	TT(A)A(A)A-N15-TATAAT	AGGAGG	42,645	<i>adhB</i>
pDGO89	0815	TT(T)A(A)A-N12-TGNTAT(T)AT	GAGG	19,843	<i>adhB</i>
pDGO90	2926		AGGAGG	15,731	<i>adhB</i>
pDGO92	0307		GGAG	2764	<i>adhB</i>
pDGO95	gapDH	TTGA(A)A-N17-TA(A)AAT	AGGAGG	7372	<i>lacZ</i>
pDGO98	none			–	<i>lacZ</i>
pDGO99	0544		GGAGG	716	<i>lacZ</i>
pDGO100	gapDH_2	TTGA(A)A-N17-TA(A)AAT	AGGAGG	1154	<i>lacZ</i>
pDGO102	<i>eno</i>	TTGA(A)A-N18-(C)AT(T)AT	GGAG	32,320	<i>lacZ</i>
pDGO104	1194	TTG(T)(T)(T)-N15-TATAAT	AGG(G)GG	18,381	<i>lacZ</i>
pDGO105	0966	TTG(C)(A)(T)-N15-TGNTATAAT	AGGA	1734	<i>lacZ</i>
pDGO106	2638	TT(A)A(A)A-N15-TATAAT	AGGAGG	6441	<i>lacZ</i>
pDGO107	0815	TT(T)A(A)A-N12-TGNTAT(T)AT	GAGG	2091	<i>lacZ</i>
pDGO108	2926	TGNTA(A)(T)AT	AGGAGG	7656	<i>lacZ</i>
pDGO109	2463		AGGAGG	8067	<i>lacZ</i>
pDGO110	0307		GGAG	858	<i>lacZ</i>
pDGO111	<i>lac</i>	TT(T)ACA-N18-TAT(G)(T)T	AGGA	390	<i>lacZ</i>
pDGO112	<i>lacUV5</i>	TT(T)ACA-N18-TATAAT	AGGA	390	<i>lacZ</i>
pDGO117	<i>cbp_2</i>	TTGA(A)(T)-N17-TATAAT	AGGAGG	50,011	<i>lacZ</i>
pDGO118	3011_2	TTGAC(T)-N17-TATAAT	AGGAGG	5978	<i>lacZ</i>

^a See Section 2.10 for details about calculation of translation initiation efficiency (TIE).

2.3. Making cell-free extracts

To prepare cell-free extracts (CFE), *C. thermocellum* cells were grown to mid-log phase. 10 ml of cells was centrifuged and the pellet was resuspended in B-PER buffer (Thermo Scientific, IL). The cells were lysed by addition of Lysonase according to the manufacturer's directions (EMD Millipore, MA). After centrifugation for 1 min at 15,000g to remove cell debris, the resulting supernatant was used for protein and enzyme assays. *E. coli* cells were also lysed using B-PER buffer and Lysonase enzyme. Protein concentration of the CFE was measured using the Coomassie Plus Bradford Assay (Thermo Scientific, IL) using bovine serum albumin as the standard. Absorbance was measured at 595 nm and 450 nm and the ratio of the two was used to determine concentration.

2.4. LacZ activity assay

The activity of the LacZ enzyme was determined by measuring the formation of the yellow cleavage product of o-nitrophenyl-β-D-galactoside (ONPG) as described by Miller (1972) with modifications as described below. The 0.2 ml assay solution contained 100 mM sodium phosphate pH 7.5, 1 mM magnesium chloride, 50 mM beta mercaptoethanol, 0.655 mg/ml ONPG and varying amounts of CFE. The assay solution was incubated at 37 °C for 4 h and the rate of increase in absorbance at 420 nm was measured in a 96-well plate with a BioRad Powerwave XS spectrophotometer. Purified LacZ enzyme (G4155 from Sigma) was used as a standard. Several 2-fold dilutions of CFE were added, and activity was determined in the linear range.

2.5. AdhB activity assay

AdhB is a bifunctional secondary alcohol and aldehyde dehydrogenase from *Thermoanaerobacter pseudethanolicus* ATCC 33223 (Teth39_0218) that is NADPH-dependent (Burdette et al., 1997). The activity of the AdhB enzyme was determined by measuring the disappearance of NADPH by measuring changes in absorbance

Table 2

Description of promoter sequences. Predicted SigA/RpoD binding sites are highlighted: “-35 boxes” in green, “-10 boxes” in yellow and “extended-10 boxes” in pink. Predicted ribosome binding sites are highlighted in blue. Start codons at the end of each promoter are indicated by bold font.

Name	Native gene locus	Length (bp)	Description of native gene	Sequence (5'→3')
cbp	Clo1313_1954	430	cellobiose metabolism	TCTATTGCATTGAAATGATTAGTTATCCGTAATAATTAATCATATCATAAATTAATATCATAAATGTTT TGACGAATGAAGTTTTGGATAAATATCAAGTAAAGGAACGCTAAAAATTTGGCGTAAAAATCAAAA TGACCAC TTGAAAT TAATATGGTAAAGTAGAT ATAAAT ATTTTGGTAAACATGCCTCAGCAAGGTTAGTATAGC TGTTCCGTATAAATTAACCGTATGGTAAACGGCAGTCAGAAAAAATAGTCATAAGATTCCGTTATGAAAAAT ATACTCGGTAGTTAATAAAGAGATAGAGGTAAAGAGATCAAGATAAGAGATATAAGGTACGAATGTA TAAGATGGTGTCTTTTAGGCACATAAATAAAAAACAAATAAACGAAAAATTTT AGGAGG ACGAAAGATG
cbp_2	Clo1313_1954	621	cellobiose metabolism	GAGTCGTGACTAAGAAGCTCAAAGTAATTAACAATACAGATTTTTCTCATGCTTTTACCCCTTCATAAAT TTAATTTTATCGTTATCATAAAAAATATAGACGTTTATATTGCTTCCGGGATATAGTCTGGCATTGCTGG TGCAAAATGTTCCGGATGAAGTGGAATGATTGATTCGATGTTGATCTATTGCTGAAATGATTAGTATCCGT AAATATTAATTAATCATATAAATTAATATATAAATGTTTTGACGAATGAAGGTTTTGGATAAAT TCAAGTAAAGGAACGCTAAAAATTTGGCGTAAAAATCAAAATGACCAC TTGAAAT TAATATGGTAAAGTAG ATAAAT ATTTTGGTAAACATGCCTCAGCAAGGTTAGATTAGCTGTTCCGTATAAATTAACCGTATGGTAA AACGGCAGTCAGAAAAAATAGTCATAAGATTCCGTTATGAAAAATACTCTGGTAGTATAAATAAGAGATA TGAGGTAAGAGATCAAGATAAGAGATATAAGGTACGAATGTAAGATGGTGTCTTTTAGGCACATAAAT AAAAACAATAAACGAAAAATTT AGGAGG ACGAAAGATG
eno	Clo1313_2090	178	glycolysis	GGAAATATTAATGAAATG TTGAA AAATGTTTAAAGATGGG CATTAT GGATAAATACTACTAGTTT GCAATAAATGCTTCTAATAATGGACTTTGGTGAATATGGTGAAGGATGCAAGTGTAAATTTTAAACAT AAAAATAAGCTATATGAAG GGAG AATGGAGAATG
gapDH	Clo1313_2095	309	glycolysis	TAATTAAGTATCTCTGGCA TTGCG GGTTTTAATAAAGAT AAAAAT TATTGACTAGAAATAAAAAATGTT CCATAAATAAGTGGCAAAAAACAAGAAATTAACATCAAAAGGAAGTAAAAATCTTTGTTAAAAAATTA TTATTTTTTACTAACTA TTGAA ATGAAAAATAAATA ATAAAT GAATCATATGCAAGAGATACCTCC CAGAGGATGAATATTTACTGCTTATGCTTTATGGCAGCTAATAGAGGCTTAAATTAATTTAATTTACA AT AGGAGG CGATATAATG
gapDH_2	Clo1313_2095	263	glycolysis	AATTAATGACTAGAAATAAAAAATTTGCCATAAATTAATGGCAAAAAACAAGAAATACATCAAGGA AGATAAAAAATCTTTGTTAAAAAATTAATTTTTTAACTAAACT TTGAA ATGAAAAATAAATA ATAA ATGAAT CATAGTGAAGAGATACCTGCCAGAGGATGAATTTTACTGCATTCATGCTTTGGCAGCTAATA GAGGCATTAATTAATTTAATTTACAAT AGGAGG CGATATAATG
lac	(from <i>E. coli</i>)	190	(exogenous)	TGAAATCTTCTCATCCGCCAAAACAGCCAAGCGCAACGCAATTAATGTAGTAGTACTACTATTAGCA CCCCAGC TTTAA CTTTATGCTCCGGCTCG TATGTT GTGTGGAATTGTGAGCGGATAACAATTTACACAG GAACAGCTATG
lacUV5	(from <i>E. coli</i>)	190	(exogenous)	TGAAATCTTCTCATCCGCCAAAACAGCCAAGCGCAACGCAATTAATGTAGTAGTACTACTATTAGCA CCCCAGC TTTAA CTTTATGCTCCGGCTCG TATaa GTGTGGAATTGTGAGCGGATAACAATTTACACAG GAACAGCTATG
0307	Clo1313_0307	88	preprotein translocase	TAAAAAAGAGGGAAATAATCGGCAGCTGGGGGAAGACTTCTGATGTGCTGACAGAAAAAATTTAAA AAT GGAG ATGTGTCGAT ATG
0544	Clo1313_0544	70	VanW-family protein	GTTCATTTGGACAAGTTTAAATATATATAAATGTTAAAGCGCTTACGTTGAAAG TGGAGG GAATTCATTAT G
0815	Clo1313_0815	154	short chain dehydrogenase	ATAATATATACCGTATTTTACAATATCCATACGAATATGGATTATTTATTTGTTATATTTTAAACAAAAA TTA TTAA CTTTTCAATG GGTATTAT ATATATGGTCACAGTTCAATGAATCAAAAAATAA AGG GT TTAATATG
0966	Clo1313_0966	226	homoserine kinase or phosphoglycerate mutase	TTCTACTACTCCCTTGCAATCAGACAAAAATTTTCTCTTAACTTTCTGTAACCGTATCCGACTACTTTG TCATAAAGTAATATAGATTATTTATACCGTCAATCAAAATAAATTTTATATCGCAATCT TTGCA CCGC TTTAAAGCTCTG TTGATAAAT TATATGTTAAATATCATAAAAAATAAATAAAGTTTAA AGGAA CTGATT TTATG
1194	Clo1313_1194	300	extracellular solute binding protein	TTAATGCCGACCAAGTTGCAATCCCGTCAAATAATGATTTTTCAGCCGACGAAACAGCAAGATACT GTATGGCTATAAATGTTTTCAGGCAAGGATATATTTGCTCCGGTAAAAATTAACAATAAGCTAAAAAC TGACGTAGGATAAGCAAAACGGCGCAATTTGAGTTGTAACGTAATTTTCACTAAAAATGATTAATTTTCA TG TTGTT TTTTTATGATTAAT TATAAT ATAATTTATGATAAGCAATATCTTAATATCATTA AGGGG AA AAAAACTATG
2463	Clo1313_2463	100	guanido phosphotransferase	AAGCGGCAATATTAGAGATCAATCAGAATCTTGAAGTCAACAAAAACAATATGTAAGTGCACATTTT GGTTTTGCGAGG AGGAGG GTATGTGATG
2638	Clo1313_2638	209	peroxiredoxin	GATAACAAAGGACGGTTCAGGGCTTCTGCTACTCTACTCTGATTGTAAGAAAGTGAAGTGAATTTTAT TTAATCTTATGAAAAAATTTTGAATTCGGTTT TTAAAG AAAAAGTGGGTATAT TATAAT AGTCAAT GATTGGTTAAAAAATTTAATAAGCAACAGAAATAAACAAAAAGTA AGGAGG AATTTG TTATG
2926	Clo1313_2926	127	ATP-dependent metalloprotease	AAAAATACAAGGTTCTTGTTTTAATACCGTTA GT TAATATAATGTAATATATATTTATAATAAATATG TATGAGAGATAGTGTGTTGCTATATGCTATAAAGAAT AGGAGG GAATAGATG
3011	Clo1313_3011	387	S-layer protein	TATTGCTTAATCGGTAGGAAAAGCCGGTAAAAACCGCTTTTAAATGCACATCAAAATCATTCCAGAA AAATCAACTATATA TTGTA TTCCAGCAATCTGTAG TATAAT AGAAATCGTTGAAACTGATTTTACT AGTCATTCAACAAAAACAGTTGTTTCAATCTGTTACAATTTCTTACACTAAATCAAAAT TTGAC CACT TATAGCCCAATGG TATAAT AAGCTTGAATCAAGATAGTTGTGTGCGCAGCTTGAAGAAAAAGCAAGAT AGGTACGGTTAAAAACAAGGGATAAGGCGTAGCCTGCTTAAATTTCAAGCTGCTTCCAAAAAACA ATACA AGGAGG ATTTGAGAGTATG
3011_2	Clo1313_3011	211	S-layer protein	TTCTGTACAATTTCTTACACTAAATCAAATTT TTGAC CACTTATAGCCCAATGG TATAAT AAAGCTGAA TCAAGATAGTTGTGTGCGCAGCTTGAAGAAAAAGCAAGATAGGTCAAGTTAAAAAAACAAGGGATAA GGCGTAGCTGGCTTAATTTCAAGCTGCTTCCAAAAAACAATACA AGGAGG ATTTGAGAGTATG

at 340 nm. The 0.2 ml assay solution contained 100 mM Tris–HCl pH 7.6, 0.4 mM NADPH, 4 mM acetaldehyde and varying amounts of CFE. The assay solution was incubated at 55 °C for 1 h and the rate of decrease in absorbance at 340 nm was measured. Measurements were performed in a BioRad Powerwave XS spectrophotometer. The path length was 0.52 cm. Several 2-fold dilutions of CFE were added, and activity was determined in the linear range.

2.6. Plasmid copy number

Plasmid copy number was determined by quantitative PCR (qPCR). Copies of the plasmid genes chloramphenicol (*cat*) and B-galactosidase (*lacZ*) were normalized to levels of the *celS* gene (Clo1313_2747) levels, which is present at single copy on the *C. thermocellum* genome (Olson et al., 2010). Primer sequences are

given in Supporting information Table S4. Reactions were performed in triplicate using 2 μ l of bacterial cultures as template and SsoFast EvaGreen Supermix (BioRad) following the manufacturer's recommendation and an annealing temperature of 55 °C. Standard curves for *cat* and *celS* were generated using the linearized plasmid pDGO-28 (Olson and Lynd, 2012b), which contains a single copy of both genes. Standard curves for *lacZ* were generated using linearized plasmid pDGO98, which contains a single copy of the *lacZ* gene. All data was normalized using a 529 bp synthetic gBlock (IDT) containing the *cat*, *celS* and *lacZ* amplicons.

2.7. RNA isolation

1 ml of bacterial culture was pelleted and lysed by digestion with lysozyme (15 mg/ml) and proteinase K (20 mg/ml). RNA was isolated with an RNAeasy minikit (Qiagen) and digested with TURBO DNase (Applied Biosystems) to remove contaminating DNA. cDNA was synthesized from 500 ng of RNA using the iScript cDNA synthesis kit (BioRad).

2.8. Gene expression

qPCR was performed in triplicate using cDNA with SsoFast EvaGreen Supermix (BioRad) at an annealing temperature of 55 °C to determine RNA levels of the gene of interest, which were normalized to *recA* RNA levels. In order to confirm removal of contaminating DNA from RNA samples, for a subset of samples, a control cDNA synthesis reaction was performed in the absence of reverse transcriptase enzyme. Standard curves were generated using a 1904 bp synthetic gBlock (IDT) containing all of the amplicons.

2.9. PCR for plasmid structural stability analysis

To determine plasmid stability primers were designed that generated an approximately 6000 bp PCR product (the exact amplicon size depended upon the promoter inserted) spanning the *repB*, *cat*, *amp* and *lacZ* genes of plasmids pDGO95 through pDGO118. The PCR product was generated using 1 μ l of bacterial culture, previously transformed with the plasmid, and Quick Load 2 \times Taq (NEB). Primers were annealed at 60 °C (forward: 5'-AAGGTGCGTT-GAAGTGTGGTATGT-3', reverse: 5'-GGCTTCTGTACGCTGACCCTA-TTA-3'). PCR products were analyzed by electrophoresis on 1% agarose gels.

2.10. Predicted translation initiation efficiency (TIE) analysis

The translation initiation efficiency was predicted based on a biophysical model of translation initiation that considers the hybridization of the 16s rRNA to the 50 bp region around the translation start site. For this model, the 16s sequence from *C. thermocellum* was used (Salis et al., 2009). The calculations were performed using a freely available calculator on the Salis lab website, <https://salis.psu.edu/software/reverse>. The predicted relative rate of translation initiation is expressed in arbitrary units.

2.11. Ribosome binding site prediction and SigA promoter prediction

Conventional ribosome binding sites (RBS) were identified using the Prodigal Gene Prediction algorithm (Hyatt et al., 2010) since that is an integral part of the algorithm and part of the output. Predicted SigA binding sites were identified manually and are based on the known *Bacillus subtilis* SigA and *E. coli* RpoD (<http://dbtbs.hgc.jp/>) consensus sequence TTGACA-N16-19-TATAAT.

3. Results and discussion

Sequences of DNA upstream from genes in *C. thermocellum* were screened for promoter sequences. First these sequences were cloned 5' of a thermostable beta galactosidase gene (*lacZ*). For sequences that gave promising results with *lacZ*, we built a second set of constructs using the NADPH-linked secondary alcohol dehydrogenase from *Thermoanaerobacterium ethanolicus* (*adhB*). In total, 27 plasmid constructs were made (Table 1), using 17 promoter regions (Table 2).

3.1. *E. coli* vs. *C. thermocellum* activity

For our purposes we were interested in finding promoters with low activity in *E. coli* and high activity in *C. thermocellum*, so we measured LacZ activity in cell-free extracts of both organisms (Fig. 1, Supporting information Table S1).

It can be seen that there is a low correlation between promoters that give good activity in *C. thermocellum* vs. *E. coli*. The promoters 2638, *cbp_2*, *gapDH* and 3011_2 have the largest differential. It is interesting to note the difference between *gapDH* and *gapDH_2*, since these promoters only differ by 46 bp. To understand the cause of the substantial variability observed, we performed a second round of experiments.

Colonies from the first round were inoculated into media with different amounts of thiamphenicol (6, 24 or 96 μ g/ml). For each culture, we measured 6 parameters including: DNA copy number of the *lacZ* reporter gene, DNA copy number of the *cat* antibiotic resistance gene, mRNA expression of the *lacZ* reporter gene, mRNA expression of the native gene (i.e. the gene whose promoter was used in the construct) and specific activity of the LacZ protein (Fig. 2, Supporting information Table S2).

To understand the effect of antibiotic concentration on plasmid copy number, we grew plasmid-containing strains of *C. thermocellum* on various concentrations of thiamphenicol. Although we did see a slight correlation between antibiotic strength and the copy number of the *cat* gene (Supporting information Table S2), this did not result in higher LacZ activity. In general, the highest activity was found when cells were grown in the presence of 6 μ g/ml thiamphenicol (Fig. 2). Promoters 307, 544, 2463, *lac* and *lacUV5* did not show activity above the negative control.




Fig. 1. Comparison of LacZ activity between *E. coli* and *C. thermocellum* in units of activity per mg of cell-free extract (CFE). Error bars represent one standard deviation, $n \geq 3$.




Fig. 2. Comparison of LacZ activity for different promoters and different thiamphenicol concentrations. The thick line represents the median, the box encloses the 25th–75th percentile of data. Individual data points are plotted.




Fig. 3. Comparison of mRNA expression between *lacZ* and native gene. Thick bar represents the median. The box encompasses the 25th–75th percentile of data. Individual data points are plotted.

After these experiments had been performed, these promoters were analyzed for the presence of SigA binding sites (Table 2, green boxes). For sequences 307, 544, 2463 and 2926, these sites were not found, suggesting either that the sequence does not contain a promoter or that the sequence contains a promoter that is recognized by a sigma factor other than SigA. In addition, the four genes downstream of these sequences appear to be internal genes in operons. The 2926 sequence is interesting since it contains three potential -10 boxes (yellow), no -35 boxes (green), and a weak extended -10 box (magenta–TGNTA(A)(T)AT), is expressed well in *E. coli* but, not at all in *C. thermocellum*. We speculate that it might contain a binding site for an *E. coli* positive regulator that overcomes the lack of a good -35 box. Furthermore, this is the only sequence of the four that showed LacZ activity above the negative control.

The *lac* and *lacUV5* promoters are derived from *E. coli* and require the cAMP receptor protein (CRP). Apparently *C. thermocellum* does not have a CRP that interacts with the *lac* or *lacUV5* promoters.

Constructs with sequences 1194 and 3011_2 were difficult to transform into *C. thermocellum*, transformation efficiency was lower than expected. For 3011_2, despite numerous attempts, we only had

one successful transformation and that transformation yielded only two colonies. We speculate that these promoters may have contributed to plasmid instability for the inherently unstable plasmid.

3.2. Expression of *lacZ* vs. native gene

To test whether *lacZ* expression was similar to expression of the native gene, we measured both by RT-PCR (Fig. 3). Since the *lacZ* gene is present on a plasmid that can be present in multiple copies, *lacZ* gene expression was normalized to *lacZ* DNA copy number. In general, *lacZ* expression was higher than that of the native gene. Since these are translational fusions, we speculate that there may be differences in mRNA stability or the activity of the gene product.

To investigate whether the results obtained with *lacZ* are applicable to another reporter gene, we measured the expression of *adhB* driven by some of these promoters (Fig. 4, Supporting information Table S3). In general, there was a good correlation between constructs driving the *lacZ* or *adhB* reporter genes. Two notable exceptions are promoters 2638 and 815. For promoter 2638, the AdhB activity is about 8-fold higher than what would be




Fig. 4. Comparison of activity levels for two different reporter genes with the same promoter. Error bars represent one standard deviation, $n \geq 3$. For the *cbp* promoter, the LacZ activity was measured with the *cbp_2* promoter and the AdhB activity was measured with the *cbp* promoter. The plasmid with *cbp* driving *lacZ* could not be cloned in *E. coli*, and the plasmid with *cbp_2* driving *adhB* could not be transformed into *C. thermocellum*.

expected based on the LacZ activity. For promoter 815, the AdhB activity is about 16-fold lower than what would be expected based on the LacZ activity.

We do not have a convincing explanation for these differences, however we speculate that translation initiation may play a role. For a given promoter, the RBS sequence upstream of *lacZ* or *adhB* is identical. However differences in mRNA secondary structure near the RBS due to the different sequences of the *lacZ* and *adhB* genes can affect translation initiation (Salis et al., 2009) and mRNA stability. For each promoter and reporter gene pair, the translation initiation efficiency was predicted (Table 2). In the case of promoter 2638, the predicted RBS strength is calculated to be about 7-fold higher for the *adhB* construct than the *lacZ* construct. This explanation does not, however, explain the lower *adhB* expression with promoter 815, as the predicted RBS strength of the *adhB* construct was also higher than the *lacZ* construct (10-fold).

Another possible explanation for the differences in gene expression between *lacZ* and *adhB* with promoters 0815 and 2638 is that these promoters are affected by redox levels. Since *adhB* encodes an alcohol dehydrogenase that could affect redox levels in the cell, that would create a feedback loop and explain the differences between the reporter genes.

3.3. Plasmid instability

Although not the primary focus of this work, we observed several indications that plasmid instability is a serious problem. One indication is the variability of the DNA copy numbers of the *lacZ* and *cat* genes. Since they are both present in a single copy on the plasmid, it was expected that their copy number would remain in a 1:1 ratio. In fact, we found an excess of the *cat* gene relative to the *lacZ* gene in about a quarter of the colonies we analyzed (Supporting information Table S2). Another indication of structural instability was results seen in PCR across the region containing the *repB*, *cat*, *amp* and *lacZ* genes. Although we would have expected a ~6 kb amplicon, in about 40% of the colonies we analyzed, we found either no amplicon or an amplicon that was much shorter or longer. Plasmid instability is commonly observed in rolling-circle-replicating plasmids (Bron et al., 1991), and since the *cat* gene is upstream of the *lacZ* gene, premature termination of plasmid replication could result in a truncated

plasmid with the *cat* gene able to provide thiamphenicol resistance but lacking the *lacZ* gene.

Another interesting observation was that some promoter sequences caused problems with recovery of frozen cultures. For all of the colonies tested, about a third could not be recovered after freezing. However the problems with recovery did not seem to be distributed randomly. Rather, the sequences that resulted in high *lacZ* activity also tended to have more problems with strain recovery (data not shown). The worst results were observed with the *gapDH* promoters, where 7 of 8 colonies initially frozen could not be recovered. Interestingly, this problem only occurred when both the *gapDH* promoter and the *lacZ* reporter gene were present. In control constructs missing either the *gapDH* promoter sequence (pDGO98) or the *lacZ* reporter gene (pDGO-66), this effect was not observed, and all frozen colonies (4 of 4 in both cases) were recovered. One possible explanation is that the presence of these sequences resulted in homologous recombination and integration of this unstable plasmid at the *gapDH* locus. This integration could be toxic to cells because it disrupts expression of the native *gapDH* gene which is essential for glycolysis.

4. Conclusions

Promoters 815, 966, 2638, 2926, *eno*, *cbp* and *cbp_2* all show activity and seem to be more generally useful than the *gapDH* and *gapDH_2* promoters. Several of these promoter regions are less than 250 bp in length, which suggests that this length of promoter region is sufficient.

An important direction for future work will be understanding and eliminating the causes of plasmid structural instability observed here.

Acknowledgments

We thank Professor Jan Westpheling for critical review of the manuscript and useful discussions. The BioEnergy Science Center is a U.S. Department of Energy Bioenergy Research Center supported by the Office of Biological and Environmental Research in the DOE Office of Science subcontract number 4000115284. The DOE Office of Science did not play any role in study design, writing the report or the decision to submit the manuscript for publication.

Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.meteno.2015.03.002>.

References

- Argyros, D.A., Tripathi, S.A., Barrett, T.F., Rogers, S.R., Feinberg, L.F., Olson, D.G., Foden, J.M., Miller, B.B., Lynd, L.R., Hogsett, D.A., Caiazza, N.C., 2011. High ethanol titers from cellulose by using metabolically engineered thermophilic, anaerobic microbes. *Appl. Environ. Microbiol.* 77, 8288–8294. <http://dx.doi.org/10.1128/AEM.00646-11>.
- Bron, S., Meijer, W., Holsappel, S., Haima, P., 1991. Plasmid instability and molecular cloning in *Bacillus subtilis*. *Res. Microbiol.* 142, 875–883.
- Burdette, D.S., Secundo, F., Phillips, R.S., Dong, J., Scott, R. a, Zeikus, J.G., 1997. Biophysical and mutagenic analysis of Thermoanaerobacter ethanolicus secondary-alcohol dehydrogenase activity and specificity. *Biochem. J.* 326 (Pt 3), 717–724.
- Deng, Y., Olson, D.G., Zhou, J., Herring, C.D., Joe Shaw, A., Lynd, L.R., 2013. Redirecting carbon flux through exogenous pyruvate kinase to achieve high ethanol yields in *Clostridium thermocellum*. *Metab. Eng.* 15, 151–158. <http://dx.doi.org/10.1016/j.ymben.2012.11.006>.

- Gibson, D.G., 2011. Chapter fifteen - Enzymatic Assembly of Overlapping DNA Fragments. In: Voigt, C.A. (Ed.), *Methods in Enzymology*. Academic Press, Waltham, MA, pp. 349–361.
- Gowen, C.M., Fong, S.S., 2010. Genome-scale metabolic model integrated with RNAseq data to identify metabolic states of *Clostridium thermocellum*. *Biotechnol. J.* 5, 759–767. <http://dx.doi.org/10.1002/biot.201000084>.
- Hyatt, D., Chen, G.L., Locascio, P.F., Land, M.L., Larimer, F.W., Hauser, L.J., 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinform.* 11, 119. <http://dx.doi.org/10.1186/1471-2105-11-119>.
- Lin, P.P., Rabe, K.S., Takasumi, J.L., Kadisch, M., Arnold, F.H., Liao, J.C., 2014. Isobutanol production at elevated temperatures in thermophilic *Geobacillus thermoglucosidarius*. *Metab. Eng.* 24, 1–8. <http://dx.doi.org/10.1016/j.ymben.2014.03.006>.
- Miller, J.H., 1972. *Experiments in Molecular Genetics*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Olson, D.G., Giannone, R.J., Hettich, R.L., Lynd, L.R., 2013. Role of the CipA scaffoldin protein in cellulose solubilization, as determined by targeted gene deletion and complementation in *Clostridium thermocellum*. *J. Bacteriol.* 195, 733–739. <http://dx.doi.org/10.1128/jb.02014-12>.
- Olson, D.G., Lynd, L.R., 2012a. Chapter seventeen – transformation of *Clostridium thermocellum* by electroporation. In: Gilbert, H.J. (Ed.), *Methods in Enzymology*. Academic Press, Waltham, MA, pp. 317–330. <http://dx.doi.org/10.1016/b978-0-12-415931-0.00017-3>.
- Olson, D.G., Lynd, L.R., 2012b. Computational design and characterization of a temperature sensitive plasmid replicon for gram positive thermophiles. *J. Biol. Eng.* 6, 5. <http://dx.doi.org/10.1186/1754-1611-6-5>.
- Olson, D.G., McBride, J.E., Shaw, A.J., Lynd, L.R., 2012. Recent progress in consolidated bioprocessing. *Curr. Opin. Biotechnol.* 23, 396–405. <http://dx.doi.org/10.1016/j.copbio.2011.11.026>.
- Olson, D.G., Tripathi, S.A., Giannone, R.J., Lo, J., Caiazza, N.C., Hogsett, D.A., Hettich, R.L., Guss, A.M., Dubrovsky, G., Lynd, L.R., 2010. Deletion of the Cel48S cellulase from *Clostridium thermocellum*. *Proc. Natl. Acad. Sci. U. S. A.* 107, 17727–17732. <http://dx.doi.org/10.1073/pnas.1003584107>.
- Raman, B., McKeown Jr., C.K., Brown, M.R., Mielenz, S.D., Rodriguez, J.R., Rodriguez Jr., M., 2011. Transcriptomic analysis of *Clostridium thermocellum* ATCC 27405 cellulose fermentation. *BMC Microbiol.* 11, 134. <http://dx.doi.org/10.1186/1471-2180-11-134>.
- Riederer, A., Takasuka, T.E., Makino, S., Stevenson, D.M., Bukhman, Y. V., Elsen, N.L., Fox, B.G., 2011. Global gene expression patterns in *Clostridium thermocellum* as determined by microarray analysis of chemostat cultures on cellulose or cellobiose. *Appl. Environ. Microbiol.* 77, 1243–1253. <http://dx.doi.org/10.1128/aem.02008-10>.
- Rydzak, T., McQueen, P.D., Krokhin, O. V., Spicer, V., Ezzati, P., Dwivedi, R.C., Shamsurin, D., Levin, D.B., Wilkins, J.A., Sparling, R., 2012. Proteomic analysis of *Clostridium thermocellum* core metabolism: relative protein expression profiles and growth phase-dependent changes in protein expression. *BMC Microbiol.* 12, 214. <http://dx.doi.org/10.1186/1471-2180-12-214>.
- Salis, H.M., Mirsky, E.A., Voigt, C.A., 2009. Automated design of synthetic ribosome binding sites to control protein expression. *Nat. Biotechnol.* 27, 946–950.
- Sambrook, J., Russell, D.W., 2001. *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Tripathi, S.A., Olson, D.G., Argyros, D.A., Miller, B.B., Barrett, T.F., Murphy, D.M., McCool, J.D., Warner, A.K., Rajgarhia, V.B., Lynd, L.R., Hogsett, D.A., Caiazza, N.C., 2010. Development of pyrF-based genetic system for targeted gene deletion in *Clostridium thermocellum* and creation of a pta mutant. *Appl. Environ. Microbiol.* 76, 6591–6599. <http://dx.doi.org/10.1128/AEM.01484-10>.
- Van der Veen, D., Lo, J., Brown, S.D., Johnson, C.M., Tschaplinski, T.J., Martin, M., Engle, N.L., van den Berg, R.A., Argyros, A.D., Caiazza, N.C., Guss, A.M., Lynd, L.R., 2013. Characterization of *Clostridium thermocellum* strains with disrupted fermentation end-product pathways. *J. Ind. Microbiol. Biotechnol.* 40, 725–734. <http://dx.doi.org/10.1007/s10295-013-1275-5>.