

Superantigen reactive $V\beta 6^+$ T cells induce perforin/granzyme B mediated caspase-independent apoptosis in tumour cells

S Mürköster^{1,4}, MA Weigand^{2,3}, C Choi¹, H Walczak², V Schirmacher^{*,1} and V Umansky¹

¹Division of Cellular Immunology, G0100, Tumorimmunology Program, German Cancer Research Center, Im Neuenheimer Feld 280, 69120 Heidelberg, Germany; ²Division of Apoptosis Regulation, Tumorimmunology Program, German Cancer Research Center, Im Neuenheimer Feld 280, 69120 Heidelberg, Germany; ³Clinic of Anesthesiology, University of Heidelberg, Im Neuenheimer Feld 110, 69120 Heidelberg, Germany

The endogenous viral superantigen 7 in DBA/2 mice serves as a target antigen on syngeneic ESb-MP lymphoma cells for allogeneic graft-vs-leukaemia reactive cells. Allogeneic viral superantigen 7 reactive $V\beta 6^+$ T cells are able to transfer graft-vs-leukaemia reactivity and to kill specifically viral superantigen 7⁺ ESb-MP tumour cells *in vitro*. Here we elucidate the mechanism of this superantigen specific cell lysis. Already 10 min after co-incubation with *in vitro* stimulated $V\beta 6^+$ T cells, viral superantigen 7⁺ ESb-MP tumour cells show an apoptotic phenotype (Annexin V-positivity, DNA-fragmentation). This extremely rapid type of cell death is not mediated by the death inducing ligands CD95L, TRAIL and TNF but by perforin and granzyme B. Surprisingly, neither mitochondria nor any of the known caspases appear to be involved in this type of tumour cell killing. In contrast, nitric oxide, released by activated macrophages and endothelial cells, induces in the same tumour cells another type of apoptosis which is much slower and involves mitochondria and caspase activation. A synergistic effect between the two different effector mechanisms of superantigen reactive donor cytotoxic T lymphocytes and nitric oxide releasing host macrophages and endothelial cells might explain the effective immune rejection of even advanced metastasised cancer in this graft-vs-leukaemia animal model.

British Journal of Cancer (2002) 86, 828–836. DOI: 10.1038/sj/bjc/6600104 www.bjcancer.com

© 2002 Cancer Research UK

Keywords: endogenous superantigen; apoptosis; graft-vs-leukaemia; caspase-independent; perforin

In patients with chronic myeloid leukaemia, the adoptive transfer of allogeneic bone marrow and additionally the use of allogeneic donor lymphocytes have been shown to result in long-term survival of up to 80% of the patients (Slavin *et al*, 1998). This favourable graft-vs-leukaemia (GvL) effect is often associated with a risk for the development of graft-vs-host (GvH) disease with significant morbidity and mortality (Mackinnon *et al*, 1995; Slavin *et al*, 1998). We have established an animal model for the investigation of GvL and GvH reactivity of tumour-immune lymphocytes (Schirmacher *et al*, 1995). *In situ* activated tumour-reactive lymphocytes from the tumour-resistant mouse strain B10.D2 are transferred into 5 Gy irradiated late-stage ESb-MP tumour-bearing DBA/2 mice. This leads to complete remission of the primary tumour and to the eradication of metastases. The two strains of mice are identical at the MHC but differ in minor histocompatibility and in Mls antigens (Schirmacher *et al*, 1995, 1998). The latter represent endogenous viral superantigens (vSAGs) which are encoded by mouse mammary tumour virus proviruses (Mtv) that are integrated in murine genomes (Beutner *et al*, 1992). In contrast to conventional antigens, SAGs are presented by MHC II molecules outside the binding groove and are recognised by certain $V\beta$ -chains

of the TCR (Choi *et al*, 1996). Since these endogenous SAGs behave like self-antigens, the reactive T cells are eliminated from the T cell repertoire by intra-thymic deletion (Speiser *et al*, 1989). One of the best studied vSAGs is vSAG7 which is encoded by *Mtv7* and is expressed in DBA/2 mice and on syngeneic ESb-MP tumour cells but not in B10.D2 mice (Schirmacher *et al*, 2000). vSAG7 is recognised by $V\beta 6^+$ T cells which are present in B10.D2 but deleted in DBA/2 mice.

We previously demonstrated that $V\beta 6^+$ T cells which are transferred with other splenic lymphocytes from the donor strain B10.D2 into tumour-bearing DBA/2 mice are able to break the anti-tumour tolerance (Schirmacher *et al*, 2000) and to infiltrate the metastasised organs (Schirmacher *et al*, 1995). In the liver, these $V\beta 6^+$ T cells form close contacts with vSAG7 expressing metastasised tumour cells (vSAG7⁺ ESb-MP) and with host macrophages in the vicinity of metastases (Mürköster *et al*, 1998). Moreover, treatment of tumour-bearing DBA/2 mice with freshly isolated $V\beta 6^+$ T cells led to retarded tumour outgrowth and to prolonged survival in comparison to untreated animals. In addition, $V\beta 6^+$ cells killed specifically tumour cells expressing the endogenous vSAG7 *in vitro* (Schirmacher *et al*, 2000).

Cytotoxic T lymphocytes (CTL) kill their target cells via apoptosis by different mechanisms, for instance, via CD95-CD95L interaction (Henkart, 1994; Kägi *et al*, 1994) or via the release of cytolytic granules containing perforin and granzymes upon contact with the target cell (Kägi *et al*, 1994; Pinkoski *et al*, 2000). Other molecules which can induce apoptosis especially in tumour cells are TNF- α and TNF-related apoptosis inducing ligand (TRAIL) (Wiley *et al*, 1995). After triggering the respective death receptor a cascade of caspases is activated, which finally leads to DNA-frag-

*Correspondence: Professor V Schirmacher;
E-mail: v.schirmacher@dkfz-heidelberg.de

⁴Current address: Laboratory of Molecular Gastroenterology, I, Department of Medicine, University of Kiel, Schittenhelmstraße 12, 24105 Kiel, Germany

Received 6 September 2001; revised 22 November 2001; accepted 23 November 2001

mentation and cell death (Krammer, 1998; Pinkoski *et al*, 2000). The apoptotic process can also implicate changes in the mitochondria that leads to the loss of the mitochondrial transmembrane potential ($\Delta\psi_m$) (Ushmorov *et al*, 1999; Kroemer and Reed, 2000). Granzymes (Ebnet *et al*, 1991) which belong to the family of serine proteases are also found to induce DNA-fragmentation, for instance by activating caspase-3 (Pinkoski *et al*, 2000). Yet, several recent reports indicate that induction of apoptosis and the subsequent DNA-fragmentation do not necessarily involve the activation of caspases (Thomas *et al*, 2000).

SAGs are a collection of bacterial and viral proteins with potent immunostimulatory properties. The targeting of them to tumour cells has become an interesting new concept to augment endogenous anti-tumour reactivity but little is known so far about the mechanisms involved (Choi *et al*, 1996). This study aimed to investigate how vSAG7 activated T lymphocytes kill SAG expressing tumour cells and to obtain more insights into the mechanism of SAG specific tumour cell lysis and GvL activity.

MATERIALS AND METHODS

Animals and cell lines

B10.D2 mice (Ola, Bicester, UK) were kept under specific pathogen free conditions and were used at the age of 6–8 weeks. All animal experiments were performed according to the standards required by the UKCCCR Guidelines. Eb is a chemically induced T cell lymphoma (L5178Y/E) of DBA/2 mice. ESb-MP is a spontaneous high metastatic adhesion variant of Eb and arose most likely after fusion of Eb cells with a host macrophage (Larizza *et al*, 1984). Jurkat is a human leukaemia T cell line, BL60 is a human leukaemia B cell line and U937 is a human promonocytic cell line. Cell lines were maintained in 5% CO₂ at 37°C in RPMI-1640 containing 10% FCS (both Life Technologies, Eggenstein, Germany). In some experiments, V β 6⁺ T cells were primed by injecting 5×10^4 ESb-MP tumour cells in PBS into the ear pinna of B10.D2 mice. Animals were sacrificed 9 days after injection.

Abs and other reagents

The rat-anti-mouse V β 6 mAb was used as culture supernatant (clone 44-22-1) (Payne *et al*, 1988). The anti-APO-1 antibody (Trauth *et al*, 1989) is directed against human (hu) CD95. hu TRAIL-R2-Fc, hu TNF-R2-Fc and hu CD95-Fc were used in the blocking experiments (Walczak *et al*, 1997, 1999). Leucine Zipper (LZ)-TRAIL, TNF- α and LZ-CD95L were used for induction of apoptosis (Walczak *et al*, 1999). All human reagents are also reactive to murine cells (Walczak *et al*, 1999). Human IgG was purchased from Sigma (Deisenhofen, Germany). Recombinant (rec.) human IFN- γ was obtained from Biomol (Hamburg, Germany). Annexin V-FITC was obtained from R&D systems (Wiesbaden, Germany). The following caspase inhibitors were purchased from Bachem (Heidelberg, Germany): ZVAD-fmk (Z-Val-Ala-DL-Asp-fluoromethylketone) and IETD-CHO (Ac-Ile-Glu-Thr-Asp-aldehyde (pseudo acid). JC1 (5,5',6,6'-tetrachloro-1,1',3,3'-tetraethyl-benzimidazol-carbocyanine iodide) was from Molecular Probes (Eugene, OR, USA). Concanamycin A and granzyme B inhibitor (Z-AAD-CMK) were obtained from Calbiochem (Bad Soden, Germany). To evaluate apoptotic effects of nitric oxide (NO), glycerol trinitrate (GTN; Merck; Darmstadt, Germany) was used.

Isolation of V β 6⁺ T cells via magnetic beads

V β 6⁺ T cells were isolated as previously described (Mürköster *et al*, 1998). Spleen cells from B10.D2 mice were incubated with rat-anti-V β 6 mAbs and with magnetic beads-conjugated anti-rat IgG isotype Abs (Dynal, Hamburg, Germany). Cells were separated via a Dynal-

magnet. The purity of the positively selected V β 6⁺ T cells was >88%. In some experiments, CD4⁺ or CD8⁺ T cells were isolated by depletion of CD8⁺ or CD4⁺ T cells by Dynabeads-conjugated anti-mouse CD8 or Dynabeads-conjugated anti-mouse CD4 (both Dynal) respectively. Then CD4⁺V β 6⁺ and CD8⁺V β 6⁺ T cells were isolated as previously described (Mürköster *et al*, 1998). After isolation, CD8⁺V β 6⁺ T cell population contained 0.4% CD4⁺ cells and the population of CD4⁺V β 6⁺ T cells 0.3% CD8⁺ cells.

Mixed lymphocyte tumour cell culture for induction of cytotoxic activity

Isolated V β 6⁺ T cells from normal or immunised B10.D2 mice were incubated with 100 Gy γ -irradiated (Gammacell 1000, Ottawa, Canada) V β 6 negative spleen cells (used as APC) and ESb-MP or Eb tumour cells for 4 days. Cytotoxic activity of *in vitro* stimulated V β 6⁺ T cells were tested either by a ⁵¹Cr-release-assay or by FACS analysis.

⁵¹Cr-release-assay ESb-MP and Eb target cells were labelled with 0.2 μ Ci ⁵¹Cr sodium chromate (Amersham, Braunschweig, Germany) in RPMI-1640 medium with 30% FCS for 90 min at 37°C. V β 6⁺ effector T cells were incubated with target cells (effector:target ratio 10:1) in 96-well round bottom plates (Renner, Dannstadt, Germany) for 4 h at 37°C. Radioactivity released in supernatants was measured by a γ -counter (LKB-Wallac, Freiburg, Germany) as described (Schirmmacher *et al*, 2000). In all experiments, an effector:target ratio of 10:1 was used. This ratio has been previously shown to be an optimal (Schirmmacher *et al*, 2000).

Determination of cytotoxicity via flow cytometry

Effector cells (V β 6⁺ T cells) were incubated with target ESb-MP cells (10:1 ratio) in 6-well plates for different periods of time (10–240 min). To determine which death receptors could be involved in the induction of apoptosis, ESb-MP cells were treated with 1 or 10 μ g ml⁻¹ LZ-TRAIL, 1 μ g ml⁻¹ LZ-CD95L, TNF- α or 10 μ g ml⁻¹ control IgG for 4 h without or with 24 h pre-incubation with IFN- γ . As a control for induction of apoptosis, either ESb-MP cells were incubated with 0.5 mM GTN for 24 h or Jurkat cells were treated with 1 μ g ml⁻¹ anti-APO-1/Protein A (10 ng ml⁻¹, Sigma) or 1 μ g ml⁻¹ LZ-CD95L, BL60 cells were treated with 1 μ g ml⁻¹ LZ-TRAIL and U937 cells were treated with TNF- α overnight to induce apoptosis. To block apoptosis in ESb-MP cells, the following reagents were used: hu TRAIL-R2-Fc (10 μ g ml⁻¹), hu CD95-Fc (10 μ g ml⁻¹), hu TNF-R2-Fc (10 μ g ml⁻¹), hu IgG1 as isotype control (10 μ g ml⁻¹), Concanamycin A (10 nM), granzyme B inhibitor (20 μ M), ZVAD-fmk (25 μ M) and IETD-CHO (25 μ M). Treatment of effector and target cells with the respective reagent started 15–60 min before the co-incubation. As a positive control, 1–30 μ g ml⁻¹ CD95-Fc, TNF-R2-Fc and TRAIL-R2-Fc were used to block apoptosis in Jurkat, BL60 and U937 cells. After removal of effector cells, ESb-MP cells were washed, detached from the plates, and cytotoxicity was determined by staining with 1 μ g ml⁻¹ propidium iodide (PI) which was added to the cells 5 min before measurement.

Target cell apoptosis

Apoptosis was assessed either by Annexin V/PI staining (Koopman *et al*, 1994) or by determining DNA-fragmentation (Nicoletti *et al*, 1991). After co-incubation with effector cells, 5×10^5 ESb-MP target cells were either resuspended in binding buffer (R&D systems), 50 ng ml⁻¹ Annexin V and 1 μ g ml⁻¹ PI or treated with a hypotonic solution (0.1% sodium citrate, 0.1% Triton X-100) containing 50 μ g ml⁻¹ PI overnight. Jurkat cells were incubated in the hypotonic solution containing PI and DNA-fragmentation was measured as described above. Target cell death and apoptosis were analyzed by flow cytometry using a FACScan analyzer with CellQuest software (BD Bioscience, Heidelberg, Germany).

Cytofluorometric analysis of mitochondrial transmembrane potential ($\Delta\psi_m$)

$\Delta\psi_m$ was measured with JC1 as described (Salvioli *et al*, 1997). JC1 is a cyanine dye which accumulates in the mitochondrial matrix under the influence of the $\Delta\psi_m$. In the presence of a high $\Delta\psi_m$, JC1 forms aggregates which have characteristic absorption and emission spectra so that cells are detectable as FL1⁻ and FL2⁺. 5×10^5 ESb-MP tumour cells were treated with $5 \mu\text{g ml}^{-1}$ JC1 for 20 min at 4°C followed by FACScan analysis. As a control, Jurkat cells were treated with $1 \mu\text{g ml}^{-1}$ anti-APO-1 mAb cross-linked by 10 ng ml^{-1} Protein A for 12 h followed by incubation with JC1 as described above.

Detection of perforin in $V\beta 6^+$ T cells

Isolated $V\beta 6^+$ T cells were stimulated either with ESb-MP or with Eb cells and APCs or were left without any stimulation. After 4 days supernatants were taken and centrifuged by 500 r.p.m. for 5 min. The pellet containing $V\beta 6^+$ T cells was washed with PBS and centrifuged on siliconised glass slides (Sigma). After drying, slides were fixed in acetone for 10 min at room temperature followed by washing in PBS. To avoid nonspecific binding, slides

were treated with 1% normal goat serum for 15 min followed by incubation with the rat-anti-mouse perforin mAb (Alexis, Grünberg, Germany) for 45 min. After washing, slides were treated with a goat-anti-rat antibody conjugated with alkaline phosphatase (AP). Then slides were washed with water, counterstained with hemalaun (Sigma) and mounted with glycerol-gelatin (Merck). The substrate for the development of AP consisted of $6.3 \mu\text{l}$ 5% Neufuchsin (Sigma) or 2 mg Fast Blue (Sigma) in $16 \mu\text{l}$ 4% sodium nitrite (Fluka, Buchs, Switzerland), 2 mg naphthol-As-Bi-phosphate (Sigma) in $20 \mu\text{l}$ N,N-dimethylformamide (Merck) and 3 ml of 0.05 M l^{-1} Tris-HCl-buffer, pH 8.7 containing 1 mM levamisole (Sigma). As negative control either the first antibody was omitted or an isotype matched control antibody was used. Both stainings revealed no staining.

RESULTS

Activated $V\beta 6^+$ T cells kill vSAG7⁺ ESb-MP tumour cells by induction of apoptosis

$V\beta 6^+$ T cells, stimulated for 4 days with tumour cells and spleen cells as APC were able to kill vSAG7⁺ ESb-MP tumour cells while vSAG7⁻ Eb tumour cells were hardly lysed (Figure 1A). Experi-




Figure 1 Cytotoxicity towards vSAG7⁺ tumour cells generated by stimulated $V\beta 6^+$ T cells *in vitro*. **(A)** $V\beta 6^+$ T cells were isolated from B10.D2 mice which were either immunised with ESb-MP cells or remained untreated (1° *in vivo*). Then, $V\beta 6^+$ T cells were stimulated for 4 days either with irradiated vSAG7⁻ Eb or with vSAG7⁺ ESb-MP cells and APC (2° *in vitro*) and tested for cytotoxicity towards Eb or ESb-MP target cells (effector: target ratio 10:1) in a 4 h ⁵¹Cr-release assay. The mean \pm standard deviation of triplicates of two independent experiments are shown. **(B)** $CD4^+ V\beta 6^+$, $CD8^+ V\beta 6^+$ and total $V\beta 6^+$ T cells were isolated from non-immunised B10.D2 mice, purified and stimulated with irradiated ESb-MP and APCs for 4 days. Then effector cells were co-cultured with target ESb-MP cells for 2 h, and tumour cell lysis was determined by PI staining and FACScan analysis. The mean \pm standard deviation from 2–4 independent experiments are shown.

ments with the separation of $V\beta 6^+$ T cells into $CD4^+$ and $CD8^+$ T cells revealed that both T cell subpopulations induced cell death in ESb-MP target cells (Figure 1B).

CTLs directed to conventional antigens consisting of peptide/MHC complexes kill their target cells via apoptosis (Henkart, 1994; Kägi *et al*, 1994). To determine whether vSAG7 reactive $V\beta 6^+$ T cells are able to induce apoptosis, we assessed the death of vSAG7⁺ ESb-MP cells after co-incubation with stimulated $V\beta 6^+$ T cells by two different methods. Figure 2A shows the kinetics of apoptosis measured by Annexin V staining/PI staining. After 10 min, 21% of the ESb-MP cells were Annexin V⁺ indicating that they were in the early phase of apoptosis. Additionally, at this time point, 26% were Annexin V⁺/PI⁺, a phenotype specific for the late stage of apoptosis. The maximal amount of apoptotic cells (63%) was detected after 20 min and remained at this level until 240 min of co-incubation. Changes in the forward/side scatter which are also typical for apoptosis started as well after 10 min of co-incubation of effector and target cells (data not shown). When measuring oligonucleosomal DNA-fragmentation, 45% of apoptotic ESb-MP cells were observed after 10 min of coculture (Figure 2B). The maximal level of apoptotic cells was reached after 30 min. Taken together, these observations reveal that $V\beta 6^+$ T cells are able to rapidly induce apoptosis in vSAG7⁺ ESb-MP tumour cells.

Induction of apoptosis in ESb-MP cells is caspase-independent and does not involve mitochondrial damage

Caspases are known as main effector enzymes responsible for the initiation of DNA-fragmentation and the typical morphological changes in apoptosis (Henkart, 1994; Krammer, 1998). To analyse the involvement of caspases in this rapid vSAG7-mediated tumour cell lysis, we added either ZVAD-fmk (a broad spectrum caspase inhibitor) or IETD-CHO (a specific inhibitor of caspase 8) to the coculture of $V\beta 6^+$ T cells and ESb-MP tumour cells. As shown in Figure 3, ZVAD-fmk and IETD-CHO blocked almost completely the death of Jurkat cells which were treated with anti-CD95 mAbs. In addition, ZVAD-fmk also inhibited NO-mediated apoptosis in ESb-MP cells. In contrast, ZVAD-fmk and IETD-CHO did not influence significantly the $V\beta 6^+$ T cell mediated apoptosis in ESb-MP cells. Thus $V\beta 6^+$ T cells were able to induce apoptosis in vSAG7⁺ ESb-MP tumour cells in a caspase-independent manner.

The dye JC1 is known as a marker for changes in the $\Delta\psi_m$ which can be associated with apoptosis (Salvioli *et al*, 1997). To evaluate whether mitochondria are involved in the induction of vSAG7 mediated apoptosis in ESb-MP cells, we performed JC1 staining of ESb-MP cells after co-incubation with $V\beta 6^+$ T cells

for different periods of time. In cells with intact mitochondria and high $\Delta\psi_m$, JC1 forms so called J-aggregates that are associated with a large shift in emission (590 nm=Fl 1). In Jurkat cells which were treated with anti-CD95 mAbs for 12 h the involvement of mitochondria could be clearly detected (Figure 4). In contrast no changes in $\Delta\psi_m$ were observed during 4 h co-culture of $V\beta 6^+$ T cells with ESb-MP tumour cells (Figure 4). Staining with 3,3'-dihexyloxycarbocyanine iodide (DiOC₆) and dihydroethidine (HE), which are used to determine changes in $\Delta\psi_m$ and the production of reactive oxygen species (ROS), respectively (Ushmorov *et al*, 1999) also showed no mitochondrial alterations during $V\beta 6^+$ T cell-mediated apoptosis in ESb-MP tumour cells (data not shown). In contrast, NO, another inducer of apoptosis in these cells, stimulated mitochondrial changes (data not shown).




Figure 2 Stimulated $V\beta 6^+$ T cells induce apoptosis in vSAG7⁺ ESb-MP tumour cells. *In vitro* stimulated $V\beta 6^+$ T cells were co-incubated with ESb-MP cells in a ratio of 10:1 for different periods of time and stained either with Annexin V-FITC and $1 \mu\text{g ml}^{-1}$ PI (A) or were treated with a hypotonic solution containing $50 \mu\text{g ml}^{-1}$ PI overnight to determine DNA-fragmentation (B). Untreated cells showed no changes in the amount of apoptotic cells during co-incubation. Apoptosis was determined by FACS-can analysis. One representative experiment out of four (A) or three (B) independent experiments is shown.




Figure 3 $V\beta 6^+$ T cell induced apoptosis in ESb-MP cells is caspase-independent. Stimulated $V\beta 6^+$ T cells were co-incubated with ESb-MP tumour cells (effector:target ratio 10:1) for 4 h or treated with 0.5 mM GTN (NO). Jurkat cells were stimulated with an anti-APO-1 (CD95) mAb/Protein A for 12 h. In some cultures, $25 \mu\text{M}$ ZVAD-fmk or $25 \mu\text{M}$ IETD-CHO were added. As a control, ESb-MP cells and Jurkat cells were cultured without $V\beta 6^+$ T cells and reagents. Cell death was determined by PI staining and FACSscan analysis. The mean \pm standard deviation from three independent experiments are shown.

TRAIL, CD95L and TNF- α are not involved in the induction of apoptosis in ESb-MP cells

Several death receptors and their respective ligands have been shown to be involved in the induction of apoptosis (Kägi *et al*, 1994; Wiley *et al*, 1995). The most common are TRAIL-R1+2/TRAIL, CD95/CD95L and TNF-R1+2/TNF. We tested whether ESb-MP cells were sensitive towards the apoptosis-inducing potential of CD95L, TRAIL and TNF- α . Figure 5A shows the positive controls: CD95L was able to induce apoptosis in Jurkat cells, TRAIL induced apoptosis in BL60 and TNF- α induced apoptosis in U937 cells. In each cell line, apoptosis could be blocked using soluble Fc-fusion proteins against the respective ligands. In contrast, ESb-MP tumour cells were not sensitive towards any of these ligands, even after IFN- γ stimulation for 24 h since cytotoxicity was not increased in comparison to untreated ESb-MP cells (Figure 5B). Furthermore, blocking with TRAIL-R2-Fc, CD95-Fc and TNF-R2-Fc did not significantly affect cell lysis of ESb-MP target cells by stimulated V β 6⁺ T cells (Figure 5C). Treatment with the isotype matched Ig control increased slightly but not significantly the number of apoptotic ESb-MP cells. We conclude that V β 6⁺ T cells induce apoptosis in ESb-MP tumour cells independently of CD95L, TRAIL and TNF- α .

V β 6⁺ T cells induce cytotoxicity in ESb-MP cells through perforin/granzyme B

Perforin/granzymes are able to induce apoptosis in a death receptor-independent manner (Kägi *et al*, 1994; Trapani *et al*, 1998; Pinkoski *et al*, 2000). Figure 6A shows that V β 6⁺ T cells stimulated with vSAG7⁺ ESb-MP cells for 4 days expressed perforin while no expression of perforin was found in V β 6⁺ T cells from control cultures without stimulation or stimulated with vSAG7⁻ Eb cells. Figure 6B shows the effects of treatment of V β 6⁺ T effector cells

and ESb-MP target cells with Concanamycin A, an inhibitor of perforin or of treatment with an inhibitor of granzyme B or a combination of both. Concanamycin A alone did not inhibit target cell lysis while treatment with the granzyme B inhibitor led to a significant reduction of apoptosis. Treatment with both inhibitors almost completely prevented the lysis of ESb-MP cells (from 35 to 5% apoptosis, Figure 6B). Together, these data indicate that vSAG7 stimulated expression of perforin in V β 6⁺ T cells. These activated cells induced apoptosis in ESb-MP tumour cells via released perforin and granzyme B.

DISCUSSION

We have recently identified a tumour associated endogenous vSAG7 as a new target antigen for allogeneic GvL reactivity in a murine tumour model (Schirmmacher *et al*, 2000). In DBA/2 mice, vSAG7 behaves as a strong tolerogen leading to central deletion of SAG-reactive V β 6⁺ T cells during thymic maturation. While the DBA/2 derived Eb T lymphoma cells did not express this SAG, the spontaneous metastatic variant ESb and its adhesion variant ESb-MP expressed endogenous mouse mammary tumour provirus (Mtv)- typical intracisternal A-particles (IAP), Mtv7 orf/SAG message and vSAG7 protein at the cell surface (Schirmmacher *et al*, 2000). MMTV and IAPs have been suggested to be associated with tumorigenicity and progression towards increased malignancy (Schirmmacher *et al*, 1998). We previously reported that the allogeneic MHC identical mouse strain B10.D2 in contrast to DBA/2 was able to strongly reject ESb and ESb-MP tumours and that vSAG7 reactive V β 6⁺ T cells were involved in this GvL reactivity (Schirmmacher *et al*, 2000).

In this study, we investigated the mechanism of tumour cell destruction *in vitro* by vSAG7 reactive V β 6⁺ T cells. We show here that V β 6⁺ T cells kill vSAG7⁺ tumour cells via release of perforin and granzyme B. Interestingly, this form of cell death is indepen-




Figure 4 V β 6⁺ T cell-induced apoptosis in ESb-MP cells does not involve changes in $\Delta\psi_m$. ESb-MP cells remained untreated (light line) or were cocultured with stimulated V β 6⁺ T cells in an effector:target ratio of 10:1 for 4 h (dark line). As a control, Jurkat cells were left untreated (light line) or treated with anti-APO-1 mAb/Protein A for 12 h (dark line). Afterwards, ESb-MP cells and Jurkat cells were either analysed for DNA-fragmentation (FL3) or for changes in $\Delta\psi_m$ by JCI staining (FL1). Upon incubation of Jurkat cells with anti-APO-1 mAb, FL2 did not significantly change, whereas FL1, which indicates the formation of dye monomer, increased due to the reduction of $\Delta\psi_m$. This increase is taken as a measure for the loss of $\Delta\psi_m$. One representative experiment out of three is shown.

dent of the known caspases and occurs without the involvement of mitochondria. It has been previously reported that vSAG7 can induce IFN- γ production by specifically primed CD8⁺ T cells but fails to trigger cytotoxicity (Herrmann *et al*, 1992). However, in contrast to Herrmann *et al* (1992), we find that vSAG7 stimulates V β 6⁺ T cells to express perforin and to kill vSAG7⁺ target cells. This kill is SAG-specific and occurs in several tumour cells that express this endogenous viral superantigen (Schirmacher *et al*, 2000). Interestingly, this SAG specific CTL activity is mediated via both CD8⁺V β 6⁺ and CD4⁺V β 6⁺ T cells.

Bacterial SAG have been shown to trigger CD8⁺ T cell clones that are specific for other antigens (e.g. influenza virus peptides), but in these systems the exerted cytotoxicity was mediated via the CD95/CD95L system (Fuller and Braciale, 1998). Sundstedt *et*

al (1998) showed that injection of staphylococcal enterotoxin A into perforin-deficient mice led to less depletion of B-cells than in control mice. This depletion was due to the release of perforin by CD8⁺ T cells. In GvL and GvH reactivity, a role for both systems (granule exocytosis as well as CD95/CD95L) was reported (Miwa *et al*, 1999; Yasukawa *et al*, 2000). It was thus of interest to elucidate the mechanism by which allogeneic V β 6⁺ T cells kill vSAG7⁺ tumour cells.

TNF- α and CD95L play a crucial role in the activation induced cell death (AICD) of mature T cells (Dhein *et al*, 1995; Zheng *et al*, 1995) whereas TRAIL has been shown to induce apoptosis mainly in tumour cells (Wiley *et al*, 1995; Walczak *et al*, 1999) but also in normal human hepatocytes, astrocytes and keratinocytes (Jo *et al*, 2000). Esb-MP tumour cells were not sensitive towards any of




Figure 5 Esb-MP cells are resistant towards TRAIL-, CD95L- or TNF- α induced apoptosis. **(A)** Jurkat cells were treated with 1 μ g ml⁻¹ CD95L, BL60 cells were cultured with 1 μ g ml⁻¹ TRAIL and U937 cells were incubated with 1 μ g ml⁻¹ TNF- α for 12 h. Apoptosis was blocked by 1–30 μ g ml⁻¹ CD95-Fc, TRAIL-R2-Fc or TNF-R2-Fc. One representative experiment out of three is shown. **(B)** Esb-MP cells were incubated with 10 nM recombinant IFN- γ for 24 h in RPMI medium supplemented with 10% FCS followed by treatment with 1 or 10 μ g ml⁻¹ TRAIL, 1 μ g ml⁻¹ CD95L, 1 or 10 μ g ml⁻¹ TNF- α . As a control, Esb-MP cells remained untreated (Esb-MP). **(C)** Esb-MP cells were cultured for 4 h alone, with V β 6⁺ T cells (V β 6), V β 6⁺ T cells and IgG (10 μ g ml⁻¹), with V β 6⁺ T cells and TRAIL-R2-Fc (10 μ g ml⁻¹), V β 6⁺ T cells and CD95-Fc (10 μ g ml⁻¹) or with V β 6⁺ T cells and TNF-R2-Fc (10 μ g ml⁻¹). In **(B)** and **(C)** the effector:target cell ratio was 10:1. The death of Esb-MP and Jurkat cells was assessed by PI staining and FACScan analysis. The mean \pm standard deviation from three independent experiments are shown.





Figure 6 Apoptosis of ESb-MP cells is mediated by perforin and granzyme B. **(A)** Shows perforin expression in $V\beta 6^+$ T cells. $V\beta 6^+$ T cells were stimulated with vSAG7⁻ Eb or vSAG7⁺ ESb-MP cells and APCs for 4 days. Then unstimulated (control) or stimulated $V\beta 6^+$ T cells were stained with an anti perforin mAb. Cells were counterstained with hemalaun. Membrane staining indicates positively labelled cells. Black dots are magnetic beads

these death ligands and blocking of these pathways during coincubation of effector and tumour cells did not influence the cytotoxic reactivity of $V\beta 6^+$ T cells.

The perforin/granzyme system is involved in the cytotoxic reactivity of CD8⁺ T cells (Kägi *et al*, 1994; Pinkoski *et al*, 2000). Recently, perforin has been shown to play a role in AICD of T cells in an allogeneic transplantation model (Spaner *et al*, 1999). In our model, $V\beta 6^+$ T cells showed a high perforin expression after 4 days of *in vitro* stimulation. The lysis of ESb-MP tumour cells was almost completely inhibited by the perforin (concanamycin A) and granzyme B inhibitors. Interestingly, treatment with granzyme B inhibitor but not with concanamycin A alone inhibited target cell lysis. This might be due to the fact that membrane damage induced by perforin does not directly cause target cell death, but facilitates the entry of granzymes into the cell (Shresta *et al*, 1999; Trapani *et al*, 1998). Granzyme B, however, can also enter the cell via the cation-independent mannose 6-phosphate/insulin-like growth factor receptor (Motyka *et al*, 2000), which might explain our observations.

The vSAG7 mediated tumour cell lysis was very fast. This is in accordance with other observations demonstrating that granzyme B, in contrast to granzyme A, is responsible for a rapid cytotoxic effect (Drénou *et al*, 1999). We cannot exclude that granzyme A might also be active in ESb-MP cells, since application of the granzyme B inhibitor alone did not completely block apoptosis. However, there is evidence that granzyme A and B do not synergize but act independently of each other (Shresta *et al*, 1999).

The requirement of caspases as the main effector enzymes in the initiation of the apoptotic cascade is still a matter of debate (Krammer, 1998; Pinkoski *et al*, 2000). Several recent reports indicate that induction of apoptosis and subsequent DNA-fragmentation do not necessarily involve the activation of caspases (Beresford *et al*, 1999; Drénou *et al*, 1999). vSAG7 specific $V\beta 6^+$ T cells could kill ESb-MP tumour cells independently of caspases since blocking with ZVAD-fmk and IETD-CHO did not lead to an inhibition of apoptosis in ESb-MP cells. In contrast, NO-induced apoptosis in these tumour cells required caspase activity. Beresford *et al* (1999) showed that cytolysis mediated by granzyme B was caspase independent whereas DNA-fragmentation was caspase-dependent. In contrast, in our model blocking with caspase inhibitors influenced neither cytolysis nor DNA-fragmentation (data not shown).

The question arises how granzyme B may lead to DNA cleavage without activation of caspases. It was reported that granzyme B directly cleaves caspase substrates such as poly (ADP-ribose) polymerase, the catalytic subunit of DNA-dependent protein kinases and NuMA to bring about apoptotic changes in cells (Froelich *et al*, 1996; Andrade *et al*, 1998). Moreover, granzyme B induced the processing of DFF45/ICAD in a caspase-independent fashion resulting in DNase activation and DNA fragmentation (Thomas *et al*, 2000). In addition, granule-mediated cell death can proceed independently of caspases through a non-nuclear pathway (Sarin *et al*, 1997). These pathways seemed to be important for cells that are infected by viruses which are able to postpone cell death by caspase inhibitors like crmA (Zhou *et al*, 1997). This might explain

which were used for the purification and could not be removed after 4 days of stimulation. Magnification $\times 400$. **(B)** ESb-MP cells were cultured either alone (ESb-MP) or with stimulated $V\beta 6^+$ T cells at an effector:target ratio of 10:1. Separate groups of effector cells were treated either with 10 nM Concanamycin A (Concan.A), 20 μ M granzyme B inhibitor (Z-AAD-CMK) or with a combination of both inhibitors ($V\beta 6$ +Concan.A+GranzB.Inh.) for 15 min before coincubation with target cells for 4 h. Incubation of ESb-MP cells alone with Concanamycin A and granzyme B inhibitor revealed no significant difference in comparison to the ESb-MP control. Apoptosis was determined by PI incorporation and FACScan analysis. The mean \pm standard deviation from three independent experiments are shown.

why we find this kind of cell death in ESb-MP tumour cells in which virus-like particles and proviral genes are expressed (Mürköster *et al*, 1998).

The involvement of mitochondria in apoptosis was described for different apoptotic agents (Kroemer and Reed, 2000). We reported previously that NO-mediated apoptosis in human leukaemia cells is associated with mitochondrial damage (i.e., a degradation of major mitochondrial lipid cardiolipin and cytochrome *c* release into the cytosol) followed by an activation of caspase-9 and caspase-3 (Ushmorov *et al*, 1999). Furthermore, we found that NO produced by activated Kupffer cells during adoptive immunotherapy with GvL-reactive T cells contributed to apoptosis in metastatic ESb-MP tumour cells in the liver (Mürköster *et al*, 2000). We showed also that this cell death can be induced by a mitochondria-dependent mechanism driven by CD40-CD40L interactions (Mürköster *et al* 2000). In contrast, vSAG7 mediated apoptosis in ESb-MP cells occurred independently of mitochondria since staining with JC1, DiOC₆ and HE revealed no mitochondrial damage.

REFERENCES

- Andrade F, Roy S, Nicholson D, Thornberry N, Rosen A, Casciola-Rosen L (1998) Granzyme B directly and efficiently cleaves several downstream caspase substrates: implications for CTL-induced apoptosis. *Immunity* **8**: 451–460
- Beresford PJ, Xia Z, Greenberg AH, Liebermann J (1999) Granzyme A loading induces rapid cytolysis and a novel form of DNA damage independently of caspase activation. *Immunity* **10**: 585–594
- Beutner U, Frankel WN, Cote MS, Coffin JM, Huber BT (1992) Mls-1 is encoded by the long terminal repeat open reading frame of the mouse mammary tumor provirus Mtv-7. *Proc Natl Acad Sci USA* **89**: 5432–5436
- Choi Y, Marrack P, Kappler JW (1996) Structural analysis of a mouse mammary tumor virus superantigen. *J Exp Med* **175**: 847–852
- Dhein J, Walczak H, Bäumler C, Debatin KM, Krammer PH (1995) Auto-crine T-cell suicide mediated by APO-1 (Fas/CD95). *Nature (Lond.)* **373**: 438–441
- Drénou B, Blanchereau V, Burgess DH, Fauchet R, Charron DJ, Mooney NA (1999) A caspase-independent pathway of MHC Class II antigen-mediated apoptosis of human B lymphocytes. *J Immunol* **163**: 4115–4124
- Ebnet K, Chluba-de Tapia J, Hurtenbach U, Kramer MD, Simon MM (1991) In vivo-primed mouse T cells selectively express T cell-specific serine proteinase-1 and proteinase-like molecules granzyme B and C. *Int Immunol* **3**: 9–19
- Froelich CJ, Hanna WL, Poirier GG, Duriez PJ, D'Amours D, Salvesen GS, Alnemri ES, Earnshaw WC, Shah GM (1996) Granzyme B/perforin-mediated apoptosis of Jurkat cells results in cleavage of poly(ADP-ribose) polymerase to the 89-kDa apoptotic fragment and less abundant 64-kDa fragment. *Biochem Biophys Res Commun* **227**: 658–665
- Fuller CL, Braciale VL (1998) Selective induction of CD8+ cytotoxic T lymphocytes effector function by Staphylococcus Enterotoxin B. *J Immunol* **161**: 5179–5186
- Henkart PA (1994) Lymphocyte-mediated cytotoxicity: two pathways and multiple effector molecules. *Immunity* **1**: 343–346
- Herrmann T, Waander GA, Chvatchko Y, MacDonald HR (1992) The viral superantigen Mls-1^a induces interferon- γ secretion by specifically primed CD8+ cells but fails to trigger cytotoxicity. *Eur J Immunol* **22**: 2789–2793
- Jo M, Kim TH, Seol DW, Esplen JE, Dorko K, Billiar TR, Strom SC (2000) Apoptosis induced in normal human hepatocytes by tumor necrosis factor-related apoptosis-inducing ligand. *Nature Med* **6**: 564–567
- Kägi D, Vignaux F, Ledermann B, Bürki K, Depraetere V, Nagata S, Hengartner H, Golstein P (1994) Fas and perforin pathways as major mechanisms of T-cell mediated cytotoxicity. *Science* **265**: 528–530
- Koopman G, Reutelingsperger CP, Kuijten GA, Keehnen, RM, Pals ST, van Oers MH (1994) Annexin V for flow cytometric detection of phosphatidylserine expression on B cells undergoing apoptosis. *Blood* **84**: 1415–1420
- Krammer PH (1998) CD95 (Apo-1/Fas)-mediated apoptosis: Live and let die. *Adv Immunol* **71**: 163–207
- Kroemer G, Reed JC (2000) Mitochondrial control of cell death. *Nature Med* **6**: 513–519
- Larizza L, Schirmmacher V, Pflüger E (1984) Acquisition of high metastatic capacity after in vitro fusion of a non-metastatic tumor line with a bone-marrow derived macrophage. *J Exp Med* **160**: 1579–1584
- Mackinnon S, Papadopoulos EB, Carabasi MH, Reich L, Collins NH, Boulad F, Castro-Malaspina H, Childs BH, Gillio AP, Kernan NA (1995) Adoptive immunotherapy evaluating escalating doses of donor leukocytes for relapse of chronic myeloid leukemia after bone marrow transplantation: separation of graft-versus-leukemia responses from graft-versus-host disease. *Blood* **86**: 1261–1268
- Miwa K, Hashimoto H, Yatomi T, Nakamura N, Nagata S, Suda T (1999) Therapeutic effect of an anti-Fas ligand mAb on lethal graft-versus-host disease. *Int Immunol* **11**: 925–931
- Motyka B, Korbitt G, Pinkowski MJ, Heilbein JA, Caputo A, Hobman M, Barry M, Shostak I, Sawchuk T, Holmes CFB, Gauldie J, Bleackley RC (2000) Mannose 6-Phosphate/Insulin-like Growth factor II receptor is a death receptor for granzyme B during cytotoxic T cell-induced apoptosis. *Cell* **103**: 491–500
- Mürköster S, Wachowski O, Zerban H, Schirmmacher V, Umansky V, Rocha M (1998) Graft-versus-leukemia reactivity involves cluster formation between superantigen-reactive donor T lymphocytes and host macrophages. *Clin Cancer Res* **4**: 3095–3106
- Mürköster S, Laman JD, Rocha M, Umansky V, Schirmmacher V (2000) Functional and *in situ* evidence for nitric oxide production driven by CD40-CD40L interactions in graft-versus-leukemia reactivity. *Clin Cancer Res* **6**: 1988–1996
- Nicoletti I, Migliorati G, Pagliacci MC, Grignani F, Riccardi C (1991) A rapid and simple method for measuring thymocyte apoptosis by propidium iodide staining and flow cytometry. *J Immunol Method* **139**: 271–279
- Payne J, Huber BT, Cannon NA, Schneider R, Schilham MW, Acha-Orbea H, Mac-Donald HR, Hengartner H (1988) Two monoclonal rat antibodies with specificity for the β -chain variable region V β 6 of the murine T-cell receptor. *Proc Natl Acad Sci USA* **85**: 7695–7698
- Pinkoski MJ, Heilbein JA, Barry M, Bleackley RC (2000) Nuclear translocation of granzyme B in target cell apoptosis. *Cell Death Diff* **7**: 17–24
- Salvioli S, Ardizzoni A, Franceschi C, Cossarizza A (1997) JC-1, but not DiOC₆(3) or rhodamine 123, is a reliable fluorescent probe to assess $\Delta\psi$ changes in intact cells: implications for studies on mitochondrial functionality during apoptosis. *FEBS Lett* **411**: 77–82
- Sarin A, Williams MS, Alexander-Miller MA, Berzofsky JA, Zacharchuk CM, Henkart PA (1997) Target cell lysis by CTL granule exocytosis is independent of ICE/Ced-3 family proteases. *Immunity* **6**: 209–215
- Schirmmacher V, Beckhove P, Krüger A, Rocha M, Umansky V, Fichtner KP, Hull WE, Zangemeister-Wittke U, Griesbach A, Jurianz K, von Hoegen P (1995) Effective immune rejection of advanced metastasized cancer. *Int J Oncol* **6**: 505–521
- Schirmmacher V, Beutner U, Bucur M, Umansky V, Rocha M, von Hoegen P (1998) Loss of endogenous mouse mammary tumor virus superantigen increases tumor resistance. *J Immunol* **161**: 563–570

- Schirmacher V, Mürköster S, Bucur M, Umansky V, Rocha M (2000) Breaking tolerance to a tumor-associated viral superantigen as a basis for graft versus leukemia (GvL) reactivity. *Int J Cancer* **87**: 695–706
- Shresta S, Graubert TA, Thomas DA, Raptis SZ, Ley TJ (1999) Granzyme A initiates an alternative pathway for granule-mediated apoptosis. *Immunity* **10**: 595–605
- Slavin S, Nagler A, Naparstek E, Kapelushnik Y, Aker M, Cividalli G, Varadi G, Kirschbaum M, Ackerstein A, Samuel S, Amar A, Brautbar C, Ben-Tal O, Eldor A, Or R (1998) Nonmyeloablative stem cell transplantation and cell therapy as an alternative to conventional bone marrow transplantation with lethal cytoreduction for the treatment of malignant and nonmalignant hematologic diseases. *Blood* **91**: 756–765
- Spaner D, Raju K, Rabinovich B, Miller RG (1999) A role for perforin in activation-induced T cell death in vivo: Increased expansion of allogeneic perforin-deficient T cells in SCID mice. *J Immunol* **162**: 1192–1199
- Speiser DE, Schneider R, Hengartner H, MacDonald HR, Zinkernagel RM (1989) Clonal deletion of self-reactive T cells in irradiation bone marrow chimeras and neonatally tolerant mice. Evidence for intercellular transfer of Mls-1^a. *J Exp Med* **170**: 559–600
- Sundstedt A, Grundstrom S, Dohlsten M (1998) T cell- and perforin-dependent depletion of B cells in vivo by staphylococcal enterotoxin A. *Immunology* **95**: 76–82
- Thomas DA, Du C, Xu M, Wang X, Ley TJ (2000) DFF45/ICAD can be directly processed by granzyme B during the induction of apoptosis. *Immunity* **12**: 621–632
- Trapani JA, Jans PJ, Smyth MJ, Froelich CJ, Williams EA, Sutton VR, Jans DA (1998) Perforin-dependent nuclear entry of granzyme B precedes apoptosis, and is not a consequence of nuclear membrane dysfunction. *Cell Death Diff* **5**: 488–496
- Trauth BC, Klas C, Peters AMJ, Matzku S, Möller P, Falk W, Debatin KM, Krammer PH (1989) Monoclonal antibody-mediated tumor regression by induction of apoptosis. *Science* **245**: 301–305
- Ushmorov A, Ratter F, Lehmann V, Dröge W, Schirmacher V, Umansky V (1999) Nitric oxide-induced apoptosis in human leukemic lines requires mitochondrial degradation and cytochrome C release. *Blood* **93**: 2342–2352
- Walczak H, Degli-Esposti MA, Johnson RS, Smolak PJ, Waugh JY, Boiani N, Timour TS, Gerhart MJ, Schooley KA, Smith KA, Goodwin RG, Rauch CT (1997) TRAIL-R2: a novel apoptosis-mediating receptor for TRAIL. *EMBO J* **16**: 5386–5397
- Walczak H, Miller RE, Ariail K, Gliniak B, Griffith TS, Kubin M, Chin W, Jones J, Woodward A, Le T, Smith C, Smolak P, Goodwin RG, Rauch CT, Schuh JC, Lynch DH (1999) Tumoricidal activity of tumor necrosis factor-related apoptosis-inducing ligand in vivo. *Nature Med* **5**: 155–162
- Wiley SR, Schooley K, Smolak PJ, Din WS, Huang C-P, Nicholl JK, Sutherland GR, Smith TD, Rauch CT, Smith CA (1995) Identification and characterization of a new member of the TNF family that induces apoptosis. *Immunity* **3**: 673–682
- Yasukawa M, Ohminami H, Kasahara Y, Ishida Y, Fujita S (2000) Granule exocytosis, and not the Fas/Fas ligand system, is the main pathway of cytotoxicity mediated by alloantigen-specific CD4⁺ as well as CD8⁺ cytotoxic T lymphocytes in humans. *Blood* **95**: 2352–2355
- Zheng L, Fisher G, Miller RE, Peschon J, Lynch DH, Lenardo MJ (1995) Induction of apoptosis in mature T cells by tumor necrosis factor. *Nature* **377**: 348–351
- Zhou Q, Snipas S, Orth K, Muzio M, Dixit VM, Salvesen GS (1997) Target protease specificity of the viral serpin CrmA. Analysis of five caspases. *J Biol Chem* **272**: 7797–7800