



Genome Sequence of *Campylobacter fetus* subsp. *venerealis* NW_ED23, Isolated from Bovine Sheath Wash

 Mpinda Edoaurd Tshipamba,^a  Stephen Abiola Akinola,^b Lubanza Ngoma,^a  Mulunda Mwanza^a

^aDepartment of Animal Health, School of Agriculture, Faculty of Agriculture, Science, and Technology, Mafikeng Campus, North-West University, Mmabatho, South Africa

^bDepartment of Microbiology, North West University, Mmabatho, South Africa

ABSTRACT *Campylobacter fetus* subsp. *venerealis* is the causative agent of bovine genital campylobacteriosis, which is mostly characterized by reproduction problems. The strain reported in this study was isolated from bull sheath wash in South Africa.

Bovine genital campylobacteriosis belongs to list B of the notifiable diseases of the International Office of Epizootics (OIE) (1). A recent study reported that the *Campylobacter fetus* subsp. *venerealis* bv. *intermedius* genome sequence is available, but the same study confirmed that more analyses were needed to verify if there are extrachromosomal replicons (2). Furthermore, van der Graaf-van Bloois and coauthors revealed that an unassembled genome does not allow full identification of the core and accessory genome and the reconstruction of pathways and surface structures that might contribute to the pathogenicity of this biovar (3). An increase in the prevalence of this biovar has been noticed in South Africa (4). Hence, there was a need to sequence this particular strain to determine if there is any deviation from that previously reported.

The sample used for the isolation of *Campylobacter fetus* subsp. *venerealis* NW_ED23 was collected in the North-West Province, South Africa, specifically, in Mamusa District Municipality, cultured on tryptose blood agar base CM0233 (Oxoid, United Kingdom) mixed with 7% sheep blood, and supplemented with *Campylobacter* selective supplement (Skirrow) SR0068E (Oxoid). The plates were incubated at 37°C for 72 h in a 2.5-liter anaerobic jar (Oxoid) with a CampyGen CN0025A sachet (Oxoid) to create microaerophilic conditions (5). The genomic DNA (gDNA) was extracted from a single colony using a Zymo Research kit (Merck, South Africa). The PCR was conducted using MG3F and MG4R primers for the amplification of the 16S rRNA gene (6). The Basic Local Alignment Search Tool (BLAST) of the NCBI website was used to confirm identity and similarity (7), and the accession number obtained was [MT138648](https://doi.org/10.1128/MRA.00854-20). The gDNA from the pure colony based on Sanger identification confirmation was used for library preparation using a NEBNext Ultra II FS kit according to the manufacturer's instructions and was used for the whole-genome sequencing.

The extracted gDNA sample was fragmented using an enzymatic approach (NEBNext Ultra II FS kit). The resulting DNA fragment was sized (200 to 500 bp) using AMPure XP beads. The fragment was end repaired, and Illumina-specific adapter sequences were ligated to each fragment. The sample was indexed, and a second size selection step was performed. The sample was then quantified using a fluorometric method and diluted to a standard concentration (4 nM). It was then sequenced on an Illumina NextSeq platform using a NEBNext Ultra II FS 300-cycle kit according to the manufacturer's instructions to generate a total of 516,191,294 reads with 2 × 150-bp paired-end read length, and the coverage was 40×.

The sequence data were analyzed with the KBase platform with default parameters

Citation Tshipamba ME, Akinola SA, Ngoma L, Mwanza M. 2020. Genome sequence of *Campylobacter fetus* subsp. *venerealis* NW_ED23, isolated from bovine sheath wash. *Microbiol Resour Announc* 9:e00854-20. <https://doi.org/10.1128/MRA.00854-20>.

Editor J. Cameron Thrash, University of Southern California

Copyright © 2020 Tshipamba et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Mulunda Mwanza, 24059676@nwu.ac.za.

Received 22 July 2020

Accepted 28 August 2020

Published 1 October 2020

(8). The sequenced data were filtered for low-quality reads and adapter regions using Trimmomatic v0.36 (9) and FastQC v0.11.5. The genome assembly was performed using SPAdes v3.13.0 (10). Functional annotation of the whole-genome sequence was performed with the NCBI Prokaryotic Genome Annotation Pipeline (11) and Rapid Annotations using Subsystems Technology (RAST) v0.11 (12).

Campylobacter fetus subsp. *venerealis* NWU_ED23 has a total length of 1,880,748 bp assembled in 161 contigs with an average G+C content of 33.05%. The sequenced data generated contain 2,004 coding genes. The N_{50} value of the genome was 26,493 bp, while the L_{50} value was 22. The entire genome assembly contains 36 RNAs and 194 subsystems.

Data availability. The whole-genome shotgun project of *Campylobacter fetus* subsp. *venerealis* NWU_ED23 has been deposited in DDBJ/ENA/GenBank under the accession number [JACASH000000000](https://doi.org/10.1128/JCM.44.3.938-945.2006). The version published in this paper is [JACASH010000000](https://doi.org/10.1128/JCM.44.3.938-945.2006). The raw reads were also submitted to the NCBI SRA under accession number [SRX8607292](https://doi.org/10.1128/JCM.44.3.938-945.2006), BioSample number [SAMN15356083](https://doi.org/10.1128/JCM.44.3.938-945.2006), and BioProject number [PRJNA641553](https://doi.org/10.1128/JCM.44.3.938-945.2006).

ACKNOWLEDGMENT

This project was funded by the North-West University through the Animal Health Department.

REFERENCES

- McMillen L, Fordyce G, Doogan VJ, Lew AE. 2006. Comparison of culture and a novel 5' Taq nuclease assay for direct detection of *Campylobacter fetus* subsp. *venerealis* in clinical specimens from cattle. *J Clin Microbiol* 44:938–945. <https://doi.org/10.1128/JCM.44.3.938-945.2006>.
- Iraola G, Pérez R, Naya H, Paolicchi F, Harris D, Lawley TD, Rego N, Hernández M, Calleros L, Carretto L, Velilla A, Morsella C, Méndez A, Gioffre A. 2013. Complete genome sequence of *Campylobacter fetus* subsp. *venerealis* biovar *intermedius*, isolated from the prepuce of a bull. *Genome Announc* 1:e00526-13. <https://doi.org/10.1128/genomeA.00526-13>.
- van der Graaf-van Bloois L, Miller WG, Yee E, Bono JL, Rijnsburger M, Campero C, Wagenaar JA, Duim B. 2014. First closed genome sequence of *Campylobacter fetus* subsp. *venerealis* bv. *intermedius*. *Genome Announc* 2:e01246-13. <https://doi.org/10.1128/genomeA.01246-13>.
- Schmidt T, Venter EH, Picard J. 2010. Evaluation of PCR assays for the detection of *Campylobacter fetus* in bovine preputial scrapings and the identification of subspecies in South African field isolates. *J S Afr Vet Assoc* 81:87–92. <https://doi.org/10.4102/jsava.v81i2.111>.
- Acke E, McGill K, Golden O, Jones B, Fanning S, Whyte P. 2009. A comparison of different culture methods for the recovery of *Campylobacter* species from pets. *Zoonoses Public Health* 56:490–495. <https://doi.org/10.1111/j.1863-2378.2008.01205.x>.
- Schulze F, Bagon A, Müller W, Hotzel H. 2006. Identification of *Campylobacter fetus* subspecies by phenotypic differentiation and PCR. *J Clin Microbiol* 44:2019–2024. <https://doi.org/10.1128/JCM.02566-05>.
- Alayande KA, Aiyegoro OA, Nengwekhulu TM, Katata-Seru L, Ateba CN. 2020. Integrated genome-based probiotic relevance and safety evaluation of *Lactobacillus reuteri* PNW1. *PLoS One* 15:e0235873. <https://doi.org/10.1371/journal.pone.0235873>.
- Arkin AP, Cottingham RW, Henry CS, Harris NL, Stevens RL, Maslov S, Dehal P, Ware D, Perez F, Canon S, Sneddon MW, Henderson ML, Riehl WJ, Murphy-Olson D, Chan SY, Kamimura RT, Kumari S, Drake MM, Brettin TS, Glass EM, Chivian D, Gunter D, Weston DJ, Allen BH, Baumohl J, Best AA, Bowen B, Brenner SE, Bun CC, Chandonia J-M, Chia J-M, Colasanti R, Conrad N, Davis JJ, Davison BH, DeJongh M, Devoid S, Dietrich E, Dubchak I, Edirisinghe JN, Fang G, Faria JP, Frybarger PM, Gerlach W, Gerstein M, Greiner A, Gurtowski J, Haun HL, He F, Jain R, et al. 2018. KBase: the United States Department of Energy Systems Biology Knowledgebase. *Nat Biotechnol* 36:566–569. <https://doi.org/10.1038/nbt.4163>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Haft DH, DiCuccio M, Badretdin A, Brover V, Chetvernin V, O'Neill K, Li W, Chitsaz F, Derbyshire MK, Gonzales NR, Gwadz M, Lu F, Marchler GH, Song JS, Thanki N, Yamashita RA, Zheng C, Thibaud-Nissen F, Geer LY, Marchler-Bauer A, Pruitt KD. 2018. RefSeq: an update on prokaryotic genome annotation and curation. *Nucleic Acids Res* 46:D851–D60. <https://doi.org/10.1093/nar/gkx1068>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.