



Kurort Health Walking Preferentially Decreases Higher Blood Pressure and Improves Mood

Shinya Minatoguchi, MD, PhD; Taro Minagawa, MD, PhD; Kazuhiko Nishigaki, MD, PhD; Shinsuke Ojio, MD, PhD; Shinji Yasuda, MD, PhD; Kaori Osawa, BSc; Munenori Sasaki, BSc; Masashi Ogawa, BSc; Tatehiro Marumo, BSc; Shin Takano, BSc

Background: *Kurort* is a German term from the words *kur* (cure) and *ort* (area), and refers to improvements in patients' health in areas full of nature. We investigated the effect of *kurort* health walking in the 2 urban-style *kurort* health walking courses opened in Gifu City on systolic blood pressure (SBP), diastolic blood pressure (DBP), heart rate, and mood.

Methods and Results: The subjects were 454 people (136 males, 318 females; mean [±SD] age 61.7±9.9 years) taking part in *kurort* health walking for the first time. SBP, DBP, and heart rate were measured before and after *kurort* health walking. Mood was assessed using a 10-item checklist after *kurort* health walking. *Kurort* health walking significantly decreased SBP and DBP and increased heart rate. The decrease in SBP was significantly greater in the SBP ≥140 than <140 mmHg group, indicating that SBP before *Kurort* health walking was inversely correlated with the change in SBP. Similarly, the decrease in DBP was significantly greater in the DBP ≥90 than <90 mmHg group, indicating that DBP before *kurort* health walking was also inversely correlated with the change in DBP. All 10 items on the mood assessment were significantly improved after *kurort* health walking.

Conclusions: *Kurort* health walking preferentially decreases higher blood pressure and improves mood.

Key Words: Diastolic blood pressure; *Kurort* health walking; Mood; Systolic blood pressure

Hypertension and mental health status (e.g., depression) have been reported to be associated with cardiovascular events.^{1–3} Therefore, it is important to control blood pressure and to maintain a healthy mental state to reduce the risk of cardiovascular events. Although antihypertensive drugs and antidepressants are useful in preventing hypertension and depression, respectively, some lifestyle modifications may also be effective. *Kurort* is a German term from the words *kur* (cure) and *ort* (area) and refers to improvements in patients' health in areas rich in nature, such as scenic hills, forests, rivers, hot springs, and areas with a good climate.⁴ In Japan, Japanese-style *kurort*, based on the German *kurort*, has recently been developed and has become popular with the support of local governments and companies. *Kurort* health walking describes walking in a healthy area called *kurort*. On October 26, 2019, Gifu City opened 2 urban-style *kurort* health walking courses, the Mt. Kinka-Nagara River-Gifu Park course and the Mt. Dodogamine-Nagara River-Fureai Forest course, and has been encouraging citizens to participate in *kurort* health walking. These 2 courses are easily

accessible to citizens. Because these 2 *kurort* health walking courses are rich in nature, well designed, and well maintained, walking such courses may have a good effect on physical and mental health, which are related to cardiovascular disease.

The aim of the present study was to investigate the effects of *kurort* health walking in the *kurort* health walking courses in Gifu City on systolic blood pressure (SBP), diastolic blood pressure (DBP), heart rate, and mood.

Methods

Participants in the *kurort* health walking program obtained information regarding the places, dates, and times for *kurort* health walking through the public relations magazines of Gifu City. The subjects in this study were 454 first-time *kurort* health walking participants who used either of the 2 *kurort* walking courses in Gifu City between June 1, 2020 and May 30, 2021.

Because some people undertook *kurort* health walking several times, the total cumulative number of participants

Received July 27, 2021; revised manuscript received August 20, 2021; accepted September 2, 2021; J-STAGE Advance Publication released online October 6, 2021 Time for primary review: 15 days

Cardiology, Gifu Municipal Hospital, Gifu (S.M., K.N., S.O., S.Y., K.O.); Minagawa Clinic, Gifu (T. Minagawa); and Health Promotion Section, Gifu Municipal Office, Gifu (M.S., M.O., T. Marumo, S.T.), Japan

S.M. is a member of *Circulation Reports*' Editorial Team.


Mailing address: Shinya Minatoguchi, MD, PhD, Cardiology, Gifu Municipal Hospital, 7-1 Kashimacho, Gifu 500-8513, Japan.

E-mail: minatos@gifu-u.ac.jp

All rights are reserved to the Japanese Circulation Society. For permissions, please e-mail: cr@j-circ.or.jp

ISSN-2434-0790





in the program was 893; however, only those people walking the course for the first time were enrolled in the present study. All participants agreed to take part in the study and provided written informed consent before the study commenced. No one dropped out during the study.

Participants were asked to provide details regarding their age, sex, height, weight, presence of hypertension, dyslipidemia, and diabetes on a checklist before walking. SBP, DBP, and heart rate were measured using a wrist-type automatic sphygmomanometer, which is easy to use outdoors, before and after *kurort* health walking. After walking, participants were asked to complete a survey to evaluate changes in mood.


To ensure safety, participants with an SBP >180 mmHg and/or a DBP >110 mmHg are prohibited from *kurort* health walking, and *kurort* health walking is not performed in July and August because of the very hot climate in Gifu City in summer. Warming-up exercise before walking and cooling-down exercises after walking were performed under the direction of the health exercise instructors.

There are 2 courses for *kurort* health walking in Gifu City. The Mt. Kinka-Nagara River-Gifu Park course (**Figure 1**) is situated in the city center close to the Nagara River and extends to the foot of Mt. Kinka. This course is 2.3 km

long and has an elevation of 30m. The Mt. Dodogamine-Nagara River-Fureai Forest course (**Figure 2**) is situated in the northern area of the city and consists of a beautiful forest with many seasonal birds. This course is 3.2 km long and has an elevation of 80m.

To maintain target heart rate during walking, participants were asked to measure their heart rate at 5 points on the Mt. Kinka-Nagara River-Gifu Park course, and on 6 points on the Mt. Dodogamine-Nagara River-Fureai Forest course (**Figures 1,2**). Participants walked either course, accompanied by 2 health exercise instructors. The target heart rate during walking was defined as $(160 - \text{age})$ beats/min. If the heart rate increased beyond the target heart rate, participants were asked to slow the pace of walking so that heart rate was maintained under the target. Because the exercise level was maintained under the anaerobic threshold, it was considered safe for cardiac patients.⁵

This study was approved by the Ethics Committee of Gifu Municipal Hospital (Approval no. 634) and conformed with the principles outlined in the Declaration of Helsinki (*Br Med J* 1964; **ii**: 177). This study was registered with the University Hospital Medical Information Network (UMIN) Clinical Trials Registry (ID: UMIN000041617).



Measurement of SBP, DBP, and Heart Rate

SBP, DBP, and heart rate were measured before and after the completion of *kurort* health walking using a wrist-type sphygmomanometer. In analyses, SBP and DBP were each divided into 2 groups (SBP ≥ 140 and < 140 mmHg; DBP ≥ 90 and < 90 mmHg group) based on the definition of hypertension in the 2019 Japanese Society of Hypertension guidelines for the management of hypertension.⁶

Heart rate was only evaluated for the 403 participants who documented their heart rate both before and after *kurort* health walking in the checklist.

Measurement of Mood

Based on a previously reported method,⁷ a questionnaire was used in this study to determine changes in the following 10 mood items: feeling lively, feeling refreshed, a vivid feeling, feeling exhilarated, feeling relaxed, feeling calm, a fun feeling, feeling anxious, feeling irritated, and feeling tired after completing the *kurort* health walking. Participants rated each item as 'improved', 'no change', or 'worsened'. These ratings were scored as 1 for 'improved', 0 for 'no change', and -1 for 'worsened'. The effects of *kurort* health walking on each of the 10 mood items individually and the sum score for all 10 mood items

were assessed by averaging scores across all 454 participants.

Statistical Analysis

Data are presented as the mean \pm SD. The Kolmogorov-Smirnov test was used to determine the normality of data distribution. The significance of the differences in variables between groups was determined by paired and unpaired Student's t-tests. Correlation coefficients between 2 variables were obtained by linear regression analysis. Two-sided $P < 0.05$ was considered significant, and $P < 0.01$ and $P < 0.001$ were considered highly significant. All statistical analyses were performed using GraphPad Prism 7 (GraphPad Software, San Diego, CA, USA).

Results

Participants' Background

Of the 454 participants in this study, 136 were male and 318 were female. The age of participants ranged from 11 to 89 years (mean age 61.7 ± 9.9 years). Mean body mass index was 22.1 ± 2.8 kg/m². Some of participants (~30%) had past histories of hypertension ($n=84$), diabetes ($n=22$), and dyslipidemia ($n=31$).




Figure 3. (A) Systolic blood pressure (SBP), (B) diastolic blood pressure (DBP), and (C) heart rate before (pre) and after (post) *kurort* health walking. Significant decreases were seen in (A) SBP (from 130.5±18.4 to 124.4±16.8mmHg; n=454; P<0.0001, paired Student's t-test) and (B) DBP (from 81.6±11.3 to 80.6±11.8mmHg; n=454; P=0.0203, paired Student's t-test), and (C) heart rate increased significantly (from 73.1±12.1 to 75.8±12.3beats/min; n=415; P<0.0001, paired Student's t-test). Data are presented as the mean±SD.

SBP, DBP, and Heart Rate

Kurort health walking significantly decreased both SBP (from 130.5±18.4 to 124.4±16.8mmHg; P<0.0001; n=454; **Figure 3A**) and DBP (from 81.6±11.3 to 80.6±11.8mmHg; P=0.0203; n=454; **Figure 3B**). Heart rate increased significantly from 73.1±12.1 to 75.8±12.3beats/min (P<0.0001; n=415; **Figure 3C**).

In the SBP ≥140mmHg group, SBP decreased significantly from 152.2±10.2 to 139.1±14.6mmHg (P<0.0001; n=140; **Figure 4A**), whereas in the SBP <140mmHg group SBP decreased significantly from 120.9±11.9 to 117.8±13.2mmHg (P<0.0001; n=314; **Figure 4B**). The decrease in SBP was significantly greater in the SBP ≥140mmHg group (-12.9±13.9mmHg; n=140) than in the SBP <140mmHg group (-3.0±11.2mmHg; n=314; P<0.0001; **Figure 4C**). Thus, there was an inverse correlation between SBP before *kurort* health walking and the change in SBP after *kurort* health walking (P<0.0001; **Figure 4D**).

In the DBP ≥90mmHg group, DBP decreased significantly from 96.1±5.6 to 92.4±9.7mmHg (P<0.0001; n=112; **Figure 5A**). In the DBP <90mmHg group, DBP decreased from 76.8±8.2 to 76.7±9.7mmHg (n=342), but the difference did not reach statistical significance (P=0.823; **Figure 5B**). The decrease in DBP was significantly greater in the DBP ≥90mmHg group (-3.6±8.8mmHg; n=112) than in the SBP <90mmHg group (-0.1±8.9mmHg; n=342; P=0.0003; **Figure 5C**). Thus, there was an inverse correlation between


DBP before *kurort* health walking and the change in DBP after *kurort* health walking (P<0.0001; n=454; **Figure 5D**).

Mood

Kurort health walking significantly improved scores for each of the mood items, namely feeling lively (mean score after walking 0.65±0.47; P<0.0001 compared with before walking), feeling refreshed (0.87±0.33; P<0.0001), a vivid feeling (0.65±0.48; P<0.0001), feeling exhilarated (0.82±0.38; P<0.0001), feeling relaxed (0.77±0.41; P<0.0001), feeling calm (0.65±0.47; P<0.0001), a fun feeling (0.74±0.43; P<0.0001), feeling anxious (0.44±0.49; P<0.0001), feeling irritated (0.51±0.50; P<0.0001), and feeling tired (0.40±0.52; P<0.0001). In addition, *kurort* health walking significantly improved the sum score of the 10 mood items (mean score after walking 6.5±3.2 out of perfect score of 10; P<0.0001 compared with before walking; **Figure 6**).

Discussion

The major findings of the present study are that: (1) *kurort* health walking in the courses in Gifu City decreased both SBP and DBP; (2) the decrease in SBP was significantly greater in the SBP ≥140 than <140mmHg group, and the change in SBP was inversely correlated with SBP before walking; (3) the decrease in DBP was significantly greater in the DBP ≥90 than <90mmHg group, and the change in




DBP was inversely correlated with DBP before walking; and (4) *kurort* health walking improved mood.

Recently in Japan, the popularity of *kurort* health walking has gradually increased with the support of local governments and companies. The 2 *kurort* health walking courses in Gifu City are well designed and well maintained by the Gifu City administration; thus, walking through these courses may improve participants' physical condition and mood by relieving physical and mental tension by attenuating augmented sympathetic nerve activity. In the present study, *kurort* health walking decreased SBP and DBP (Figure 3A,B) and significantly increased heart rate (Figure 3C). On average, *kurort* health walking decreased SBP by 6.1 mmHg, decreased DBP by 1 mmHg, and increased heart rate by 2.7 beats/min. The increase in heart rate after exercise such as walking is a physiologically normal response, and an increase in heart rate of 2.7 beats/min is very small and can be regarded as safe (Figure 3C). Checking participants' heart rate during *kurort* health walking to ensure that it was maintained under the target rate (calculated as $[160 - \text{age}]$ beats/min) meant that *kurort* health walking was performed safely.

When participants were divided into 2 groups based on SBP before *kurort* health walking (i.e., SBP ≥ 140 and < 140 mmHg), the decrease in SBP after *kurort* health walk-

ing was significantly greater in the SBP ≥ 140 than < 140 group (-12.9 ± 13.9 vs. -3.0 ± 11.2 mmHg, respectively; $P < 0.0001$), indicating that the higher the SBP, the greater the decrease in SBP after *kurort* health walking (Figure 4D). When participants were divided into 2 groups based on DBP before *kurort* health walking (i.e., DBP ≥ 90 and < 90 mmHg), the decrease in DBP after *kurort* health walking was significantly greater in the DBP ≥ 90 than < 90 mmHg group (-3.6 ± 8.8 vs. -0.1 ± 8.9 mmHg, respectively; $P = 0.0003$), indicating that the higher the DBP, the greater the decrease in DBP (Figure 5D). Furthermore, as shown in Figures 4D and 5D, there is an inverse correlation between SBP and DBP before *kurort* health walking and the change in SBP and DBP after *kurort* health walking. *Kurort* health walking preferentially decreased higher than lower blood pressure and was safely performed even by hypertensive patients with SBP ≥ 140 mmHg and DBP ≥ 90 mmHg, decreasing their high blood pressure (Figures 4,5). Based on these findings, *kurort* health walking may be a useful strategy to reduce SBP and DBP in hypertensive patients. It has previously been reported that aerobic exercise decreases both SBP and DBP in both hypertensive patients and normotensive subjects.⁸⁻¹¹ The precise mechanisms by which aerobic exercise decreases blood pressure have not been fully clarified; however, some possible mech-



anisms have been suggested, including attenuation of plasma norepinephrine and epinephrine concentrations and enhancement of plasma prostaglandin E concentrations by aerobic exercise.⁸

Kurort health walking also improved all 10 mood items (Figure 6). Of the 10 items evaluated, scores for feeling refreshed (0.87 ± 0.33), feeling exhilarated (0.82 ± 0.38), feeling relaxed (0.77 ± 0.41), and a fun feeling (0.74 ± 0.43) were higher, suggesting that these 4 feelings were greatly improved by *kurort* health walking. Furthermore, the sum of all 10 mood items was 6.5 ± 3.2 (out of perfect score of 10); hence, more than 65% of participants showed an improvement in the sum of all 10 mood items following *kurort* health walking. These results suggest that *kurort* health walking may relieve mental tension and bring about an improvement in mood. Based on these results, *kurort* health walking may be a therapeutic strategy for improving mental health status. Previous studies found that aerobic exercise improved mental health status in patients with diabetes and depressive disorders,^{12,13} and others have reported that people who do not walk outside their home show more depressive symptoms or a greater likelihood of clinical depression.¹⁴ These reports may explain the beneficial effect of the *kurort* health walking in Gifu City, which consisted of walking outdoors, on mental health status. Because hypertension

and mental health status have been reported to be associated with cardiovascular events,¹⁻³ *kurort* health walking may be effective in reducing the risk of cardiovascular events because it decreases SBP and DBP and improves mood.

The advantages of *kurort* walking in Gifu City compared with normal walking may be that: (1) participants can walk in areas rich in nature, such as scenic hills, forests, rivers, and hot springs, and in a good climate,³ resulting in good mental feelings; (2) participants are accompanied by 2 health exercise instructors who provide appropriate advice during the walk; (3) blood pressure is measured before and after walking; and (4) heart rate during walking is maintained below the target rate, defined as $(160 - \text{age})$ beats/min, by checking heart rate at various points over the course to ensure safety (Figures 1,2).

Study Limitations

This study only showed the short-term effects of a single bout of exercise. The long-term effects of *kurort* health walking on blood pressure and mood remain to be investigated. Because the participants in the *kurort* health walking program are general citizens of Gifu City who happened to take part in the program, we only recorded whether they had hypertension, diabetes, and dyslipidemia, which are

Effect of kurort health walking on changes in mental feelings




Figure 6. Effects of *kurort* health walking on changes in mood. Improvements were seen in all 10 mood items evaluated on the checklist after *kurort* health walking. Changes in mood items after *kurort* health walking were rated as improved (score=1), no change (score=0), or worsened (score=-1). For each item, the mean±SD scores before and after *kurort* health walking are shown, with percentages for each category (1, improved; 0, no change; -1, worsened) across all 454 participants also presented. Changes in mood items from before to after *kurort* health walking were assessed by paired Student's t-test.

risk factors for coronary artery disease, on the checklist and did not obtain information regarding a history of cardiac and pulmonary diseases or drugs used. Furthermore, it is quite difficult to take blood samples from citizens who take part in *kurort* walking in the outdoors, although a survey based on blood data, such as blood glucose level, HbA1c, and cholesterol concentrations, may be important. However, because *kurort* health walking can be regarded as a type of outdoor cardiac rehabilitation, information regarding a history of cardiac disease should have been obtained.

Clinical Perspectives

In this study, SBP and DBP decreased and mood improved in subjects who participated in the *kurort* health walking program in Gifu City. Of all the participants, approximately 30% had a history of hypertension, diabetes, and dyslipidemia, which are risk factors for coronary artery disease.¹⁵ Even in these participants, *kurort* health walking was performed safely. The present study was performed between June 1, 2020 and May 30, 2021, during the coro-

navirus pandemic, suggesting that because *kurort* health walking is mostly an outdoor activity, it can be safely performed in the coronavirus era.

In addition to effectively decreasing blood pressure and improving mood, *kurort* health walking could be an alternative tool for cardiac rehabilitation. Because cardiac rehabilitation has recently been reported to improve the prognosis of cardiac diseases such as acute myocardial infarction and heart failure,^{16,17} cardiac rehabilitation is recommended for these patients in the guidelines from the Japanese Circulation Society¹⁸ and as part of the standard cardiac rehabilitation program for heart failure from the Japanese Association of Cardiac Rehabilitation Standard Cardiac Rehabilitation Program Planning Committee.¹⁹ However, under the health insurance system in Japan, cardiac rehabilitation is limited to 20 weeks after the onset of cardiac diseases. Thus, *kurort* health walking may be a suitable option for maintaining the health of cardiac patients after they have completed 20 weeks of cardiac rehabilitation. Further investigations are warranted.

Acknowledgment

The authors are grateful for the support of the Health Promotion Section of Gifu Municipal Office, Gifu City, Japan.

Sources of Funding

This study was funded from the Sugiura Memorial Foundation, Obu City, Aichi, Japan.

Author Contributions

S.M.: organization and design of the study, data interpretation and analysis, manuscript writing, financial support, and final approval of manuscript. T.M., K.N., S.O., S.Y.: data collection and data interpretation. K.O.: data analysis. S.T., M.S., M.O., T.M.: organization of *kurort* health walking and data collection.

Disclosures

S.M. is a member of *Circulation Reports*' Editorial Team. The remaining authors have no conflicts of interest to disclose.

IRB Information

This study was approved by the Ethics Committee of Gifu Municipal Hospital (Approval no. 634).

Data Availability

The deidentified participant data will not be shared.

References

1. Stamler J, Stamler R, Neaton JD. Blood pressure, systolic and diastolic, and cardiovascular risks: US population data. *Arch Intern Med* 1993; **153**: 598–615.
2. Nielsen TJ, Vestergaard M, Christensen B, Christensen KS, Larsen KK. Mental health status and risk of new cardiovascular events or death in patients with myocardial infarction: A population-based cohort study. *BMJ Open* 2013; **3**: e003045.
3. Hata J, Kiyohara Y. Epidemiology of stroke and coronary artery disease in Asia. *Circ J* 2013; **77**: 1923–1932.
4. Guenther K. Exercises in therapy: Neurological gymnastics between Kurort and hospital medicine, 1880–1945. *Bull Hist Med* 2014; **88**: 102–131.
5. Wasserman K, Whipp BJ, Koyl SN, Beaver WL. Anaerobic threshold and respiratory gas exchange during exercise. *J Appl Physiol* 1973; **35**: 236–243.
6. Umemura S, Arima H, Arima S, Asayama K, Dohi Y, Hirooka Y, et al. The Japanese Society of Hypertension guidelines for the management of hypertension (JSH 2019). *Hypertens Res* 2019; **42**: 1235–1481.
7. Ueda H, Machida K, Kawamura N, Koseki N. A study on the process of mood alteration through forest walking. *Journal of The Japanese Institute of Landscape Architecture* 2013; **76**: 533–538.
8. Kiyonaga A, Arakawa K, Tanaka H, Shindo M. Blood pressure and hormonal responses to aerobic exercise. *Hypertension* 1985; **7**: 125–131.
9. Ghadieh AS, Saab B. Evidence for exercise training in the management of hypertension in adults. *Can Fam Physician* 2015; **61**: 233–239.
10. Carpio-Rivera E, Moncada-Jiménez J, Salazar-Rojas W, Solera-Herrera A. Acute effects of exercise on blood pressure: A meta-analytic investigation. *Arq Bras Cardiol* 2016; **106**: 422–433.
11. Whelton SP, Chin A, Xin X, He J. Effect of aerobic exercise on blood pressure: A meta-analysis of randomized, controlled trials. *Ann Intern Med* 2002; **136**: 493–503.
12. Gilani SRM, Feizabad AK. The effects of aerobic exercise training on mental health and self-esteem of type 2 diabetes mellitus patients. *Health Psychol Res* 2019; **7**: 6576.
13. Morres ID, Hatzigeorgiadis A, Stathi A, Comoutos N, Arpin-Cribbie C, Krommidas C, et al. Aerobic exercise for adult patients with major depressive disorder in mental health services: A systematic review and meta-analysis. *Depress Anxiety* 2019; **36**: 39–53.
14. Julien D, Gauvin L, Richard L, Kestens Y, Payette H. The role of social participation and walking in depression among older adults: Results from the VoisiNuAge study. *Can J Aging* 2013; **32**: 1–12.
15. Kannel WB. Some lessons in cardiovascular epidemiology from Framingham. *Am J Cardiol* 1976; **37**: 269–282.
16. Witt BJ, Jacobsen SJ, Weston SA, Killian JM, Meverden RA, Allison TG, et al. Cardiac rehabilitation after myocardial infarction in the community. *J Am Coll Cardiol* 2004; **44**: 988–996.
17. O'Connor CM, Whellan DJ, Lee KL, Keteyian SJ, Cooper LS, Ellis SJ, et al. Efficacy and safety of exercise training in patients with chronic heart failure: HF-ACTION randomized controlled trial. *JAMA* 2009; **301**: 1439–1450.
18. JCS Joint Working Group. Guidelines for rehabilitation in patients with cardiovascular disease (JCS 2012): Digest version. *Circ J* 2014; **78**: 2022–2093.
19. Izawa H, Yoshida T, Ikegame T, Izawa K, Ito Y, Okamura H, et al. Standard cardiac rehabilitation program for heart failure. *Circ J* 2019; **83**: 2394–2398.