

CORRECTION

Open Access



# Correction to: In vitro assessment of nutraceutical compounds and novel nutraceutical formulations in a liver-steatosis-based model

Antonietta Stellavato<sup>1</sup>, Anna Virginia Adriana Pirozzi<sup>1</sup>, Francesca de Novellis<sup>1</sup>, Ilaria Scognamiglio<sup>1</sup>, Valentina Vassallo<sup>1</sup>, Andrea Maria Giori<sup>2</sup>, Mario De Rosa<sup>1</sup> and Chiara Schiraldi<sup>1\*</sup>

**Correction to: *Lipids Health Dis* 17, 24 (2018)**  
<https://doi.org/10.1186/s12944-018-0663-2>

Following the publication of the original article [1], the authors noticed an inaccuracy related to western blotting actin bands in Figs. 5 and 6.

In this respect, the authors apologize for uploading the western blotting (actins bands) that could be misleading. The actins obtained for the filters incubated for SOD-2 antibody were incorrectly duplicated for filters incubated with PPAR- $\gamma$ . Presented here are the correct Figs. 5 and 6 with the correct actins references.

---

The original article can be found online at <https://doi.org/10.1186/s12944-018-0663-2>.


\* Correspondence: [chiara.schiraldi@unicampania.it](mailto:chiara.schiraldi@unicampania.it)


<sup>1</sup>Department of Experimental Medicine, Section of Biotechnology, Medical Histology and Molecular Biology, Università della Campania "Luigi Vanvitelli", Naples, Italy

Full list of author information is available at the end of the article



© The Author(s). 2022 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.





**Fig. 6** PPARα and PPARγ protein expression levels on HepG2 cells determined by Western blotting. Actin is used to normalize the results

#### Author details

<sup>1</sup>Department of Experimental Medicine, Section of Biotechnology, Medical Histology and Molecular Biology, Università della Campania "Luigi Vanvitelli", Naples, Italy. <sup>2</sup>R&D - IBSA Farmaceutici Italia, Lodi, Italy.

Published online: 06 April 2022

#### Reference

1. Stellavato A, Pirozzi AVA, de Novellis F, et al. In vitro assessment of nutraceutical compounds and novel nutraceutical formulations in a liver-steatosis-based model. *Lipids Health Dis.* 2018;17:24. <https://doi.org/10.1186/s12944-018-0663-2>.