

Comments on “Double-Blind Comparison of Ultrasonic and Conventional Osteotomy in Terms of Early Postoperative Edema and Ecchymosis”

Can Alper Çağıcı, MD

Aesthetic Surgery Journal
2018, Vol 38(2) NP29–NP30
© 2017 The American Society for
Aesthetic Plastic Surgery, Inc.
This is an Open Access article
distributed under the terms of the
Creative Commons Attribution
Non-Commercial License ([http://
creativecommons.org/licenses/
by-nc/4.0/](http://creativecommons.org/licenses/by-nc/4.0/)), which permits non-
commercial re-use, distribution,
and reproduction in any medium,
provided the original work is
properly cited. For commercial
reuse, please contact journals.
permissions@oup.com.
DOI: 10.1093/asj/sjw232
www.aestheticsurgeryjournal.com

OXFORD
UNIVERSITY PRESS

Editorial Decision date: November 1, 2016; online publish-ahead-of-print December 21, 2017.

I read the manuscript by Ilhan et al¹ with great interest. The authors performed a randomized study comparing two types of osteotomy in rhinoplasty: ultrasonic and conventional osteotomy. They found that ultrasonic osteotomy results in less edema and ecchymosis. Their patients were blind to the study and not informed about which osteotomy technique was applied to them. The same surgeon performed both types of surgery and was not blinded. The postoperative evaluation of the surgical results was performed by blinded examiners. Although the authors called their study a “double-blinded” comparison, I do not agree with them.

I agree that blinding of the examiners was important, because their attitude about the technique might affect their assessment. I did not understand how a contribution was expected from patient blindness in this study. Although they did not discuss the effect of the patient blindness on the study, I hope that such an effect is minimal. On the other hand, the blinding of surgeon, evaluator, and biostatistician might be important in this clinical trial to overcome transfer of their attitudes toward or against an intervention during the study. In a “double-blinded” study like this the readers probably expect that the surgeon and evaluators are blinded, not the patients.

The authors considered their study double-blinded because the patients and evaluators were blinded. I prefer that the surgeon instead of the patient is blinded in a “double-blinded” study because unblinded clinicians may transfer their attitude to the treatment.² However, I am aware that it is impossible to design a study with surgeon blindness. Two options are available in such cases: the lack of the surgeon’s blindness and the resultant potential bias may be discussed in the paper, or the study may be performed

using an expertise-based design. In an expertise-based design, the patients are randomized to paired surgeons with expertise in only one procedure.³ The expertise-based design is advised to overcome transfer of the surgeon’s attitude toward or against an intervention during the surgery. I agree that it is difficult to randomize patients who initially presented to a particular aesthetic surgeon. In this situation, a double-blind study is not required. If a double-blind study is still desired, it is possible to carry out but the drawbacks of the study must be addressed in the Discussion section.

Randomization is important because it eliminates selection bias. All persons associated with the study (participants, surgeons, data collectors, outcome adjudicators, and data analysts) should be blinded as much as possible.² The term “double-blinded” should be avoided in surgical trials.³ Its meaning changes according to the individual using it. It is more meaningful to define the persons who are blinded to the study in the Methods section instead of calling the study “double-blinded.”³

Disclosures

The author declared no potential conflicts of interest with respect to the research, authorship, and publication of this article.

Dr Çağıcı is an Associate Professor, ENT Department, Baskent University Adana Seyhan Hospital, Seyhan/Adana, Turkey.

Corresponding Author:

Dr Can Alper Çağıcı, ENT Department, Baskent University Adana Seyhan Hospital, Gazipaşa Mahallesi Baraj Caddesi No. 7, Seyhan/Adana 01150, Turkey.

E-mail: ccagici@hotmail.com

Funding

The author received no financial support for the research, authorship, and publication of this article.

REFERENCES

1. Ilhan AE, Cengiz B, Caypinar Eser B. Double-blind comparison of ultrasonic and conventional osteotomy in terms of early postoperative edema and ecchymosis. *Aesthet Surg J*. 2016;36(4):390-401.
2. Karanicolas PJ, Farrokhyar F, Bhandari M. Practical tips for surgical research: blinding: who, what, when, why, how? *Can J Surg*. 2010;53(5):345-348.
3. Farrokhyar F, Karanicolas PJ, Thoma A, et al. Randomized controlled trials of surgical interventions. *Ann Surg*. 2010;251(3):409-416.

Putting Patient Safety First Benefits Everyone

PERFORMANCE • EFFICIENCY • PATIENT SATISFACTION • REFERRALS • REVENUE

The Aesthetic Society's Patient Safety Committee would like to remind you that an increased focus on patient safety leads to enhanced surgical performance and efficiency. When a surgeon puts safety first, patients are better satisfied, resulting in more referrals, which ultimately impacts your bottom line.

Culture
of
Safety

"Primum non nocere"
First do no harm



THE AMERICAN SOCIETY FOR
AESTHETIC PLASTIC SURGERY, INC.

800-364-2147 • 562-799-2356
www.surgery.org