



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Emergency department screening for multisystem inflammatory syndrome (MIS-C) in children



1. Introduction

Multisystem inflammatory syndrome in children (MIS-C) was defined by the Centers for Disease Control and Prevention (CDC) in May 2020 [1]. An approach to the identification of this uncommon yet life-threatening illness among well-appearing febrile children presenting for medical care is urgently needed as the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection continues to affect millions worldwide [2,3]. Thus, the aims of this report are to: (1) describe our institutional approach to screening for MIS-C; (2) report the test characteristics of our approach in excluding MIS-C; and (3) provide recommendations for screening for MIS-C among well appearing children with fever.

2. Methods

Screening for MIS-C at an urban, tertiary care pediatric emergency department (PED) located in a geographic area with a high burden of SARS-CoV-2 infections [2] was implemented on May 7, 2020. For well appearing children presenting with fever ≥ 48 h with rash, gastrointestinal symptoms, or clinical features of Kawasaki Disease [4], the initial recommended screening laboratory studies include: complete blood count with differential (CBC), comprehensive metabolic panel (CMP), C-reactive protein (CRP), ferritin, d-dimer, and high-sensitivity troponin I (troponin). We retrospectively reviewed the PED logs for May 7 to May 31, 2020 to identify patients < 21 years of age with a relevant chief complaint (e.g. fever, diarrhea, rash) who had a ferritin, d-dimer or troponin performed (as CBC, CMP, and CRP are not specific to our screening algorithm). Demographic and clinical characteristics, including final diagnoses, and laboratory values were abstracted by clinicians using a standardized data collection tool and entered into Research Electronic Data Capture (REDCap) [5].

For analysis, patients were dichotomized as MIS-C or not (see Table 1). Means/standard deviations, medians/interquartile ranges, or frequencies/95% confidence intervals (CI) were calculated for patient characteristics. Univariate logistic regression (LR) was used to assess the association of clinical data with the primary outcome (non-MIS-C); those with $p < 0.1$ were assessed for independent association using multivariable LR. Co-linearity of predictors was assessed using variance inflation factor. As our goal was to identify characteristics associated with a low probability of MIS-C, composite predictor variables were created, with candidates selected from the multivariable analysis or “by meaning”. Continuous predictors (except age) were dichotomized with a cut-point determined by

Youden's J statistic. Meeting institutional or CDC clinical criteria for screening was also included in the composite variable. Final model selection was based on comparison of Akaike's Information Criteria, area under the receiver operating characteristic curve (AUC), and CI width (narrower considered more clinically informative). Two-tailed significance level was set at 0.05. Statistical analysis was performed using SAS v9.4 (Cary, NC).

3. Results

Among 2536 patients presenting during the study period, 196 (8%) were included in the analysis and 12 (0.5%) were categorized as MIS-C. Demographics and screening laboratory results are reported in Table 1. On multivariable analysis, troponin, d-dimer, ferritin, CRP, and age had significant inverse associations with the primary outcome. Composite laboratory value predictor variables performed differently depending on whether institutional or CDC clinical screening criteria were used. Final models for each, along with independent contribution of age, are presented in Table 2.

4. Conclusions

Although limited by sample size, this timely report is the first to provide clinicians with a preliminary approach to screening febrile children for MIS-C. While larger validation studies are needed, children meeting the CDC clinical criteria [1] should have a CRP, troponin, and ferritin performed. Patients with results below the identified cut-points are at lower risk for MIS-C and can be discharged home with close primary care follow-up. Given its association with greater likelihood of disease, increased age should be incorporated into clinical decision making among children being screened for MIS-C.

Funding support

None.

Contributors statement

DeLaroche - conceptualization, data curation, project administration, software supervision, validation, writing - original draft, writing - review and editing. Stankovic & Ruffing: conceptualization, data curation, writing - review and editing. Ehrman - formal analysis, writing - original draft, writing - review and editing. Noble, Arora, Maksimowski - data curation, writing - review and editing.

Declaration of Competing Interest

The authors have no conflicts of interest relevant to this article to disclose.

Table 1
Demographic and clinical characteristics of the study cohort

Characteristic	All Cases <i>n</i> = 196	Non MIS-C <i>n</i> = 184	MIS-C* <i>n</i> = 12
Demographics, <i>n</i> (%)			
Age, median (IQR), years	2.0 (6.0)	1.5 (6.0)	8.5 (6.5)
Female gender	103 (53)	94 (51)	9 (75)
Race			
Caucasian	48 (25)	46 (25)	2 (17)
African-American	95 (49)	88 (48)	7 (58)
Other	53 (27)	50 (27)	3 (25)
Index visit	167 (85)	159 (86)	8 (67)
PED Disposition			
Admission	129 (66)	117 (64)	12 (100)
Discharge	67 (34)	67 (36)	0 (0)
PICU admission during hospitalization	20 (16)	10 (9)	10 (83)
Presenting Clinical Features by System, <i>n</i> (%)			
Fever [§]	182 (93)	171 (93)	11 (92)
Respiratory	78 (40)	71 (39)	7 (58)
Gastrointestinal	163 (83)	152 (83)	11 (92)
Neurological	23 (12)	17 (9)	6 (50)
Cardiac	12 (6)	11 (6)	1 (8)
Musculoskeletal	13 (7)	12 (7)	1 (8)
Mucocutaneous	78 (40)	71 (39)	7 (58)
Screening Laboratory Study Results, median (IQR)			
White blood cell count (reference range: 4.1–10.1 × 10 ⁹ /L)	8.50 (6.50) [195]	8.40 (6.40) [183]	10.95 (8.70) [12]
Neutrophil count (reference range: 1.58–7.13 × 10 ⁹ /L)	4.20 (5.50) [190]	4.00 (5.20) [178]	9.00 (7.20) [12]
Lymphocyte count (reference range: 0.8–5.0 × 10 ⁹ /L)	2.50 (2.90) [190]	2.70 (3.10) [178]	1.05 (0.70) [12]
Hemoglobin (reference range: 11.7–15.9 g/dL)	11.90 (1.40) [195]	11.90 (1.30) [183]	11.35 (1.90) [12]
Platelet count (reference range: 130–450 × 10 ⁹ /L)	260 (119) [194]	262 (113) [182]	202 (216) [12]
CRP (reference range: <5.0 mg/L)	14.65 (63.90) [194]	11.55 (52.90) [182]	139.65 (76.55) [12]
Ferritin (reference range: 11.0–306.8 µg/L)	73.40 (93.50) [191]	70.80 (83.80) [179]	318.15 (418.25) [12]
Albumin (reference range: 3.8–5.1 g/dL)	4.40 (0.50) [192]	4.40 (0.40) [180]	3.90 (0.75) [12]
ALT (reference range: 7–52 U/L)	17.00 (12.00) [192]	17.00 (12.00) [180]	17.50 (45.00) [12]
Troponin I, high sensitivity (reference range: 3–17 ng/L)	7.00 (7.00) [194]	7.00 (6.00) [182]	36.50 (97.00) [12]
D-dimer (reference range: <0.50 mg/L FEU)	0.71 (0.87) [189]	0.69 (0.79) [177]	1.95 (1.60) [12]

Abbreviations: ALT, alanine aminotransferase; CRP, C-reactive protein; PED, pediatric emergency department; PICU, pediatric intensive care unit; FEU, fibrinogen equivalent units; IQR, interquartile range; MIS-C, multisystem inflammatory syndrome in children; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.

SI conversion factors: To convert hemoglobin to g/L multiply by 10; to convert albumin to g/L multiply by 10; to convert ALT to µkat/L multiply by 0.0167.

Two patients with MIS-C met CDC criteria except for laboratory confirmed SARS-CoV-2 infection or an epidemiological link to SARS-CoV-2. One patient with MIS-C met CDC criteria except for fever, which was absent at home but had reported chills at home and a documented fever in the PED.

* Patients included in the MIS-C group had a presumptive diagnosis of MIS-C early in their hospital course made by a specialist in pediatric infectious disease and all were treated with intravenous immunoglobulin. Final discharge diagnoses were MIS-C (11) and hemophagocytic lymphohistiocytosis (1). Eight patients with MIS-C met CDC criteria.

§ Fever included subjective or documented fever prior to PED arrival.

Table 2
OR, AUC, sensitivity, and specificity of screening algorithms for the exclusion of MIS-C among pediatric patients presenting to the PED

Algorithm	OR	95% CI	AUC (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)	Age [#] OR (95% CI)
A. CDC clinical criteria* met plus composite laboratory criteria 1 [§]	17.44	3.67–82.75	0.81 (0.69–0.92)	0.78 (0.72–0.84)	0.83 (0.62–1.00)	0.86 (0.76–0.97)
B. Institutional clinical criteria [∇] met plus composite laboratory criteria 2 [^]	19.65	5.35–72.08	0.78 (0.65–0.93)	0.91 (0.87–0.95)	0.67 (0.40–0.93)	0.87 (0.77–0.99)
C. CDC clinical criteria* met plus composite laboratory criteria 2 [^]	60.71	12.10–304.66	0.88 (0.77–0.99)	0.92 (0.89–0.96)	0.83 (0.62–1.00)	0.91 (0.79–1.04)
D. Institutional clinical criteria [∇] met plus composite laboratory criteria 1 [§]	5.83	1.68–20.25	0.71 (0.56–0.85)	0.74 (0.68–0.81)	0.67 (0.40–0.93)	0.84 (0.75–0.95)

Abbreviations: AUC, area under the curve; CDC, Centers for Disease Control and Prevention; CI, confidence interval; CRP, c-reactive protein; PED, pediatric emergency department; MIS-C, multisystem inflammatory syndrome in children; OR, odds ratio.

* CDC clinical criteria defined as fever ≥24 h plus ≥2 systems involved at the time of PED presentation.

§ Lab criteria 1: CRP & high-sensitivity Troponin I & ferritin all above cut-points, defined as CRP ≥ 87.5 mg/L; troponin I ≥ 17 ng/L; ferritin ≥121.6 ng/mL.

∇ Institutional clinical criteria defined as fever ≥48 h plus one of the following: gastrointestinal features (abdominal pain, vomiting, diarrhea), mucocutaneous features (rash, conjunctivitis, erythema of oral/pharyngeal mucosa), or lymphadenopathy.

^ Lab criteria 2: [CRP & d-dimer & ferritin] OR troponin I above the optimal cut-point, defined as CRP ≥ 87.5 mg/L; troponin I, high sensitivity ≥17 ng/L; ferritin ≥121.6 ng/mL; d-dimer ≥1.07 mg/L fibrinogen equivalent units (FEU).

Effect per 1 year increase in age when added to corresponding clinical/laboratory model.

References

- [1] Centers for Disease Control and Prevention Emergency Preparedness and Response: Multisystem Inflammatory Syndrome in Children (MIS-C) Associated with Coronavirus Disease 2019 (COVID-19). Published May 14, 2020. Accessed May 16, 2020 <https://emergency.cdc.gov/han/2020/han00432.asp>; 2020.
- [2] Feldstein LR, Rose EB, Horwitz SM, et al. Multisystem inflammatory syndrome in U.S. children and adolescents. *N Engl J Med*. 2020. <https://doi.org/10.1056/NEJMoa2021680> [Online ahead of print].
- [3] Center for Systems Science and Engineering at Johns Hopkins University. COVID-19 Dashboard. Accessed July 14, 2020. Available at: <https://coronavirus.jhu.edu/map.html>; 2020.
- [4] McCrindle BW, Rowley AH, Newburger JW, et al. American Heart Association Rheumatic Fever, Endocarditis, and Kawasaki Disease Committee of the Council on Cardiovascular Disease in the Young; Council on Cardiovascular and Stroke Nursing; Council on Cardiovascular Surgery and Anesthesia; and Council on Epidemiology and Prevention. Diagnosis, treatment, and long-term management of Kawasaki disease: a scientific statement for health professionals for the American Heart Association. *Circulation*. 2017;135(17):e927–99. <https://doi.org/10.1161/CIR.0000000000000484>.
- [5] Harris PA, Taylor R, Thielke R, et al. Research electronic data capture (REDCap) – a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42:377–81.

Amy M. DeLaroche MBBS

Division of Emergency Medicine, Department of Pediatrics, Children's Hospital of Michigan, Detroit, MI, United States of America

Corresponding author at: Division of Emergency Medicine, Department of Pediatrics, Children's Hospital of Michigan, 3901 Beaubien Street, Detroit, MI 48201, United States of America.

E-mail address: adelaroc@dmc.org

Curt Stankovic MD, MBA

Division of Emergency Medicine, Department of Pediatrics, Children's Hospital of Michigan, Detroit, MI, United States of America

Robert R. Ehrman MD, MS

Department of Emergency Medicine, Wayne State University School of Medicine, Detroit, MI, United States of America

Department of Emergency Medicine, Detroit Medical Center/Sinai-Grace Hospital, Detroit, MI, United States of America

Jennifer Noble MD

Division of Emergency Medicine, Department of Pediatrics, Children's Hospital of Michigan, Detroit, MI, United States of America

Rajan Arora MD

Division of Emergency Medicine, Department of Pediatrics, Children's Hospital of Michigan, Detroit, MI, United States of America

Karolina Maksimowski MD

Division of Emergency Medicine, Department of Pediatrics, Children's Hospital of Michigan, Detroit, MI, United States of America

Ronald P. Ruffing MD, MPH

Division of Emergency Medicine, Department of Pediatrics, Children's Hospital of Michigan, Detroit, MI, United States of America

22 September 2020