

underlying neuro-behavioural effects in this model. Although the results of the morphometric analysis suggest a causal link between behavioural changes with altered synaptic plasticity, the mechanisms and involved signalling remain to be defined.

Disclosure: No significant relationships.

Keywords: animal models of autism; dendritic spines; tuberous sclerosis complex; enriched environment

EPP0201

Burnout or anxiety?! A medical trial regarding resident doctors in a first line University Hospital during COVID 19 pandemic.

C. Sapaniuc^{1,*}, G. Lacatusu¹, I. Mihai², M. Rascanu³, D. Bran³ and D. Manciu^{1,2}

¹Hospital of Infectious Diseases "Sf. Parascheva", Infectious Diseases, Iasi, Romania; ²University of Medicine and Pharmacy "Gr. T. Popa", Infectious Diseases, Iasi, Romania and ³Hospital of Infectious Diseases "Sf. Parascheva", Psychology, Iasi, Romania

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.507

Introduction: Coronavirus Disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has been reported as a worldwide emergency. Due to the extensiveness of spread and death, it has been declared as a pandemic.

Objectives: To highlight how COVID-19 pandemic psycho-emotional affects the medical staff of a frontline University Hospital in the "battle" with new coronavirus.

Methods: We employed a cross-sectional survey of 71 resident doctors from a frontline Hospital after a one-year pandemic and analyzed the prevalence and associated factors with work-related psychological distress among our study group.

Results: Out of the hospital resident doctors, 71 participated and completed the questionnaire, offering an overall response rate of 100%. The majority of participants were women (86% - 61). The average age was 29 years. Most respondents were unmarried. A total of 67% of participants were non-smokers, 5% stated that they occasionally consumed alcohol, none of them used drugs. As a result of the qualitative and quantitative analysis of the data, aspects related to anxiety (21.12% - 15), exhaustion (15.49% - 11), and depression (11.26% - 8) are highlighted. In our study, no people were identified who would reach extreme exhaustion in the work process, due to good resilience and due to a well thought out program of work and rest during the pandemic.

Conclusions: The psychological pressure at work, as well as the one felt after limiting and restricting mobility for shorter or longer periods, had an impact on the psycho-emotional state of health care workers, requiring further psychological reassessments and psychological support.

Disclosure: No significant relationships.

Keywords: burnout; anxiety; pandemic; covid 19; resident doctor; hospital

EPP0202

Psychiatric care in Croatia during COVID-19 lockdown and earthquake: significant decrease in admissions to University Psychiatric Hospital Vrapce in Zagreb

D. Polšek^{1,*}, A. Botica², T. Sabo¹, M. Baković³, V. Marinović¹ and G. Arbanas^{4,5}

¹University Psychiatric Hospital Vrapče, Department For Psychotic Disorders, Zagreb, Croatia; ²University Hospital Centre Split, Dpt. Of Psychiatry, Split, Croatia; ³Psychiatric Hospital Ugljan, Dpt. Of Psychiatry, Ugljan, Croatia; ⁴University of Rijeka, Medical Department, Rijeka, Croatia and ⁵University Psychiatric Hospital Vrapče, Department For Forensic Psychiatry, Zagreb, Croatia

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.508

Introduction: The COVID-19 pandemic has had an enormous impact on both physical and mental health of people around the world.

Objectives: The aim of the study was to evaluate the number and characteristics of people seeking emergency psychiatric help during combined psychosocial stressful events in March 2020.

Methods: Data for 3927 patients seeking emergency psychiatric help were collected and analyzed for the months preceding, during and after lockdown due to COVID-19 pandemic and concomitant earthquake that took place on 22nd March 2020 in Zagreb, and compared with the same months of 2019.

Results: A significant decrease in both the number of visits and admissions to the hospital was found for the month of lockdown. There was a significant decrease in the number of out-patients visits and day hospital admissions. Compared with other months, more women and younger patients sought help. There was a significant rise in the number of patients presenting with suicidal thoughts, as well as a larger percentage of involuntary admissions.

Conclusions: Overall less people sought psychiatric help in the face of an unpredictable acute threat, which was interpreted in the light of prioritizing fear of infection over mental health issues. Alternatively, it is possible that people threatened with immediate danger mobilize short-term compensatory psychological resources which help deal with or put off mental illness. This research was conducted as part of the project of the Croatian Science Foundation CORONA-04-2086 Life in the time of COVID-19-social implications on the security and well-being of vulnerable groups in the European context.

Disclosure: No significant relationships.

Keywords: COVID-19 pandemic; Emergency psychiatric care; Hospital admission; lockdown

EPP0203

Mechanisms and Effectiveness of Online CB-ART Interventions in Reducing Covid-19-related Distress

D. Segal-Engelchin*, V. Daichman, E. Huss and O. Sarid Ben-Gurion University of the Negev, Social Work, Beer-Sheva, Israel

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.509

Introduction: The combination of cognitive behavioral interventions and art therapy provides a unique tool for image transformation as a strategy for managing distress in extremely stressful