

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2,2'-Dihydroxy-*N,N'*-(ethane-1,2-diyl)-dibenzamide

Daniel Pereira da Costa,^a Sabrina Madruga Nobre,^{a*}
 Bruna Gonçalves Lisboa,^a Juliano Rosa de Menezes
 Vicenti^a and Davi Fernando Back^b

^aEscola de Química e Alimentos, Universidade Federal do Rio Grande, Av. Itália, km 08, Campus Carreiros, 96203-900 Rio Grande-RS, Brazil, and ^bDepartamento de Química, Universidade Federal de Santa Maria, Av. Roraima Campus, 97105-900 Santa Maria-RS, Brazil

Correspondence e-mail: sabrinanobre@furg.br


Received 20 December 2012; accepted 27 December 2012

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.049; wR factor = 0.143; data-to-parameter ratio = 15.4.

The asymmetric unit of the title compound, $\text{C}_{16}\text{H}_{16}\text{N}_2\text{O}_4$, contains one half-molecule, the whole molecule being generated by an inversion center located at the mid-point of the C—C bond of the central ethane group. An intramolecular O—H...O hydrogen bond forms an $S(6)$ ring motif. In the crystal, molecules are connected *via* N—H...O hydrogen bonds, generating infinite chains along $[1\bar{1}0]$.

Related literature

For the synthesis of bisamides, see: Apurba *et al.* (2002); Fry *et al.* (2010). For similar bisamide crystal structures, see: Booyesen *et al.* (2009). For applications of bisamides as catalysts, see: Maurya *et al.* (2003); Liu *et al.* (2011).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{16}\text{N}_2\text{O}_4$
 $M_r = 300.31$
 Triclinic, $P\bar{1}$
 $a = 4.6311$ (3) Å
 $b = 5.0435$ (3) Å

$c = 15.2957$ (9) Å
 $\alpha = 89.091$ (4)°
 $\beta = 83.315$ (4)°
 $\gamma = 85.956$ (4)°
 $V = 353.94$ (4) Å³

$Z = 1$
 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹

$T = 296$ K
 $0.34 \times 0.24 \times 0.11$ mm

Data collection

Bruker APEXII CCD
 diffractometer
 Absorption correction: gaussian
 (SADABS; Bruker, 2009)
 $T_{\min} = 0.966$, $T_{\max} = 0.989$

9142 measured reflections
 1535 independent reflections
 924 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.033$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.143$
 $S = 1.01$
 1535 reflections

100 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.13$ e Å⁻³
 $\Delta\rho_{\min} = -0.16$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O1}-\text{H1}\cdots\text{O2}$	0.82	1.76	2.495 (2)	148
$\text{N}-\text{H0}\cdots\text{O1}^i$	0.86	2.21	2.993 (2)	151

Symmetry code: (i) $x + 1, y - 1, z$.

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: DIAMOND (Brandenburg, 2006); software used to prepare material for publication: publCIF (Westrip, 2010).

The authors gratefully acknowledge Professor Manfredo Hörner, Department of Chemistry, Universidade Federal de Santa Maria, Brazil, for his support and help with the X-ray measurements. They also thank PRONEX FAPERGS/CNPq 10/0009-2 and PRONEM FAPERGS/CNPq 11/2026-4 for funding this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LR2096).

References

- Apurba, K., Patra, A. K., Olmstead, M. M. & Mascharak, P. K. (2002). *Inorg. Chem.* **41**, 5403–5409.
- Booyesen, I., Gerber, T. I. A., Hosten, E. & Mayer, P. (2009). *Acta Cryst.* **E65**, o850.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2009). *APEX2*, *SAINTE* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Fry, N. L., Rose, M. J., Rogow, D. L., Nyitray, C., Kaur, M. & Mascharak, P. K. (2010). *Inorg. Chem.* **49**, 1487–1495.
- Liu, H. Y., Wang, K., Fu, H. Y., Yuan, M. L. & Chen, H. (2011). *Chin. Chem. Lett.* **22**, 738–740.
- Maurya, M. R., Titinchi, S. J. J. & Chand, S. (2003). *Catal. Lett.* **89**, 219–227.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2013). E69, o201 [doi:10.1107/S1600536812051963]

2,2'-Dihydroxy-*N,N'*-(ethane-1,2-diyl)dibenzamide

Daniel Pereira da Costa, Sabrina Madruga Nobre, Bruna Gonçalves Lisboa, Juliano Rosa de Menezes Vicenti and Davi Fernando Back

Comment

Nitrogenated compound amines, imines and amides are part of a broad class of molecules having a pharmacological and technological profile (Fry *et al.* 2010). Bisamides can be synthesized from a reaction between a diamine and an acylhalide under reflux (Apurba *et al.* 2002 ; Fry *et al.* 2010). Some synthesized bisamides have had their crystal structure determined (Booyesen *et al.* 2009). However, others have been synthesized and applied to several reactions, such as the hydroformylation of phenol (Maurya *et al.* 2003) and due to the simplicity of the synthesis of these compounds, they have been applied to coupling reactions, such as the Suzuki reaction (Liu *et al.* 2011). The asymmetric unit is formed by one half of the molecule and the whole molecule is generated through a crystallographic center of symmetry placed in the mid-point of the C-C bond of the ethane bridging moiety (Fig. 1). Intramolecular and intermolecular classical hydrogen bond interactions are observed in the solid state. Each molecule participates in intermolecular hydrogen bond interactions with two other neighboring molecules through $N^{\text{ii}}-H0^{\text{ii}}\cdots O1$ with a distance of 2.210 Å (symmetry code: (ii) $-1 + x, 1 + y, z$). The intramolecular hydrogen bonding systems forms six membered rings, showing a shorter distance of 1.763 Å ($O2\cdots H1-O1$). These hydrogen bonding may contribute to the stabilization of the crystal structure (Fig. 2).

Experimental


Ethylenediamine (0.1 mol, 6.6 ml) was added to a solution containing methyl salicylate (0.2 mol, 25.7 ml). The mixture was refluxed for 7 h. The reaction product was washed with ethyl acetate (3 times in 30 ml) and dried over $CaCl_2$. After recrystallization from hexane, a yellow solid was obtained (23.1 g, 77%).

Refinement


H atoms attached to aromatic C atoms were positioned with idealized geometry and were refined isotropic with $U_{\text{eq}}(\text{H})$ set to 1.2 times of the $U_{\text{eq}}(\text{C})$ using a riding model with $C-H = 0.93$ Å. H atoms attached to N atoms and to the ethylene fragment were located in difference Fourier maps. Their positional and isotropic displacement parameters were refined.

Computing details

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT* (Bruker, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *pubCIF* (Westrip, 2010).


Figure 1

Molecular projection showing the asymmetric unit represented in dark colors. Ellipsoid probability: 50%. Symmetry codes: (i) $2-x, 1-y, 2-z$


Figure 2

Packing diagram view showing the intramolecular and intermolecular hydrogen bond interactions. Some hydrogen atoms were omitted for clarity. (ii) $-1+x, 1+y, z$

2,2'-Dihydroxy-*N,N'*-(ethane-1,2-diyl)dibenzamide

Crystal data

$C_{16}H_{16}N_2O_4$

$M_r = 300.31$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 4.6311(3)\ \text{\AA}$

$b = 5.0435(3)\ \text{\AA}$

$c = 15.2957(9)\ \text{\AA}$

$\alpha = 89.091(4)^\circ$

$\beta = 83.315(4)^\circ$

$\gamma = 85.956(4)^\circ$

$V = 353.94(4)\ \text{\AA}^3$

$Z = 1$

$F(000) = 158$

$D_x = 1.409\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 1575 reflections

$\theta = 2.7\text{--}25.8^\circ$

$\mu = 0.10\ \text{mm}^{-1}$

$T = 296\ \text{K}$

Block, colourless

$0.34 \times 0.24 \times 0.11\ \text{mm}$

Data collection

Bruker APEXII CCD diffractometer	9142 measured reflections
Radiation source: fine-focus sealed tube	1535 independent reflections
Graphite monochromator	924 reflections with $I > 2\sigma(I)$
φ and ω scans	$R_{\text{int}} = 0.033$
Absorption correction: gaussian (SADABS; Bruker, 2009)	$\theta_{\text{max}} = 27.1^\circ$, $\theta_{\text{min}} = 2.7^\circ$
$T_{\text{min}} = 0.966$, $T_{\text{max}} = 0.989$	$h = -5 \rightarrow 5$
	$k = -6 \rightarrow 6$
	$l = -19 \rightarrow 19$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.049$	H-atom parameters constrained
$wR(F^2) = 0.143$	$w = 1/[\sigma^2(F_o^2) + (0.0644P)^2 + 0.0596P]$
$S = 1.01$	where $P = (F_o^2 + 2F_c^2)/3$
1535 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
100 parameters	$\Delta\rho_{\text{max}} = 0.13 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.16 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.4041 (3)	1.1108 (3)	0.79344 (10)	0.0718 (5)
H1	0.4677	1.0900	0.8411	0.108*
O2	0.7318 (3)	0.9317 (3)	0.90207 (9)	0.0629 (5)
N	1.0113 (3)	0.5534 (3)	0.87856 (10)	0.0507 (5)
H0	1.0709	0.4248	0.8433	0.061*
C1	0.7459 (4)	0.7361 (3)	0.76153 (11)	0.0426 (5)
C2	0.5357 (4)	0.9263 (4)	0.73646 (12)	0.0496 (5)
C3	0.4584 (5)	0.9234 (5)	0.65190 (15)	0.0676 (6)
H3	0.3184	1.0495	0.6352	0.081*
C4	0.5845 (5)	0.7384 (4)	0.59253 (14)	0.0649 (6)
H4	0.5312	0.7405	0.5357	0.078*
C5	0.7895 (5)	0.5493 (5)	0.61620 (14)	0.0655 (6)
H5	0.8742	0.4220	0.5760	0.079*
C6	0.8674 (5)	0.5507 (4)	0.69979 (13)	0.0589 (6)
H6	1.0068	0.4226	0.7156	0.071*
C7	0.8293 (4)	0.7466 (4)	0.85175 (12)	0.0448 (5)
C8	1.1112 (4)	0.5513 (4)	0.96491 (12)	0.0523 (5)

H8A	1.1517	0.7304	0.9796	0.063*
H8B	1.2913	0.4402	0.9636	0.063*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0882 (11)	0.0606 (9)	0.0622 (9)	0.0348 (8)	-0.0120 (8)	-0.0122 (7)
O2	0.0780 (10)	0.0543 (9)	0.0545 (9)	0.0202 (7)	-0.0125 (7)	-0.0214 (7)
N	0.0591 (10)	0.0484 (10)	0.0439 (9)	0.0123 (8)	-0.0115 (7)	-0.0116 (7)
C1	0.0458 (10)	0.0388 (10)	0.0426 (10)	0.0022 (8)	-0.0050 (8)	-0.0043 (8)
C2	0.0569 (11)	0.0402 (10)	0.0501 (11)	0.0053 (9)	-0.0049 (9)	-0.0011 (9)
C3	0.0806 (15)	0.0620 (14)	0.0601 (13)	0.0147 (12)	-0.0198 (12)	0.0036 (11)
C4	0.0828 (16)	0.0679 (15)	0.0459 (12)	-0.0029 (12)	-0.0165 (11)	-0.0007 (11)
C5	0.0820 (15)	0.0642 (14)	0.0490 (12)	0.0113 (12)	-0.0094 (11)	-0.0158 (10)
C6	0.0696 (13)	0.0567 (13)	0.0488 (12)	0.0176 (10)	-0.0111 (10)	-0.0120 (10)
C7	0.0458 (10)	0.0420 (10)	0.0456 (10)	0.0040 (8)	-0.0041 (8)	-0.0085 (8)
C8	0.0540 (11)	0.0563 (12)	0.0477 (11)	0.0058 (9)	-0.0154 (8)	-0.0091 (9)

Geometric parameters (\AA , $^\circ$)

O1—C2	1.349 (2)	C3—C4	1.365 (3)
O1—H1	0.8206	C3—H3	0.9300
O2—C7	1.245 (2)	C4—C5	1.372 (3)
N—C7	1.334 (2)	C4—H4	0.9300
N—C8	1.449 (2)	C5—C6	1.369 (3)
N—H0	0.8600	C5—H5	0.9300
C1—C6	1.383 (2)	C6—H6	0.9300
C1—C2	1.400 (3)	C8—C8 ⁱ	1.511 (4)
C1—C7	1.478 (2)	C8—H8A	0.9700
C2—C3	1.382 (3)	C8—H8B	0.9700
C2—O1—H1	109.5	C6—C5—C4	119.07 (19)
C7—N—C8	122.58 (15)	C6—C5—H5	120.5
C7—N—H0	118.6	C4—C5—H5	120.5
C8—N—H0	118.8	C5—C6—C1	122.21 (18)
C6—C1—C2	117.95 (17)	C5—C6—H6	118.9
C6—C1—C7	123.62 (16)	C1—C6—H6	118.9
C2—C1—C7	118.42 (15)	O2—C7—N	120.36 (17)
O1—C2—C3	119.21 (17)	O2—C7—C1	120.87 (16)
O1—C2—C1	121.37 (17)	N—C7—C1	118.78 (15)
C3—C2—C1	119.42 (17)	N—C8—C8 ⁱ	112.0 (2)
C4—C3—C2	120.98 (19)	N—C8—H8A	109.2
C4—C3—H3	119.5	C8 ⁱ —C8—H8A	109.2
C2—C3—H3	119.5	N—C8—H8B	109.2
C3—C4—C5	120.4 (2)	C8 ⁱ —C8—H8B	109.2
C3—C4—H4	119.8	H8A—C8—H8B	107.9
C5—C4—H4	119.8		
C6—C1—C2—O1	178.94 (18)	C2—C1—C6—C5	0.3 (3)
C7—C1—C2—O1	-2.0 (3)	C7—C1—C6—C5	-178.7 (2)

C6—C1—C2—C3	-0.3 (3)	C8—N—C7—O2	-1.8 (3)
C7—C1—C2—C3	178.77 (19)	C8—N—C7—C1	178.24 (17)
O1—C2—C3—C4	-179.4 (2)	C6—C1—C7—O2	173.61 (19)
C1—C2—C3—C4	-0.2 (4)	C2—C1—C7—O2	-5.4 (3)
C2—C3—C4—C5	0.6 (4)	C6—C1—C7—N	-6.5 (3)
C3—C4—C5—C6	-0.6 (4)	C2—C1—C7—N	174.55 (17)
C4—C5—C6—C1	0.2 (4)	C7—N—C8—C8 ⁱ	80.7 (3)

Symmetry code: (i) $-x+2, -y+1, -z+2$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
O1—H1...O2	0.82	1.76	2.495 (2)	148
N—H0...O1 ⁱⁱ	0.86	2.21	2.993 (2)	151

Symmetry code: (ii) $x+1, y-1, z$.