

ARTICLE

Received 21 Mar 2016 | Accepted 17 Aug 2016 | Published 27 Sep 2016

DOI: 10.1038/ncomms12933

OPEN

Carbene-catalysed reductive coupling of nitrobenzyl bromides and activated ketones or imines via single-electron-transfer process

Bao-Sheng Li^{1,*}, Yuhuang Wang^{1,*}, Rupert S.J. Proctor¹, Yuexia Zhang¹, Richard D. Webster¹, Song Yang², Baoan Song² & Yonggui Robin Chi^{1,2}

Benzyl bromides and related molecules are among the most common substrates in organic synthesis. They are typically used as electrophiles in nucleophilic substitution reactions. These molecules can also be activated via single-electron-transfer (SET) process for radical reactions. Representative recent progress includes α -carbon benzylation of ketones and aldehydes via photoredox catalysis. Here we disclose the generation of (nitro)benzyl radicals via *N*-heterocyclic carbene (NHC) catalysis under reductive conditions. The radical intermediates generated via NHC catalysis undergo formal 1,2-addition with ketones to eventually afford tertiary alcohol products. The overall process constitutes a formal polarity-inversion of benzyl bromide, allowing a direct coupling of two initially electrophilic carbons. Our study provides a new carbene-catalysed reaction mode that should enable unconventional transformation of (nitro)benzyl bromides under mild organocatalytic conditions.

¹Division of Chemistry & Biological Chemistry, Nanyang Technological University, School of Physical & Mathematical Sciences, Singapore 637371, Singapore.

²Laboratory Breeding Base of Green Pesticide and Agricultural Bioengineering, Key Laboratory of Green Pesticide and Agricultural Bioengineering, Ministry of Education, Guizhou University, Huaxi District, Guiyang 550025, China. * These authors contributed equally to this work. Correspondence and requests for materials should be addressed to Y.R.C. (email: robinchi@ntu.edu.sg).

Alkyl halides, such as benzyl bromides, are readily available and inexpensive raw materials in chemical synthesis. Due to the electronegative nature of the halide atoms, these molecules are typically used as electrophiles to react with nucleophilic atoms in substitution reactions^{1,2}. Alkyl halides can also be activated to form radical intermediates for various reactions. Indeed, the use of alkyl halides to generate radical intermediates^{3–6} for α -carbon alkylation of aldehydes^{7–10} and ketones¹¹ has recently received intensive attention (Fig. 1a)¹². For instance, MacMillan has pioneered the use of ruthenium photoredox catalysts to enable single-electron-transfer (SET) processes under light to form radicals that can then react with enamines catalytically generated from aldehydes and amine catalysts^{7,8}. Melchiorre has realized asymmetric α -carbon alkylation of aldehydes via a radical process driven by the formation of a donor-acceptor complex and light^{9,10}.

Our laboratory is interested in using *N*-heterocyclic carbene (NHC) organic catalysts to develop new reaction modes for rapid synthesis of functional molecules^{13,14}. In our recent efforts in developing NHC-enabled radical reactions, we have found that the presence of a nitro group (in either substrates¹⁵ or reagents¹⁶) is important to initiate a SET process under NHC catalysis. Here we report that benzyl bromides, bearing a nitro substituent on the benzene ring, can undergo an NHC-mediated SET process using an aldehyde as a formal reductant (Fig. 1a). The catalytically generated benzyl radical intermediate then undergoes a formal 1,2-addition to a ketone to eventually form a tertiary alcohol product. The overall reaction is a formal reductive coupling of nitrobenzyl bromide **1** and ketone **2** (Fig. 1b).

Notably, the use of alkyl halides to react with another electrophile typically need to go through pre-formed organometallic reagents (such as Grignard reagents)^{17,18}. Catalytic variants of the approaches with organometallic reagents have also been developed for the coupling of alkyl halides with another electrophilic substrate^{19–21}. However, sensitive functional groups such as nitro and ketone substituents often cannot survive under conditions with organometallic reagents. In the area of organic catalysis, NHCs have mainly been used in electron-pair-transfer reactions^{22–29}. Despite the fact that *vitamin B1* (an NHC precursor)-mediated reactions in biological systems go through SET processes³⁰, NHC-mediated radical reactions in synthetic chemistry are much less developed. Studer³¹ reported

the oxidation of aldehydes to carboxylic esters using TEMPO as a single-electron oxidant via NHC catalysis. We developed the dimerisation of nitroalkenes via NHC-mediated generation of radical anion intermediates from nitroalkenes¹⁵. Rovis³² and our¹⁶ laboratories independently reported β -hydroxylation of unsaturated aldehydes through SET processes³³ under NHC catalysis. Our present study in catalytically turning (nitro)benzyl bromides to radical intermediates for unusual reaction offers a new reaction mode of NHC catalysis that should provide unique opportunities in chemical synthesis.

Results

Reaction optimization. Key results of initial studies are summarized in Table 1 (the detailed conditions please see the Supplementary Table 1). The 4-nitrobenzyl bromide **1a**, and α,β -unsaturated ketoester **2a** were used as model substrates. Optimal conditions for the formation of alcohol product **4a** (in 75% isolated yield) were found when triazolium salt **A**³⁴ was used as an NHC precursor, aryl aldehyde **3a** as a reductant, and *N,N*-diisopropylethylamine as a base (entry 1). The reactions under darkness (the reaction flask was covered with aluminium foil) or normal room lighting conditions gave nearly the same outcomes, indicating that the generation of radical intermediates under our system was not driven by light. Both the NHC catalyst (entry 2) and the aryl aldehyde reductant (entry 3) were required for the reductive coupling reaction to proceed. The nitro group in substrate **1a** was critical for this reaction to occur, and its removal or replacement with other substituents (for example, **1b**, **1c** or **1d**) led to no detectable formation of **4a** (entry 4). The substituents on the aryl aldehyde reductant could also affect the reaction efficiency (entries 5, 6). When aryl aldehyde such as **3b** and **3c** were used as the reductant, **4a** was formed in much lower yield. In these cases (entries 4–6), most of the benzyl bromide and




Figure 1 | NHC-catalysed reduction coupling of nitrobenzyl bromide with activated carbonyl compounds. (a) Formation of benzyl radical via photoredox or NHC catalysis. (b) Reductive coupling of nitrobenzyl bromides and ketones.

Table 1 | Key results of condition optimizations.

Entry*	Benzyl bromide	Ar'CHO	NHC	4a
1	1a	3a	A	78 (75)*
2		As entry 1, no NHC		ND
3		As entry 1, no Ar'CHO		ND
4		As entry 1, benzyl bromide = 1b , 1c or 1d		ND
5		As entry 1, Ar'CHO = 3b		36
6		As entry 1, Ar'CHO = 3c		27
7		As entry 1, NHC = B		35
8		As entry 1, NHC = C		29
9		As entry 1, NHC = D		23
10		As entry 1, NHC = E		ND

DIPEA, *N,N*-diisopropylethylamine; Mes, 1,3,5-trimethylbenzene; ND, not detected.
 ***1a** to **1d** (0.15 mmol), **2a** (0.10 mmol, 19.0 mg), NHC pre-catalyst **A** to **E** (5 mol%), aldehyde **3** (0.15 eq.), DIPEA (0.15 mmol, 25 μ l), CH₃OH (1.0 ml), 8 h. Yields were estimated via ¹H NMR analysis. Isolated yield after SiO₂ chromatography. Bold entries indicate optimal condition.

ketoester substrates remained unreacted, and the aryl aldehyde substrate was consumed to form benzoin adducts and the corresponding carboxylic ester. The *N*-aryl substituents in the triazolium-based NHC pre-catalysts (entries 1, 7 and 8) could affect the reaction outcomes. Imidazolium NHC pre-catalyst **D** could also mediate this reaction, albeit with a low yield (23%, entry 9). A thiazolium pre-catalyst (**E**) could not mediate this reaction (entry 10). When catalyst **B** or **C** were used, a significant amount of benzoin condensation products of the aldehyde were observed (yields of benzoin adduct: <5% with catalyst **A**; 45% with catalyst **B**; 40% with catalyst **C**).

Substrates scope with α -keto esters. With an acceptable condition in hand, we examined the scope of the reductive coupling reaction (Fig. 2). Examples of nitrobenzyl bromides that could react effectively (with ketoester **2a** as a model ketone substrate)

are given in Fig. 2a. The use of benzyl bromides with the NO₂ group at the para (**4a**) or ortho (**4b**) position of the benzene ring both worked well, while meta-nitro benzyl bromide could not react to give the proposed reductive coupling product. Placing additional substituents on the para-nitro benzyl bromides (**4c**, **4d**) and ortho-nitro benzyl bromides (**4e–j**) were all tolerated. For example, having an ester (**4f**) or ketone (**4g**) moiety on the ortho-nitrobenzyl bromide did not affect the reaction outcomes. It should be noted that such electrophilic functional groups (for example, esters, ketones) are likely problematic in conventional approaches using organometallic reagents^{17,18}. Secondary and tertiary nitrobenzyl bromides (**4k**, **4l**) are also effective substrates. Examples of ketoester substrates are given in Fig. 2b. Different substituents and substitution patterns on the phenyl ring of unsaturated ketoester **2a** were tolerated (**4m–r**). The phenyl unit of **2a** could be replaced with a thiofuran (**4s**), alkene (**4t**) or alkyl (**4u**) unit. Phenyl ketoester **4v** could effectively react with




Figure 2 | α -Ketoester as substrates and its transformations. Scope of the reaction (**a,b**) and synthetic transformation of the reductive coupling product (**c**). The reactions were performed on 0.1 mmol scale (condition as in Table 1, entry 1). Isolated yields base on ketoesters **2**. *The reaction was carried out at 5 mmol (ketoester) scale. **Relative configuration. MCPBA, 3-chloroperoxybenzoic acid.

nitrobenzyl bromide **1a**, while the use of alkyl ketoesters (such as methyl 2-oxobutanoate) did not lead to the proposed products. The carbon-carbon double bond of unsaturated ketoester **2a** could be replaced with a carbon-carbon triple bond (**4w-z**), with little influence on the reaction yields. The reaction, carried out under mild conditions, can be easily scaled up. For example, product **4b** could be prepared in gram scale without loss in yield.

The reductive coupling adduct from our reactions can be readily converted to other functional molecules (Fig. 2c). For example, reduction of the NO₂ group in **4b** followed by an intramolecular ester-to-amide exchange could give lactam **4b-1** that bears a quinolone core widely found in natural products and bioactive molecules³⁵. The alkene in allylic alcohol **4b** could be easily converted to epoxide **4b-2** in 87% yield with excellent stereo-selectivity (only one diastereoisomer was obtained). The hydroxyl group in **4b** could undergo a formal 1,3-migration under acidic conditions to form γ -hydroxyl α , β -unsaturated ester **4b-3** in 81% yield.

Substrates scope with isatins. To further expand the scope of our reductive coupling reactions, we next examined isatin derivatives as ketone substrates to react with the nitrobenzyl bromides

(Fig. 3). To our delight, the standard conditions used above for the ketoester substrates (Table 1, entry 1) worked effectively for the isatin substrates without any further optimization. The reaction yields here were slightly higher than those reactions using ketoesters. Nitrobenzyl bromides with different substituents and substitution patterns were well tolerated, giving the desired alcohol products in 68–81% yields (**7a-j**, Fig. 3a). The substituents on the isatin had little effect on the reaction yields (**7k-t**, Fig. 3b). The isatin without an *N*-substituent (**7u**) reacted effectively as well. Further synthetic transformations of the reductive coupling products can provide a shortcut to functional molecules (Fig. 3c). For example, reduction and cyclization of **7u** followed by a facile methylation gave neocryptolepine **7u-2**, a natural product isolated from the bark of the roots of *Cryptolepis sanguinolenta*^{36,37}. Neocryptolepine and its analogs exhibit a strong antiparasmodial activity against *Plasmodium falciparum* chloroquine-resistant strains³⁸.

Substrates scope with imines. Imines as reaction partners also worked effectively under our catalytic condition (Fig. 4). The results further illustrated the broad applicability of our NHC-mediated SET approach.




Figure 3 | Isatin as substrates and its application. Isatin as the substrates (**a,b**), and transformation of **7u** to natural product (**c**). The condition as in Table 1, entry 1. Isolated yields base on isatin (**6**). * The reaction was carried out at 5 mmol (ketoester) scale.




Figure 4 | Imines as the substrates. The condition as in Table 1, entry 1. Isolated yields base on imines (**8**).

Discussion

To understand the reaction pathway (as postulated in Fig. 5f), we analysed the side reactions and performed multiple control experiments (Fig. 5, additional control experiments see Supplementary Methods). Under the standard conditions (Table 1, entry 1), a debromination product (**10a**) of nitrobenzyl bromide **1a** was observed in 42% yield (Fig. 5a; the yield of **10a** is based on ketone ester **2a**—the limiting reagent). This side product (**10a**) likely come from a further reduction of nitrobenzyl radical intermediate **B2** (Fig. 5a). Specifically, the reduction of **B2** formed a nitrobenzyl anion that underwent spontaneous protonation in CH₃OH as the solvent and proton source to form **10a**. The nitrotoluene substrate **10a** could not react with our ketone substrate **2a** (Fig. 5b,c), suggesting that the formation of **4a** in our catalytic reaction unlikely came from a nucleophilic addition of nitrobenzyl anion to the ketone substrate. A possible reduction³⁹ of the ketone substrate **2a** to form α -hydroxyl ester **11a** was not observed. To rule out the possibility that **9a** was formed *in situ* and quickly reacted with the bromide substrate **1a**. We prepared **11a** and evaluated its reaction with **1a** (Fig. 5d). No formation of **4a** was observed under various conditions (Fig. 5d).

We have also measured the reduction potential of compounds **1a**, **2a** and **6a** via cyclic voltammetry (see Supplementary Method and Supplementary Fig. 130). The experiments showed that compound **1a** ($E_p^{\text{red}} = -1.18$ V vs) has a higher reduction potential than compounds **2a** ($E_p^{\text{red}} = -1.47$ V vs) and **6a** ($E_p^{\text{red}} = -1.38$ V vs). The results suggested that most likely **1a** is easier to be reduced, supporting our mechanistic proposal (Fig. 5f) that the SET process first occurs on nitrobenzyl bromide **1a**. Finally, the addition of radical trapping agent 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) to our reaction diminished the formation of **4a** (Fig. 5e). A radical–radical coupling adduct **12a** between TEMPO and nitrobenzyl radical **B2** was observed in 7% yield (Fig. 5e). Similar observations in trapping of benzyl radical intermediates with TEMPO under photocatalysis were reported by Melchiorre⁹ and Meggers¹¹, respectively.

A postulated reaction pathway is illustrated in Fig. 5f. Aldehyde reagent **3** reacts with the NHC catalyst to form a Breslow intermediate⁴⁰ **A1** that then serves as a single-electron

reductant to convert nitrobenzyl bromide **1a** into radical anion intermediate **B1**. Breslow intermediate **A1** is oxidized to a radical cation intermediate³¹ **A2** in this process. The leaving of a bromide anion (Br^-) from **B1** affords nitrobenzyl radical intermediate **B2**. On the one hand, addition of radical intermediate **B2** to ketone^{41–44} substrate **2** forms radical intermediate **B3**. Subsequent reduction of **B3** by Breslow intermediate-derived radical cation **A2** forms **B4**, which can then be protonated to form alcohol product **4**. The oxidized **A2** becomes azolium ester intermediate **A3** that can then be trapped by CH₃OH to form ester **5** with regeneration of the NHC catalyst. Alternatively, reduction of radical intermediate **B2** to form intermediate **B5** with an anionic benzylic carbon may occur. Nucleophilic addition of **B5** to ketone substrate **2** can also lead to product **4**. At this point, this ionic pathway (via intermediate **B5**) cannot be ruled out. In the overall process, the initially electrophilic nitrobenzyl bromide is catalytically activated to directly react with another electrophilic substrate (the ketone substrate).

In summary, we have developed a new organic catalytic method for the generation of nitrobenzyl radical intermediates from nitrobenzyl bromides under mild conditions. The catalytically formed radical intermediates react with ketones, resulting in a formal reductive coupling of initially electrophilic nitrobenzyl bromides and ketones. The products from our catalytic reactions can be readily transformed to functional molecules and natural products. Our study provides a new reaction mode that can likely allow for unique design in organic synthesis. It may also encourage further adventures in developing novel organocatalytic activations by single-electron-transfer processes. Mechanistic studies and additional development of this reaction mode for challenging substrates are in progress.

Methods

General strategy of reductive coupling reactions. To a reaction tube equipped with a stirrer bar were added NHC catalyst salt **A** (0.005 mmol), aldehyde **3** (0.15 mmol), ketone (**2**, **6** or **8**, 0.1 mmol), and benzyl bromide **1** (0.15 mmol). The tube was sealed, evacuated and purged, and a solution of *N,N*-diisopropylethylamine (0.15 mmol, 19.4 mg, 25 μ l) in methanol (1.0 ml) was added. The reaction was stirred for 8 h. The solvent was removed *in vacuo* and the residue was purified by flash column chromatography.




Figure 5 | Experiments to elucidate the reaction mechanism and possible reaction pathways. (a) The experiment indicates that the α -ketoester has no effect for the generation of nitrobenzyl radical. (b) The result suggest that the 1,2-addition product unlikely come from 4-nitro toluene in the current reaction system. (c) The coupling product could not be produced in the presence of NaOMe. (d) The final product could not be obtained via the alkylation of α -hydroxyl ester. (e) The result indicates that the nitro benzyl radical is existent in current reaction system. (f) The nitrobenzyl radical **B2** could be formed via oxidation of Breslow intermediate **A1** and leaving of bromide anion from **B1**. Subsequently, the carbon-carbon bond can be formed via 1,2-addition of nitrobenzyl radical intermediate **B3** or ionic intermediate **B5** to carbonyl compounds.

Data availability. For ^1H , ^{13}C NMR and high-performance liquid chromatography spectra of compounds in this manuscript, see Supplementary Figs 1–129. For Oak Ridge thermal ellipsoid plot (ORTEP) of products **4w** and **7l**, see Supplementary Information. The X-ray crystallographic coordinates for structures

reported in this article have been deposited at the Cambridge Crystallographic Data Centre (**4w**: CCDC 1451838; **7l**: CCDC 1451840). These data could be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

References

- Semb, J. & McElvain, S. M. Reaction of organic halides with piperidine. I. Alkyl bromide. *J. Am. Chem. Soc.* **53**, 690–696 (1931).
- Evans, A. G. Reaction of organic halides in solution. *Nature* **157**, 438–439 (1946).
- Moreno, S. N. J., Schreiber, J. & Mason, R. Nitrobenzyl radical metabolites from microsomal reduction of nitrobenzyl chlorides. *J. Biol. Chem.* **261**, 7811–7815 (1986).
- Lawless, J. G., Bartak, D. E. & Hawley, M. D. The electrochemical reduction of nitrobenzyl halides in acetonitrile. *J. Am. Chem. Soc.* **91**, 7121–7127 (1969).
- Bays, J. P., Blumer, S. T., Baral-Tosh, S., Behar, D. & Neta, P. Intramolecular electron transfer and dehalogenation of nitroaromatic anion radicals. *J. Am. Chem. Soc.* **105**, 320–324 (1983).
- Narayanam, J. M. R., Tucker, J. W. & Stephenson, C. R. J. Electron-transfer photoredox catalysis: development of a tin-free reductive dehalogenation reaction. *J. Am. Chem. Soc.* **131**, 8756–8757 (2009).
- Nicewicz, D. A. & MacMillan, D. W. C. Merging photoredox catalysis with organocatalysis: the direct asymmetric alkylation of aldehydes. *Science* **322**, 77–80 (2008).
- Shih, H.-W., Vander Wal, M. N., Grange, R. L. & MacMillan, D. W. C. Enantioselective β -benzylation of aldehydes via photoredox organocatalysis. *J. Am. Chem. Soc.* **132**, 13600–13603 (2010).
- Arceo, E., Jurberg, I. D., Álvarez-Fernández, A. & Melchiorre, P. Photochemical activity of a key donor–acceptor complex can drive stereoselective catalytic-alkylation of aldehydes. *Nature Chem.* **5**, 750–756 (2013).
- Arceo, E., Bahamonde, A., Bergonzini, G. & Melchiorre, P. Enantioselective direct α -alkylation of cyclic ketones by means of photo-organocatalysis. *Chem. Sci.* **5**, 2438–2442 (2014).
- Huo, H. *et al.* Asymmetric photoredox transition-metal catalysis activated by visible light. *Nature* **515**, 100–103 (2015).
- Du, J. N., Skubi, K. L., Schultz, D. M. & Yoon, T. P. A dual-catalysis approach to enantioselective [2 + 2] photocycloadditions using visible light. *Science* **344**, 392–396 (2014).
- Li, B.-S. *et al.* Carbon–carbon bond activation of cyclobutenones enabled by the addition of chiral organocatalyst to ketone. *Nat. Commun.* **6**, 6207 (2015).
- Fu, Z., Xu, J., Zhu, T., Leong, W. W. Y. & Chi, Y. R. β -Carbon activation of saturated carboxylic esters through N-heterocyclic carbene organocatalysis. *Nature Chem.* **5**, 835–839 (2013).
- Du, Y. *et al.* N-heterocyclic carbene organocatalytic reductive β,β -coupling reactions of nitroalkenes via radical intermediates. *Org. Lett.* **16**, 5678–5681 (2014).
- Zhang, Y. *et al.* N-heterocyclic carbene-catalyzed radical reactions for highly enantioselective β -hydroxylation of enals. *J. Am. Chem. Soc.* **137**, 2416–2419 (2015).
- Boudier, A., Bromm, L. O., Lotz, M. & Knochel, P. New applications polyfunctional organometallic compounds in organic synthesis. *Angew. Chem. Int. Ed.* **39**, 4414–4435 (2000).
- Terao, J. & Kambe, N. Cross-coupling reaction of alkyl halides with grignard reagents catalyzed by Ni, Pd, or Cu complexes with π -carbon Ligand(s). *Acc. Chem. Res.* **41**, 1545–1554 (2008).
- Knochel, P. *et al.* Highly functionalized organomagnesium reagents prepared through halogen–metal exchange. *Angew. Chem. Int. Ed.* **42**, 4302–4320 (2003).
- Kambe, N., Iwasakia, T. & Terao, J. Pd-catalyzed cross-coupling reactions of alkyl halides. *Chem. Soc. Rev.* **40**, 4937–4947 (2011).
- Dudnik, A. S. & Fu, G. C. Nickel-catalyzed coupling reactions of alkyl electrophiles, including unactivated tertiary halides, to generate carbon–boron bonds. *J. Am. Chem. Soc.* **134**, 10693–10697 (2012).
- Flanigan, D. M., Michailidis-Romanov, F., White, N. A. & Rovis, T. Organocatalytic reactions enabled by N-heterocyclic carbenes. *Chem. Rev.* **115**, 9307–9387 (2015).
- Enders, D., Niemeier, O. & Henseler, A. Organocatalysis by N-heterocyclic carbenes. *Chem. Rev.* **207**, 5606–5655 (2007).
- Ryan, S. J., Candish, L. & Lupton, D. W. Acyl anion free N-heterocyclic carbene organocatalysis. *Chem. Soc. Rev.* **42**, 4906–4917 (2013).
- De Sarkar, S., Biswas, A., Samanta, R. C. & Studer, A. Catalysis with N-heterocyclic carbenes under oxidative conditions. *Chem. Eur. J.* **19**, 4664–4678 (2013).
- Cohen, D. T. & Scheidt, K. A. Cooperative Lewis acid/N-heterocyclic carbene catalysis. *Chem. Sci.* **3**, 53–57 (2012).
- Hopkinson, M. N., Richter, C., Schedler, M. & Glorius, F. An overview of N-heterocyclic carbenes. *Nature* **510**, 485–496 (2014).
- Mahatthananchai, J. & Bode, J. W. On the mechanism of N-heterocyclic carbene-catalyzed reactions involving acyl azoliums. *Acc. Chem. Res.* **47**, 696–707 (2014).
- Nair, V. *et al.* Employing homoenolate generated by NHC in carbon–carbon bond-forming reactions: state of the art. *Chem. Soc. Rev.* **40**, 5336–5346 (2011).
- Ragsdale, S. W. Pyruvate ferredoxin oxidoreductase and its radical intermediate. *Chem. Rev.* **103**, 2333–2346 (2003).
- Guin, J., Sarkar, S. D., Grimme, S. & Studer, A. Biomimetic carbene-catalyzed oxidations of aldehydes using TEMPO. *Angew. Chem., Int. Ed.* **47**, 8727–8730 (2008).
- White, N. A. & Rovis, T. Enantioselective N-heterocyclic carbene-catalyzed β -hydroxylation of enals using nitroarenes: an atom transfer reaction that proceeds via single electron transfer. *J. Am. Chem. Soc.* **136**, 14674–14677 (2014).
- White, N. A. & Rovis, T. Oxidatively initiated NHC-catalyzed enantioselective synthesis of 3,4-disubstituted cyclopentanones from enals. *J. Am. Chem. Soc.* **137**, 10112–10115 (2015).
- Kerr, M. S., de Alaniz, J. R. & Rovis, T. An efficient synthesis of achiral and chiral 1,2,4-triazolium salts: bench stable precursors for N-heterocyclic carbenes. *J. Org. Chem.* **70**, 5725–5728 (2005).
- Simonetti, S. O., Larghi, E. L. & Kaufman, T. S. A convenient approach to an advanced intermediate toward the naturally occurring, bioactive 6-substituted 5-hydroxy-4-aryl-1H-quinolin-2-ones. *Org. Biomol. Chem.* **14**, 2625–2636 (2016).
- Sharaf, M. H. M., Schiff, Jr P. L., Tackie, A. N., Phoebe, Jr C. H. & Martin, G. E. Two new indoloquinoline alkaloids from *Cryptolepis sanguinolenta*: cryptosanguinolentine and cryptotackieine. *J. Heterocycl. Chem.* **33**, 239–243 (1996).
- Cimanga, K., De Bruyne, T., Pieters, L., Claeys, M. & Vlietinck, A. New alkaloids from *Cryptolepis sanguinolenta*. *Tetrahedron Lett.* **37**, 1703–1706 (1996).
- Cimanga, K., De Bruyne, T., Pieters, L., Claeys, M. & Vlietinck, A. *In vitro* and *in vivo* antiplasmodial activity of cryptolepine and related alkaloids from *Cryptolepis sanguinolenta*. *J. Nat. Prod.* **60**, 688–691 (1997).
- Chan, A. & Scheidt, K. A. Hydroacylation of activated ketones catalyzed by N-heterocyclic carbenes. *J. Am. Chem. Soc.* **128**, 4558–4559 (2006).
- Breslow, R. On the mechanism of thiamine action. IV. I Evidence from studies on model systems. *J. Am. Chem. Soc.* **80**, 3719–3726 (1958).
- Halland, N. & Jørgensen, K. A. Intermolecular addition of alkyl radicals to imines in the absence and in the presence of a Lewis acid. *J. Chem. Soc., Perkin Trans. 1*, 1290–1295 (2001).
- Friestad, G. K. Addition of carbon-centered radicals to imines and related compounds. *Tetrahedron* **57**, 5461–5496 (2001).
- Kawamoto, T. *et al.* Radical addition of alkyl halides to formaldehyde in the presence of cyanoborohydride as a radical mediator. a new protocol for hydroxymethylation Reaction. *J. Am. Chem. Soc.* **134**, 875–877 (2012).
- Henry, D. J., Coote, M., Gómez-Balderas, R. & Radom, L. Comparison of the kinetics and thermodynamics for methyl radical addition to C=C, C=O, and C=S double bonds. *J. Am. Chem. Soc.* **2004**, 1732–1740 (2004).

Acknowledgements

We acknowledge support by Singapore National Research Foundation (NRF), Ministry of Education, Singapore Economic Development Board (EDB), GlaxoSmithKline (GSK) and Nanyang Technological University (NTU); and China's National Key program for Basic Research (No. 2010CB 126105), Thousand Talent Plan, National Natural Science Foundation of China (No.21132003; No. 21472028), the Returned Oversea Student Science and Technology Activity Program of Guizhou Province, The Guizhou Science and Technology Department's Key Program and Guizhou University. We thank Dr R. Ganguly and Y. Li (NTU) for assistance with X-ray structure analysis. We thank Prof. Hua Zhang and Dr Huan Xu (MSE, NTU) for help with cyclic voltammetry experiments. We thank Yinguo Liu for his help in preparing several substrates, and Xingxing Wu for helpful discussions.

Author contributions

B.-S.L. and Y.W. contributed equally to this work. B.-S.L. proposed the reductive coupling reactions of benzyl bromides with ketones, and performed most of the experiments; Y.W. initially proposed NHC-mediated generation of nitrobenzyl radicals; R.S.J.P. performed part of the experiments; R.D.W. carried out EPR studies attempting to understand the reaction mechanism; Y.Z., S.Y. and B.-A.S. participated in designing part of the experiments; Y.R.C. conceptualized and directed the project, and drafted the manuscript with the assistance from all co-authors. All authors contributed to discussions.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Li, B.-S. *et al.* Carbene-catalysed reductive coupling of nitrobenzyl bromides and activated ketones or imines via single-electron-transfer process. *Nat. Commun.* 7, 12933 doi: 10.1038/ncomms12933 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016