Open Access Full Text Article

LETTER

Commentary on the Paper by Yang et al: Development and Validation of Nomogram Prediction Model for Postoperative Sleep Disturbance in Patients Undergoing Non-Cardiac Surgery: A Prospective Cohort Study [Letter]

Jiaojiao Yang, Huihui Miao, Hui Qiao, Tianzuo Li

Department of Anesthesiology, Beijing Shijitan Hospital, Capital Medical University, Beijing, 100038, People's Republic of China

Correspondence: Huihui Miao; Tianzuo Li, Department of Anesthesiology, Beijing Shijitan Hospital, Capital Medical University, 10th Tieyi Road, Haidian District, Beijing, 100038, People's Republic of China, Email miaohuihui@bjsjth.cn; litz@bjsjth.cn

Dear editor

We read with great interest the article by Yang et al,¹ which focused on postoperative sleep disturbance (PSD) in patients who underwent non-cardiac surgery and evaluated the perioperative risk factors associated with PSD, including sex, anxiety, satisfaction of ward environment, combined regional nerve block, postoperative nausea and vomiting, longer duration of stay in the postanesthesia care unit, a higher dose of midazolam or sufentanil, and a higher postoperative NRS score. However, while reading the article, we felt that some methodological aspects of the study should be discussed thoroughly.

First, although this study excluded patients with a long-term usage of sedatives, or those under antidepressants, drugs, or alcohol dependence, it did not consider if patients took hypnotic drugs from the day of admission to the night before surgery, including benzodiazepines, non-benzodiazepines, and melatonin. If patients in a clinic have sleep disturbance, doctors usually prescribe hypnotics for adequate rest before surgery. Few patients even opt for hypnosis by themselves. If this aspect is discussed, it may have a significant impact on the results of the study.

Second, during the preoperative period, majority of the patients experience a certain degree of sleep deprivation a night before surgery due to ward environment, fear of death, potential risks of surgery, postoperative recovery, and economic burden, all of which lead to preoperative sleep disturbance. Moreover, patients with preoperative sleep deprivation have a higher incidence of severe postoperative pain. Perioperative mental disease, sleep disturbance, and post-operative pain can result in a vicious circle for the patient.² Therefore, evaluating the patient's sleep quality before surgery is necessary. However, in the study, the patient's sleep quality one night before the surgery was not measured. Although PSQI was performed before surgery, it mainly measured the sleep quality of the patient in the past one month.³

Third, this study did not consider the impact of the timing of surgery on the circadian rhythm. The effects of surgery on the quality of postoperative sleep at different time periods are different. Song et al found that patients who underwent surgery in the evening had more severe sleep disorders than those who underwent surgery in the morning.⁴ Moreover, different time periods of medication administration have different effects on postoperative sleep quality. For example, Song et al found that the intraoperative use of dexmedetomidine during surgery in the day could improve sleep quality compared with its use at night.⁵

Finally, the information in the Abstract and text does not match; for example, the Abstract stated "a higher dose of midazolam and sufentanil is a perioperative risk factor associated with PSD." However, the text stated that "a lower dose of midazolam was associated with an increased probability of PSD in non-cardiac surgery patients." The nomogram also

© 2022 Yang et al. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms.php you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (http://www.dovepress.com/terms.php). confirmed that the lower the dose of midazolam and sufentanil, the more likely it is to cause postoperative sleep disturbance.

Disclosure

The authors report no conflicts of interest in this communication.

References

- 1. Yang S, Zhang Q, Xu Y, et al. Development and validation of nomogram prediction model for postoperative sleep disturbance in patients undergoing non-cardiac surgery: a prospective cohort study. *Nat Sci Sleep*. 2021;13:1473–1483. doi:10.2147/NSS.S319339
- 2. Dandan L, Xiao H, Sun Y, et al. Perioperative sleep disorder: a review. Front Med (Lausanne). 2021;8:640416. doi:10.3389/fmed.2021.640416
- 3. Buysse DJ, Reynolds CF, Monk TH, et al. The Pittsburgh sleep quality index: a new instrument for psychiatric practice and research. *Psychiatry Res.* 1989;28(2):193–213. doi:10.1016/0165-1781(89)90047-4
- 4. Bijia S, Yang L, Xiufei T, et al. Comparison of morning and evening operation under general anesthesia on intraoperative anesthetic requirement, postoperative sleep quality, and pain: a randomized controlled trial. *Nat Sci Sleep.* 2020;12:467–475. doi:10.2147/NSS.S257896
- 5. Song B, Li Y, Teng X, et al. The effect of intraoperative use of dexmedetomidine during the daytime operation vs the nighttime operation on postoperative sleep quality and pain under general anesthesia. *Nat Sci Sleep.* 2019;11:207–215. doi:10.2147/NSS.S225041

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Nature and Science of Sleep 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Nature and Science of Sleep editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Nature and Science of Sleep

Dovepress

Publish your work in this journal

Nature and Science of Sleep is an international, peer-reviewed, open access journal covering all aspects of sleep science and sleep medicine, including the neurophysiology and functions of sleep, the genetics of sleep, sleep and society, biological rhythms, dreaming, sleep disorders and therapy, and strategies to optimize healthy sleep. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/nature-and-science-of-sleep-journal

https://doi.org/10.2147/NSS.S357538