

Review Article

Using nuclear envelope mutations to explore age-related skeletal muscle weakness

Edmund Battey^{1,2}, Matthew J. Stroud² and  Julien Ochala^{1,3,4}

¹Centre of Human and Applied Physiological Sciences, School of Basic and Medical Biosciences, Faculty of Life Sciences and Medicine, King's College London, London, SE1 1UL, U.K.; ²British Heart Foundation Centre of Excellence, School of Cardiovascular Medicine and Sciences, King's College London, London, U.K.; ³Randall Centre for Cell and Molecular Biophysics, School of Basic and Medical Biosciences, Faculty of Life Sciences and Medicine, Guy's Campus, King's College London, SE1 1UL, U.K.; ⁴Department of Biomedical Sciences, University of Copenhagen, Blegdamsvej 3, 2200 Copenhagen N, Denmark

Correspondence: Julien Ochala (julien.ochala@kcl.ac.uk)



Skeletal muscle weakness is an important determinant of age-related declines in independence and quality of life but its causes remain unclear. Accelerated ageing syndromes such as Hutchinson–Gilford Progerin Syndrome, caused by mutations in genes encoding nuclear envelope proteins, have been extensively studied to aid our understanding of the normal biological ageing process. Like several other pathologies associated with genetic defects to nuclear envelope proteins including Emery–Dreifuss muscular dystrophy, Limb–Girdle muscular dystrophy and congenital muscular dystrophy, these disorders can lead to severe muscle dysfunction. Here, we first describe the structure and function of nuclear envelope proteins, and then review the mechanisms by which mutations in genes encoding nuclear envelope proteins induce premature ageing diseases and muscle pathologies. In doing so, we highlight the potential importance of such genes in processes leading to skeletal muscle weakness in old age.

Introduction

The human lifespan has increased substantially over the past half-century and this trend is projected to continue well into the 21st century [1]. This extension of the lifespan, however, has not been accompanied by an equivalent extension of the healthspan in old age; instead, morbidity has been extended, and independence and quality of life attenuated [2]. This increasing dependence on healthcare services has associated economic costs. Thus, a ‘managed compression of morbidity’ is necessary to address social and economic issues associated with an extending lifespan [3].

One important aspect of deteriorated healthspan and morbidity is skeletal muscle weakness, which is detrimental for independence and quality of life [4,5]. The causes of such progressive age-related generalised muscle dysfunction remain poorly understood, limiting development of potential therapeutic interventions to improve healthspan (pharmacologically or physically via personalised exercise regimens).

Interestingly, in recent years, premature ageing syndromes caused by genetic defects to nuclear envelope proteins (estimated prevalence of 1:4,000,000 to 1:10,000,000) have been increasingly studied as models to reveal the causes of the normal biological ageing process [6–8]. Among these is the Hutchinson–Gilford Progerin Syndrome (HGPS), which results in severe pathologies including heart disease, arteriosclerosis and an average life expectancy of ~13 years [9,10]. HGPS and other related disorders such as Emery–Dreifuss muscular dystrophy, Limb–Girdle muscular dystrophy and congenital muscular dystrophy can also lead to severe muscle pathologies resembling muscle weakness in old age, suggesting how modifications in the nuclear envelope may alter skeletal muscle development and function. Such deterioration in skeletal muscle behaviour is often not discussed in the clinical literature, partly because of the severity of the phenotype and the early age of death of patients, which makes the study of skeletal muscle function challenging; hence, its extent remains unclear.

Received: 14 April 2020
Revised: 04 August 2020
Accepted: 17 August 2020

Version of Record published:
26 August 2020




Figure 1. A muscle nucleus connected to sarcomeres through nuclear membrane and cytoskeletal proteins

Chromatin (grey) is organised into exposed, transcriptionally active sections (euchromatin) and tightly packed, transcriptionally repressed sections located at nucleoli and the nuclear periphery (heterochromatin). Heterochromatin associates with DNA interaction proteins BAF and HDAC3, which associate with inner nuclear membrane (INM) proteins such as Emerin, and the nuclear lamina, which interacts with SUN 1/2. SUN 1/2 bind to Nesprins to form the LINC complex, which links the nucleus to actin, as well as microtubules and Desmin via Kinesin and Plectin, respectively. Desmin binds to the Z-disk of sarcomeres, completing the connection between nuclei and myofibrils. Through this network of proteins, transcriptional activity is responsive to cytoskeletal changes; INM, inner nuclear membrane, ONM, outer nuclear membrane.

In this review, we start by briefly summarising what is known about nuclear membrane proteins. We then describe the mechanisms by which mutations in genes encoding these nuclear envelope proteins induce premature ageing diseases and muscle pathologies. Finally, we suggest how this knowledge could be used to unveil the key determinants of skeletal muscle weakness in old age.

The nuclear envelope and mechanotransduction

As the site of DNA transcription, the nucleus is responsible for orchestrating cell structure, function and adaptive responses [11]. Each nucleus is surrounded by an envelope, termed the nuclear envelope (NE), which sets a barrier between the cytoplasm and nuclear contents and consists of outer and inner nuclear membranes (ONM and




Figure 3. Hypothetical age-related defects in mechanotransduction in skeletal muscle fibres

(**Left panel**) In young, healthy skeletal muscle fibres, the LINC complex and its associated nuclear envelope proteins effectively transduce cytoskeletal forces to the nucleus to regulate signalling pathways, normal chromatin organisation and gene expression. (**Right panel**) In aged skeletal muscle, the content or distribution of the LINC complex and its associated proteins may be altered, leading to defective mechanotransduction and a gene transcription that may affect expression of contractile proteins.

Mutations in nuclear envelope proteins and related genetic diseases

Hutchinson–Gilford Progerin Syndrome (HGPS) is the most common premature ageing disorder [9,10]. HGPS is notably associated with mutations in the *LMNA* gene, which encodes the nuclear lamina components Lamin A/C, and the *ZMPSTE24* gene, which encodes the zinc metalloprotease 24 enzyme essential for Lamin A/C maturation [8,48]. These mutations disrupt Lamin A expression and function in nuclei of all cell types and can lead to altered forms known as Progerin or Prelamin A [8,49–52]. Mouse models of premature ageing where Lamin A/C and Prelamin A contents are modulated and are able to recapitulate patient phenotypes (i.e. cardiac and arteriosclerosis problems) as well as causing striking generalised skeletal muscle weakness [6,10,53–55]. Indeed, muscle-specific overexpression of human Progerin in mice significantly decreased muscle mass and myofibre size and halved grip strength [55]. Similarly, absence of the *ZMPSTE24* gene, resulting in the inability to form mature Lamin A from Prelamin A, leads to weaker and atrophic lower limb muscles and a reduction in the intrinsic force-generating capacity of myofibres and myofilaments [6,53].

Mutations in the *LMNA* gene also cause diseases not related to HGPS, collectively referred to as laminopathies, such as Emery–Dreifuss muscular dystrophy (EDMD), Limb–Girdle muscular dystrophy and *LMNA*-congenital muscular dystrophy. These are relatively well characterised clinically by skeletal muscular dystrophies and cardiomyopathies [56–61]. Other missense and nonsense mutations in genes encoding LINC complex and associated NE proteins have been identified in humans and have similar cardiac and skeletal muscle phenotypes. For instance, mutations have been found in *EMD*, encoding Emerin; *TMEM43* encoding Luma; and *SYNE1* and *SYNE2*, encoding Nesprin-1 and Nesprin-2, respectively [62–71].

Mechanisms underlying skeletal muscle weakness associated with mutations in genes encoding LINC complex and associated proteins

Lamin A-related disruption in nuclear architecture and mechanotransduction

The mechanisms of skeletal muscle dysfunction associated with mutations in genes for NE proteins are complex and in-depth mechanistic studies are currently lacking. The absence of Lamin A/C is usually characterised by aberrant nuclear morphology, nuclear mechanics and mechanotransduction [42,48,61,72]. Lamin A/C-deficient fibroblasts display increased nuclear deformation and altered expression of mechanosensitive genes *egr-1* and *iex-1* in response to mechanical strain and altered NF- κ B signalling [42]. When myoblasts from Lamin A-deficient mice are cultured to form *in vitro* muscle fibres, nuclear deformations, elongation, and rupture are evident and accompanied by DNA damage [73]. Additionally, cultured myoblasts from a human patient with a muscular dystrophy-causing mutation in the *LMNA* gene exhibit altered nuclear morphology, gene expression and increased cellular senescence [74].

Recently, it was shown that *LMNA*-related congenital muscular dystrophy patient fibroblasts and mouse myoblasts displayed altered Lamin A/C localisation, associated with altered expression and localisation of nuclear envelope proteins [61]. In contrast with control and EDMD cells, Lamin A/C was predominantly localised in the nucleoplasm, rather than forming the nuclear lamina at the nuclear periphery. Absence of peripheral Lamin A/C and its accumulation in the nucleoplasm was associated with attenuated expression and mislocalisation of several nuclear envelope transmembrane proteins, and mislocalisation of Nesprin-1 α . The authors postulated that the Lamin A/C mislocalisation and associated misregulation of nuclear envelope proteins may explain the greater severity of *LMNA*-related congenital muscular dystrophy compared with EDMD [61] (Figure 3).

Cells expressing Progerin, a truncated, permanently farnesylated form of Lamin A, which lacks the cleavage site for the enzyme *ZMPSTE24* to generate mature Lamin A, show evidence of impaired mechanotransduction, although the specific effects on skeletal muscle cells remain to be determined (Figure 2). Nuclei from Progerin-expressing fibroblasts display aberrant morphology, increased mechanical sensitivity and stiffness leading to senescence and cell death, indicating altered mechanotransduction [48,75]. Interestingly, it has been shown that replacement of mature Lamin A/C by Progerin in fibroblasts and iPSC-derived smooth muscle cells directly triggers premature senescence upon cell differentiation [76,77]. Progerin overexpression has been achieved in human myogenic cells, but whether they displayed a phenotype was not investigated [78].

Cells accumulating Prelamin A, a permanently farnesylated premature form of Lamin A, also exhibit aberrant nuclear morphology in fibroblasts and cardiomyocytes [54,79,80] (Figure 2). It is worth noting altered nuclear morphology has not directly been shown in skeletal muscle and this is an area for further research. However, aberrant nuclear morphology is associated with muscle degeneration, altered transcriptional activity, impaired contractility and muscle weakness, suggesting the accelerated ageing phenotype observed in HGPS fibroblasts could also occur in skeletal muscle cells [6,53,54,81]. A potential mechanism for this muscle weakness is reduced myonuclear number, resulting in a volume of cytoplasm too large for the transcriptional capability of each nucleus (termed the myonuclear domain theory, see [77–79,82–84]). Indeed, reduced myonuclear number was associated with reduced transcriptional activity and myosin content in mice muscle fibres with Prelamin A accumulation [53]. To compound the reduction in myonuclear number, altered nuclear integrity may contribute to defective mechanotransduction, further impairing the transcription of contractile proteins.

Lamin A-related alteration in Ca²⁺ metabolism

Recently, it was reported that Lamin A and Progerin interact with endoplasmic reticulum based proteins involved in Ca²⁺ transport and alter Ca²⁺ metabolism [55]. Immunoprecipitation in HGPS fibroblasts has shown that both Lamin A and Progerin interact with Sarcolipin, a protein involved in thermogenesis and Ca²⁺ homeostasis; however, Progerin binding is more potent [55,85,86]. Overexpression of Progerin in C2C12 myoblasts resulted in elevated cytosolic Ca²⁺ concentration and altered control of store operated Ca²⁺ entry [55]. Strikingly, mice lacking Lamin A/C upregulate Sarcolipin and skeletal muscle in mice expressing human Progerin have markedly altered Sarcolipin function and Ca²⁺ metabolism, together with ruffled nuclear morphology [55]. Since Ca²⁺ is essential component of muscle contraction, these studies implicate a role of Ca²⁺ metabolism in striated muscle tissue defects caused by mutations in the *LMNA* gene, an area which requires further research [87].

LINC complex protein-related changes in nuclear behaviour and mechanotransduction

As mentioned earlier, the LINC complex and its associated NE proteins are involved in muscle pathologies and may contribute to skeletal muscle defects with age. For example, fibroblasts expressing muscular dystrophy-associated variants of SUN1 and SUN2 have significant nuclear mispositioning, and this has also been shown in C2C12 myotubes expressing SUN1 variants [88]. In myotubes generated from a human patient with heterozygous SUN1 mutations, myonuclear organisation was similarly defective [88]. In fibroblasts from EDMD patients with Nesprin-1 and -2 mutations, nuclei exhibit aberrant morphologies [66]. Additionally, Nesprin-1 mutant C2C12 cells have defects in myoblast differentiation, specifically reduced fusion index, and down-regulated expression of myogenic transcription factors MyoD and Myogenin, and Myosin Heavy Chain [67]. Ablation of both Nesprin-1 and the muscle-specific isoform, Nesprin-1 α 2, affects expression of muscle differentiation genes, nuclear number and their positioning, increasing the myonuclear domain and impairing contractile protein expression [53,89–93].

Similarly, Emerin-deficient fibroblasts exhibit abnormal nuclear shape, stability and defective mechanotransduction [72]. In C2C12 cells, Emerin has been shown to interact with A- and B-type lamins, influencing nuclear architecture and stability and ultimately compromising mechanotransduction [94]. Furthermore, loss of Emerin in mice results in nuclear fragility, MAPK signalling activation and delayed induction of MyoD-related genes during muscle regeneration, although overall muscle function and regeneration were minimally affected [27,38,95,96]. The lack of effect on muscle function in Emerin-null mice is unlike the human EDMD phenotype, and may be explained by Lamina-associated polypeptide 1 (LAP1) compensation; LAP1 is highly expressed in mouse compared with human skeletal muscle, and reducing LAP1 expression in Emerin-null mice induced muscle abnormalities [97].

Thus, alterations to Emerin, Nesprin-1/2 and SUN1 in striated muscle tissue can result in nuclear mislocalisation, altered nuclear morphology and mechanotransduction, and impaired development and function. These findings emphasise the role of the LINC complex and associated NE proteins in muscle structure and function, and may therefore have implications for normal physiological ageing, discussed next.

Implications for normal muscle ageing

The above evidence suggests that alterations in the cytoskeletal-nuclear network could result in structural changes to the nucleus and impact gene expression in skeletal muscle. With age, compromised nuclear integrity, through alterations to LINC complex and associated proteins, may impair mechanotransduction. This may in turn lead to reduced expression of contractile proteins, resulting in a loss of muscle mass and function with age (Figure 3). To date, the only study which has investigated LINC complex proteins in human ageing showed SUN1 levels were increased in

fibroblasts from older (84–91 years old) compared with pre-pubescent (3–10 years old) individuals [98]. In line with these data, SUN1 accumulation in *LMNA* mutant mouse fibroblasts and human HGPS patient fibroblasts was associated with nuclear defects and cellular senescence, which were corrected upon reduction of the accumulated SUN1 [99]. Accumulation of SUN1 in human skeletal muscle may strengthen the association between the nucleus and the cytoskeleton, resulting in oversensitivity to cytoskeletal forces thereby causing aberrant mechanotransduction and negatively impacting cell function [98]. Alternatively, SUN1 mislocalisation, as shown in *LMNA*^{-/-} fibroblasts, could impair force transmission from the cytoskeleton to the nucleus [99]. Thus, altered SUN1 expression or distribution in skeletal muscle could impair mechanotransduction, elevating atrophic gene expression and suppressing the expression of genes encoding contractile proteins.

Future research

Current studies mainly focus on alterations to LINC complex components in fibroblasts. Future work using human skeletal muscle biopsies from young and old individuals will be pertinent to investigate mechanisms relating to both skeletal muscle function and dysfunction. Specifically:

- (1) Establishing whether protein levels and localisation of LINC complex and NE proteins are altered in skeletal muscle from young and old adults, to reveal potential associations between NE function and skeletal muscle ageing.
- (2) Understanding the effects of exercise on nuclear shape and LINC complex function.
- (3) Elucidating functional changes to nuclear mechanics in human skeletal muscle to provide insight into the role of NE proteins in aged skeletal muscle.

Furthermore, approaches to model muscular dystrophy in culture using cells from patients to create induced-pluripotent stem cell-derived artificial 3D skeletal muscle may shed new light on pathophysiological mechanisms underlying nuclear envelopopathies [100–102].

Concluding remarks

In this review, we have discussed the mechanisms by which mutations in genes encoding NE proteins induce premature ageing diseases and muscle pathologies, and suggested how this information may be used to unveil the key determinants of muscle weakness in old age. In these diseases, mutations in the LINC complex and its associated proteins cause aberrant nuclear morphology, defective mechanotransduction and muscle weakness. This has been extensively demonstrated in mouse and human fibroblasts, but to a limited degree in muscle cells. Whether changes to NE proteins contribute to defective mechanotransduction and muscle weakness in normal but aged muscle is understudied and is an exciting avenue for future research (Figure 3).

Competing Interests

The authors declare that there are no competing interests associated with the manuscript.

Funding

J.O. and E.B. are funded by the Medical Research Council of the UK [grant number MR/S023593/1]. M.J.S. is supported by British Heart Foundation Intermediate Fellowship [FS/17/57/32934] and King's BHF Centre for Award Excellence [RE/18/2/34213].

Open Access

Open access for this article was enabled by the participation of King's College London in an all-inclusive *Read & Publish* pilot with Portland Press and the Biochemical Society under a transformative agreement with JISC.

Acknowledgements

We thank the Medical Research Council (MR/S023593/1) and British Heart Foundation (FS/17/57/32934 and RE/18/2/34213) for support.

Abbreviations

BAF, barrier to autointegration factor; EDMD, Emery–Dreifuss muscular dystrophy; HDAC3, histone deacetylase 3; HGPS, Hutchinson–Gilford progerin syndrome; INM, inner nuclear membrane; KASH, Klarsicht, ANC-1, Syne Homology; LAP1,

lamina-associated polypeptide 1; LEM, lamina-associated polypeptide 2, Emerin, MAN1; LINC, linker of nucleoskeleton and cytoskeleton; NE, nuclear envelope; Nesprin, nuclear envelope spectrin repeat proteins; ONM, outer nuclear membrane; SUN, Sad1 and UNC-84 domain containing.

References

- United Nations (2019) World population prospects.
- Roser, M. (2019) Life Expectancy — Our World in Data.
- Brown, G.C. (2015) Living too long. *EMBO Rep.* **16**, 137–141, <https://doi.org/10.15252/embr.201439518>
- Trombetti, A., Reid, K.F., Hars, M., Herrmann, F.R., Pasha, E., Phillips, E.M. et al. (2016) Age-associated declines in muscle mass, strength, power, and physical performance: impact on fear of falling and quality of life. *Osteoporos. Int.* **27**, 463–471, <https://doi.org/10.1007/s00198-015-3236-5>
- McGregor, R.A., Cameron-Smith, D. and Poppitt, S.D. (2014) It is not just muscle mass: A review of muscle quality, composition and metabolism during ageing as determinants of muscle function and mobility in later life. *Longev. Heal.* **3**, 9, <https://doi.org/10.1186/2046-2395-3-9>
- Greising, S.M., Call, J.A., Lund, T.C., Blazar, B.R., Tolar, J. and Lowe, D.A. (2012) Skeletal muscle contractile function and neuromuscular performance in *Zmpste24*^{-/-} mice, a murine model of human progeria. *Age (Omaha)* **34**, 805–819, <https://doi.org/10.1007/s11357-011-9281-x>
- Maggi, L., Carboni, N. and Bernasconi, P. (2016) Skeletal Muscle Laminopathies: A Review of Clinical and Molecular Features. *Cells* **5**, 33, <https://doi.org/10.3390/cells5030033>
- Eriksson, M., Brown, W.T., Gordon, L.B., Glynn, M.W., Singer, J., Scott, L. et al. (2003) Recurrent de novo point mutations in lamin A cause Hutchinsonin-Gilford progeria syndrome. *Nature* **423**, 293–298, <https://doi.org/10.1038/nature01629>
- Hennekam, R.C.M. (2006) Hutchinson-Gilford progeria syndrome: Review of the phenotype. *Am. J. Med. Genet. Part A* **140**, 2603–2624, <https://doi.org/10.1002/ajmg.a.31346>
- Merideth, M.A., Gordon, L.B., Clauss, S., Sachdev, V., Smith, A.C.M., Perry, M.B. et al. (2008) Phenotype and Course of Hutchinsonin-Gilford Progeria Syndrome. *N. Engl. J. Med.* **358**, 592–604, <https://doi.org/10.1056/NEJMoa0706898>
- Wansink, D.G., Schul, W., Van der Kraan, I., Van Steensel, B., Van Driel, R. and De Jong, L. (1993) Fluorescent labeling of nascent RNA reveals transcription by RNA polymerase II in domains scattered throughout the nucleus. *J. Cell Biol.* **122**, 283–293, <https://doi.org/10.1083/jcb.122.2.283>
- Gerace, L. and Burke, B. (1988) Functional organization of the nuclear envelope. *Annu. Rev. Cell Biol.* **4**, 335–374, <https://doi.org/10.1146/annurev.cb.04.110188.002003>
- Wang, N., Tytell, J.D. and Ingber, D.E. (2009) Mechanotransduction at a distance: Mechanically coupling the extracellular matrix with the nucleus. *Nat. Rev. Mol. Cell Biol.* **10**, 75–82, <https://doi.org/10.1038/nrm2594>
- Chambliss, A.B., Khatau, S.B., Erdenberger, N., Robinson, D.K., Hodzic, D., Longmore, G.D. et al. (2013) The LINC-anchored actin cap connects the extracellular milieu to the nucleus for ultrafast mechanotransduction. *Sci. Rep.* **3**, 1–9, <https://doi.org/10.1038/srep01087>
- Fletcher, D.A. and Mullins, R.D. (2010) Cell mechanics and the cytoskeleton. *Nature* **463**, 485–492, <https://doi.org/10.1038/nature08908>
- Maniotis, A., Chen, C. and Ingber, D. (1997) Demonstration of mechanical connections between integrins, cytoskeletal filaments, and nucleoplasm that stabilize nuclear structure. **94**, 849–854
- Crisp, M., Liu, Q., Roux, K., Rattner, J.B., Shanahan, C., Burke, B. et al. (2006) Coupling of the nucleus and cytoplasm: Role of the LINC complex. *J. Cell Biol.* **172**, 41–53, <https://doi.org/10.1083/jcb.200509124>
- Rajgor, D. and Shanahan, C.M. (2013) Nesprins: from the nuclear envelope and beyond. *Expert Reviews in Molecular Medicine* **15**, 1–17
- Lombardi, M.L., Jaalouk, D.E., Shanahan, C.M., Burke, B., Roux, K.J. and Lammerding, J. (2011) The interaction between nesprins and sun proteins at the nuclear envelope is critical for force transmission between the nucleus and cytoskeleton. *J. Biol. Chem.* **286**, 26743–26753, <https://doi.org/10.1074/jbc.M111.233700>
- Banerjee, I., Zhang, J., Moore-Morris, T., Pfeiffer, E., Buchholz, K.S., Liu, A. et al. (2014) Targeted Ablation of Nesprin 1 and Nesprin 2 from Murine Myocardium Results in Cardiomyopathy, Altered Nuclear Morphology and Inhibition of the Biomechanical Gene Response. *PLoS Genet.* **10**, e1004114, <https://doi.org/10.1371/journal.pgen.1004114>
- Padmakumar, V.C., Libotte, T., Lu, W., Zaim, H., Abraham, S., Noegel, A.A. et al. (2005) The inner nuclear membrane protein Sun1 mediates the anchorage of Nesprin-2 to the nuclear envelope. *J. Cell Sci.* **118**, 3419–3430, <https://doi.org/10.1242/jcs.02471>
- Haque, F., Smallwood, D.T., Trembath, R.C., Dent, C.L., Shanahan, C.M., Shackleton, S. et al. (2006) SUN1 Interacts with Nuclear Lamin A and Cytoplasmic Nesprins To Provide a Physical Connection between the Nuclear Lamina and the Cytoskeleton. *Mol. Cell. Biol.* **26**, 3738–3751, <https://doi.org/10.1128/MCB.26.10.3738-3751.2006>
- Starr, D.A. (2011) KASH and SUN proteins. *Curr. Biol.* **21**, R414–R415, <https://doi.org/10.1016/j.cub.2011.04.022>
- Ivorra, C., Kubicek, M., González, J.M., Sanz-González, S.M., Álvarez-Barrientos, A., O'Connor, J.E. et al. (2006) A mechanism of AP-1 suppression through interaction of c-Fos with lamin A/C. *Genes Dev.* **20**, 307–320, <https://doi.org/10.1101/gad.349506>
- Ho, C.Y., Jaalouk, D.E., Vartiainen, M.K. and Lammerding, J. (2013) Lamin A/C and emerin regulate MKL1-SRF activity by modulating actin dynamics. *Nature* **497**, 507–511, <https://doi.org/10.1038/nature12105>
- Dechat, T., Pflieger, K., Sengupta, K., Shimi, T., Shumaker, D.K., Solimando, L. et al. (2008) Nuclear lamins: Major factors in the structural organization and function of the nucleus and chromatin. *Genes Dev.* **22**, 832–853, <https://doi.org/10.1101/gad.1652708>
- Muchir, A., Pavlidis, P., Bonne, G., Hayashi, Y.K. and Worman, H.J. (2007) Activation of MAPK in hearts of EMD null mice: Similarities between mouse models of X-linked and autosomal dominant Emery - Dreifuss muscular dystrophy. *Hum. Mol. Genet.* **16**, 1884–1895, <https://doi.org/10.1093/hmg/ddm137>
- Muchir, A., Wu, W., Choi, J.C., Iwata, S., Morrow, J., Homma, S. et al. (2012) Abnormal p38 α mitogen-activated protein kinase signaling in dilated cardiomyopathy caused by lamin A/C gene mutation. *Hum. Mol. Genet.* **21**, 4325–4333, <https://doi.org/10.1093/hmg/dds265>

- 29 Senior, A. and Gerace, L. (1988) Integral membrane proteins specific to the inner nuclear membrane and associated with the nuclear lamina. *J. Cell Biol.* **107**, 2029–2036, <https://doi.org/10.1083/jcb.107.6.2029>
- 30 Foisner, R. (2001) Inner nuclear membrane proteins and the nuclear lamina. *J. Cell Sci.* **114**, 3791–3792
- 31 Goldberg, M., Harel, A. and Gruenbaum, Y. (1999) The nuclear lamina: Molecular organization and interaction with chromatin. *Crit. Rev. Eukaryot. Gene Expr.* **9**, 285–293, <https://doi.org/10.1615/CritRevEukarGeneExpr.v9.i3-4.130>
- 32 Berk, J.M., Tiffit, K.E. and Wilson, K.L. (2013) The nuclear envelope LEM-domain protein emerin. *Nucleus* **4**, 298–314, <https://doi.org/10.4161/nucl.25751>
- 33 Wagner, N. and Krohne, G. (2007) LEM-Domain Proteins: New Insights into Lamin-Interacting Proteins. *Int. Rev. Cytol.* **261**, 1–46, [https://doi.org/10.1016/S0074-7696\(07\)61001-8](https://doi.org/10.1016/S0074-7696(07)61001-8)
- 34 Herrada, I., Bourgeois, B., Samson, C., Buendia, B., Worman, H.J. and Zinn-Justin, S. (2016) Purification and Structural Analysis of LEM-Domain Proteins. *Methods Enzymol.*, vol. 569, pp. 43–61, Academic Press Inc
- 35 Montes de Oca, R., Shoemaker, C.J., Gucek, M., Cole, R.N. and Wilson, K.L. (2009) Barrier-to-Autointegration Factor Proteome Reveals Chromatin-Regulatory Partners. *PLoS ONE* **4**, e7050, <https://doi.org/10.1371/journal.pone.0007050>
- 36 Demmerle, J., Koch, A.J. and Holaska, J.M. (2012) The nuclear envelope protein emerin binds directly to histone deacetylase 3 (HDAC3) and activates HDAC3 activity. *J. Biol. Chem.* **287**, 22080–22088, <https://doi.org/10.1074/jbc.M111.325308>
- 37 Holaska, J.M. and Wilson, K.L. (2007) An emerin “Proteome”: Purification of distinct emerin-containing complexes from HeLa cells suggests molecular basis for diverse roles including gene regulation, mRNA splicing, signaling, mechanosensing, and nuclear architecture. *Biochemistry* **46**, 8897–8908, <https://doi.org/10.1021/bi602636m>
- 38 Demmerle, J., Koch, A.J. and Holaska, J.M. (2013) Emerin and histone deacetylase 3 (HDAC3) cooperatively regulate expression and nuclear positions of MyoD, Myf5, and Pax7 genes during myogenesis. *Chromosom Res.* **21**, 765–779, <https://doi.org/10.1007/s10577-013-9381-9>
- 39 Muchir, A., Pavlidis, P., Decostre, V., Herron, A.J., Arimura, T., Bonne, G. et al. (2007) Activation of MAPK pathways links LMNA mutations to cardiomyopathy in Emery-Dreifuss muscular dystrophy. *J. Clin. Invest.* **117**, 1282–1293, <https://doi.org/10.1172/JCI29042>
- 40 Tapia, O., Fong, L.G., Huber, M.D., Young, S.G. and Gerace, L. (2015) Nuclear envelope protein Lem2 is required for mouse development and regulates MAP and AKT kinases. *PLoS ONE* **10**, 1–20, <https://doi.org/10.1371/journal.pone.0116196>
- 41 Markiewicz, E., Tilgner, K., Barker, N., Van De Wetering, M., Clevers, H., Dorobek, M. et al. (2006) The inner nuclear membrane protein Emerin regulates β -catenin activity by restricting its accumulation in the nucleus. *EMBO J.* **25**, 3275–3285, <https://doi.org/10.1038/sj.emboj.7601230>
- 42 Lammerding, J., Schulze, P., Takahashi, T., Kozlov, S., Sullivan, T., Kamm, R. et al. (2004) Lamin A/C deficiency causes defective nuclear mechanics and mechanotransduction. *J. Clin. Invest.* **113**, 370–378, <https://doi.org/10.1172/JCI200419670>
- 43 Houben, F., Ramaekers, F.C.S., Snoeckx, L.H.E.H. and Broers, J.L.V. (2007) Role of nuclear lamina-cytoskeleton interactions in the maintenance of cellular strength. *Biochim. Biophys Acta - Mol. Cell Res.* **1773**, 675–686, <https://doi.org/10.1016/j.bbamcr.2006.09.018>
- 44 Zwerger, M., Jaalouk, D.E., Lombardi, M.L., Isermann, P., Mauer mann, M., Dialynas, G. et al. (2013) Myopathic lamin mutations impair nuclear stability in cells and tissue and disrupt nucleo-cytoskeletal coupling. *Hum. Mol. Genet.* **22**, 2335–2349, <https://doi.org/10.1093/hmg/ddt079>
- 45 Poh, Y.C., Shevtsov, S.P., Chowdhury, F., Wu, D.C., Na, S., Dundr, M. et al. (2012) Dynamic force-induced direct dissociation of protein complexes in a nuclear body in living cells. *Nat. Commun.* **3**, 810–866, <https://doi.org/10.1038/ncomms1873>
- 46 Dupont, S. (2016) Role of YAP/TAZ in cell-matrix adhesion-mediated signalling and mechanotransduction. *Exp. Cell Res.* **343**, 42–53, <https://doi.org/10.1016/j.yexcr.2015.10.034>
- 47 Ramage, L., Nuki, G. and Salter, D.M. (2009) Signalling cascades in mechanotransduction: Cell-matrix interactions and mechanical loading. *Scand J. Med. Sci. Sport* **19**, 457–469, <https://doi.org/10.1111/j.1600-0838.2009.00912.x>
- 48 Goldman, R.D., Shumaker, D.K., Erdos, M.R., Eriksson, M., Goldman, A.E., Gordon, L.B. et al. (2004) Accumulation of mutant lamin A causes progressive changes in nuclear architecture in Hutchinson-Gilford progeria syndrome. *Proc. Natl Acad. Sci. U.S.A.* **101**, 8963–8968, <https://doi.org/10.1073/pnas.0402943101>
- 49 Young, S.G., Meta, M., Yang, S.H. and Fong, L.G. (2006) Prelamin A farnesylation and progeroid syndromes. *J. Biol. Chem.* **281**, 39741–39745, <https://doi.org/10.1074/jbc.R600033200>
- 50 De Sandre-Giovannoli, A., Bernard, R., Cau, P., Navarro, C., Amiel, J., Boccaccio, I. et al. (2003) Lamin A truncation in Hutchinson-Gilford progeria. *Science (80-)* **300**, 2055, <https://doi.org/10.1126/science.1084125>
- 51 Aebi, U., Cohn, J., Buhle, L. and Gerace, L. (1986) The nuclear lamina is a meshwork of intermediate-type filaments. *Nature* **323**, 560–564, <https://doi.org/10.1038/323560a0>
- 52 Davies, B.S.J., Fong, L.G., Yang, S.H., Coffinier, C. and Young, S.G. (2009) The Posttranslational Processing of Prelamin A and Disease. *Annu. Rev. Genomics Hum. Genet.* **10**, 153–174, <https://doi.org/10.1146/annurev-genom-082908-150150>
- 53 Levy, Y., Ross, J.A., Niglas, M., Snetkov, V.A., Lynham, S., Liao, C.-Y. et al. (2018) Prelamin A causes aberrant myonuclear arrangement and results in muscle fiber weakness. *JCI Insight* **3**, 1–18, <https://doi.org/10.1172/jci.insight.120920>
- 54 Pendás, A.M., Zhou, Z., Cadiñanos, J., Freije, J.M.P., Wang, J., Hultenby, K. et al. (2002) Defective prelamin A processing and muscular and adipocyte alterations in Zmpste24 metalloproteinase-deficient mice. *Nat. Genet.* **31**, 94–99, <https://doi.org/10.1038/ng871>
- 55 Wang, W.P., Wang, J.Y., Lin, W.H., Kao, C.H., Hung, M.C., Teng, Y.C. et al. (2020) Progerin in muscle leads to thermogenic and metabolic defects via impaired calcium homeostasis. *Aging Cell* **19**, e13090, <https://doi.org/10.1111/acer.13090>
- 56 Di Barletta, M.R., Ricci, E., Galluzzi, G., Tonali, P., Mora, M., Morandi, L. et al. (2000) Different mutations in the LMNA gene cause autosomal dominant autosomal recessive Emery-Dreifuss muscular dystrophy. *Am. J. Hum. Genet.* **66**, 1407–1412, <https://doi.org/10.1086/302869>
- 57 Emery, A.E.H. (1989) Emery-Dreifuss syndrome. *J. Med. Genet.* **26**, 637–641, <https://doi.org/10.1136/jmg.26.10.637>
- 58 Raharjo, W.H., Enarson, P., Sullivan, T., Stewart, C.L. and Burke, B. (2001) Nuclear envelope defects associated with LMNA mutations cause dilated cardiomyopathy and Emery-Dreifuss muscular dystrophy. *J. Cell Sci.* **114**, 4447–4457

- 59 Muchir, A. (2000) Identification of mutations in the gene encoding lamins A/C in autosomal dominant limb girdle muscular dystrophy with atrioventricular conduction disturbances (LGMD1B). *Hum. Mol. Genet.* **9**, 1453–1459, <https://doi.org/10.1093/hmg/9.9.1453>
- 60 Maggi, L., D'Amico, A., Pini, A., Sivo, S., Pane, M., Ricci, G. et al. (2014) LMNA-associated myopathies: The Italian experience in a large cohort of patients. *Neurology* **83**, 1634–1644, <https://doi.org/10.1212/WNL.0000000000000934>
- 61 Bertrand, A.T., Brull, A., Azibani, F., Benarroch, L., Chikhaoui, K., Stewart, C.L. et al. (2020) Lamin A/C Assembly Defects in LMNA-Congenital Muscular Dystrophy Is Responsible for the Increased Severity of the Disease Compared with Emery-Dreifuss Muscular Dystrophy. *Cells* **9**, 844, <https://doi.org/10.3390/cells9040844>
- 62 Manilal, S., Nguyen, T.M., Sewry, C.A. and Morris, G.E. (1996) The Emery-Dreifuss muscular dystrophy protein, emerin, is a nuclear membrane protein. *Hum. Mol. Genet.* **5**, 801–808, <https://doi.org/10.1093/hmg/5.6.801>
- 63 Nagano, A., Koga, R., Ogawa, M., Kurano, Y., Kawada, J., Okada, R. et al. (1996) Emerin deficiency at the nuclear membrane in patients with Emery-Dreifuss muscular dystrophy. *Nat. Genet.* **12**, 254–259, <https://doi.org/10.1038/ng0396-254>
- 64 Beaudin, M., Gamache, P.-L., Gros-Louis, F. and Dupré, N. (1993) *SYNE1 Deficiency*, University of Washington, Seattle
- 65 Chen, Z., Ren, Z., Mei, W., Ma, Q., Shi, Y., Zhang, Y. et al. (2017) A novel SYNE1 gene mutation in a Chinese family of Emery-Dreifuss muscular dystrophy-like. *BMC Med. Genet.* **18**, 63, <https://doi.org/10.1186/s12881-017-0424-5>
- 66 Zhang, Q., Bethmann, C., Worth, N.F., Davies, J.D., Wasner, C., Feuer, A. et al. (2007) Nesprin-1 and -2 are involved in the pathogenesis of Emery - Dreifuss muscular dystrophy and are critical for nuclear envelope integrity. *Hum. Mol. Genet.* **16**, 2816–2833, <https://doi.org/10.1093/hmg/ddm238>
- 67 Zhou, C., Li, C., Zhou, B., Sun, H., Koullourou, V., Holt, I. et al. (2017) Novel nesprin-1 mutations associated with dilated cardiomyopathy cause nuclear envelope disruption and defects in myogenesis. *Hum. Mol. Genet.* **26**, 2258–2276, <https://doi.org/10.1093/hmg/ddx116>
- 68 Stroud, M.J., Banerjee, I., Veevers, J. and Chen, J. (2014) Linker of Nucleoskeleton and Cytoskeleton Complex Proteins in Cardiac Structure, Function, and Disease. *Circ. Res.* **114**, 538–548, <https://doi.org/10.1161/CIRCRESAHA.114.301236>
- 69 Stroud, M.J., Fang, X., Zhang, J., Guimarães-Camboa, N., Veevers, J., Dalton, N.D. et al. (2018) Luma is not essential for murine cardiac development and function. *Cardiovasc. Res.* **114**, 378–388, <https://doi.org/10.1093/cvr/cvx205>
- 70 Christensen, A., Andersen, C., Tybjærg-Hansen, A., Haunso, S. and Svendsen, J. (2011) Mutation analysis and evaluation of the cardiac localization of TMEM43 in arrhythmogenic right ventricular cardiomyopathy. *Clin. Genet.* **80**, 256–264, <https://doi.org/10.1111/j.1399-0004.2011.01623.x>
- 71 Merner, N.D., Hodgkinson, K.A., Haywood, A.F.M., Connors, S., French, V.M., Drenckhahn, J.D. et al. (2008) Arrhythmogenic Right Ventricular Cardiomyopathy Type 5 Is a Fully Penetrant, Lethal Arrhythmic Disorder Caused by a Missense Mutation in the TMEM43 Gene. *Am. J. Hum. Genet.* **82**, 809–821, <https://doi.org/10.1016/j.ajhg.2008.01.010>
- 72 Lammerding, J., Hsiao, J., Schulze, P.C., Kozlov, S., Stewart, C.L. and Lee, R.T. (2005) Abnormal nuclear shape and impaired mechanotransduction in emerin-deficient cells. *J. Cell Biol.* **170**, 781–791, <https://doi.org/10.1083/jcb.200502148>
- 73 Earle, A.J., Kirby, T.J., Fedorchak, G.R., Isermann, P., Patel, J., Iruvanti, S. et al. (2018) Mutant lamins cause mechanically-induced nuclear envelope rupture, DNA damage, and DNA-PK activation in muscle. *BioRxiv*, 364778
- 74 Kandert, S., Wehnert, M., Müller, C.R., Buendia, B. and Dabauvalle, M.C. (2009) Impaired nuclear functions lead to increased senescence and inefficient differentiation in human myoblasts with a dominant p.R545C mutation in the LMNA gene. *Eur. J. Cell Biol.* **88**, 593–608, <https://doi.org/10.1016/j.ejcb.2009.06.002>
- 75 Verstraeten, V.L.R.M., Ji, J.Y., Cummings, K.S., Lee, R.T. and Lammerding, J. (2008) Increased mechanosensitivity and nuclear stiffness in Hutchinson-Gilford progeria cells: Effects of farnesyltransferase inhibitors. *Aging Cell* **7**, 383–393, <https://doi.org/10.1111/j.1474-9726.2008.00382.x>
- 76 Cao, K., Blair, C.D., Faddah, D.A., Kieckhafer, J.E., Olive, M., Erdos, M.R. et al. (2011) Progerin and telomere dysfunction collaborate to trigger cellular senescence in normal human fibroblasts. *J. Clin. Invest.* **121**, 2833–2844, <https://doi.org/10.1172/JCI43578>
- 77 Liu, G.H., Barkho, B.Z., Ruiz, S., Diep, D., Qu, J., Yang, S.L. et al. (2011) Recapitulation of premature ageing with iPSCs from Hutchinson-Gilford progeria syndrome. *Nature* **472**, 221–227, <https://doi.org/10.1038/nature09879>
- 78 Luo, Y.B., Mitrant, C., Adams, A.M., Johnsen, R.D., Fletcher, S., Mastaglia, F.L. et al. (2014) Antisense oligonucleotide induction of progerin in human myogenic cells. *PLoS ONE* **9**, 98306, <https://doi.org/10.1371/journal.pone.0098306>
- 79 Brayson, D., Frustaci, A., Verardo, R., Chimenti, C., Russo, M.A., Hayward, R. et al. (2019) Prelamin A mediates myocardial inflammation in dilated and HIV-Associated cardiomyopathies. *JCI Insight* **4**, e126315, <https://doi.org/10.1172/jci.insight.126315>
- 80 Young, S.G., Fong, L.G. and Michaelis, S. (2005) Prelamin A, Zmpste24, Misshapen Cell Nuclei, and Progeria—New Evidence Suggesting that Protein Farnesylation Could Be Important for Disease Pathogenesis. *J. Lipid Res.* **46**, 2531–2558, <https://doi.org/10.1194/jlr.R500011-JLR200>
- 81 Bergo, M.O., Gavino, B., Ross, J., Schmidt, W.K., Hong, C., Kendall, L.V. et al. (2002) Zmpste24 deficiency in mice causes spontaneous bone fractures, muscle weakness, and a prelamin A processing defect. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 13049–13054, <https://doi.org/10.1073/pnas.192460799>
- 82 Allen, D.L., Roy, R.R. and Reggie Edgerton, V. (1999) Myonuclear domains in muscle adaptation and disease. *Muscle Nerve* **22**, 1350–1360, [https://doi.org/10.1002/\(SICI\)1097-4598\(199910\)22:10%3c1350::AID-MUS3%3e3.0.CO;2-8](https://doi.org/10.1002/(SICI)1097-4598(199910)22:10%3c1350::AID-MUS3%3e3.0.CO;2-8)
- 83 Hall, Z.W. and Ralston, E. (1989) Nuclear domains in muscle cells. *Cell* **59**, 771–772, [https://doi.org/10.1016/0092-8674\(89\)90597-7](https://doi.org/10.1016/0092-8674(89)90597-7)
- 84 Pavlath, G.K., Rich, K., Webster, S.G. and Blau, H.M. (1989) Localization of muscle gene products in nuclear domains. *Nature* **337**, 570–573, <https://doi.org/10.1038/337570a0>
- 85 Bal, N., Maurya, S. and Sopariwala, D. (2012) Medicine SS-N, 2012 undefined. Sarcolipin is a newly identified regulator of muscle-based thermogenesis in mammals. *Nat Med.* **18**, <https://doi.org/10.1038/nm.2897>
- 86 Pant, M., Bal, N.C. and Periasamy, M. (2016) Sarcolipin: A Key Thermogenic and Metabolic Regulator in Skeletal Muscle. *Trends Endocrinol. Metab.* **27**, 881–892, <https://doi.org/10.1016/j.tem.2016.08.006>
- 87 Rivera-Torres, J., Calvo, C.J., Llach, A., Guzmán-Martínez, G., Caballero, R., González-Gómez, C. et al. (2016) Cardiac electrical defects in progeroid mice and Hutchinson-Gilford progeria syndrome patients with nuclear lamina alterations. *Proc. Natl. Acad. Sci. U.S.A.* **113**, E7250–E7259, <https://doi.org/10.1073/pnas.1603754113>

- 88 Meinke, P., Mattioli, E., Haque, F., Antoku, S., Columbaro, M., Straatman, K.R. et al. (2014) Muscular Dystrophy-Associated SUN1 and SUN2 Variants Disrupt Nuclear-Cytoskeletal Connections and Myonuclear Organization. *PLoS Genet.* **10**, e1004605, <https://doi.org/10.1371/journal.pgen.1004605>
- 89 Stroud, M.J., Feng, W., Zhang, J., Veevers, J., Fang, X., Gerace, L. et al. (2017) Nesprin 1 α 2 is essential for mouse postnatal viability and nuclear positioning in skeletal muscle. *J. Cell Biol.* **216**, 1915–1924, <https://doi.org/10.1083/jcb.201612128>
- 90 Gimpel, P., Lee, Y.L., Sobota, R.M., Calvi, A., Koullourou, V., Patel, R. et al. (2017) Nesprin-1 α -Dependent Microtubule Nucleation from the Nuclear Envelope via Akap450 Is Necessary for Nuclear Positioning in Muscle Cells. *Curr. Biol.* **27**, 2999.e9–3009.e9, <https://doi.org/10.1016/j.cub.2017.08.031>
- 91 Zhang, J., Felder, A., Liu, Y., Guo, L.T., Lange, S., Dalton, N.D. et al. (2009) Nesprin 1 is critical for nuclear positioning and anchorage. *Hum. Mol. Genet.* **19**, 329–341, <https://doi.org/10.1093/hmg/ddp499>
- 92 Zhang, X., Xu, R., Zhu, B., Yang, X., Ding, X., Duan, S. et al. (2007) Syne-1 and Syne-2 play crucial roles in myonuclear anchorage and motor neuron innervation. *Development* **134**, 901–908, <https://doi.org/10.1242/dev.02783>
- 93 Puckelwartz, M.J., Kessler, E., Zhang, Y., Hodzic, D., Randles, K.N., Morris, G. et al. (2009) Disruption of nesprin-1 produces an Emery Dreifuss muscular dystrophy-like phenotype in mice. *Hum. Mol. Genet.* **18**, 607–620, <https://doi.org/10.1093/hmg/ddn386>
- 94 Fairley, E.A.L., Kendrick-Jones, J. and Ellis, J.A. (1999) The Emery-Dreifuss muscular dystrophy phenotype arises from aberrant targeting and binding of emerin at the inner nuclear membrane. *J. Cell Sci.* **112**, 4801
- 95 Melcon, G., Koslov, S., Cutler, D.A., Sullivan, T., Hernandez, L., Zhao, P. et al. (2006) Loss of emerin at the nuclear envelope disrupts the Rb1/E2F and MyoD pathways during muscle regeneration. *Hum. Mol. Genet.* **15**, 637–651, <https://doi.org/10.1093/hmg/ddi479>
- 96 Ozawa, R., Hayashi, Y.K., Ogawa, M., Kurokawa, R., Matsumoto, H., Noguchi, S. et al. (2006) Emerin-lacking mice show minimal motor and cardiac dysfunctions with nuclear-associated vacuoles. *Am. J. Pathol.* **168**, 907–917, <https://doi.org/10.2353/ajpath.2006.050564>
- 97 Shin, J.Y., Méndez-López, I., Wang, Y., Hays, A.P., Tanji, K., Lefkowitz, J.H. et al. (2013) Lamina-associated polypeptide-1 interacts with the muscular dystrophy protein emerin and is essential for skeletal muscle maintenance. *Dev. Cell* **26**, 591–603, <https://doi.org/10.1016/j.devcel.2013.08.012>
- 98 Chang, W., Wang, Y., Gant Luxton, G.W., Östlund, C., Worman, H.J. and Gundersen, G.G. (2019) Imbalanced nucleocytoskeletal connections create common polarity defects in progeria and physiological aging. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 3578–3583, <https://doi.org/10.1073/pnas.1809683116>
- 99 Chen, C.Y., Chi, Y.H., Mutalif, R.A., Starost, M.F., Myers, T.G., Anderson, S.A. et al. (2012) Accumulation of the inner nuclear envelope protein Sun1 is pathogenic in progeric and dystrophic laminopathies. *Cell* **149**, 565–577, <https://doi.org/10.1016/j.cell.2012.01.059>
- 100 Maffioletti, S.M., Sarcar, S., Henderson, A.B.H., Mannhardt, I., Pinton, L., Moyle, L.A. et al. (2018) Three-Dimensional Human iPSC-Derived Artificial Skeletal Muscles Model Muscular Dystrophies and Enable Multilineage Tissue Engineering. *Cell Rep.* **23**, 899–908, <https://doi.org/10.1016/j.celrep.2018.03.091>
- 101 van der Wal, E., Herrero-Hernandez, P., Wan, R., Broeders, M., in 't Groen, S.L.M., van Gestel, T.J.M. et al. (2018) Large-Scale Expansion of Human iPSC-Derived Skeletal Muscle Cells for Disease Modeling and Cell-Based Therapeutic Strategies. *Stem Cell Rep.* **10**, 1975–1990, <https://doi.org/10.1016/j.stemcr.2018.04.002>
- 102 Zhou, T., Benda, C., Duzinger, S., Huang, Y., Li, X., Li, Y. et al. (2011) Generation of induced pluripotent stem cells from urine. *J. Am. Soc. Nephrol.* **22**, 1221–1228, <https://doi.org/10.1681/ASN.2011010106>