

## Original Article



# Association of Early Favipiravir Use with Reduced COVID-19 Fatality among Hospitalized Patients

Ercan Karatas <sup>1</sup>, Lacin Aksoy <sup>2</sup>, and Ersin Ozaslan <sup>3</sup>

<sup>1</sup>Department of Internal Medicine, Tuzla State Hospital, Istanbul, Turkey

<sup>2</sup>Department of Family Medicine, Marmara University, Istanbul, Turkey

<sup>3</sup>Department of Medical Oncology, Acibadem Kayseri Hospital, Kayseri, Turkey

## OPEN ACCESS

Received: Dec 21, 2020

Accepted: Mar 23, 2021

### Corresponding Author:

Ercan Karatas, MD

Department of Internal Medicine, Tuzla State Hospital, Istanbul 34959, Turkey.

Tel: +905354405744

Fax: +902164940959

E-mail: canertaskara@hotmail.com

Copyright © 2021 by The Korean Society of Infectious Diseases, Korean Society for Antimicrobial Therapy, and The Korean Society for AIDS

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

### ORCID iDs

Ercan Karatas

<https://orcid.org/0000-0002-3133-4199>

Lacin Aksoy

<https://orcid.org/0000-0002-4105-0595>

Ersin Ozaslan

<https://orcid.org/0000-0002-8333-676X>

### Conflict of Interest

No conflicts of interest.

### Author Contributions

Conceptualization: EK. Data curation: EK. Formal analysis: LA. Investigation: LA. Methodology: EÖ. Project administration: EK. Resources: EK. Software: LA. Supervision: EÖ. Validation: LA. Visualization: EK. Writing - original draft: EK. Writing - review & editing: LA.

## ABSTRACT

**Background:** The antiviral agent favipiravir is an RNA-dependent RNA polymerase (RdRp) inhibitor.

**Materials and Methods:** We examined patients with a clinical, laboratory, and radiological diagnosis of severe coronavirus disease 2019 (COVID-19) pneumonia. We investigated the effect of administering enteral favipiravir at a 2 × 1,600 mg loading dose and 2 × 600 mg maintenance dose for 5 days in addition to the standard COVID-19 treatment.

**Results:** In total, 180 patients, who were hospitalized at the Istanbul Tuzla State Hospital and received favipiravir treatment between March 20, 2020 and May 30, 2020, were examined. Of these, 47 patients died. Thirty-three of the patients who died were aged over 65 years (70%), indicating that fatality was higher in elderly patients. Most of those who died had at least one comorbidity. Of the 101 patients who initiated favipiravir within ≤3 days of hospitalization, 17 died (17%). Of the 79 patients who initiated favipiravir after >3 days of hospitalization, 30 died (38%) ( $P = 0.002$ ).

**Conclusion:** We found that initiation of favipiravir within the first 72 h after the onset of disease symptoms reduced fatality in patients with COVID-19.

**Keywords:** COVID-19; Favipiravir; Time-to-treatment; Hospitalization

## INTRODUCTION

The novel severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has spread globally and caused an unprecedented pandemic, with 150 million cases and over 3 million deaths in a year. SARS-CoV-2 is an enveloped, positive-sense, single-stranded RNA virus responsible for causing coronavirus disease 2019 (COVID-19) [1]. Although various strategies have been developed to treat COVID-19, there is still no standard treatment [2]. Antivirals that have been used to treat other viral infections are also being tried against COVID-19 [3].

Favipiravir is an oral broad-spectrum anti-viral agent that selectively and potently inhibits the RNA-dependent RNA polymerase (RdRp) of RNA viruses [4]. Favipiravir has shown potent *in vitro* activity against SARS-CoV-2 [5]. Consequently, favipiravir has been added to the options available for treating COVID-19 [6]. It was determined that favipiravir treatment reduced viral

clearance and provided clinical and radiological improvement [7]. In Turkey, Ministry of Health suggested the usage of favipiravir as an antiviral according to clinical researches in Europe.

In our study, we investigated whether the time when favipiravir is initiated influences the risk of fatality in hospitalized patients with severe COVID-19 pneumonia.

## MATERIAL AND METHODS

### 1. Patients

This study was carried out at Tuzla State Hospital, Istanbul, Turkey, with approval of the Marmara University Faculty of Medicine Ethics Committee after obtaining permission from the Scientific Research Unit of the Ministry of Health and the Istanbul Provincial Health Directorate (Marmara University Faculty of Medicine Research Ethics Committee; approval no.: E.70737436-050.01.04-2000202899; date: September 29, 2020). Patients with severe COVID-19 pneumonia, who were admitted to Tuzla State Hospital between March 20, 2020 and May 30, 2020 and treated with favipiravir at any stage of their treatment, were examined retrospectively. All patients with their or their family consent used favipiravir. The study conducted according to their health records with no personal information. Therefore, informed consent not asked to the patients whether their families. The study included patients aged over 18 years who were not pregnant, had severe pneumonia (respiratory rate  $>30/\text{min}$ ) with or without comorbidities and/or severe respiratory distress (dyspnea or use of extra respiratory muscles) and/or fingertip oxygen saturation  $<90\%$  ( $\text{PaO}_2/\text{FiO}_2 <300$  in patients receiving oxygen), were treated with favipiravir at any stage of their treatment, and had bilateral multi-lobar ground glass opacities observed in a computed tomography (CT) scan of the lungs. During the first half of the study period, according to the COVID-19 treatment protocol followed in our country, favipiravir could be administered to patients in whom standard COVID-19 treatment was not successful. Therefore, favipiravir was started later in some of our patients. However, the treatment protocol changed in the second half of our study, and favipiravir was used from the beginning of the treatment.


### 2. Statistical Analysis

The research data were collected retrospectively through the Tuzla State Hospital registration system. Descriptive statistics were used to analyze categorical variables, represented as numbers, and continuous variables, represented as mean  $\pm$  SD or median and interquartile range. The results were compared using *t*-test with the software SPSS 22.0 Statistics (IBM, Armonk, NY, USA).

## RESULTS

We examined 180 patients who were hospitalized with severe pneumonia in the Istanbul Tuzla State Hospital and treated with favipiravir. The median age of the patients was  $59 \pm 17.4$ . Of these, 108 patients were aged  $\leq 65$  years. Fifteen of these patients died, indicating a fatality rate of 12% (15/108). In addition, 32 of 72 patients who were aged  $>65$  years died; the fatality rate of these patients was 45% (32/72;  $P < 0.001$ ). This indicates that fatality increased with an increase in age. In total, 47 patients died, of whom 30 were male and 17 were female.


We observed that as the number of comorbidities increased, the fatality rate also increased. In our study, 113 of 180 patients (63% of all patients) had at least one comorbidity (Fig. 1). The



**Figure 1.** Relationship between comorbidities and survival (X: Types of comorbidities, Y: Number of patients). HT, hypertension; CAD, coronary artery disease; PAH, peripheral artery disease; HL, hyperlipidemia; CHF, congestive heart failure; DM, diabetes mellitus; COPD, chronic obstructive pulmonary disease.

main comorbidities were hypertension (79 patients), hyperlipidemia (57 patients), diabetes mellitus (52 patients), and coronary artery disease (33 patients). Of the 47 patients who died, 42 had at least one comorbid disease (89%). Cough (94 patients, 52%), dyspnea (86 patients, 47%), and fever (82 patients, 45%) were the topmost initial clinical findings (Fig. 2).

Among the hematological parameters, lymphopenia (lymphocyte count  $<1 \times 10^9/L$ ), low hemoglobin levels (hemoglobin  $<12 \text{ g/dL}$ ), leukocytosis (leukocyte count  $>7.5 \times 10^9/L$ ) ( $P = 0.001$ ) and neutrophilia (neutrophil count  $>7.5 \times 10^9/L$ ) ( $P = 0.003$ ) are associated with disease severity. In this study, disease severity and platelet count were not correlated ( $P > 0.5$ ). C-reactive protein (CRP) levels were increased in the patients; survivors had mean CRP values of about 85 mg/L, while non-survivors had mean values of 122 mg/L, which correlates with disease severity and prognosis ( $P = 0.004$ ). Other predictors studied were the serum ferritin and lactate dehydrogenase (LDH) levels. High levels of ferritin ( $>200 \mu\text{g/L}$ ) and LDH ( $>260 \text{ units/L}$ ) are associated with disease severity ( $P = 0.001$ ). Survivors showed significantly higher plasma D-dimer levels than the non-survivors. Among the coagulation parameters studied, D-dimer level  $>1,000 \text{ ng/mL}$  was a predictor of fatality ( $P = 0.0003$ ). Clinicians should consider low lymphocyte, leukocyte, and hemoglobin counts as well as the CRP, LDH, ferritin, and



**Figure 2.** Clinical findings (X: Number of patients, Y: Symptoms).

**Table 1.** Laboratory findings

	Survivors (n = 133)	Non-survivors (n = 47)	Normal range	P-value
Age, mean (IQR), years	54 (45 – 67)	71 (60 – 82)		
Male	77	30		
Female	56	17		
Duration of symptoms, days	3.11 (1 – 5)	5.72 (2 – 7)		
White blood cells, median (IQR), $\times 10^9/L$	7.4 (4.8 – 8.8)	9.2 (6.3 – 11.2)	4 – 11	0.001
Hemoglobin count, median (IQR), g/dL	12.7 (11.6 – 13.9)	9.2 (9.8 – 12.5)	Males: 13 – 18 Females: 11.5 – 16.5	0.001
Lymphocyte count, median (IQR), $\times 10^9/L$	1.17 (0.8 – 1.4)	0.75 (0.4 – 1)	1.5 – 4.5	0.001
Neutrophil count, median (IQR), $\times 10^9/L$	5.5 (3 – 6.5)	7.94 (4.7 – 10.1)	2.0 – 7.5	0.003
Platelet count, median (IQR), $\times 10^9/L$	230 (151 – 274)	236 (175 – 256)	150 – 450	>0.5
C-reactive protein, median (IQR), mg/L	85 (23 – 108)	122 (111 – 233)	<8.0 mg/L	0.004
Ferritin, median (IQR), $\mu g/L$	330 (94 – 339)	812 (218 – 1,043)	Females: 10 – 200 $\mu g/L$ ; Males: 30 – 300 $\mu g/L$	0.001
Lactate dehydrogenase, median (IQR), units/L	333 (222 – 369)	528 (368 – 640)	110 – 210	0.001
Creatinine, mg/dL	0.9 (0.7 – 1.0)	1.3 (0.7 – 1.7)	<1.2	0.004
ALT, U/L	31 (22 – 39)	33 (17 – 35)	<24	>0.5
Na, mEq/L	135 (133 – 139)	136 (133 – 140)	135 – 145	>0.5
D-Dimer, median (IQR), ng/mL	830 (430 – 1,210)	2,897 (1,070 – 3,185)	<500	0.0003


P-values were calculated using Pearson's chi-square test.  
IQR, interquartile range; ALT, alanine aminotransferase.

**Table 2.** Respiratory support

Favipiravir initiation time	$\leq 72$ h (n = 133)	$> 72$ h (n = 47)	P-value
Supplemental oxygen	104	2	0.0001
High-flow nasal oxygen	2	7	0.02
Non-invasive mechanical ventilation	4	3	0.3
Invasive ventilation	23	35	0.001

D-dimer serum levels. These serum values can be used in risk stratification to predict severe and fatal COVID-19 in hospitalized patients (Table 1).

Moreover, the need for advanced respiratory support increased in patients who initiated favipiravir after  $> 72$  h (Table 2). When we evaluated the time-to-treatment after hospitalization, we found that the fatality rate was lower in those who received favipiravir treatment within the first 72 h of hospitalization ( $P = 0.002$ ; Fig. 3).



**Figure 3.** Relationship between the time of initiating favipiravir and survival (X: Number of survivors and non-survivors, Y: Number of patients).

## DISCUSSION

COVID-19 is a multisystemic inflammatory disease [8]. Currently, there is no universal standard for drug treatment, and various combination therapies are being administered [9]. Our standard treatment involves the administration of antivirals (favipiravir), broad spectrum antibiotics, low-molecular weight heparin, antithrombotic drugs, corticosteroids, and oxygen therapy [10]. For severe COVID-19 pneumonia, cytokine storm, macrophage activation syndrome, and acute respiratory distress syndrome, convalescent plasma, interleukin-6 inhibitors, and anakinra, an interleukin-1 receptor antagonist, are added to the treatment [11]. Combination therapy is currently customized for each patient.

Based on other viral infections, early antiviral therapy is considered more useful in the treatment of COVID-19. The benefit of early antiviral therapy in the treatment of influenza has been observed previously [12, 13]. SARS, Middle East respiratory syndrome (MERS), and COVID-19 are caused by viruses with similar genetic structures [14]. The benefit of early antiviral treatment has also been reported in SARS and MERS [15]. Several clinical trials on COVID-19 have demonstrated that early antiviral therapy is necessary and beneficial.

For instance, Magleby et al. conducted a retrospective cohort study in 678 hospitalized patients with COVID-19 in New York. They demonstrated that high viral load was independently associated with mortality, and that early antiviral therapy prevents increased viral load in the lower respiratory tract by inactivating the viruses in the upper respiratory tract [16]. Weiss et al. determined that the viral load in the upper respiratory tract peaks early in mild cases of COVID-19 (4 days) compared to that in moderate to severe cases (8 days). Similarly, viral load in the lower respiratory tract peaks early in mild cases (6 days) compared to that in moderate to severe cases (11 days). This indicates the importance of early diagnosis and the time to initiate treatment [17].

Goyal et al. reported that if a patient receives antiviral therapy in the early phase of infection, there are high chances that the duration of shedding and intensity of the effector immune response may decrease; however, there may be a limited impact on viral area under the curve, possibly owing to higher levels of early SARS-CoV-2 replication [18]. Early antiviral therapy can effectively reduce virus clearance time and prevent the rapid progression of COVID-19. Therefore, COVID-19 patients should receive combination treatment with antiviral agents at an early stage. By reducing the rate, quantity, or duration of viral replication, the degree of direct viral injury to the respiratory epithelium, vasculature, and organs may be minimized. Additionally, secondary processes that depend on viral stimulation, including the activation of inflammatory cells, cytokines, and coagulation, could potentially be minimized if the viral replication is attenuated [19].

According to Saber-Ayad et al., antiviral drugs administered shortly after the onset of symptoms can shorten the course of clinical illness and reduce its infectivity by lowering viral shedding [20]. In another study conducted in Wuhan, Yu et al. divided 129 hospitalized patients with mild to moderate COVID-19, who were treated with antiviral drugs, into the early antiviral treatment and late antiviral treatment groups. The group receiving early antiviral treatment recovered in 7 days less than the group receiving late antiviral treatment [21].

Wu J et al. found that older adults and patients with underlying diseases are more likely to experience severe COVID-19 progression. It is recommended to start antiviral treatment in

time to slow the progression of the disease and improve the prognosis [22]. Early antiviral therapy, along with the combination therapy, increases viral clearance and reduces viral load, resulting in fatal complications including cytokine storm, macrophage activation syndrome, and acute respiratory distress syndrome [23]. According to a previous multivariate analysis, one comorbidity and the time from illness to antiviral treatment are statistically significantly associated with the severity of disease [24]. Similarly, since it is not known precisely which patients will have a more severe course of viral infection, antiviral treatment should be initiated without delay, especially in the elderly patients, patients with multiple chronic diseases, and patients who meet the clinical and laboratory criteria for severe COVID-19 at the time of onset [25]. Murohashi et al. demonstrated the importance of the early-stage use of a combination of favipiravir and methylprednisolone in severe cases, to achieve a favorable clinical outcome [26]. Doi et al. described the successful treatment of patients with COVID-19 using favipiravir to rapidly and substantially clear SARS-CoV-2 from nasal secretions irrespective of whether it was started relatively early or later within the first week of infection [27].

Similar to the above-mentioned studies, we found that early antiviral therapy is beneficial and reduces fatality in patients with COVID-19.

The primary defense mechanism against viral infections is the response formed by the humoral immune system. However, in some infectious diseases, such as COVID-19, the immune system may not be able to neutralize the virus. In these cases, it is necessary to use treatments that will reduce the rate of replication of the virus. Early initiation of antiviral drugs will reduce the rate of viral replication and decrease the time required for the humoral defense mechanism to neutralize the virus.

In conclusion, we found that initiation of the antiviral agent favipiravir, an RdRp inhibitor, within the first 72 h of the onset of disease symptoms significantly reduced fatality. Nevertheless, the research has limitations, such as the fact that the study was conducted with the participation of a small group of patients.

## REFERENCES

1. Sallenave JM, Guillot L. Innate immune signaling and proteolytic pathways in the resolution or exacerbation of SARS-CoV-2 in Covid-19: key therapeutic Targets? *Front Immunol* 2020;11:1229. [PUBMED](#) | [CROSSREF](#)
2. Helmy YA, Fawzy M, Elawad A, Sobieh A, Kenney SP, Shehata AA. The COVID-19 pandemic: a comprehensive review of taxonomy, genetics, epidemiology, diagnosis, treatment, and control. *J Clin Med* 2020;9:1225. [PUBMED](#) | [CROSSREF](#)
3. Han W, Quan B, Guo Y, Zhang J, Lu Y, Feng G, Wu Q, Fang F, Cheng L, Jiao N, Li X, Chen Q. The course of clinical diagnosis and treatment of a case infected with coronavirus disease 2019. *J Med Virol* 2020;92:461-3. [PUBMED](#) | [CROSSREF](#)
4. Furuta Y, Komeno T, Nakamura T. Favipiravir (T-705), a broad spectrum inhibitor of viral RNA polymerase. *Proc Jpn Acad Ser B Phys Biol Sci* 2017;93:449-63. [PUBMED](#) | [CROSSREF](#)
5. Joshi S, Parkar J, Ansari A, Vora A, Talwar D, Tiwaskar M, Patil S, Barkate H. Role of favipiravir in the treatment of COVID-19. *Int J Infect Dis* 2021;102:501-8. [PUBMED](#) | [CROSSREF](#)
6. Du YX, Chen XP. Favipiravir: Pharmacokinetics and Concerns About Clinical Trials for 2019-nCoV Infection. *Clin Pharmacol Ther* 2020;108:242-7. [PUBMED](#) | [CROSSREF](#)



7. Cai Q, Yang M, Liu D, Chen J, Shu D, Xia J, Liao X, Gu Y, Cai Q, Yang Y, Shen C, Li X, Peng L, Huang D, Zhang J, Zhang S, Wang F, Liu J, Chen L, Chen S, Wang Z, Zhang Z, Cao R, Zhong W, Liu Y, Liu L. Experimental treatment with favipiravir for COVID-19: an open-label control study. *Engineering (Beijing)* 2020;6:1192-8.  
[PUBMED](#) | [CROSSREF](#)
8. Yang L, Liu S, Liu J, Zhang Z, Wan X, Huang B, Chen Y, Zhang Y. COVID-19: immunopathogenesis and Immunotherapeutics. *Signal Transduct Target Ther* 2020;5:128.  
[PUBMED](#) | [CROSSREF](#)
9. Vijayvargiya P, Esquer Garrigos Z, Castillo Almeida NE, Gurram PR, Stevens RW, Razonable RR. Treatment considerations for COVID-19: a critical review of the evidence (or lack thereof). *Mayo Clin Proc* 2020;95:1454-66.  
[PUBMED](#) | [CROSSREF](#)
10. Xu X, Ong YK, Wang Y. Role of adjunctive treatment strategies in COVID-19 and a review of international and national clinical guidelines. *Mil Med Res* 2020;7:22.  
[PUBMED](#) | [CROSSREF](#)
11. Shakoory B, Carcillo JA, Chatham WW, Amdur RL, Zhao H, Dinarello CA, Cron RQ, Opal SM. Interleukin-1 receptor blockade is associated with reduced mortality in sepsis patients with features of macrophage activation syndrome: reanalysis of a prior phase III trial. *Crit Care Med* 2016;44:275-81.  
[PUBMED](#) | [CROSSREF](#)
12. Wang Y, Wang Y, Chen Y, Qin Q. Unique epidemiological and clinical features of the emerging 2019 novel coronavirus pneumonia (COVID-19) implicate special control measures. *J Med Virol* 2020;92:568-76.  
[PUBMED](#) | [CROSSREF](#)
13. Hsu J, Santesso N, Mustafa R, Brozek J, Chen YL, Hopkins JP, Cheung A, Hovhannisyian G, Ivanova L, Flottorp SA, Saeterdal I, Wong AD, Tian J, Uyeki TM, Akl EA, Alonso-Coello P, Smaill F, Schünemann HJ. Antivirals for treatment of influenza: a systematic review and meta-analysis of observational studies. *Ann Intern Med* 2012;156:512-24.  
[PUBMED](#) | [CROSSREF](#)
14. Uyeki TM, Bernstein HH, Bradley JS, Englund JA, File TM, Fry AM, Gravenstein S, Hayden FG, Harper SA, Hirshon JM, Ison MG, Johnston BL, Knight SL, McGeer A, Riley LE, Wolfe CR, Alexander PE, Pavia AT. Clinical practice guidelines by the Infectious Diseases Society of America: 2018 update on diagnosis, treatment, chemoprophylaxis, and institutional outbreak management of seasonal influenza. *Clin Infect Dis* 2019;68:e1-47.  
[PUBMED](#) | [CROSSREF](#)
15. Chan KS, Lai ST, Chu CM, Tsui E, Tam CY, Wong MM, Tse MW, Que TL, Peiris JS, Sung J, Wong VC, Yuen KY. Treatment of severe acute respiratory syndrome with lopinavir/ritonavir: a multicentre retrospective matched cohort study. *Hong Kong Med J* 2003;9:399-406.  
[PUBMED](#)
16. Magleby R, Westblade LF, Trzebucki A, Simon MS, Rajan M, Park J, Goyal P, Safford MM, Satlin MJ. Impact of SARS-CoV-2 viral load on risk of intubation and mortality among hospitalized patients with coronavirus disease 2019. *Clin Infect Dis* 2020:ciaa851.  
[PUBMED](#) | [CROSSREF](#)
17. Weiss A, Jellingsø M, Sommer MOA. Spatial and temporal dynamics of SARS-CoV-2 in COVID-19 patients: a systematic review and meta-analysis. *EBioMedicine* 2020;58:102916.  
[PUBMED](#) | [CROSSREF](#)
18. Goyal A, Lurie Y, Meissner EG, Major M, Sansone N, Uprichard SL, Cotler SJ, Dahari H. Modeling HCV cure after an ultra-short duration of therapy with direct acting agents. *Antiviral Res* 2017;144:281-5.  
[PUBMED](#) | [CROSSREF](#)
19. McCullough PA, Kelly RJ, Ruocco G, Lerma E, Tumlin J, Wheelan KR, Katz N, Lepor NE, Vijay K, Carter H, Singh B, McCullough SP, Bhambi BK, Palazzuoli A, De Ferrari GM, Milligan GP, Safder T, Tecson KM, Wang DD, McKinnon JE, O'Neill WW, Zervos M, Risch HA. Pathophysiological basis and rationale for early outpatient treatment of SARS-CoV-2 (COVID-19) infection. *Am J Med* 2021;134:16-22.  
[PUBMED](#) | [CROSSREF](#)
20. Saber-Ayad M, Saleh MA, Abu-Gharbieh E. The Rationale for Potential Pharmacotherapy of COVID-19. *Pharmaceuticals (Basel)* 2020;13:96.  
[PUBMED](#) | [CROSSREF](#)
21. Yu T, Tian C, Chu S, Zhou H, Zhang Z, Luo S, Hu D, Fan H. COVID-19 patients benefit from early antiviral treatment: A comparative, retrospective study. *J Med Virol* 2020;92:2675-83.  
[PUBMED](#) | [CROSSREF](#)

22. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *JAMA* 2020;323:1239-42.  
[PUBMED](#) | [CROSSREF](#)
23. Kim JS, Lee JY, Yang JW, Lee KH, Effenberger M, Szpirt W, Kronbichler A, Shin JI. Immunopathogenesis and treatment of cytokine storm in COVID-19. *Theranostics* 2021;11:316-29.  
[PUBMED](#) | [CROSSREF](#)
24. Wu J, Li W, Shi X, Chen Z, Jiang B, Liu J, Wang D, Liu C, Meng Y, Cui L, Yu J, Cao H, Li L. Early antiviral treatment contributes to alleviate the severity and improve the prognosis of patients with novel coronavirus disease (COVID-19). *J Intern Med* 2020;288:128-38.  
[PUBMED](#) | [CROSSREF](#)
25. Kalil AC. Treating COVID-19-off-label drug use, compassionate use, and randomized clinical trials during pandemics. *JAMA* 2020;323:1897-8.  
[PUBMED](#) | [CROSSREF](#)
26. Murohashi K, Hagiwara E, Kitayama T, Yamaya T, Higa K, Sato Y, Otoshi R, Shintani R, Okabayashi H, Ikeda S, Niwa T, Nakazawa A, Oda T, Okuda R, Sekine A, Kitamura H, Baba T, Komatsu S, Iwasawa T, Kaneko T, Ogura T. Outcome of early-stage combination treatment with favipiravir and methylprednisolone for severe COVID-19 pneumonia: a report of 11 cases. *Respir Investig* 2020;58:430-4.  
[PUBMED](#) | [CROSSREF](#)
27. Doi Y, Hibino M, Hase R, Yamamoto M, Kasamatsu Y, Hirose M, Mutoh Y, Homma Y, Terada M, Ogawa T, Kashizaki F, Yokoyama T, Koba H, Kasahara H, Yokota K, Kato H, Yoshida J, Kita T, Kato Y, Kamio T, Kodama N, Uchida Y, Ikeda N, Shinoda M, Nakagawa A, Nakatsumi H, Horiguchi T, Iwata M, Matsuyama A, Banno S, Koseki T, Teramachi M, Miyata M, Tajima S, Maeki T, Nakayama E, Taniguchi S, Lim CK, Saijo M, Imai T, Yoshida H, Kabata D, Shintani A, Yuzawa Y, Kondo M. A Prospective, randomized, open-label trial of early versus late favipiravir therapy in hospitalized patients with COVID-19. *Antimicrob Agents Chemother* 2020;64:e01897-20.  
[PUBMED](#) | [CROSSREF](#)