

Keeping you healthy: BK channel activation by omega-3 fatty acids

Ramon Latorre and Gustavo Contreras

Centro Interdisciplinario de Neurociencia de Valparaíso, Facultad de Ciencias, Universidad de Valparaíso, Valparaíso 2366103, Chile

In the exuberant world of K⁺ channels, the Ca²⁺- and voltage-activated K⁺ (BK, MaxiK, Slo1) channel stands alone. It is coded by a single gene (*slo1* or *KCNMA1*), and the pore-forming α subunit has seven transmembrane segments instead of six as found in voltage-dependent K⁺ channels (Atkinson et al., 1991; Adelman et al., 1992; Butler et al., 1993; Wallner et al., 1996). Being activated by depolarizing voltages and cytoplasmic Ca²⁺, the BK channel is the perfect molecular machine to retard or to simply stop excitatory signals. The negative feedback (hyperpolarization) created by the opening of these K⁺ channels is caused by the perfect tuning between Ca²⁺ and voltage sensors. The communication between these two types of sensors is allosterically established, that is, voltage or internal Ca²⁺ alone is able to open the BK channel, but channel opening is increasingly facilitated as more Ca²⁺ and voltage sensors are activated (Horrigan and Aldrich, 2002; Horrigan, 2012) (Fig. 1 A). Another important difference between BK channels and Kv channels, where opening is tightly coupled to voltage-sensor activation (Soler-Llavina et al., 2003) (Fig. 1 B), is that, albeit with a very low probability, BK channels can open in a voltage sensor- and Ca²⁺-independent manner (reaction C \leftrightarrow O defined by the equilibrium constant *L* in Fig. 1 A).

Despite being coded by a single gene, BK channel diversity is large. Alternative splicing, posttranslational modifications, and/or the presence of modulatory β or γ subunits create this diversity (Orío et al., 2002; Salkoff et al., 2006; Yan and Aldrich, 2010, 2012). In particular, modifications induced in BK gating kinetics by the β 1, β 2, and β 4 subunits proved to be of crucial importance in many physiological processes, ranging from shaping neuronal excitability and neurosecretion to smooth muscle tone, and in others not so physiological, such as alcohol tolerance (Brenner et al., 2000; Hu et al., 2001; Gollasch et al., 2002; Grimm and Sansom, 2007; Martin et al., 2008). The expression of β subunits is highly tissue specific; β 1 is the only β subunit expressed in smooth muscle, and β 4 is mainly expressed in the nervous system (Orío et al., 2002; Wu and Marx, 2010). In vascular smooth muscle cells (SMCs), the presence of β 1 plays a vital role in vasoregulation, and its lack leads to hypertension (Brenner et al., 2000; Fernández-Fernández et al., 2004; Nelson and Bonev, 2004). β 1 and β 2 have been observed to dramatically slow down activation and deactivation kinetics as well as

increase the apparent Ca²⁺ sensitivity of the BK channel. Although β 4 also decelerates BK activation and deactivation kinetics, even more so than β 1, Ca²⁺ sensitivity of channels formed by α/β 4 subunits is complex. Channels are less sensitive to Ca²⁺ at low internal Ca²⁺ concentrations (<10 μ M) than channels formed by the α subunit alone. However, Ca²⁺ is more effective in activating α/β 4 channels at higher Ca²⁺ concentrations (Ha et al., 2004; Wang et al., 2006).

In addition to their effects on channel gating, β subunits grant BK channels sensitivity to several physiologically important compounds, thus making these subunits targets for possible pharmacological interventions. For example, β 1-containing BK channels but not channels formed by the α subunit alone appear to be the target of 17 β -estradiol and other compounds such as estrogen analogues, anti-estrogens, and the bile salt component lithocholic acid (Valverde et al., 1999; Dick et al., 2001; Bukiya et al., 2009; Maher et al., 2013). The activation of BK channels by 17 β -estradiol has been proposed as the possible mechanism that mediates the acute relaxation of vascular smooth muscle induced by the hormone (White et al., 1995; Ruehlmann et al., 1998). On the other hand, stress steroids activate channels formed by the α/β 4 complex but not by α/β 2 (King et al., 2006). Polyunsaturated fatty acids such as arachidonic acid (AA) are also able to directly activate BK channels, but in this case, AA enhances BK current in the presence of either β 2 or β 3 (Sun et al., 2007). Findings by Sun et al. (2007) also show that AA is able to remove inactivation, suggesting that this fatty acid is interacting with the β 2-inactivating peptide. Tissue specificity of β subunits and their particular capacity to endow BK channels with different pharmacological profiles have greatly increased the importance of BK channels in maintaining the adequate cellular electrical homeostasis in different tissues.

Docosahexaenoic acid (DHA), an omega-3 fatty acid known to be associated with beneficial cardiovascular effects, was reported to be a potent activator of BK currents in rat coronary artery SMCs and to promote dilation of isolated small coronary arteries (Lai et al., 2009;

© 2013 Latorre and Contreras This article is distributed under the terms of an Attribution-Noncommercial-Share Alike-No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms>). After six months it is available under a Creative Commons License (Attribution-Noncommercial-Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>).

Wang et al., 2011). DHA, an omega-3 fatty acid found in fish oil (salmon, sardines, herring, etc.) and also in plant seeds, is the most abundant omega-3 fatty acid in the brain. By studying the Greenland Inuit tribe, which consumes large amounts of fat from fish, the conclusion was reached that high levels of omega-3 fatty acids consumed by the Inuit were the cause of their reduced triglycerides and blood pressure. Hence, this study underscores the benefits of consuming this type of lipid (Dyerberg et al., 1975). Below-normal levels of this fatty acid have been also associated with cognitive decline and increase in neural cell death (Serhan et al., 2004; Lukiw et al., 2005). However, the detailed mechanisms underlying the mode of action of this important fatty acid remain unclear.

The work done by Hoshi et al. (2013b) adds to the numerous beneficial effects of DHA by including the possibility that this fatty acid could be clinically relevant if targeted to BK channels, because of its blood pressure-lowering effects. In wild-type mice, but not in Slo1 knockout (*SLO1*^{-/-}) mice, DHA injections have been observed to reversibly reduce blood pressure and produce a significant increase in BK-mediated K⁺ currents in isolated aortic vascular SMCs, a current enhancement that was absent in SMCs dissociated from *SLO1*^{-/-} mice. When applied to the intracellular side of inside-out membrane patches, DHA was able to quickly activate $\alpha/\beta1$ channels in a reversible manner and with an EC₅₀ of ~500 nM. These findings indicate that the omega-3 fatty acid directly acts on the $\alpha/\beta1$ complex with an affinity that is ~20-fold greater than the affinity to a G-coupled receptor associated to the antiinflammatory properties of this fatty acid (Oh et al., 2010). Other important fatty acids like, for example, omega-6 fatty acids, α -linoleic and eicosapentanoic acid or AA, are also able to activate BK channels, albeit with a lower potency. Because DHA BK channel activation can be elicited with all the voltage sensors at

rest and in the absence of internal Ca²⁺, Hoshi's group arrived at the conclusion that DHA acts directly on the C \leftrightarrow O equilibrium (Fig. 1 A) and destabilizes the channel closed conformation of the pore gate.

The next step in this BK channel saga was to identify the molecular determinants in the $\beta1$ subunit that confer BK its ability to be activated by the omega-3 fatty acid (Hoshi et al., 2013a). The effects of DHA were first tested in channel with various subunits ($\beta1$, $\beta2$ [inactivation removed], $\beta4$, and $\gamma1$ [LRRC26]), with the result that robust channel activation by DHA was only observed in channels formed by $\alpha/\beta1$ and $\alpha/\beta4$. β subunits consist of two transmembrane domains connected by a large external loop and with N and C termini oriented toward the cytoplasm. The $\beta1$ phenotype can be recovered by creating a chimera containing the N terminus and the N-terminal half of the first transmembrane segment (TM1) of $\beta1$ in a $\beta2$ background. Two $\beta1$ amino acid residues, one in the amino terminus (R11) and the other in TM1 (C18), proved to be enough to recover the full effect of DHA when replacing the corresponding amino acids in $\beta2$ (A42, L49). On the other hand, the corresponding amino acids in $\beta4$ are E12 and R19. As in the case of the double mutant $\beta2$ A42R:L49C, BK channels formed by $\alpha/\beta2$ A42E:L49R subunits have very similar responses to DHA as $\alpha/\beta4$ channels. Mechanistically, it is still unclear how these of amino acid residues in $\beta1$ and $\beta4$ confer DHA sensitivity to the BK channel. However, the fact that the $\alpha/\beta1$ as well as the $\alpha/\beta4$ channel can be activated by DHA opens the possibility that DHA regulation of neuronal BK channel activity may play an important role in the nervous system.

In this issue of *JGP*, Hoshi et al. continue dissecting the effects of DHA, and their queries have led them to the identification of a single amino acid residue in the BK S6 transmembrane domain, which has been seen to establish the sensitivity to the omega-3 fatty acid.




Figure 1. Sequential versus allosteric models. (A) Allosteric model. In this model, P_o is given by the expression: $P_o = [L(1 + JD + KC + JKCD)^4] / [L(1 + JD + KC + JKCD)^4 + (1 + J + K + JKE)^4]$; $J = e^{(-z_1 F(V - V_h) / RT)}$, with $z_1 = 0.6 e_0$, $V_h = 168$ mV; $L = Lo e^{(zL FV / RT)}$, with $zL = 0.3 e_0$ and $Lo = 10^{-6}$. The values for the allosteric factors were: $D = 19$, $C = 14$, and $E = 3.8$, and the constant $K_d = 8.2$ μ M defines the Ca²⁺-binding reaction $X \leftrightarrow XCa$ (Orio and Latorre, 2005). (B) A sequential model of the Hodgkin and Huxley type. In this type of model, the open probability (P_o) is given by the expression: $P_o = (1 / (1 + J))^4$, where the equilibrium constant that defines the reaction Resting \leftrightarrow Active of the voltage sensors is defined as: $J = e^{(-z_1 F(V - V_h) / RT)}$, with $z_1 = 1.2 e_0$ and $V_h = 20$ mV.

The finding that DHA is able to activate BK channels in the absence of β subunits, albeit with about a fivefold loss in potency, led Hoshi et al. (2013c) to search for the molecular determinants of this β subunit-independent mode of action of the lipid, for which the fruit fly came to the rescue. *Drosophila melanogaster* BK channels turned out to be insensitive to DHA, and chimeras created by mixing regions of *Drosophila* and human BK channels made it possible to identify the pore domain (PD; S5-P loop–S6) as the region that is necessary and sufficient to recover the full effect of DHA in the absence of β subunits (about a fourfold increase in ionic currents). From all the point mutations made in the PD, Y318S was the only one capable of dramatically decreasing the DHA effect. Y318 is located toward the C terminus end of S6. However, before continuing with the DHA issue, an overview of certain BK activation gate characteristics having direct implications on the DHA mode of action must be offered. The BK internal vestibule is much wider than the intracellular mouth of Kv channels (Li and Aldrich, 2004; Brelidze and Magleby, 2005; Geng et al., 2011; Zhou et al., 2011). Additionally, large quaternary ammonium ions and the Shaker “ball” peptide can block closed channels, implying that the bundle crossing does not hinder the passage of ions (Wilkins and Aldrich, 2006; Thompson and Begenisich, 2012). Although this evidence strongly suggests that the BK activation gate is not cytoplasmic like in Kv channels and may reside in the selectivity filter, there is ample evidence that certain key residues in the S6 transmembrane segment can control channel gating (Wu et al., 2009; Chen and Aldrich, 2011). In particular, some of the point mutations introduced in S6 (e.g., L312Q) produced permanently open channels at all the potentials studied. The fact that the S332Y mutation located near the C terminus of S6 renders the BK channel insensitive to DHA draws attention to the importance of the S6 transmembrane segment in modulating BK channel gating. Recalling the finding that DHA is able to potentiate BK currents in the absence of activated voltage and/or Ca^{2+} sensors (Hoshi et al., 2013a), Hoshi et al., (2013c) advance the hypothesis that this omega-3 lipid modifies the closed–open equilibrium of the ion conduction gate, possibly by increasing the forward rate constant with the $\beta 1$ subunit playing an amplifying role within the functional effects of DHA binding. How this potentiating effect of the $\beta 1$ subunit develops is not clear, as its modulatory effects on BK channels gating are caused by a stabilization of the active configuration of the voltage sensor and to a large increase in the energy barrier that separates the closed from the open channel configuration (Bao and Cox, 2005; Contreras et al., 2012). One possibility, however, could be that the interaction between α and $\beta 1$ produces structural changes that favor DHA binding. Whether serine 332 is a key residue in a DHA binding pocket or if it is part of the coupling system that

transforms DHA binding energy into the pore opening is an open question. Perhaps the answer will emerge in the next paper by Hoshi’s group.

This work was supported by a grant from the Fondo Nacional de Ciencia y Tecnología (Fondecyt 1110430). The Centro Interdisciplinario de Neurociencia de Valparaíso is a Millennium Institute supported by the Millennium Initiative of the Ministerio de Economía, Fomento y Turismo.

Kenton J. Swartz served as editor.

REFERENCES

- Adelman, J.P., K.Z. Shen, M.P. Kavanaugh, R.A. Warren, Y.N. Wu, A. Lagrutta, C.T. Bond, and R.A. North. 1992. Calcium-activated potassium channels expressed from cloned complementary DNAs. *Neuron*. 9:209–216. [http://dx.doi.org/10.1016/0896-6273\(92\)90160-F](http://dx.doi.org/10.1016/0896-6273(92)90160-F)
- Atkinson, N.S., G.A. Robertson, and B. Ganetzky. 1991. A component of calcium-activated potassium channels encoded by the *Drosophila slo* locus. *Science*. 253:551–555. <http://dx.doi.org/10.1126/science.1857984>
- Bao, L., and D.H. Cox. 2005. Gating and ionic currents reveal how the BK_{Ca} channel’s Ca^{2+} sensitivity is enhanced by its $\beta 1$ subunit. *J. Gen. Physiol.* 126:393–412. <http://dx.doi.org/10.1085/jgp.200509346>
- Brelidze, T.I., and K.L. Magleby. 2005. Probing the geometry of the inner vestibule of BK channels with sugars. *J. Gen. Physiol.* 126:105–121. <http://dx.doi.org/10.1085/jgp.200509286>
- Brenner, R., G.J. Pérez, A.D. Bonev, D.M. Eckman, J.C. Kosek, S.W. Wiler, A.J. Patterson, M.T. Nelson, and R.W. Aldrich. 2000. Vasoregulation by the beta1 subunit of the calcium-activated potassium channel. *Nature*. 407:870–876. <http://dx.doi.org/10.1038/35038011>
- Bukiya, A.N., T. Vaithianathan, L. Toro, and A.M. Dopico. 2009. Channel $\beta 2$ -4 subunits fail to substitute for $\beta 1$ in sensitizing BK channels to lithocholate. *Biochem. Biophys. Res. Commun.* 390:995–1000. <http://dx.doi.org/10.1016/j.bbrc.2009.10.091>
- Butler, A., S. Tsunoda, D.P. McCobb, A. Wei, and L. Salkoff. 1993. mSlo, a complex mouse gene encoding “maxi” calcium-activated potassium channels. *Science*. 261:221–224. <http://dx.doi.org/10.1126/science.7687074>
- Chen, X., and R.W. Aldrich. 2011. Charge substitution for a deep-pore residue reveals structural dynamics during BK channel gating. *J. Gen. Physiol.* 138:137–154. <http://dx.doi.org/10.1085/jgp.201110632>
- Contreras, G.F., A. Neely, O. Alvarez, C. Gonzalez, and R. Latorre. 2012. Modulation of BK channel voltage gating by different auxiliary β subunits. *Proc. Natl. Acad. Sci. USA*. 109:18991–18996. <http://dx.doi.org/10.1073/pnas.1216953109>
- Dick, G.M., C.F. Rossow, S. Smirnov, B. Horowitz, and K.M. Sanders. 2001. Tamoxifen activates smooth muscle BK channels through the regulatory beta 1 subunit. *J. Biol. Chem.* 276:34594–34599. <http://dx.doi.org/10.1074/jbc.M104689200>
- Dyerberg, J., H.O. Bang, and N. Hjorne. 1975. Fatty acid composition of the plasma lipids in Greenland Eskimos. *Am. J. Clin. Nutr.* 28:958–966.
- Fernández-Fernández, J.M., M. Tomás, E. Vázquez, P. Orio, R. Latorre, M. Sentí, J. Marrugat, and M.A. Valverde. 2004. Gain-of-function mutation in the KCNMB1 potassium channel subunit is associated with low prevalence of diastolic hypertension. *J. Clin. Invest.* 113:1032–1039.
- Geng, Y., X. Niu, and K.L. Magleby. 2011. Low resistance, large dimension entrance to the inner cavity of BK channels determined by changing side-chain volume. *J. Gen. Physiol.* 137:533–548. <http://dx.doi.org/10.1085/jgp.201110616>

- Gollasch, M., J. Tank, F.C. Luft, J. Jordan, P. Maass, C. Krasko, A.M. Sharma, A. Busjahn, and S. Bähring. 2002. The BK channel beta1 subunit gene is associated with human baroreflex and blood pressure regulation. *J. Hypertens.* 20:927–933. <http://dx.doi.org/10.1097/00004872-200205000-00028>
- Grimm, P.R., and S.C. Sansom. 2007. BK channels in the kidney. *Curr. Opin. Nephrol. Hypertens.* 16:430–436. <http://dx.doi.org/10.1097/MNH.0b013e32826fbc7d>
- Ha, T.S., M.S. Heo, and C.S. Park. 2004. Functional effects of auxiliary beta4-subunit on rat large-conductance Ca²⁺-activated K⁺ channel. *Biophys. J.* 86:2871–2882. [http://dx.doi.org/10.1016/S0006-3495\(04\)74339-8](http://dx.doi.org/10.1016/S0006-3495(04)74339-8)
- Horrigan, F.T. 2012. Perspectives on: Conformational coupling in ion channels: Conformational coupling in BK potassium channels. *J. Gen. Physiol.* 140:625–634. <http://dx.doi.org/10.1085/jgp.201210849>
- Horrigan, F.T., and R.W. Aldrich. 2002. Coupling between voltage sensor activation, Ca²⁺ binding and channel opening in large conductance (BK) potassium channels. *J. Gen. Physiol.* 120:267–305. <http://dx.doi.org/10.1085/jgp.20028605>
- Hoshi, T., Y. Tian, R. Xu, S.H. Heinemann, and S. Hou. 2013a. Mechanism of the modulation of BK potassium channel complexes with different auxiliary subunit compositions by the omega-3 fatty acid DHA. *Proc. Natl. Acad. Sci. USA.* 110:4822–4827. <http://dx.doi.org/10.1073/pnas.1222003110>
- Hoshi, T., B. Wissuwa, Y. Tian, N. Tajima, R. Xu, M. Bauer, S.H. Heinemann, and S. Hou. 2013b. Omega-3 fatty acids lower blood pressure by directly activating large-conductance Ca²⁺-dependent K⁺ channels. *Proc. Natl. Acad. Sci. USA.* 110:4816–4821. <http://dx.doi.org/10.1073/pnas.1221997110>
- Hoshi, T., R. Xu, S. Hou, S.H. Heinemann, and Y. Tian. 2013c. A point mutation in the human Slo1 channel that obliterates its sensitivity to omega-3 docosahexaenoic acid. *J. Gen. Physiol.* 142:507–522.
- Hu, H., L.R. Shao, S. Chavoshy, N. Gu, M. Trieb, R. Behrens, P. Laake, O. Pongs, H.G. Knaus, O.P. Ottersen, and J.F. Storm. 2001. Presynaptic Ca²⁺-activated K⁺ channels in glutamatergic hippocampal terminals and their role in spike repolarization and regulation of transmitter release. *J. Neurosci.* 21:9585–9597.
- King, J.T., P.V. Lovell, M. Rishniw, M.I. Kotlikoff, M.L. Zeeman, and D.P. McCobb. 2006. Beta2 and beta4 subunits of BK channels confer differential sensitivity to acute modulation by steroid hormones. *J. Neurophysiol.* 95:2878–2888. <http://dx.doi.org/10.1152/jn.01352.2005>
- Lai, L.H., R.X. Wang, W.P. Jiang, X.J. Yang, J.P. Song, X.R. Li, and G. Tao. 2009. Effects of docosahexaenoic acid on large-conductance Ca²⁺-activated K⁺ channels and voltage-dependent K⁺ channels in rat coronary artery smooth muscle cells. *Acta Pharmacol. Sin.* 30:314–320. <http://dx.doi.org/10.1038/aps.2009.7>
- Li, W., and R.W. Aldrich. 2004. Unique inner pore properties of BK channels revealed by quaternary ammonium block. *J. Gen. Physiol.* 124:43–57. <http://dx.doi.org/10.1085/jgp.200409067>
- Lukiw, W.J., J.G. Cui, V.L. Marcheselli, M. Bodker, A. Botkjaer, K. Gotlinger, C.N. Serhan, and N.G. Bazan. 2005. A role for docosahexaenoic acid-derived neuroprotectin D1 in neural cell survival and Alzheimer disease. *J. Clin. Invest.* 115:2774–2783. <http://dx.doi.org/10.1172/JCI25420>
- Maher, J., A.C. Hunter, J.G. Mabley, J. Lippiat, and M.C. Allen. 2013. Smooth muscle relaxation and activation of the large conductance Ca²⁺-activated K⁺ (BK_{Ca}) channel by novel oestrogens. *Br. J. Pharmacol.* 169:1153–1165. <http://dx.doi.org/10.1111/bph.12211>
- Martin, G.E., L.M. Hendrickson, K.L. Penta, R.M. Friesen, A.Z. Pietrzykowski, A.R. Tapper, and S.N. Treistman. 2008. Identification of a BK channel auxiliary protein controlling molecular and behavioral tolerance to alcohol. *Proc. Natl. Acad. Sci. USA.* 105:17543–17548. <http://dx.doi.org/10.1073/pnas.0801068105>
- Nelson, M.T., and A.D. Bonev. 2004. The beta1 subunit of the Ca²⁺-sensitive K⁺ channel protects against hypertension. *J. Clin. Invest.* 113:955–957.
- Oh, D.Y., S. Talukdar, E.J. Bae, T. Imamura, H. Morinaga, W. Fan, P. Li, W.J. Lu, S.M. Watkins, and J.M. Olefsky. 2010. GPR120 is an omega-3 fatty acid receptor mediating potent anti-inflammatory and insulin-sensitizing effects. *Cell.* 142:687–698. <http://dx.doi.org/10.1016/j.cell.2010.07.041>
- Orio, P., and R. Latorre. 2005. Differential effects of β1 and β2 subunits on BK channel activity. *J. Gen. Physiol.* 125:395–411. <http://dx.doi.org/10.1085/jgp.200409236>
- Orio, P., P. Rojas, G. Ferreira, and R. Latorre. 2002. New disguises for an old channel: MaxiK channel beta-subunits. *News Physiol. Sci.* 17:156–161.
- Ruehlmann, D.O., J.R. Steinert, M.A. Valverde, R. Jacob, and G.E. Mann. 1998. Environmental estrogenic pollutants induce acute vascular relaxation by inhibiting L-type Ca²⁺ channels in smooth muscle cells. *FASEB J.* 12:613–619.
- Salkoff, L., A. Butler, G. Ferreira, C. Santi, and A. Wei. 2006. High-conductance potassium channels of the SLO family. *Nat. Rev. Neurosci.* 7:921–931. <http://dx.doi.org/10.1038/nrn1992>
- Serhan, C.N., K. Gotlinger, S. Hong, and M. Arita. 2004. Resolvins, docosatrienes, and neuroprotectins, novel omega-3-derived mediators, and their aspirin-triggered endogenous epimers: an overview of their protective roles in catabasis. *Prostaglandins Other Lipid Mediat.* 73:155–172. <http://dx.doi.org/10.1016/j.prostaglandins.2004.03.005>
- Soler-Llavina, G.J., M. Holmgren, and K.J. Swartz. 2003. Defining the conductance of the closed state in a voltage-gated K⁺ channel. *Neuron.* 38:61–67. [http://dx.doi.org/10.1016/S0896-6273\(03\)00157-0](http://dx.doi.org/10.1016/S0896-6273(03)00157-0)
- Sun, X., D. Zhou, P. Zhang, E.G. Moczydlowski, and G.G. Haddad. 2007. Beta-subunit-dependent modulation of hSlo BK current by arachidonic acid. *J. Neurophysiol.* 97:62–69. <http://dx.doi.org/10.1152/jn.00700.2006>
- Thompson, J., and T. Begegnisich. 2012. Selectivity filter gating in large-conductance Ca²⁺-activated K⁺ channels. *J. Gen. Physiol.* 139:235–244. <http://dx.doi.org/10.1085/jgp.201110748>
- Valverde, M.A., P. Rojas, J. Amigo, D. Cosmelli, P. Orío, M.I. Bahamonde, G.E. Mann, C. Vergara, and R. Latorre. 1999. Acute activation of Maxi-K channels (hSlo) by estradiol binding to the beta subunit. *Science.* 285:1929–1931. <http://dx.doi.org/10.1126/science.285.5435.1929>
- Wallner, M., P. Meera, and L. Toro. 1996. Determinant for beta-subunit regulation in high-conductance voltage-activated and Ca²⁺-sensitive K⁺ channels: an additional transmembrane region at the N terminus. *Proc. Natl. Acad. Sci. USA.* 93:14922–14927. <http://dx.doi.org/10.1073/pnas.93.25.14922>
- Wang, B., B.S. Rothberg, and R. Brenner. 2006. Mechanism of β4 subunit modulation of BK channels. *J. Gen. Physiol.* 127:449–465. <http://dx.doi.org/10.1085/jgp.200509436>
- Wang, R.X., Q. Chai, T. Lu, and H.C. Lee. 2011. Activation of vascular BK channels by docosahexaenoic acid is dependent on cytochrome P450 epoxygenase activity. *Cardiovasc. Res.* 90:344–352. <http://dx.doi.org/10.1093/cvr/cvq411>
- White, R.E., D.J. Darkow, and J.L. Lang. 1995. Estrogen relaxes coronary arteries by opening BKCa channels through a cGMP-dependent mechanism. *Circ. Res.* 77:936–942. <http://dx.doi.org/10.1161/01.RES.77.5.936>
- Wilkins, C.M., and R.W. Aldrich. 2006. State-independent block of BK channels by an intracellular quaternary ammonium. *J. Gen. Physiol.* 128:347–364. <http://dx.doi.org/10.1085/jgp.200609579>
- Wu, R.S., and S.O. Marx. 2010. The BK potassium channel in the vascular smooth muscle and kidney: α- and β-subunits. *Kidney Int.* 78:963–974. <http://dx.doi.org/10.1038/ki.2010.325>

- Wu, Y., Y. Xiong, S. Wang, H. Yi, H. Li, N. Pan, F.T. Horrigan, Y. Wu, and J. Ding. 2009. Intersubunit coupling in the pore of BK channels. *J. Biol. Chem.* 284:23353–23363. <http://dx.doi.org/10.1074/jbc.M109.027789>
- Yan, J., and R.W. Aldrich. 2010. LRRC26 auxiliary protein allows BK channel activation at resting voltage without calcium. *Nature.* 466:513–516. <http://dx.doi.org/10.1038/nature09162>
- Yan, J., and R.W. Aldrich. 2012. BK potassium channel modulation by leucine-rich repeat-containing proteins. *Proc. Natl. Acad. Sci. USA.* 109:7917–7922. <http://dx.doi.org/10.1073/pnas.1205435109>
- Zhou, Y., X.M. Xia, and C.J. Lingle. 2011. Cysteine scanning and modification reveal major differences between BK channels and Kv channels in the inner pore region. *Proc. Natl. Acad. Sci. USA.* 108:12161–12166. <http://dx.doi.org/10.1073/pnas.1104150108>