

ARTICLE

Received 27 Sep 2012 | Accepted 3 Dec 2012 | Published 15 Jan 2013

DOI: 10.1038/ncomms2354

OPEN

# Transgenerational gene silencing causes gain of virulence in a plant pathogen

Dinah Qutob<sup>1</sup>, B. Patrick Chapman<sup>1</sup> & Mark Gijzen<sup>1</sup>

Avirulence (*Avr*) genes of plant pathogens encode effector proteins that trigger immunity in plants carrying appropriate resistance (*R*) genes. The *Phytophthora sojae Avr3a* gene displays allelic variation in messenger RNA transcript levels. *P. sojae* strains with detectable *Avr3a* gene transcripts are avirulent on plants carrying the *R*-gene *Rps3a*, whereas strains lacking *Avr3a* mRNA escape detection by *Rps3a* and are virulent. Here we show non-Mendelian interactions between naturally occurring *Avr3a* alleles that result in transgenerational gene silencing, and we identify small RNA molecules of 25 nucleotides that are abundant in gene-silenced strains but not in strains with *Avr3a* mRNA. This example of transgenerational gene silencing is exceptional because it is naturally occurring and results in gain of virulence in a pathogenic organism.

<sup>1</sup>Agriculture and Agri-Food Canada, 1391 Sandford Street, London, Ontario N5V 4T3, Canada. Correspondence and requests for materials should be addressed to M.G. (email: mark.gijzen@agr.gc.ca).

Plant pathogens secrete effector proteins to enable disease, but evolution favors host plant immune systems to recognize these factors as cues for activating rapid defence responses that arrest the infection<sup>1</sup>. Effector-triggered immunity in plants is the consequence of signalling events that are determined by resistance (*R*) gene products interacting, directly or indirectly, with pathogen avirulence (*Avr*) effectors.

*Phytophthora sojae* causes stem and root rot of soybean and is a major disease problem that plagues this crop<sup>2</sup>. The *P. sojae* *Avr3a* gene encodes a predicted protein of 111 amino acids that includes a signal peptide, a host-targeting motif and a carboxy-terminal effector domain, which are common features of oomycete *Avr* effectors<sup>3–6</sup>. *Avr3a* resides in a 10.8 kilobase (kb) DNA segment that displays copy number variation among *P. sojae* strains<sup>5,6</sup>. Four haplotypes of *Avr3a* are known and gain of virulence on the *R*-gene *Rps3a* is caused by messenger RNA transcript differences between strains. In crosses between *P. sojae* strains P6497 (*Avr3a*<sup>P6497</sup>/*Avr3a*<sup>P6497</sup>) and P7064 (*Avr3a*<sup>P7064</sup>/*Avr3a*<sup>P7064</sup>), avirulence towards *Rps3a* plants segregates as a dominant Mendelian trait conditioned by the presence of *Avr3a*<sup>P6497</sup>. Transcripts of *Avr3a*<sup>P6497</sup> but not *Avr3a*<sup>P7064</sup> mRNA are detectable in each of the parental strains and their progeny<sup>5,6</sup>.

Sequencing of *Avr3a* alleles and analysis of mRNA transcripts in 17 different *P. sojae* strains revealed four different haplotypes<sup>6</sup>, but the inheritance behaviour of each of these haplotypes in segregating populations has not been investigated. The present study was intended to test the inheritance *Avr3a* in *P. sojae* strains with haplotypes that are different from strains P6497 and P7064. To our surprise, we discovered that genetic outcrosses between *P. sojae* strains ACR10 and P7076 resulted in transgenerational gene silencing of *Avr3a* and gain of virulence on soybean plants carrying the *Rps3a* gene.

## Results

**Transgenerational gene silencing of *Avr3a*.** To determine the inheritance of other *Avr3a* alleles, we created crosses between *P. sojae* strains P7076 (*Avr3a*<sup>P7076</sup>/*Avr3a*<sup>P7076</sup>) and ACR10 (*Avr3a*<sup>ACR10</sup>/*Avr3a*<sup>ACR10</sup>). The strain P7076 is avirulent, whereas strain ACR10 is virulent on *Rps3a* plants, and mRNA transcripts are detectable for *Avr3a*<sup>P7076</sup> but not *Avr3a*<sup>ACR10</sup>, as we have previously shown<sup>5,6</sup> (Fig. 1a). To select hybrid progeny and follow segregation of *Avr3a* alleles we used co-dominant DNA markers derived from cleaved amplified polymorphic (CAP) fragments, because *Avr3a*<sup>P7076</sup> and *Avr3a*<sup>ACR10</sup> differ in sequence<sup>5,6</sup>. A total of 28 F<sub>1</sub> hybrid (*Avr3a*<sup>P7076</sup>/*Avr3a*<sup>ACR10</sup>) progeny were isolated, scored for virulence on *Rps3a* plants and assessed for the presence of *Avr3a* mRNA transcripts by reverse transcriptase polymerase chain reaction (RT-PCR) (Table 1, Fig. 2, Supplementary Fig. S1). All 28 F<sub>1</sub> progeny were virulent on *Rps3a* plants and were lacking *Avr3a* mRNA transcripts. This result was unexpected because heretofore avirulent alleles of *Avr3a* that accumulate mRNA transcripts were dominant to virulent alleles that lack transcripts, such as in crosses between *P. sojae* strains P6497 and P7064. To further explore the segregation of *Avr3a* alleles, virulence and mRNA transcript accumulation in this cross, we generated three different F<sub>2</sub> populations from separate F<sub>1</sub> individuals. In all, 139 F<sub>2</sub> progeny were tested for segregation of *Avr3a* alleles and mRNA transcripts and scored for virulence on *Rps3a* plants. The *Avr3a*<sup>P7076</sup> and *Avr3a*<sup>ACR10</sup> alleles segregated normally in each F<sub>2</sub> population but remarkably all 139 F<sub>2</sub> progeny lacked *Avr3a* mRNA transcripts and were virulent on *Rps3a* plants (Table 1). A sampling of eight F<sub>3</sub> individuals, derived from homozygous *Avr3a*<sup>P7076</sup>/*Avr3a*<sup>P7076</sup> F<sub>2</sub> cultures, indicates that gene silencing is maintained and meiotically stable in these inbred lines

(Supplementary Fig. S2). These results demonstrate heritable transgenerational gene silencing of *Avr3a* in crosses between *P. sojae* strains ACR10 and P7076.


**Distinctive profile of small RNA (sRNA) in silenced strains.** To explore the cause of transgenerational silencing of *Avr3a*, we performed deep sequencing of sRNA from each parental *P. sojae* strain ACR10 and P7076, and from a sample of F<sub>2</sub> cultures with the genotype *Avr3a*<sup>P7076</sup>/*Avr3a*<sup>P7076</sup>. Results from this analysis show that the profile of sRNA coverage across the *Avr3a* region differs between ACR10 and P7076 (Fig. 1b). The sRNA sequences matching to the *Avr3a* region are far more abundant in strain ACR10 compared with P7076 (Table 2). The most prevalent of these sRNA molecules are 24–26 nucleotides in length (Fig. 1c). In strain ACR10, there is a nearly continuous coverage of sRNA sequences that align to a 3.7 kb span of the *Avr3a* interval including the 5' and 3' intergenic segments but not extending into flanking genes. In contrast, in strain P7076 coverage of matching sRNA is limited to a 0.3-kb segment occurring in the 5' intergenic region. Sequencing of sRNA from homozygous *Avr3a*<sup>P7076</sup>/*Avr3a*<sup>P7076</sup> F<sub>2</sub> cultures resulted in a profile of coverage that is nearly identical to that from strain ACR10 (*Avr3a*<sup>ACR10</sup>/*Avr3a*<sup>ACR10</sup>) rather than that from P7076 (*Avr3a*<sup>P7076</sup>/*Avr3a*<sup>P7076</sup>). The sRNA profiles were replicated from independent samples of ACR10, P7076, F<sub>1</sub> and F<sub>2</sub> cultures, and extended to include samples from additional *P. sojae* strains P6497 and ACR16 (Table 2; Supplementary Fig. S3). The results consistently show a pattern of extensive sRNA coverage of *Avr3a* occurs in gene-silenced strains.

## Discussion

Previous studies have shown that transformation of organisms, including species of *Phytophthora*, with recombinant DNA or RNA molecules can cause internuclear or transgenerational gene silencing that resembles our observations of naturally occurring silencing of *Avr3a* (refs 7,8). Moreover, results from comparative genome sequencing of species within the *Phytophthora infestans* clade suggest a role for epigenetic mechanisms in the rapid evolution of *Phytophthora* species<sup>9</sup>. Thus, we propose that transgenerational gene silencing of *Avr3a* in *P. sojae* is the result of epigenetic changes that modulate the expression state of this gene, and that are transmitted to progeny in a non-Mendelian fashion.

Given the emerging functions of sRNAs in controlling epigenetic marks and gene expression states in numerous experimental systems, it is noteworthy that we found an association between sRNAs and gene silencing at the *Avr3a* locus. In plants, sRNA of 24–26 nucleotides are involved in directing chromatin changes including DNA methylation, whereas shorter ones of 21–22 nucleotides participate in mRNA degradation<sup>10</sup>. Heterochromatic silencing in yeast is initiated by an interaction of sRNA and RNA polymerase II, which guides histone modifications that control chromatin structure and gene expression<sup>11</sup>. Thus, the evidence suggests a role for sRNA in directing the transgenerational gene expression changes of the *P. sojae* *Avr3a* gene.

Our observations of transgenerational gene silencing of *Avr3a* provide a remarkable example of pathogen adaptation to host immune surveillance. The interactions between pathogens and their hosts are often interpreted as an evolutionary race, where pathogen virulence and host resistance are at the forefront<sup>12</sup>. Transgenerational gene silencing of an *Avr* effector locus represents a novel method of host immune evasion that offers the ability to spread quickly in pathogen populations, but yet is potentially reversible as the sequence of the *Avr* gene itself



**Figure 1 | Characteristics of *P. sojae* *Avr3a* alleles and deep sequencing of sRNA.** (a) A schematic illustration of *Avr3a* alleles from four different strains of *P. sojae*. The segment of DNA harbouring *Avr3a* and four other predicted genes is shown. Predicted open reading frames are indicated by bold arrows. Shown for each allele is the copy number of the DNA segment, presence (+) or absence (-) of *Avr3a* mRNA transcripts and disease outcome on *Rps3a* plants as virulent (V) or avirulent (A). (b) Profiles of sRNA from *P. sojae* strains ACR10 and P7076, and from a pool of F<sub>2</sub> cultures selected for their *Avr3a*<sup>P7076</sup>/*Avr3a*<sup>P7076</sup> genotype. Average depth of coverage across the 10.8-kb genome segment containing *Avr3a* was normalized to the total number of trimmed reads for each sample. (c) Size of sRNA aligning to the 10.8-kb DNA segment. The total number of counts of RNA sequence reads and the size in nucleotides (nt) is shown for *P. sojae* strain ACR10.

**Table 1 | Segregation of *Avr3a* alleles and virulence phenotypes in F<sub>1</sub> and F<sub>2</sub> populations (ACR10 × P7076) of *Phytophthora sojae*.**


Population	n	<i>Avr3a</i> allele genotype			P	Virulence		mRNA	
		<i>Avr3a</i> <sup>ACR10</sup> / <i>Avr3a</i> <sup>ACR10</sup>	<i>Avr3a</i> <sup>ACR10</sup> / <i>Avr3a</i> <sup>P7076</sup>	<i>Avr3a</i> <sup>P7076</sup> / <i>Avr3a</i> <sup>P7076</sup>		A	V	(+)	(-)
F <sub>1</sub>	28	0	28	0	1.0	0	28	0	28
F <sub>2</sub> -1	51	16	26	9	0.38	0	51	0	51
F <sub>2</sub> -2	44	14	16	14	0.19	0	44	0	44
F <sub>2</sub> -3	44	10	22	12	0.91	0	44	0	44

The number of progeny (n) in each population and the P value from a  $\chi^2$ -test for Mendelian segregation of genotypes is shown. Virulent (V) and avirulent (A) phenotypes were scored by inoculation of plants, and the presence (+) or absence (-) of mRNA transcripts of *Avr3a* as determined by RT-PCR.

remains unchanged. We predict that other eukaryotic pathogens of plants, humans and animals will employ similar mechanisms to escape detection by host immune systems.

**Methods**

***Phytophthora sojae* strains and crosses.** The origin of *P. sojae* strains and methods of culture, propagation and oospore isolation have been described<sup>5,6,13-15</sup>.



**Figure 2 | Photographs of *P. sojae*-infected soybean plants showing disease outcomes.** Virulence assays of *P. sojae* cultures were performed on greenhouse-grown seedlings of soybean cultivar Williams (*rps3a*) and the isoline L83-570 (*Rps3a*) by hypocotyl inoculation with *P. sojae* mycelia. Shown are disease outcomes 6 days after inoculation of plants with the parental strains, the F<sub>1</sub> hybrids used to construct each of the three F<sub>2</sub> populations, or with representative F<sub>2</sub> individuals illustrating each *Avr3a* genotype from the three F<sub>2</sub> populations tested. Pots are 10 cm in diameter.

*P. sojae* is an oomycete and diploid microorganism that reproduces sexually through the development of oogonia (female gametophyte) and antheridia (male gametophyte). *P. sojae* is homothallic and is not known to have mating types,

therefore oospores can develop from self-fertilization or from out-crossing when different strains of the organism are grown together<sup>2</sup>. To perform sexual crosses of *P. sojae*, parental strains are co-cultivated and oospores are isolated from culture

**Table 2 | Deep sequencing of sRNA from *Phytophthora sojae*.\***

Sample <sup>†</sup>	<i>Avr3a</i> genotype	<i>Avr3a</i> mRNA <sup>‡</sup>	Virulence on <i>Rps3a</i>	Total number of trimmed sRNA sequence reads	Matches to <i>Avr3a</i> segment <sup>§</sup>	Normalized matches per 10 <sup>6</sup> reads
<i>Parental strains</i>						
P7076 (rep1)	<i>Avr3a</i> <sup>P7076</sup> / <i>Avr3a</i> <sup>P7076</sup>	+	A	4,640,922	306	65
P7076 (rep2)	<i>Avr3a</i> <sup>P7076</sup> / <i>Avr3a</i> <sup>P7076</sup>	+	A	13,040,733	659	50
ACR10 (rep1)	<i>Avr3a</i> <sup>ACR10</sup> / <i>Avr3a</i> <sup>ACR10</sup>	–	V	3,338,276	1,715	513
ACR10 (rep2)	<i>Avr3a</i> <sup>ACR10</sup> / <i>Avr3a</i> <sup>ACR10</sup>	–	V	13,164,585	9,488	720
<i>Progeny</i>						
F <sub>1</sub> -2	<i>Avr3a</i> <sup>ACR10</sup> / <i>Avr3a</i> <sup>P7076</sup>	–	V	48,505,836	10,603	218
F <sub>2</sub> -2-14	<i>Avr3a</i> <sup>ACR10</sup> / <i>Avr3a</i> <sup>ACR10</sup>	–	V	50,286,928	30,809	612
F <sub>2</sub> -2-26	<i>Avr3a</i> <sup>ACR10</sup> / <i>Avr3a</i> <sup>P7076</sup>	–	V	45,921,916	17,812	387
F <sub>2</sub> -2-35	<i>Avr3a</i> <sup>P7076</sup> / <i>Avr3a</i> <sup>P7076</sup>	–	V	45,141,247	12,883	285
F <sub>2</sub> pool (rep1)	<i>Avr3a</i> <sup>P7076</sup> / <i>Avr3a</i> <sup>P7076</sup>	–	V	14,712,321	4,411	299
F <sub>2</sub> pool (rep2)	<i>Avr3a</i> <sup>P7076</sup> / <i>Avr3a</i> <sup>P7076</sup>	–	V	20,495,465	6,204	302
<i>Other strains</i>						
P6497	<i>Avr3a</i> <sup>P6497</sup> / <i>Avr3a</i> <sup>P6497</sup>	+	A	17,919,740	345	19
ACR16	<i>Avr3a</i> <sup>ACR16</sup> / <i>Avr3a</i> <sup>ACR16</sup>	–	V	20,450,286	3,338	163

\*Samples of sRNA were purified from mycelia cultures of *P. sojae* for library construction and sequencing, as described in Methods section.

<sup>†</sup>Source of all *P. sojae* strains and sequence of *Avr3a* alleles has previously been described<sup>5,6</sup>.

<sup>‡</sup>Presence (+) or absence (–) of *Avr3a* mRNA, as determined by RT-PCR.

<sup>§</sup>Matches of sRNA sequences to 10.8 kb *Avr3a* segment, as shown in Fig. 1.

A, avirulent; rep, replicate; V, virulent.

homogenates by a series of purification steps<sup>13–15</sup>. Purified oospores are diluted and plated out on water agar to germinate. Individual germinating oospores are picked under a dissecting microscope and transferred to rich growth medium. Samples of purified DNA from the resulting cultures are tested using co-dominant DNA markers polymorphic between the parents, to determine whether the individuals result from self-fertilization or from out-crossing events between the parental strains. Such markers designed to distinguish *Avr3a*<sup>ACR10</sup> and *Avr3a*<sup>P7076</sup> alleles (described below) were used to identify F<sub>1</sub> progeny in crosses between strain ACR10 and P7076. The F<sub>2</sub> populations were generated by isolating oospores from self-fertilized F<sub>1</sub> individuals.

#### Purification of nucleic acids and scoring of markers and transcripts for *Avr3a*.

Mycelia cultures of *P. sojae* for nucleic acid isolation were grown on vegetable juice (V8) agar media overlaid with cellophane (BioRad). After growth in darkness for 7 days at 25 °C, mycelia-grown cellophane disks were peeled off the media, frozen at –80 °C and ground to a fine powder in liquid N<sub>2</sub> using a mortar and pestle. Genomic DNA was purified using conventional phenol–chloroform extraction procedures followed by precipitation with isopropanol<sup>16</sup>. Purified DNA pellets were washed with 70% (v/v) ethanol, dried and resuspended in a solution of 10 mM Tris HCl, pH 7.4, 0.1 mM EDTA. CAP markers were used to track segregation of the *Avr3a*<sup>ACR10</sup> and *Avr3a*<sup>P7076</sup> alleles. For CAPs analysis, PCR amplification was performed with 5 ng of genomic DNA as template in a volume of 25 µl under the following conditions: 0.2 mM dNTPs, 1.5 mM MgCl<sub>2</sub>, 0.4 µM of each primer (forward primer 5'-GCTGCTTCCTTCCTGGTTGC-3' and reverse primer 5'-GCTGCTGCTTTTGCTTCTC-3'), 1.25 U Taq DNA polymerase and 1 × PCR buffer as recommended by the supplier (Invitrogen, Life Technologies). The amplification was carried out in a 96-well plate (GeneAmp PCR System 9700, Applied Biosystems) under the following conditions: 94 °C for 2 min; 40 cycles of 94 °C for 40 s, 58 °C for 40 s, 72 °C for 2 min; followed by an extension step of 10 min at 72 °C. For restriction analysis of the PCR fragments, 10 µl of the amplified products were incubated overnight at 37 °C in a volume of 20 µl using 3 U of AluI (New England Biolabs). Digestion products were resolved by agarose gel electrophoresis and visualized by ethidium bromide staining according to standard methods<sup>16</sup>. The *Avr3a*<sup>ACR10</sup> sequence includes an AluI site, whereas the *Avr3a*<sup>P7076</sup> sequence lacks the restriction site and is not digested<sup>5,6</sup>.

Total RNA was extracted from frozen, pulverized mycelia samples in a solution of phenol-guanidine isothiocyanate (TRIzol, Invitrogen) according to instructions provided by the manufacturer. Approximately 1 µg total RNA was subject to DNaseI (Invitrogen) treatment before RT-PCR, as previously described<sup>5,6</sup>. Cycling conditions for detecting transcripts of *Avr3a* as well as for the *P. sojae actin* gene (forward primer 5'-CGAAATTGTGCGCGACATCAAG-3' and reverse primer 5'-GGTACCGCCGACAGCAGAT-3'), used as a control, were described as above with the exception that 25 cycles were used for transcript analysis.

**Plant inoculation and virulence assays.** Virulence assays of *P. sojae* cultures were performed on greenhouse-grown seedlings of soybean cultivar Williams (*rps*)

and the isolate L83-570 (*Rps3a*), according to standard methods<sup>5,6</sup>. Soybean seeds (20–30 seeds per pot) were sown in 10 cm pots and grown for 7 days before disease testing. For inoculation, *P. sojae* cultures were grown for 5–7 days on 0.9% (v/v) V8 agar plates. Inoculum was prepared by excising 1 cm segments of agar cut from the growing edge of mycelia colonies and macerating the cultures through a 3-ml syringe attached to an 18-gauge needle. Soybean plants were inoculated in the mid-section of each hypocotyl by making a small incision for injection of the mycelial slurry. Inoculated plants were covered with plastic bags to maintain humidity for 2 days. Disease symptoms were allowed to develop for an additional 4 days before phenotypes were scored. A minimum of three independent replicates of the disease assay were performed for each *P. sojae* culture tested.

**Deep sequencing of *P. sojae* sRNA.** To profile sRNA molecules of *P. sojae* cultures, samples of RNA were purified from mycelia cultures as described above. The size selection of sRNA, attachment of adaptor molecules and library construction was performed using prepared supplies and reagents (Illumina small RNA v1.5), following instructions provided by the manufacturer. Sequencing was performed on a flow cell instrument (Illumina GAII). After trimming to remove adaptor sequences and low-quality sequences, the depth of sRNA coverage of the *Avr3a* region was determined by alignment, and normalized to the number of reads per library. Average depth of coverage across the 10.8-kb genome segment containing *Avr3a* was determined using a 50-base pair moving interval and values normalized to the total number of trimmed reads for each sample. The number of trimmed reads obtained for each of the sequenced sRNA libraries is proved in Table 2.

#### References

- Jones, J. D. & Dangl, J. L. The plant immune system. *Nature* **444**, 323–329 (2006).
- Erwin, D. C. & Ribeiro, O. K. *Phytophthora Diseases Worldwide* (APS Press, 1997).
- Tyler, B. M. Entering and breaking: virulence effector proteins of oomycete plant pathogens. *Cell Microbiol.* **11**, 13–20 (2009).
- Oliva, R. et al. Recent developments in effector biology of filamentous plant pathogens. *Cell Microbiol.* **12**, 705–715 (2010).
- Qutob, D. et al. Copy number variation and transcriptional polymorphisms of *Phytophthora sojae* RXLR effector genes *Avr1a* and *Avr3a*. *PLoS ONE* e50666 (2009).
- Dong, S. et al. Sequence variants of the *Phytophthora sojae* RXLR effector *Avr3a/5* are differentially recognized by *Rps3a* and *Rps5* in soybean. *PLoS ONE* e20172 (2011).
- van West, P., Kamoun, S., van't Klooster, J. W. & Govers, F. Internuclear gene silencing in *Phytophthora infestans*. *Mol. Cell.* **3**, 339–348 (1999).
- van West, P. et al. Internuclear gene silencing in *Phytophthora infestans* is established through chromatin remodelling. *Microbiology* **154**, 1482–1490 (2008).
- Raffaele, S. et al. Genome evolution following host jumps in the Irish potato famine pathogen lineage. *Science* **330**, 1540–1543 (2010).
- Molnar, A., Melnyk, C. & Baulcombe, D. C. Silencing signals in plants: a long journey for small RNAs. *Genome Biol.* **12**, 215 (2011).

11. Zaratiegui, M. *et al.* RNAi promotes heterochromatic silencing through replication-coupled release of RNA Pol II. *Nature* **479**, 135–137 (2011).
12. Woolhouse, M. E., Webster, J. P., Domingo, E., Charlesworth, B. & Levin, B. R. Biological and biomedical implications of the co-evolution of pathogens and their hosts. *Nat. Genet.* **32**, 569–577 (2002).
13. MacGregor, T. *et al.* Genetic and physical mapping of *Avr1a* in *Phytophthora sojae*. *Genetics* **160**, 949–959 (2002).
14. Tyler, B. M., Forster, H. & Coffey, M. D. Inheritance of avirulence factors and restriction fragment length polymorphism markers in outcrosses of the oomycete *Phytophthora sojae*. *Mol. Plant-Microbe Interact.* **8**, 515–523 (1995).
15. Gijzen, M. & Qutob, D. in *Oomycete Genetics and Genomics: Diversity, Interactions and Research Tools* (eds Lamour, K. & Kamoun, S.) 303–329 (Wiley-Blackwell, 2009).
16. Sambrook, J., Fritsch, E. F. & Maniatis, T. *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Press, 1989).

## Acknowledgements

We thank The Centre for Applied Genomics (Toronto) and the BC Cancer Agency (Vancouver) for sRNA sequencing; Aldona Gaidauskas-Scott, Irina Mutiu, Kufrom Kufllu and Alex Molnar for contributions to the work; and Brett Tyler and Krzysztof

Szczyglowski for comments on the manuscript. This work was funded by the Agriculture and Agri-Food Canada Crop Genomics Initiative.

## Author contributions

D.Q. and M.G. conceived and designed the experiments. D.Q. performed the experiments. D.Q., B.P.C. and M.G. analysed the data. M.G. wrote the paper.

## Additional information

**Supplementary Information** accompanies this paper at <http://www.nature.com/naturecommunications>

**Competing financial interests:** The authors declare no competing financial interests.

**Reprints and permission** information is available online at <http://npg.nature.com/reprintsandpermissions/>

**How to cite this article:** Qutob, D. *et al.* Transgenerational gene silencing causes gain of virulence in a plant pathogen. *Nat. Commun.* **4**:1349 doi: 10.1038/ncomms2354 (2013).



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>