

RETRACTION NOTE

Open Access



Retraction Note To: Functional activation of Proline-rich Tyrosine Kinase2 (PYK2) in peripheral blood mononuclear cells from patients with systemic lupus erythematosus

Meiying Wang^{1†}, Hongsheng Sun^{1†}, Wei Zhang² and Yuanchao Zhang^{1*}

Retraction

This article [1] has been retracted by the editors of BMC Musculoskeletal Disorders due to extensive overlap with previously published work [2]. We apologise for the inconvenience caused.

Author details

¹Department of Rheumatology and Immunology, Provincial Hospital affiliated to Shandong University, Jinan 250021, China. ²Department of Pain Management, Hospital affiliated to Medical College of Qingdao University, Oinodao 266003, China.

Received: 7 January 2015 Accepted: 7 January 2015 Published online: 19 June 2015

References

- Wang M, Sun H, Zhang W, Zhang Y. Functional activation of Proline-rich Tyrosine Kinase2 (PYK2) in peripheral blood mononuclear cells from patients with systemic lupus erythematosus. BMC Musculoskelet Disord. 2009;10:141.
- Wang M, Zhang W, Zhang Y. Activation signal transduction by Proline-rich Tyrosine Kinase 2 (PYK2) in Peripheral blood mononuclear cells from patients with systemic lupus erythematosus. Hybridoma. 2009;28:5.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit



¹Department of Rheumatology and Immunology, Provincial Hospital affiliated to Shandong University, Jinan 250021, China Full list of author information is available at the end of the article



^{*} Correspondence: sdfs2004@sohu.com

[†]Equal contributors