

HOSTED BY



Contents lists available at ScienceDirect

## International Journal of Nursing Sciences

journal homepage: <http://www.elsevier.com/journals/international-journal-of-nursing-sciences/2352-0132>

## Original Article

## Validation of the Persian version of the Compassionate Care Assessment Tool

Aliakbar Vaisi-Raygani <sup>a,b</sup>, Asghar Dalvandi <sup>b</sup>, Kian Nourozi <sup>b</sup>, Abbas Ebadi <sup>c</sup>, Mahdi Rahgozar <sup>b</sup>, Rostam Jalali <sup>a</sup>, Nader Salari <sup>a</sup>, Alireza Abdi <sup>a,\*</sup><sup>a</sup> Department of Nursing, School of Nursing and Midwifery, Kermanshah University of Medical Sciences, Kermanshah, Iran<sup>b</sup> Nursing Department, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran<sup>c</sup> Behavioral Sciences Research Center, Life Style Institute, Baqiyatallah University of Medical Sciences, Teheran, Iran

## ARTICLE INFO

## Article history:

Received 11 April 2020

Received in revised form

4 November 2020

Accepted 8 December 2020

Available online 14 December 2020

## Keywords:

Compassionate care

Iran

Nursing care

Psychometrics

Statistical factor analysis

## ABSTRACT

**Objectives:** The present study aimed to translate and determine the psychometric properties of the Persian version of the Compassionate Care Assessment Tool (CCAT) ©.**Methods:** The study was carried out to translate and validate of the CCAT©. After securing permission from the designer of the tool and translating it, the psychometric properties were determined through examining face validity, construct validity, internal consistency, and test/retest reliability. With regard to construct validity, confirmatory factor analysis was used so that 300 patients in internal and surgery wards were selected by a simple random sampling method from three hospitals. Data were analyzed using SPSS (v.24.0) and LISREL statistical software version 8.8.**Results:** The results of the confirmatory factor analysis supported the validity and reliability of the Persian version of the CCAT© and its four factors. The reliability of the tool and internal consistency were confirmed through test/retest method with two weeks' interval. At the two areas of importance and provision of compassionate care, Cronbach's  $\alpha$  coefficient equaled to 0.918 and 0.933 and intraclass consistency equaled to 0.848 and 0.907 respectively.**Conclusion:** The results showed that the Persian version of the CCAT© was adequately valid and reliable for Iranian patients. Given the acceptable psychometric parameters of the tool, using it in future studies to measure importance and provision of compassionate nursing care to Iranian patients at internal and surgery wards is recommended.© 2020 The authors. Published by Elsevier B.V. on behalf of the Chinese Nursing Association. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## What is known?

- Nursing care of patients should be holistic and compassionate care is a fundamental part of nursing care that named as the art of nursing.
- The Compassionate Care Assessment Tool (CCAT) © by Burnell is one of the tools specially designed to assess compassionate nursing from the patient's perspective.

## What is new?

- This study confirmed the validity and reliability of the Persian version of the CCAT©, which provides an instrument to evaluate compassionate care in viewpoints of patients in Iran.

## 1. Introduction

Companionate is the nursing profession's foundation; indeed it is one of the professional standards of nursing performance [1]. The American Nurses Association (ANA) defines compassion as a vital element and feature of nursing and a requisite for improvement of nursing care quality and patients' recovery [2].

Compassion may have direct influence on the quality of care for the patient and the patients' assessment of the nursing services is usually based on the compassion they feel in the care [3–5].

\* Corresponding author.

E-mail address: [A\\_abdi61@yahoo.com](mailto:A_abdi61@yahoo.com) (A. Abdi).

Peer review under responsibility of Chinese Nursing Association.

Because of this, the terms “compassion” and “compassionate care” are usually used interchangeably in nursing literature to highlight the same meaning [6]. Emotional compassion develops as a result of witnessing others’ pain and hardship and leads to attempts to help [7]. Compassion begins with empathy and accomplishes with willingness and taking measure to decrease or remove the pain and suffering of others [5]. It is defined as perception and awareness of others’ pain and willingness to remove it [8,9].

Compassion is not a mere feeling in nursing; it is a connection between the nurse and patient that may lead the measures and decisions taken by the nurse so that the outcome will be an outstanding clinical performance by the nurse [10]. Compassionate nursing care means perceiving the pain and suffering of the patient and taking measures to decrease or remove the pain by adopting proper nursing intervention consistent with cultural backgrounds of patients and nurses [11]. Gordon notes that compassionate care is a patient-centered process to examine the situation and program the care [12]. Perry maintains that signs of compassionate nursing include being considerate and accurate in dealing with the patient’s problems, being committed to recognizing the patient, and attempting to remove the patient’s pain along with keeping a respectful relationship with the patient [13].

Thereby, compassionate nursing care is not limited to demonstrating empathy and dealing with the disease but also encompasses empowerment of patients along with providing appropriate nursing care through perceiving physical and emotional needs and emotional problems of patients [14–16]. Compassionate nursing care may lead to establishing an effective relationship between the provider and receiver of the care, which results in faster recovery of the patient and better therapeutic results [17,18]. It may also lead to higher satisfaction in the patients, safer care services, time and cost-saving, satisfaction in the personnel, sense of usefulness, higher self-confidence, and development of coping behavior in patients [17,19]. Lack of compassion in providing care is an imperfection and a sign of the low quality of health care [20]. Some consider it as the best and most valuable gift that a nurse can give a patient in health systems [9,21]. It entails recognizing and perceiving the patients’ needs and expectations with compassionate care [22–24]. Without patients’ feedback, nurses risk providing care only based on their professional perception and presumptions and neglecting patients’ attitudes and viewpoints [25]. Without feedback from patients as the receivers of care and failing to determine characteristics and behaviors perceived as compassionate nursing, it is not possible to supervise and assess compassionate nursing or determine its effectiveness [5,26,27]. The Royal College of Nursing in the UK and the American Nursing Association (ANA) have emphasized supervising and assessing compassionate care provision as a priority for professional development, even if it challenges the status quo [2,28].

Wear and Zarconi believed that rather than the patient, nowadays, more emphasis is put on the diseases, and naturally, this approach leads to negligence of attitudes and needs of the patient [29]. As recommended by research works, despite the importance and role of compassionate care in nursing, today’s nursing is unfortunately mostly based on quantitative evidences and clinical skills, while ethics and compassion are less emphasized [30–32].

Therefore, this aspect of nursing has been neglected and deemphasized as moral and arbitrary aspects of nursing [33]; so that this key need of the patient has been overshadowed [34]. In this spirit, about 50% of patients in the USA in 2010 noted that compassion is a missing element of care services [10].

There are several hurdles in the way of promoting compassionate care and probably one of the most important of them is the absence of a valid and reliable tool to assess and monitor provision of compassionate care [34,35]. It is essential to develop a valid and

reliable tool to assess and monitor compassionate care in clinical settings [28,35]. Several tools are available to assess compassion [2]; however, there are a limited number of tools to measure compassion in health personnel and provided care and in particular the care provided by nurses. One of the tools specially designed to assess compassionate nursing from the patient’s perspective is the Compassionate Care Assessment Tool (CCAT) © by Burnell. The reason for choosing this tool was that it is specifically focused on assessing the importance and provision of compassionate care from the viewpoint of patients hospitalized in internal and surgery wards. In addition, the subscales of the tool provide better coverage of the concept of compassionate nursing care [34,35]. The tool was designed by Burnell in 2011 with 28 statements to assess compassionate nursing care in cancer patients. Afterward, Burnell and Agen (2013) examined the tool’s validity for patients in the USA in internal and surgery wards. The new tools consist of 20 statements and four factors, including meaningful connection (eight statements), patients’ expectations (five statements), caring attributes (four statements), and capable practitioner (three statements) [9].

Despite the evidence about the direct relationship between low quality of care and lack of compassion and the weight of compassionate care as a construct in the nursing field that can be manipulated [34], it has been rarely dealt with in Iran. Explanations for this include the subjective and multi-aspect nature of the concept and the fact that is a function of cultural ground of the society. In addition, Iranian nurses are not completely familiarized with the concept, and there is no normalized and valid tool to measure it in Iran. This highlights the need for a valid and reliable tool to assess and survey the provision of compassionate nursing care in clinical settings. The present study is an attempt to translate and validate of the CCAT© for Iranian patients hospitalized in internal and surgery wards.

## 2. Material and methods

The study was carried out during May 2017 to September 2018 as a methodological work to translate the CCAT© into Persian and validate the tool for the patients hospitalized in internal and surgery wards. The study population was the patients hospitalized in three educational Kermanshah-based hospitals affiliated with Kermanshah University of Medical Sciences. Kermanshah city is home to the largest hospitals in the west of Iran and hosts patients from four neighboring provinces. To examine the validity of the tool, 300 patients were selected through random cluster method from three hospitals in Kerman city.

Inclusion criteria were age range 18–65 years, minimum hospitalization term of three days, consent to participate in the study, Persian speaking and understanding abilities, no pain, and stable condition. The questionnaires not fully completed were excluded.

The proposed method by Wild et al. was followed to translate and validate the tool [36]. At first, a written permission from the designer of the tool was secured, and then two translators translated it from English into Persian. Then the two versions of the questionnaire were compared and combined to obtain on translation. Afterward, the Persian version was translated into English by three translators (one of them was a native English speaker). The three English versions were compared by a supervisor and consistency with the original tool was confirmed. The designer also received a copy of the translated tool for confirmation. With confirmation by the designer of the CCAT©, the Persian version of the tool was examined in terms of face validity, construct validity, internal consistency, and test/retest reliability. Qualitative face validity of the tool was examined based on cognitive questions from 15 patients hospitalized in Kermanshah-based hospitals. The

translated tool was provided to 15 patients, including both genders who had been hospitalized for at least three days in internal and surgery wards. They were asked to state their opinions about clarity, ease of understanding, and understandability of the statements. The Persian version passes this stage with minor alterations before confirmatory factor analysis.

For construct validity stage, the patients were first briefed about the objective and method of the study and an informed letter of consent was signed by them. Afterward, they were asked to fill out a demographical information form (age, gender, education, marital status, ward, the term of hospitalization, and hospitalization occasions in the past year). Then they were asked to fill out the CCAT©; each questionnaire was filled out in 25–30 min. The participants were asked to rank each one of 20 statements of the tool in terms of importance and provision of cares from the aspect covered by the statements. The statements were designed based on Likert's four-point scale (not important till extremely important in term of importance and never to all of the time for provision of care). Confirmatory factor analysis was performed for both aspects of importance and provision of care.

Construct validity of the tool was examined in LISREL statistical software version 8.8 using confirmatory factor analysis to examine the goodness of fit of the tool's factors. Several goodnesses of fit indices are available for this purpose, and it is recommended to use more than one index [37,38]. Here, Squared-chi, Minimum Discrepancy Function by degrees of freedom divided (CMIN/DF), root mean square error of approximation (RMSEA), adjusted goodness of fit index (AGFI), incremental fit index (IFI), and comparative fit index (CFI) were used.

Internal consistency and test/retest reliability were examined in SPSS (v.24). To determine internal consistency, Cronbach's  $\alpha$  coefficient was computed for the whole tool and the subscales. To examine test/retest reliability and interclass consistency coefficient, 30 patients hospitalized in internal and surgery wards of the three educational hospitals in Kermanshah City who met the inclusion criteria were selected to fill the tool two times in two weeks' interval. These participants did not participate in the structural validity stage.

### 3. Results

Following translation and confirmation of the translation by the designer, face and structural validity construct validity were examined. In the case of face validity, the tool was examined by the patients in terms of clarity and simplicity. In construct validity, the collected data from 300 patients hospitalized in internal and surgery wards at three educational hospitals in Kermanshah city were analyzed. The mean age of the patients was  $43.6 \pm 13.66$  years with age range of 18–65. Among the patients, 63.7% were men, 56.3% did not have high school diplomas, and 51.0% were at the internal ward. Demographic characteristics of the participants are listed in Table 1.

About construct validity and confirmatory factor analysis for a tool with four factors in two areas of importance and provision of the services, the results showed that squared-chi to the degree of freedom ( $df$ ) ratios were 2.768 and 2.464 respectively ( $P < 0.001$ ) and RMSEA values were 0.083 and 0.070 respectively. For the rest of the indices including AGFI, CFI, NFI, IFI, and SRMR, goodness of fit was at acceptable level. Notably, these indices were higher in the area of importance of the services. Concerning AGFI, the goodness of fit was not acceptable in none of the areas (Table 2). The conceptual model of provision and importance of compassionate care is provided in Figs. 1 and 2.

Examining internal consistency of the Persian version of the CCAT© in the two areas of importance and provision of compassionate services from the participants' point of view indicated good

internal consistency of the whole tool in the area of importance of services (Cronbach's  $\alpha = 0.918$ ). All the aspects except for "competent practitioner" had a high internal consistency, higher than 0.7; this might be due to the small number of statements (three statements). In the area of provision of services, the tool had an excellent internal consistency (Cronbach's  $\alpha = 0.933$ ). In addition, all the three subscales had good internal consistency higher than 0.7 (Table 3).

The results of test/retest with two weeks' interval showed intraclass consistency coefficient in the areas of importance and provision of compassionate cares were 0.848 and 0.907 respectively (Table 4).

### 4. Discussion

In this study, the validity and reliability of the Persian version of the CCAT© were supported. Confirmatory factor analysis results for the CCAT© supported acceptable goodness of fit of the tool in the two areas of importance and provision of services. Thus, the Persian version of the CCAT© with four factors and 20 statements is supported in terms of structural validity.

With regard to the reliability of the tool, stratified internal consistency index – i.e., an estimate of consistency of scores from two or more evaluations to measure a quantitative index – indicated that the tool and the four factors had an acceptable consistency in the two areas of importance and provision of services from the patients' point of view. In addition, the internal consistency of the whole CCAT© and its four factors was acceptable based on Cronbach's  $\alpha$  coefficient in the two areas of importance and provision of services. These findings are consistent with Burnell's study (2013), which obtained Cronbach's  $\alpha > 0.7$  for the whole tool and four factors [9].

Sinkler et al. (2017) evaluated the compassionate assessment tools in health services and showed that one of the weaknesses of the available tools is the lack of validation in different societies and cultures [34]. The present study deals with this disadvantage. As the results showed, the CCAT© is a reliable, valid, and replicable tool. Since providing care with compassion is part of nurses' job, provision of compassionate services is a key element in satisfaction of patients with nursing cares and provision of quality health services to patients [39–42]. Assessing patients' satisfaction with compassionate nursing care is a reliable way to enable nurses to understand patients' needs and expectations with regard to cares. This measure constitutes the centerpiece in the provision of proper and timely services [43–45].

A literature review showed that compassion is a function of culture [46], and the dominant culture in society affects how compassionate behaviors are formed [47]. Therefore, nurses' behaviors are a function of culture so that compassionate behaviors

**Table 1**  
Demographic characteristics of the participants (n = 300).


Variables		n (%)
Gender	Male	191 (63.7)
	Female	109 (36.3)
Age group (year)	18–35	95 (31.7)
	36–45	53 (17.7)
	46–55	70 (23.3)
	56–65	82 (27.3)
Marital status	Married	226 (75.3)
	Unmarried	74 (24.7)
Education level	Below diploma	169 (56.3)
	Diploma	86 (28.7)
	Academic degree	45 (15.0)
Ward	Internal	153 (51.0)
	Surgery	147 (49.0)

**Table 2**  
Goodness of fit indices of confirmatory factor analysis of the Persian version of the Compassionate Care Assessment Tool.

Goodness of fit indices	RMSEA (90%CI)	CFI	NFI	IFI	AGFI	sRMR	$\chi^2$	P
Provision	0.070 (0.061–0.079)	0.98	0.96	0.98	0.85	0.047	2.464	<0.001
Importance	0.083 (0.075–0.091)	0.96	0.95	0.96	0.82	0.046	2.768	<0.001

and behavioral specifications that demonstrate compassion in nurses are affected by traditions, customs, culture, values, beliefs, and sociopolitical structure of the society [18,33,48,49]. The dominant culture in Iran is Iranian-Islamic culture, and nursing is assumed as worshipping. There are verses in the Holy Quran and hadiths about the value of looking after patients, which is the main job of nurses. That is, when a nurse provides cares with religious intentions to a patient, their job becomes a holy and heavenly job and resemblance of the best deeds. Therefore, it is highly fruitful to

examine and perceive the aspects and specifications of compassionate nursing care in clinical settings based on cultural and social differences between Iranian-Islamic and Western societies and highlight the deep effects of these differences on the perception and expectation of patients with compassionate care. This highlights the need for and recommends conducting qualitative studies to better understand different and unknown aspects of the concept of compassionate nursing. Among the limitations of the present study was the lack of a suitable tool for Iranian society to conduct



**Fig. 1.** The conceptual model of the importance of compassionate care. F = Factor. IM=Importance.




Fig. 2. The conceptual model of the provision of compassionate care. F = Factor. PR=Provision.

**Table 3**  
Internal consistency of the Persian version of the Compassionate Care Assessment Tool and subscales in the area of provision and importance of compassionate care.

Subscale	Number of items	Cronbach's $\alpha$	
		Importance	Provision
Meaningful connection	8	0.799	0.835
Patients' expectations	5	0.745	0.818
Caring attributes	4	0.745	0.821
Capable practitioner	3	0.657	0.716
Total	20	0.918	0.933

criterion-related validity examination.

### 5. Conclusion

This study confirmed the cultural validation of the CCAT. As to the advantages of the tool examined in this work, it is notable that provision of compassionate nursing care was evaluated by the patients at internal and surgery wards. Thus, the collected information helps us gain deeper understanding of the needs and expectations of patients with reception of compassionate cares and the amount of such services to meet the needs and expectations of patients.

### Ethics approval and consent to participate

The study is confirmed by research ethics committee of University of Social Welfare and Rehabilitation. (IR.USWR.REC.1396.250).

**Table 4**  
Intra-class correlation of the Persian version of the Compassionate Care Assessment Tool and subscale.

Subscale		ICC	95%CI
Meaningful connection	Importance	0.889	0.767–0.947
	Provision	0.935	0.863–0.969
Patients' expectations	Importance	0.847	0.679–0.927
	Provision	0.905	0.801–0.955
Caring attributes	Importance	0.852	0.688–0.929
	Provision	0.933	0.859–0.968
Competent practitioner	Importance	0.848	0.610–0.928
	Provision	0.729	0.430–0.871
Total score of tool	Importance	0.848	0.682–0.928
	Provision	0.907	0.804–0.956

Note: ICC= Intra-class correlation.

## Availability of data and material

Data available by contacting the corresponding author.

## Funding

The study was funded by the University of Social Welfare and Rehabilitation.

## CRedit authorship contribution statement

**Aliakbar Vaisi-Raygai:** Conceptualization, Data Curation, Methodology, Writing Original Draft, Writing-Review & Editing. **Asghar Dalvandi:** Supervision, Conceptualization, Methodology, Writing Original Draft. **Kian Nourozi:** Conceptualization, Methodology, Writing Original Draft. **Abbas Ebadi:** Conceptualization, Methodology, Writing Original Draft, Validation, Formal Analysis. **Mahdi Rahgozar:** Conceptualization, Methodology, Writing Original Draft, Validation. **Rostam Jalali:** Conceptualization, Methodology, Writing Original Draft, Validation. **Nader Salari:** Conceptualization, Methodology, Writing Original Draft, Formal Analysis, Software. **Alireza Abdi:** Conceptualization, Methodology, Writing Original Draft, Data Curation, Writing- Review & Editing, Software.

## Declaration of competing interest

The authors declare that they have no conflict of interest.

## Acknowledgement

The present paper is part of a PhD dissertation in nursing approved by the University of Rehabilitation Medical Sciences. The authors wish to express their gratitude to all participating patients for their contribution and the Department of Research of Tehran University of Social Welfare and Rehabilitation for their support of the study plan.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ijnss.2020.12.004>.

## References

- [1] Brown SJ. Evidence-based nursing. The research-practice connection. Sudbury, MA: Jones & Bartlett Learning; 2012.
- [2] American Nurses Association. Nursing scope and standards of practice. American Nurses Association; 2010.
- [3] Archer KR, Castillo RC, Wegener ST, Abraham CM, Obremskey WT. Pain and satisfaction in hospitalized trauma patients: the importance of self-efficacy and psychological distress. *J Trauma Acute Care Surg* 2012;72(4):1068–77. <https://doi.org/10.1097/JA.0b013e3182452df5>.
- [4] Papastavrou E, Andreou P, Tsangari H, Merkouris A. Linking patient satisfaction with nursing care: the case of care rationing - a correlational study. *BMC Nurs* 2014;13(1):26. <https://doi.org/10.1186/1472-6955-13-26>.
- [5] Kret DD. The qualities of a compassionate nurse according to the perceptions of medical-surgical patients. *Medsurg Nurs* 2011;20(1):29.
- [6] Bloomfield J, Pegram A. Care, compassion and communication. *Nurs Stand* 2015;29(25):45–50. <https://doi.org/10.7748/ns.29.25.45.e7653>.
- [7] DeSteno D. Compassion and altruism: how our minds determine who is worthy of help. *Curr Opin Behav Sci* 2015;3:80–3. <https://doi.org/10.1016/j.cobeha.2015.02.002>.
- [8] Chochinov HM. Dignity and the essence of medicine: the A, B, C, and D of dignity conserving care. *BMJ* 2007;335(7612):184–7. <https://doi.org/10.1136/bmj.39244.650926.47>.
- [9] Burnell L, Agan DL. Compassionate care: can it be defined and measured? The development of the compassionate care assessment tool. *Int J Caring Sci* 2013;6(2):180–7.
- [10] Papadopoulos I, Taylor G, Ali SL, Aagard M, Akman O, Alpers LM, et al. Exploring nurses' meaning and experiences of compassion: an international online survey involving 15 countries. *J Transcult Nurs* 2017;28(3):286–95. <https://doi.org/10.1177/1043659615624740>.
- [11] Papadopoulos I. Courage, compassion and cultural competence. Amsterdam: The 13th Anna Reynvaan Lecture; 2011. 2011.
- [12] Gordon S. Beyond the bedside: what do nurses really do? *Ohio Nurses Rev* 2006;81(4):1. PMID: 16910198.
- [13] Perry B. Conveying compassion through attention to the essential ordinary. *Nurs Older People* 2009;21(6):14–21. <https://doi.org/10.7748/nop2009.07.21.6.14.c7137>.
- [14] Boyle DA. Countering compassion fatigue: a requisite nursing agenda. *Online J Issues Nurs* 2011;16(1):2. <https://doi.org/10.3912/ojin.vol16no01man02>.
- [15] Davison N, Williams K. Compassion in nursing. 1: defining, identifying and measuring this essential quality. *Nursing Times* 2009;105(36):16–8. PMID: 19835194.
- [16] Dewar B, Cook F. Developing compassion through a relationship centred appreciative leadership programme. *Nurse Educ Today* 2014;34(9):1258–64. <https://doi.org/10.1016/j.nedt.2013.12.012>.
- [17] Youngson R. Compassion in Healthcare: the missing dimension of healthcare reform. Caregiver stress and staff support in illness, dying, and bereavement. Oxford University Press; 2011. p. 49–61.
- [18] Meyer RML, Li A, Klaristenfeld J, Gold JL. Pediatric novice nurses: examining compassion fatigue as a mediator between stress exposure and compassion satisfaction, burnout, and job satisfaction. *J Pediatr Nurs* 2015;30(1):174–83. <https://doi.org/10.1016/j.pedn.2013.12.008>.
- [19] Lown BA, Rosen J, Marttila J. An agenda for improving compassionate care: a survey shows about half of patients say such care is missing. *Heal Aff* 2011;30(9):1772–8. <https://doi.org/10.1377/hlthaff.2011.0539>.
- [20] Blomberg K, Griffiths P, Wengström Y, May C, Bridges J. Interventions for compassionate nursing care: a systematic review. *Int J Nurs Stud* 2016;62:137–55. <https://doi.org/10.1016/j.ijnurstu.2016.07.009>.
- [21] Papadopoulos I, Shea S, Taylor G, Pezzella A, Foley L. Developing tools to promote culturally competent compassion, courage, and intercultural communication in healthcare. *J Compassionate Heal Care* 2016;3(1):1–10. <https://doi.org/10.1186/s40639-016-0019-6>.
- [22] van der Cingel M. Compassion: the missing link in quality of care. *Nurse Educ Today* 2014;34(9):1253–7. <https://doi.org/10.1016/j.nedt.2014.04.003>.
- [23] Zamanzadeh V, Valizadeh L, Rahmani A, van der Cingel M, Ghafourifard M. Factors facilitating nurses to deliver compassionate care: a qualitative study. *Scand J Caring Sci* 2018;32(1):92–7. <https://doi.org/10.1111/scs.12434>.
- [24] Winch S, Henderson A, Jones J. Recognizing the dialectic of compassionate care in the workplace: feedback from nurse educators. *J Continuing Educ Nurs* 2015;46(5):228–32. <https://doi.org/10.3928/00220124-20150420-03>.
- [25] Curtis K, Horton K, Smith P. Student nurse socialisation in compassionate practice: a Grounded Theory study. *Nurse Educ Today* 2012;32(7):790–5. <https://doi.org/10.1016/j.nedt.2012.04.012>.
- [26] Straughair C. Exploring compassion: implications for contemporary nursing. Part 2. *Br J Nurs* 2012;21(4):239–44. <https://doi.org/10.12968/bjon.2012.21.4.239>.

- [27] Badger K, Royse D. Describing compassionate Care: the burn survivor's perspective. *J Burn Care Res* 2012;33(6):772–80. <https://doi.org/10.1097/BCR.0b013e318254d30b>.
- [28] Sturgeon D. Measuring compassion in nursing. *Nurs Stand* 2008;22(46):42–3. <https://doi.org/10.7748/ns2008.07.22.46.42.c6599>.
- [29] Wear D, Zarconi J. Can compassion be taught? let's ask our students. *J Gen Intern Med* 2008;23(7):948–53. <https://doi.org/10.1007/s11606-007-0501-0>.
- [30] Lee Y, Seomun G. Development and validation of an instrument to measure nurses' compassion competence. *Appl Nurs Res* 2016;30:76–82. <https://doi.org/10.1016/j.apnr.2015.09.007>.
- [31] Baljani E, Azimi N, Hosseinloo A. A survey on nurses' perception of the importance of caring behaviors and factors affecting its provision. *Evidence Based Care* 2012;2(1):13–21. <https://doi.org/10.22038/EBCJ.2012.377>.
- [32] Henderson A, Jones J. Developing and maintaining compassionate care in nursing. *Nurs Stand* 2017;32(4):60–9. <https://doi.org/10.7748/ns.2017.e10895>.
- [33] Nurs Forum. In: Schantz ML, editor. *Compassion: a concept analysis*, vol. 42; 2007. p. 48–55. <https://doi.org/10.1111/j.1744-6198.2007.00067.x>.
- [34] Sinclair S, Russell LB, Hack TF, Kondejewski J, Sawatzky R. Measuring compassion in healthcare: a comprehensive and critical review. *Patient - Patient - Centered Outcomes Res* 2017;10(4):389–405. <https://doi.org/10.1007/s40271-016-0209-5>.
- [35] Papadopoulos I, Ali SL. Measuring compassion in nurses and other healthcare professionals: an integrative review. *Nurse Educ Pract* 2016;16(1):133–9. <https://doi.org/10.1016/j.nepr.2015.08.001>.
- [36] Wild D, Grove A, Martin M, Eremenco S, McElroy S, Verjee-Lorenz A, et al. Principles of good practice for the translation and cultural adaptation process for patient-reported outcomes (PRO) measures: report of the ISPOR task force for translation and cultural adaptation. *Value Heal* 2005;8(2):94–104. <https://doi.org/10.1111/j.1524-4733.2005.04054.x>.
- [37] Seo DC, Torabi MR, Blair EH, Ellis NT. A cross-validation of safety climate scale using confirmatory factor analytic approach. *J Saf Res* 2004;35(4):427–45. <https://doi.org/10.1016/j.jsr.2004.04.006>.
- [38] Bentler PM. Comparative fit indexes in structural models. *Psychol Bull* 1990;107(2):238–46. <https://doi.org/10.1037/0033-2909.107.2.238>.
- [39] Bradshaw A. Measuring nursing care and compassion: the McDonaldised nurse? *J Med Ethics* 2009;35(8):465–8. <https://doi.org/10.1136/jme.2008.028530>.
- [40] Cornwell J, Donaldson J, Smith P. Nurse Education Today: special issue on compassionate care. *Nurse Educ Today* 2014;34(9):1188–9. <https://doi.org/10.1016/j.nedt.2014.06.001>.
- [41] Hill R. Compassion, quality and standards of care. *Nurse Prescr* 2010;8(3):100–1. <https://doi.org/10.12968/npre.2010.8.3.100>.
- [42] Mooney H. Can you measure compassion. *Nursing Times* 2009 (Online)(accessed 2007 15) Available from: <http://www.nursingtimes.net/whats-new-in-nursing/management/can-you-measure-compassion/5000543.article>.
- [43] Curtis K. 21st Century challenges faced by nursing faculty in educating for compassionate practice: embodied interpretation of phenomenological data. *Nurse Educ Today* 2013;33(7):746–50. <https://doi.org/10.1016/j.nedt.2013.05.007>.
- [44] Lown BA, Muncer SJ, Chadwick R. Can compassionate healthcare be measured? The Schwartz center compassionate care Scale™. *Patient Educ Couns* 2015;98(8):1005–10. <https://doi.org/10.1016/j.pec.2015.03.019>.
- [45] McCaffrey G, McConnell S. Compassion: a critical review of peer-reviewed nursing literature. *J Clin Nurs* 2015;24(19–20):3006–15. <https://doi.org/10.1111/jocn.12924>.
- [46] Spandler H, Stickley T. No hope without compassion: the importance of compassion in recovery-focused mental health services. *J Ment Heal* 2011;20(6):555–66. <https://doi.org/10.3109/09638237.2011.583949>.
- [47] Cornelius E. The motivation and limits of compassion. *HTS Theologisches Studien/Theol Stud* 2013;69(1):1–7. <https://doi.org/10.4102/hts.v69i1.1189>.
- [48] Dewar B, Adamson E, Smith S, Surfleet J, King L. Clarifying misconceptions about compassionate care. *J Adv Nurs* 2014;70(8):1738–47. <https://doi.org/10.1111/jan.12322>.
- [49] Burnell L. *Compassionate care: the patient perspective*. ProQuest Dissertations and Theses. University of San Diego; 2011. <https://search.proquest.com/docview/892725774?accountid=27797>.