

## Research Article

# Type IV Secretion System Is Not Involved in Infection Process in Citrus

Tiago Rinaldi Jacob,<sup>1</sup> Marcelo Luiz de Laia,<sup>2</sup> Leandro Marcio Moreira,<sup>3</sup>  
Janaína Fernandes Gonçalves,<sup>2</sup> Flavia Maria de Souza Carvalho,<sup>4</sup>  
Maria Inês Tiraboschi Ferro,<sup>1</sup> and Jesus Aparecido Ferro<sup>1</sup>

<sup>1</sup> Faculdade de Ciências Agrárias e Veterinárias de Jaboticabal, Departamento de Tecnologia, Universidade Estadual Paulista (UNESP), 14.884-900 Jaboticabal, SP, Brazil

<sup>2</sup> Departamento de Engenharia Florestal, Faculdade de Ciências Agrárias, Universidade Federal dos Vales do Jequitinhonha e Mucuri, 39.100-000 Diamantina, MG, Brazil

<sup>3</sup> Departamento de Ciências Biológicas (DECBI), Instituto de Ciências Exatas e Biológicas (ICEB) and Núcleo de Pesquisas em Ciências Biológicas (NUPEB), Universidade Federal de Ouro Preto, 35.400-000 Ouro Preto, MG, Brazil

<sup>4</sup> Embrapa Informática Agropecuária, CNPTIA-EMBRAPA, Campinas, SP, Brazil

Correspondence should be addressed to Jesus Aparecido Ferro; [jesus@fcav.unesp.br](mailto:jesus@fcav.unesp.br)

Received 17 August 2013; Accepted 10 January 2014; Published 23 February 2014

Academic Editor: Joseph Falkinham

Copyright © 2014 Tiago Rinaldi Jacob et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The type IV secretion system (T4SS) is used by Gram-negative bacteria to translocate protein and DNA substrates across the cell envelope and into target cells. *Xanthomonas citri* subsp. *citri* contains two copies of the T4SS, one in the chromosome and the other is plasmid-encoded. To understand the conditions that induce expression of the T4SS in *Xcc*, we analyzed, *in vitro* and *in planta*, the expression of 18 ORFs from the T4SS and 7 hypothetical flanking genes by RT-qPCR. As a positive control, we also evaluated the expression of 29 ORFs from the type III secretion system (T3SS), since these genes are known to be expressed during plant infection condition, but not necessarily in standard culture medium. From the 29 T3SS genes analyzed by qPCR, only *hrpA* was downregulated at 72 h after inoculation. All genes associated with the T4SS were downregulated on *Citrus* leaves 72 h after inoculation. Our results showed that unlike the T3SS, the T4SS is not induced during the infection process.

## 1. Introduction

*Xanthomonas citri* subsp. *citri* (*Xcc*) is a Gram-negative plant pathogenic bacteria that causes severe disease in many economically important citrus plants [1]. The citrus canker induced by *Xcc* is a destructive disease characterized by canker lesions on leaves, stems, and fruits; furthermore the pathogen induces defoliation, which results in reduced yield and premature fruit drop [2]. Control is difficult in areas where the disease is already established, and plant eradication is the only effective way to control and prevent the disease spread. Recurrent and severe attacks of the pathogen are responsible for serious economic losses in citrus groves around the world [2].

The focus of the present study is aimed at understanding the role of bacterial secretory systems, specifically the type IV secretory system (T4SS). Seven secretory systems are known and described in prokaryotic organisms, and each one is related to a physiological process [3–5]. Of these, the best studied is the type III secretory system (T3SS), which enables bacterial pathogens to deliver effector proteins into eukaryotic cells [6]. Some bacterial pathogens, including species from *Chlamydia*, *Xanthomonas*, *Pseudomonas*, *Ralstonia*, *Shigella*, *Salmonella*, *Escherichia*, and *Yersinia*, depend on the T3SS to induce damage to the host. In contrast, other organisms including *Agrobacterium tumefaciens*, *Helicobacter pylori*, and *Legionella pneumophila* depend on the type IV secretory system (T4SS) for virulence induction [7].

The T3SS contains at least twenty distinct proteins, which are subdivided into three parts. The basal body of the system [8–11] is associated with an ATPase and likely facilitates the entry of substrates into the secretion apparatus [12]. This basal body is also associated with a protein needle (the second part of the T3SS), which binds the bacterium to host cells and acts as a conduit for effectors secretion. The third part of the T3SS is composed of three proteins that are exported through the bacterial needle and form a pore on the surface of the host cell, which facilitates the export of toxins into the target cytoplasm [13].

On the other hand, the T4SS is a multiprotein complex that consists of a protein channel (encoded by *virB* and *virD*) through which proteins or protein-DNA complexes can be translocated between bacteria (cell-to-cell communication) and into host cells [14]. Translocation is driven by a number of cytoplasmic ATPases that potentially energize large conformational changes in the translocation complex [15]. There are three functional types of T4SS. The first type, found in many Gram-positive and Gram-negative bacteria and some Archaea, functions in conjugation and in the transfer of T-DNA into plant cells by *A. tumefaciens*. A second type of T4SS mediates DNA uptake in the transformation process (found in *H. pylori*). A third type of T4SS is used to transfer toxic effector proteins or protein complexes into the cytoplasm of host cells (found in *Bordetella pertussis*, *Legionella pneumophila*, *Pseudomonas aeruginosa*, and *Xcc*) [15].

Substrates transported by the T4SS modulate various cellular processes including apoptosis, vesicular traffic, and ubiquitination; furthermore, the number of type IV effector proteins continues to increase [16–19]. However, most of these substrates have not been functionally characterized, and their role in bacterial pathogenesis remains unknown [20].

In many bacteria, these two secretory systems types are well characterized; however, in *Xanthomonas* spp. only the T3SS has been intensively studied [1], in contrast to the limited number of studies involving the T4SS [21, 22]. In *Xcc* the T4SS deserves more attention, especially since the genome contains two copies of this system (chromosomal and plasmid-borne) [23]. The plasmid copy is homologous to others *Xanthomonas* that contain the T4SS on extra-chromosomal DNA. However, only *Xcc* and *Xanthomonas campestris* contain the chromosomal copy of the T4SS.

To understand the conditions that stimulate expression of T4SS genes in *Xcc* we used qPCR to analyze the *in vitro* and *in planta* expression of 18 ORFs from the T4SS (both chromosomal and plasmid copies) and 7 hypothetical ORFs flanking the T4SS. To validate the qPCR data, we compared expression of 29 ORFs from the T3SS, which are known to be expressed *in planta*.

Our results showed that the T4SS is not induced during the infection process in *Xcc*, but may be very important in cell-to-cell communication.

## 2. Materials and Methods

**2.1. Microbiological Procedures.** *Xcc* strain 306 was previously sequenced by da Silva et al. [23]. *Xcc* was maintained in sterile tap water and grown on nutrient agar (NA) medium

containing 3 g beef extract, 5 g peptone, and 15 g agar in 1 L of distilled water at 28°C. After 24 h, three single colonies were transferred to new NA plates and incubated for 12 h at 28°C. These *Xcc* solid cultures were used as pre-inoculum for *in planta* and *in vitro* studies. For *in planta* studies, pre-inoculum cultures were aseptically transferred to sterile flasks containing 50 mL of nutrient broth (NB) and the bacterial concentration was adjusted to 10<sup>8</sup> CFU/mL (OD<sub>600</sub> = 0.3). This bacterial suspension was taken up in a 1 mL needleless syringe and used to infiltrate orange leaves (*Citrus sinensis* cv. pera) grown in 20 L capacity pots. A total of nine plants were inoculated (20 leaves per plant) and maintained for 72 h in a growth room at 28°C, with a 12 h photoperiod and light intensity ~2,000 lux. Three plants were used for each incubation period. After multiplication, inoculated leaves were collected, sliced into thin strips with a razor blade, and placed in a beakers with sterile distilled water in an ice bath with gentle agitation. Samples from each plant were placed in separate beakers for bacterial exudation. After 5 min, leaf debris was removed by filtration through gauze, and bacterial cells were recovered by centrifugation at 5,000 ×g for 5 min at 4°C. Total RNA was extracted immediately, as described below. For *in vitro* studies, 1 mL of each *Xcc* suspension at 10<sup>8</sup> CFU/mL were aseptically transferred to three Erlenmeyer flasks and incubated for 12 h at 28°C with shaking (200 rpm). After incubation, bacterial cells were harvested by centrifugation at 5,000 ×g for 5 min. Total RNA was extracted as described below. Thus, we obtained three independent biological replicates for *Xcc* growth *in vitro* and *in planta*. All cultivation media were obtained from Difco Chemical Co., Detroit, USA. For confirmatory RT-qPCR, the same procedures were followed except the plant incubation period was only 72 h, due to RNA quality and concentration sufficient to yield reliable results. The time point 72 hours was the shortest period to allow obtaining satisfactory bacterial mass to the proposed analyzes involving gene expression. The bacterial cells concentration in short time (12 or 24 hours) don't allows this study. For this reason, some works are using culture media that mimic vegetable conditions to investigate the infectious process under these time scales [24, 25].

**2.2. Extraction of RNA from *Xcc*.** Each inoculated flask represented one independent biological replicate. RNA was extracted using an Illustra-RNAspin Mini RNA Isolation Kit (Amersham Biosciences) following the manufacturer's instructions. To ensure that samples did not contain DNA, PCR was performed using RNA samples treated with DNase I as a template. PCR conditions consisted of an initial denaturing step of 94°C for 3 min, followed by 35 cycles of a denaturation at 94°C for 30 s, annealing at 60°C for 30 s, and elongation at 72°C for 2 min. A final elongation step of 72°C for 4 min was performed, and then samples were maintained at 4°C until needed. The amplification reaction was conducted in a total volume of 25 µL containing 200 ng RNA in 2.5 µL, PCR buffer (2.5 µL, Invitrogen), 1.5 mM MgCl<sub>2</sub>, 0.2 mM dNTP, 300 nM 16S rRNA primer [26], and 1 U Taq DNA polymerase (Invitrogen). The products were electrophoresed in a 1% agarose gel with TAE buffer, stained

with ethidium bromide, and visualized using a UV transilluminator. No products were observed (data not shown). To verify the quality of extracted RNA, the samples were analyzed by electrophoresis in a 1% agarose gel using TAE buffer followed by ethidium bromide staining. The A260/280 ratio of RNA samples was measured, and RNA was quantified using a NanoDrop ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA). RNA samples were stored at  $-80^{\circ}\text{C}$  until needed.

**2.3. Gene Expression Using RT-qPCR.** First strand cDNA synthesis and all RT-qPCR reactions were done using the SuperScript III First-Strand Synthesis SuperMix for RT-qPCR (Invitrogen) as recommended by the manufacturer's specifications except for the amount of cDNA in each reaction, which was 20 ng. All PCR was performed with SYBR Green on a 7500 Real-Time PCR instrument (Applied Biosystems) using three biological replicates and three technical replicates (one for each biological replicate). PCR was for 2 min at  $50^{\circ}\text{C}$ , 10 min at  $95^{\circ}\text{C}$ , followed by 40 cycles of 15 s at  $95^{\circ}\text{C}$ , and 1 min at  $60^{\circ}\text{C}$ . To determine PCR efficiency, standard curves were generated using cDNA samples at five dilutions and measured in triplicate. *rpoB*, *atpD*, and *gyrB* were used as reference genes in all experiments [26]. To perform relative expression analysis, we used the  $2^{-\Delta\Delta\text{CT}}$  method [27]. Primer features are presented in Table 1.

### 3. Results and Discussion

**3.1. qPCR Results for the T3SS.** The results obtained from *Xcc* growing in *planta* or *in vitro* are presented in Figure 1. From the 29 T3SS genes analyzed by qPCR, only *hrpA* was downregulated at 72 h after inoculation (Figure 1(b)). *hpa1*, *hpaB*, *hrpD6*, *hrpE*, and *hrpF* showed the highest rates of expression, whereas *phaE*, *hrpG*, *hrpX*, and *hrcA* showed the lowest levels of expression. Interestingly, *hrpB4* and *hrpXct*, which were shown to be necessary for *Xcc* pathogenesis [28], were not among the most upregulated genes in this study.

**3.2. qPCR Results for the T4SS.** The results for the expression levels of T4SS genes in *planta* and *in vitro* are shown in Figure 2. All genes associated with the T4SS were downregulated on *Citrus* leaves 72 h after inoculation (Figure 2(c)). We also analyzed the expression of hypothetical genes located near *virB* in chromosomal DNA (XAC2606, XAC2611, XAC2613, and XAC2622) and in plasmid pXAC64 (XACb0035, XACb0042, XACb0043, XACb0048, and XACb0049) (Figure 2(d) and Table 1). ORFs representing hypothetical genes in both the chromosome and plasmid were down-regulated when *Xcc* was cultivated in *Citrus sinensis* for 72 h. An exception was XAC2611, where expression was similar when *Xcc* was cultivated in culture medium or *in planta* (Figure 2(d)).

Our results clearly show different gene expression profiles for the T3SS and T4SS in *Xcc* during *Citrus* infection. The T4SS was previously described in *A. tumefaciens*, where it mediates the transfer of DNA and protein substrates to plants and other organisms via a cell contact-dependent mechanism

[14]. In *Xcc*, the genome sequence revealed the presence of two *virB* operons, one on the chromosome and a second copy in the 64 kb plasmid pXAC64 [23]. The chromosomal genes *virB1*, *virB3*, *virB4*, *virB8*, *virB9*, and *virB11* and the plasmid-encoded genes *virB1*, *virB2*, *virB4*, *virB6*, *virB9*, and *virB11* were all down-regulated in *planta* (Figure 2(c)). This was somewhat surprising because the T4SS is essential for a successful infection in many Gram-negative pathogenic bacteria [29]. The fact that ORFs representing hypothetical genes in both the chromosome and plasmid near the *virB* genes were also down-regulated under the same condition (Figure 2(d)) indicates that they could be related to the T4SS system in *Xcc*. Recent studies in *Brucella suis* [30] and in *A. tumefaciens* [31] showed that alterations in *VirB8* result in protein dimerization, a process that modifies the T4SS structure and affects bacterial virulence. In *Streptococcus suis* [32], the knockout of the *virD4-89K* and *virB4-89K* of the T4SS eliminated the lethality of a highly virulent strain and impaired its ability to trigger host immune responses in mice. Recent studies continue to demonstrate the importance of the T4SS and its components in virulence; thus the lack of induction of *Xcc* T4SS genes in the present study is intriguing. However, in *Xanthomonas campestris* pv. *campestris* T4SS-deletion mutant displayed the same virulence as wild type and authors conclude that T4SS is not involved in pathogenicity in that *Xanthomonas* [33].

Wang et al. [34] presented evidence that *in planta* transfer of a 37 kb plasmid (pXcB) from *Xanthomonas aurantifolii* to *Xanthomonas citri* can occur via T4SS. Thus, at least the T4SS copy present in *Xcc* plasmid can play a role in horizontal gene transfer.

All but one of the twenty-nine genes from the T3SS of *Xcc* were up-regulated in *planta*. Laia et al. [28] showed that *Xcc* mutants containing mutations in *hrpB4* or *hrpXct* failed to cause disease and growth in citrus leaves was lower than the wild-type *Xcc* strain 306. These two genes, which are not among the most up-regulated genes in the present study, are part of the *hrp* (hypersensitive reaction and pathogenicity) system and comprise part of the T3SS [35]. In the related pathogen, *Xanthomonas campestris* pv. *vesicatoria* (*Xcv*), a *hrpB4* mutant was unable to cause disease in susceptible pepper plants or the hypersensitive reaction in pepper plants carrying the respective compatible R gene [36]. Previously, *hrpXv* was shown to be necessary for transcriptional activation of five *hrp* genes (loci *hrpB* to *hrpF*) [37], and *hrpB4* was required for complete functionality of the T3SS in *Xcv* [36]. Thus it is apparent that a gene does not have to be strongly up-regulated during infection to play an important role in virulence. Multiple genes, including *hpaA*, *hpaE*, *hpaF*, *hpaP*, *hrpB*, *hrpB1*, *hrpB2*, *hrpB7*, *hrpD5*, *hrpM*, *hrpW*, *hrcC*, *hrcQ*, *hrcS*, *hrcT*, and *hrcV*, were expressed at levels similar to *hrpXct* and *hrpB4* and are good candidates for further studies into their roles in citrus canker disease.

Among the most up-regulated genes, *hrpD6* warrants further attention. A *hrpD6* mutant of *Xanthomonas oryzae* pv. *oryzae* (*Xoo*) failed to trigger a hypersensitive response in tobacco and was nonpathogenic in rice because the mutation in *hrpD6* impacts the secretion of T3SS effectors, such as *Hpa1*, which is translocated through the T3SS [38].

TABLE 1: General features of the T3SS and T4SS primer sequences.

XAC ID	Gene	Gene ID	Product (NCBI Annotation)	Primers (Forward)	Primers (Reverse)	Amplicon (bp)	System
XAC0293	<i>hrpB</i>	1154364	ATP-dependent RNA helicase	GCCGTATCCCGTTGACGTT	ACGGCCCCCTTCATCTCTTTT	81	T3SS
XAC0393	<i>hpaF</i>	1154464	HpaF protein	GGCGAGCGTTTTGACAA	CCGGAGATCCTGGCAGTTT	57	T3SS
XAC0394	<i>hrpF</i>	1154465	HrpF protein	GCCGGACGCGAGTTG	CGGTGATCGAGTTGTTCAA	62	T3SS
XAC0396	<i>hpaB</i>	1154467	HpaB protein	GTCCGTGTACCGCAAGAC	TGACGCCGGCATCATG	53	T3SS
XAC0397	<i>hrpE</i>	1154468	HrpE protein	ACGAGCCCCAGAAGTCCAT	CCACGTTGAAGTCCAGATCATTT	66	T3SS
XAC0398	<i>hrpD6</i>	1154469	HrpD6 protein	ACCGATACGGTCACCCAAGA	GATCCGCCGCTTGGT	55	T3SS
XAC0399	<i>hrpD5</i>	1154470	HrpD5 protein	GCCCTGAAGTGGTTCGTTA	TCACGGACGGTCTTGCT	54	T3SS
XAC0400	<i>hpaA</i>	1154471	HpaA protein	CGCCACCCCAACAGAAAA	ACGGCATCAGCTCCAICAA	59	T3SS
XAC0401	<i>hrcS</i>	1154472	HrcS protein	CGACGATCTAGTGGGATTACCT	CGACCACCGGCAAGGA	68	T3SS
XAC0402	<i>hrcR</i>	1154473	Protein T3SS	ACACGGGAACCGGAAAAA	CCTTGGGCCAGATCTGTTGT	58	T3SS
XAC0403	<i>hrcQ</i>	1154474	HrcQ protein	GCAGCTGGAAGTGGACCAA	GGCTGCAAAACCCGACAAC	57	T3SS
XAC0404	<i>hpaP</i>	1154475	HpaP protein	TGTCGATGGCACGAAATACC	CCCAACCGGGCCATT	58	T3SS
XAC0405	<i>hrcV</i>	1154476	HrcV protein	GCCGCAAGTTGTGAGTT	CTCTGCCATATCGCGATTCC	58	T3SS
XAC0406	<i>hrcU</i>	1154477	HrcU protein T3SS	AGCGGGTGTCTTTG	TAGTCGCCGGCCGATTT	51	T3SS
XAC0407	<i>hrpB1</i>	1154478	HrpB1 protein	CTGCGGCCAGAGCTGAAG	ACCGCGCTTGATGGAATC	60	T3SS
XAC0408	<i>hrpB2</i>	1154479	HrpB2 protein	CAGCGCAGCAGATCAAGTTG	CGCCGACGCTGACATTG	69	T3SS
XAC0409	<i>hrcJ</i>	1154480	HrcJ protein	GAGCGGCACGCCAATC	TGTCGAGATCAAGCCATCCTT	54	T3SS
XAC0410	<i>hrpB4</i>	1154481	HrpB4 protein	GGACAACACGCGGATCGA	GTCCGTCCAGCGCTGAA	52	T3SS
XAC0411	<i>hrpB5</i>	1154482	Protein T3SS HrpB	CGCACGCCCTGGGATATG	AGGCCGGATTCTGTTCCA	63	T3SS
XAC0412	<i>hrcN</i>	1154483	T3SS ATPase protein	TCGATGGGCACCTGATTCTC	CGTCGATTGCCGGGTAFT	63	T3SS
XAC0413	<i>hrpB7</i>	1154484	HrpB7 protein	CCGGGTGATGGACAAG	GCTGCAGGATCGTTCGATA	60	T3SS
XAC0414	<i>hrcT</i>	1154485	HrcT protein	ATCCAGCAATCCGACAGCAT	CACCGGTGGGACAGTTT	54	T3SS
XAC0415	<i>hrcC</i>	1154486	HrcC protein	CAGATGGGGTGGATGTG	CCGTCCACGGTGTGGAT	59	T3SS
XAC0416	<i>hpaI</i>	1154487	HpaI protein	GCAGCAGGCCGGTCACT	TTCATCAGCATCTGGGTGTAITG	57	T3SS
XAC0459	<i>phaE</i>	1154530	monovalent cation/H <sup>+</sup> antiporter subunit E	CCATCCCCGGCTTCATCTG	GCGATGCCGTGAATGTTG	54	T3SS
XAC0618	<i>hrpM</i>	1154689	Glucosyltransferase MdoH	CAAGGGCCTGCACTGGAT	GCATCAACAGACCCCAATC	86	T3SS
XAC1265	<i>hrpG</i>	1155336	HrpG protein	CGGCAAGCCGATCGTATT	GAAAAGAGCAGCCAGGCAAT	54	T3SS
XAC1266	<i>hrpXct</i>	1155337	HrpX protein	GCCTACAGCTACATGATCACCAAT	TGCGGCCACTTCGTTGA	63	T3SS

TABLE 1: Continued.

XAC ID	Gene	Gene ID	Product (NCBI Annotation)	Primers (Forward)	Primers (Reverse)	Amplicon (bp)	System
XAC1520	<i>hrcA</i>	1155591	Heat-inducible transcriptional repressor	AGTTCGGGTTTCGGCATAATC	CGGTTCTGCACCTCGTTGT	91	T3SS
XAC1994	<i>hrpX</i>	1156064	HrpX-like protein	CTGGTCTGATGGAGCTTTCT	GCGTACTCGTCCACGATCAA	90	T3SS
XAC2047	<i>phaE</i>	1156118	PHA synthase subunit	AGCAGGGTGCATCGAAGAAG	TTTTTCGGAGCCGCCCTTTT	63	T3SS
XAC2606	—	1156677	Hypothetical protein	CCCTCAGTGTGGGCCAGAT	AGGAGGGCGGTTGAGGGGATA	58	H_T4SS_C
XAC2611	—	1156682	Hypothetical protein	GGGAGCCCCCAATGTG	GGAGGCGTGGGAATATCATCT	59	H_T4SS_C
XAC2612	<i>virB6</i>	1156683	VirB6 protein	TGCAGCAGGGAGGATTAGGA	CGGCGGTGCGGTGAT	56	T4SS_C
XAC2613	—	1156684	Hypothetical protein	CGAAGGGAGATGGAAGCAGTT	GCTCAACAGATCGGCTGGAA	59	H_T4SS_C
XAC2614	<i>virB4</i>	1156685	VirB4 protein	GACGTCGCTTGGTGGAGAAA	TTCCCTGGGCGCAATCT	53	T4SS_C
XAC2615	<i>virB3</i>	1156686	VirB3 protein	GCCCCGCCATGTTTTTG	CGCCCGCACCAATGAA	54	T4SS_C
XAC2616	<i>virB2</i>	1156687	VirB2 protein	TGCGGATGCCAAAATG	CAGCAACGAAATGAATAGCAATG	62	T4SS_C
XAC2617	<i>virB1</i>	1156688	VirB1 protein	CGTATCGAGTCGTGCGGTAAT	CACCAAGGCGTCCACCAA	57	T4SS_C
XAC2618	<i>virB11</i>	1156689	VirB11 protein	CGCTGGTCAACCACATTC	CCTGGCGTCTCAATCGT	56	T4SS_C
XAC2619	<i>virB10</i>	1156690	VirB10 protein	CCATTGCAGCTCTTTGATTTCTC	GAATCTCCCCGCTTTCG	61	T4SS_C
XAC2620	<i>virB9</i>	1156691	VirB9 protein	GCTGAGCCCCAACGAAAAA	GCTCCATCCACCAAGTAA	59	T4SS_C
XAC2621	<i>virB8</i>	1156692	VirB8 protein	TCGTCATGGCAGATGCGTAT	CCGAAGTTGGGCTCAAGCT	61	T4SS_C
XAC2622	—	1156693	Hypothetical protein	GGCGGATTCCAAATATGCAGTT	TGGCCCCGATCAAGGTGTAG	59	H_T4SS_C
XAC2623	<i>virD4</i>	1156694	VirD4 protein	TCGCTGGAATCCATTGACCTA	TCAAATCCGAAACGCGAAAT	59	T4SS_C
XAC2922	<i>hrpW</i>	1156993	HrpW protein	GGCGGACACACCGACATC	TGCCITTTACAGGGTGGAGTCTT	58	T3SS
XAC3122	<i>hrpA</i>	1157193	ATP-dependent RNA helicase	TGCGCGCTCTGAAATCG	GGACTGGGTAAAGATCTTCAATGCA	76	T3SS
XACb0035	—	1158522	Hypothetical protein	CGAGACGAGAACGGCCATT	TGTCACGCAGCAATTCGAA	55	H_T4SS_P
XACb0036	<i>virB1</i>	1158523	VirB1 protein	CGTTACAACCTCGCCAAATAATG	CTGCCCGCTTGAGATTG	74	T4SS_P
XACb0037	<i>virB11</i>	1158524	VirB11 protein	CCCCGAAAGCGGTTGAGT	CGTGAGCGACCCCTTGTTG	58	T4SS_P
XACb0038	<i>virB10</i>	1158525	VirB10 protein	TCTACGTGAACAAGACCTCGAAT	CGCTTGGCTCCGTCGAT	64	T4SS_P
XACb0039	<i>virB9</i>	1158526	VirB9 protein	GCTCACGGCGGCAAGTT	GCACTTGTGTGACAGCATCATC	58	T4SS_P
XACb0040	<i>virB8</i>	1158527	VirB8 protein	CTCGAACAAAGTCAACACAGGAA	AACAGCAGCAACCCGATGA	60	T4SS_P
XACb0041	<i>virB6</i>	1158528	VirB6 protein	GCGCAGCTCGTCAATCGT	CGCGGACACCCCAACATG	55	T4SS_P
XACb0042	—	1158529	Hypothetical protein	CCGGGAACCCCGCTTTTA	GGAAGCCACTGCCGAAACT	52	H_T4SS_P



TABLE 1: Continued.

XAC ID	Gene	Gene ID	Product (NCBI Annotation)	Primers (Forward)	Primers (Reverse)	Amplicon (bp)	System
XACb0043	—	1158530	Hypothetical protein	TTCGGGGCCCTCATCTC	AGGGCGACTTGTGTGAACCTTGTT	58	H_T4SS_P
XACb0044	<i>virB5</i>	1158531	VirB5 protein	CAGGCCGCTACGAGAAG	GGGATTGGCGAACACAGGTCTT	57	T4SS_P
XACb0045	<i>virB4</i>	1158532	VirB4 protein	GGAACAGGCAAAAGACGTCAAC	CCCGACAAGTTTGAACGA	57	T4SS_P
XACb0046	<i>virB3</i>	1158533	VirB3 protein	TCTTGGGATGAGCTTTTATTCC	CCCAAAGTGATCGCGTGATT	59	T4SS_P
XACb0047	<i>virB2</i>	1158534	VirB2 protein	CGATCTGGCGGCTAACGT	ATGCGAGCTGCGGAAGAA	56	T4SS_P
XACb0048	—	1158535	Hypothetical protein	GGCGGATGACGTTGAAGCT	CGCGGTAACGACCTCATAAAC	56	H_T4SS_P
XACb0049	—	1158536	Hypothetical protein	ACCACGGATCCTGGCAAAT	AGCAGCCCCCGTGAACA	51	H_T4SS_P

This table provides primers sequences, ORF's identification ID, gene's name, and respective *Xanthomonas citri* subsp. *citri* secretory systems. T4SS: type IV secretion system; T3SS: type III secretion system; C: chromosomal copy; P: plasmid copy; H: hypothetical genes associated with the T4SS.




FIGURE 1: T3SS genes in *Xanthomonas citri* subsp. *citri* and their expression levels. Schematic representation of T3SS genes in *Xanthomonas citri* subsp. *citri* and their expression levels. (a) shows functional maps of T3SS genes on the chromosome. The expression levels of genes in the T3SS (b) are indicated. Solid green bars indicate *in vivo* expression with respect to their expression *in vitro* (black bars). Error bars indicate standard deviation of the replicates.




FIGURE 2: T4SS genes in *Xanthomonas citri* subsp. *citri* and their expression levels. (a) and (b) show functional maps of T4SS genes on the chromosome (a) and plasmid pXAC64 (b), respectively. The expression levels of genes in the T4SS (c) and ORFs localized near the *virB* genes (d) are indicated. Solid green bars indicate *in vivo* expression with respect to their expression *in vitro* (black bars). Error bars indicate standard deviation of the replicates.

Our qPCR data showed that *hpaI* was the most up-regulated gene during *Citrus sinensis* infection by *Xcc*, followed by *hrpD6* (Figure 1(b)). Conversely, *hrpA*, which encodes an important component of the T3SS pilus [39], was down-regulated in our study. Previously, *hrpA* mutants of *Pseudomonas syringae* pv. *tomato* DC3000 showed reduced accumulation of structural components of the *hrp* pilus [39]. Haapalainen et al. [40] showed that soluble plant cell signals induce the expression of *hrp/hrc* genes and specifically upregulate *hrpA*. Interestingly, they found that HrpA does not accumulate to high levels intracellularly, suggesting that some kind of feedback regulation takes place between secretion and production of HrpA, perhaps through degradation of intracellular HrpA. Our work shows that this may also be applicable to *hrpA* gene in *Xcc*: its mRNA may be degraded faster, just as happen with HrpA in *Pseudomonas syringae*.

#### 4. Conclusions

Our finds confirm that the T3SS genes are induced in the successful infection of *Citrus sinensis* by *Xcc*, and therefore are excellent targets to help control or decrease the severity of citrus canker, whereas T4SS seems not have any relation with virulence. On the other hand, our results show that T4SS is not induced under the same infection conditions, which could indicate that it is not necessary for infection but only for cell-to-cell communication.

#### Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

#### Acknowledgments

This work was supported by grants from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Fundo de Defesa da Citricultura (FUNDECITRUS), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for the biological tests and fellowship Grants, and by Grants from Fundação de Amparo à Pesquisa do Estado de Minas Gerais (CBB-APQ-04425-10) for *Xcc* functional database creation. The funding agencies had no role in study design, data collection and analysis, decision to publish, or preparation of the paper.

#### References

- [1] J. Boch and U. Bonas, "Xanthomonas AvrBs3 family-type III effectors: discovery and function," *Annual Review of Phytopathology*, vol. 48, pp. 419–436, 2010.
- [2] J. H. Graham, T. R. Gottwald, J. Cubero, and D. S. Achor, "Xanthomonas axonopodis pv. citri: factors affecting successful eradication of citrus canker," *Molecular Plant Pathology*, vol. 5, no. 1, pp. 1–15, 2004.
- [3] W. Bitter, E. N. G. Houben, J. Luirink, and B. J. Appelmelk, "Type VII secretion in mycobacteria: classification in line with cell envelope structure," *Trends in Microbiology*, vol. 17, no. 8, pp. 337–338, 2009.
- [4] D. Büttner and U. Bonas, "Regulation and secretion of Xanthomonas virulence factors," *FEMS Microbiology Reviews*, vol. 34, no. 2, pp. 107–133, 2010.
- [5] E. J. M. Stoop, W. Bitter, and A. M. van der Sar, "Tubercle bacilli rely on a type VII army for pathogenicity," *Trends in Microbiology*, vol. 20, no. 10, pp. 477–484, 2012.
- [6] P. Troisfontaines and G. R. Cornelis, "Type III secretion: more systems than you think," *Physiology*, vol. 20, no. 5, pp. 326–339, 2005.
- [7] B. Coburn, I. Sekirov, and B. B. Finlay, "Type III secretion systems and disease," *Clinical Microbiology Reviews*, vol. 20, no. 4, pp. 535–549, 2007.
- [8] J. L. Hodgkinson, A. Horsley, D. Stabat et al., "Three-dimensional reconstruction of the Shigella T3SS transmembrane regions reveals 12-fold symmetry and novel features throughout," *Nature Structural & Molecular Biology*, vol. 16, no. 5, pp. 477–485, 2009.
- [9] T. F. Moraes, T. Spreter, and N. C. Strynadka, "Piecing together the Type III injectisome of bacterial pathogens," *Current Opinion in Structural Biology*, vol. 18, no. 2, pp. 258–266, 2008.
- [10] O. Schraidt, M. D. Lefebvre, M. J. Brunner et al., "Topology and organization of the Salmonella typhimurium type III secretion needle complex components," *PLoS Pathogens*, vol. 6, no. 4, article e1000824, 2010.
- [11] T. Spreter, C. K. Yip, S. Sanowar et al., "A conserved structural motif mediates formation of the periplasmic rings in the type III secretion system," *Nature Structural and Molecular Biology*, vol. 16, no. 5, pp. 468–476, 2009.
- [12] Y. Akeda and J. E. Galán, "Chaperone release and unfolding of substrates in type III secretion," *Nature*, vol. 437, no. 7060, pp. 911–915, 2005.
- [13] P.-J. Matteï, E. Faudry, V. Job, T. Izoré, I. Attree, and A. Dessen, "Membrane targeting and pore formation by the type III secretion system translocon," *FEBS Journal*, vol. 278, no. 3, pp. 414–426, 2011.
- [14] P. J. Christie, K. Atmakuri, V. Krishnamoorthy, S. Jakubowski, and E. Cascales, "Biogenesis, architecture, and function of bacterial type IV secretion systems," *Annual Review of Microbiology*, vol. 59, pp. 451–485, 2005.
- [15] K. Wallden, A. Rivera-Calzada, and G. Waksman, "Type IV secretion systems: versatility and diversity in function," *Cellular Microbiology*, vol. 12, no. 9, pp. 1203–1212, 2010.
- [16] K. S. De Felipe, S. Pampou, O. S. Jovanovic et al., "Evidence for acquisition of Legionella type IV secretion substrates via inter-domain horizontal gene transfer," *Journal of Bacteriology*, vol. 187, no. 22, pp. 7716–7726, 2005.
- [17] H. Nagai, J. C. Kagan, X. Zhu, R. A. Kahn, and C. R. Roy, "A bacterial guanine nucleotide exchange factor activates ARF on Legionella phagosomes," *Science*, vol. 295, no. 5555, pp. 679–682, 2002.
- [18] S. Ninio and C. R. Roy, "Effector proteins translocated by Legionella pneumophila: strength in numbers," *Trends in Microbiology*, vol. 15, no. 8, pp. 372–380, 2007.
- [19] X. Pan, A. Lührmann, A. Satoh, M. A. Laskowski-Arce, and C. R. Roy, "Ankyrin repeat proteins comprise a diverse family of bacterial type IV effectors," *Science*, vol. 320, no. 5883, pp. 1651–1654, 2008.
- [20] M. I. Marchesini, C. K. Herrmann, S. P. Salcedo, J.-P. Gorvel, and D. J. Comerchi, "In search of Brucella abortus type iv secretion



- substrates: screening and identification of four proteins translocated into host cells through virB system," *Cellular Microbiology*, vol. 13, no. 8, pp. 1261–1274, 2011.
- [21] M. C. Alegria, D. P. Souza, M. O. Andrade et al., "Identification of new protein-protein interactions involving the products of the chromosome- and plasmid-encoded type IV secretion loci of the phytopathogen *Xanthomonas axonopodis* pv. *citri*," *Journal of Bacteriology*, vol. 187, no. 7, pp. 2315–2325, 2005.
- [22] D. P. Souza, M. O. Andrade, C. E. Alvarez-Martinez, G. M. Arantes, C. S. Farah, and R. K. Salinas, "A component of the Xanthomonadaceae type IV secretion system combines a VirB7 motif with a N0 domain found in outer membrane transport proteins," *PLoS Pathogens*, vol. 7, no. 5, Article ID e1002031, 2011.
- [23] A. C. R. Da Silva, J. A. Ferro, F. C. Reinach et al., "Comparison of the genomes of two *Xanthomonas* pathogens with differing host specificities," *Nature*, vol. 417, no. 6887, pp. 459–463, 2002.
- [24] A. Facincani, L. Moreira, M. Soares et al., "Comparative proteomic analysis reveals that T3SS, Tfp, and xanthan gum are key factors in initial stages of Citrus sinensis infection by *Xanthomonas citri* subsp. *citri*," *Functional & Integrative Genomics*, pp. 1–13, 2013.
- [25] M. R. Soares, A. P. Facincani, R. M. Ferreira et al., "Proteome of the phytopathogen *Xanthomonas citri* subsp. *citri*: a global expression profile," *Proteome Science*, vol. 8, article 55, 2010.
- [26] T. R. Jacob, M. L. Laia, J. A. Ferro, and M. I. T. Ferro, "Selection and validation of reference genes for gene expression studies by reverse transcription quantitative PCR in *Xanthomonas citri* subsp. *citri* during infection of Citrus sinensis," *Biotechnology Letters*, vol. 33, no. 6, pp. 1177–1184, 2011.
- [27] K. J. Livak and T. D. Schmittgen, "Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method," *Methods*, vol. 25, no. 4, pp. 402–408, 2001.
- [28] M. L. Laia, L. M. Moreira, J. DezaJacomio et al., "New genes of *Xanthomonas citri* subsp. *citri* involved in pathogenesis and adaptation revealed by a transposon-based mutant library," *BMC Microbiology*, vol. 9, article 12, 2009.
- [29] H. Nagai and T. Kubori, "Type IVB secretion systems of legionella and other gram-negative bacteria," *Frontiers in Microbiology*, vol. 2, article 136, 2011.
- [30] L. Andrieux, G. Bourg, A. Pirone, D. O'Callaghan, and G. Patey, "A single amino acid change in the transmembrane domain of the VirB8 protein affects dimerization, interaction with VirB10 and Brucella suis virulence," *FEBS Letters*, vol. 585, no. 15, pp. 2431–2436, 2011.
- [31] D. Sivanesan and C. Baron, "The dimer interface of Agrobacterium tumefaciens VirB8 is important for type IV secretion system function, stability, and association of VirB2 with the core complex," *Journal of Bacteriology*, vol. 193, no. 9, pp. 2097–2106, 2011.
- [32] Y. Zhao, G. Liu, S. Li et al., "Role of a type IV-like secretion system of Streptococcus suis 2 in the development of streptococcal toxic shock syndrome," *Journal of Infectious Diseases*, vol. 204, no. 2, pp. 274–281, 2011.
- [33] Y.-Q. He, L. Zhang, B.-L. Jiang et al., "Comparative and functional genomics reveals genetic diversity and determinants of host specificity among reference strains and a large collection of Chinese isolates of the phytopathogen *Xanthomonas campestris* pv. *campestris*," *Genome Biology*, vol. 8, no. 10, article R218, 2007.
- [34] Y. Wang, M. Xiao, X. Geng, J. Liu, and J. Chen, "Horizontal transfer of genetic determinants for degradation of phenol between the bacteria living in plant and its rhizosphere," *Applied Microbiology and Biotechnology*, vol. 77, no. 3, pp. 733–739, 2007.
- [35] G. R. Cornelis and F. Van Gijsegem, "Assembly and function of type III secretory systems," *Annual Review of Microbiology*, vol. 54, pp. 735–774, 2000.
- [36] O. Rossier, G. Van Den Ackerveken, and U. Bonas, "HrpB2 and hrpF from Xanthomonas are type III-secreted proteins and essential for pathogenicity and recognition by the host plant," *Molecular Microbiology*, vol. 38, no. 4, pp. 828–838, 2000.
- [37] K. Wengelnik and U. Bonas, "HrpXv, an AraC-type regulator, activates expression of five of the six loci in the hrp cluster of *Xanthomonas campestris* pv. *vesicatoria*," *Journal of Bacteriology*, vol. 178, no. 12, pp. 3462–3469, 1996.
- [38] X. Guo, H. Zou, Y. Li, L. Zou, and G. Chen, "HrpD6 gene determines Xanthomonas oryzae pv. oryzae to trigger hypersensitive response in tobacco and pathogenicity in rice," *Wei Sheng Wu Xue Bao*, vol. 50, no. 9, pp. 1155–1163, 2010.
- [39] E. Roine, J. Saarinen, N. Kalkkinen, and M. Romantschuk, "Purified HrpA of Pseudomonas syringae pv. tomato DC3000 reassembles into pili," *FEBS Letters*, vol. 417, no. 2, pp. 168–172, 1997.
- [40] M. Haapalainen, K. Van Gestel, M. Pirhonen, and S. Taira, "Soluble plant cell signals induce the expression of the type III secretion system of Pseudomonas syringae and upregulate the production of pilus protein HrpA," *Molecular Plant-Microbe Interactions*, vol. 22, no. 3, pp. 282–290, 2009.