

RESEARCH

Open Access



# The potential role of herbal extract Wedelolactone for treating particle-induced osteolysis: an in vivo study

Yung-Chang Lu<sup>1,2,3</sup>, Ting-Kuo Chang<sup>1,2,3</sup>, Tzu-Chiao Lin<sup>2</sup>, Shu-Ting Yeh<sup>2</sup>, Hsu-Wei Fang<sup>4</sup>,  
Chun-Hsiung Huang<sup>1,2,3,5</sup> and Chang-Hung Huang<sup>2,3,6,7\*</sup>

## Abstract

**Background:** Osteolysis is one of the most prevalent clinical complications affecting people who undergo total joint replacement (TJR). Wedelolactone (WDL) is a coumestan compound derived from the *Wedelia chinensis* plant and has been demonstrated to exhibit anti-inflammatory properties. This study aimed to investigate the oral administration of WDL as a potential treatment for particle-induced osteolysis using a well-established mice calvarial disease model.

**Methods:** Thirty-two C57BL/6 J mice were randomized into four groups: Sham, vehicle, osteolysis group with oral WDL treatment for 4 weeks (WDL 4w), and osteolysis group treated for 8 weeks (WDL 8w). Micro-CT was used to quantitatively analyze the bone mineral density (BMD), bone volume/tissue volume (BV/TV) and trabecular bone thickness (Tb.Th). Osteoclast numbers were also measured from histological slides by two investigators who were blind to the treatment used.

**Results:** The results from micro-CT observation showed that BMD in the WDL 8w group improved significantly over the vehicle group ( $p < 0.05$ ), but there was no significant difference between WDL 4w and 8w for BV/TV and Tb.Th. Osteoclast numbers in the WDL 4w group were also lower than the vehicle group ( $p < 0.05$ ), but the difference between WDL 8w and 4w groups was not significant.

**Conclusions:** Particle-induced osteolysis is an inevitable long-term complication after TJR. The results of this animal study indicate that an oral administration of WDL can help reduce the severity of osteolysis without adverse effects.

**Keywords:** Particle-induced osteolysis, Wedelolactone, Herbal medicine, Murine calvarial model

## Background

Artificial joint replacement (AJR) is considered an effective method of treating severe joint degeneration [1]. However, periprosthetic osteolysis resulting from the deposition of wear particles from the articulating joint is one of the major clinical complications following AJR [2]. Wear particles can stimulate inflammatory responses

and osteoclastic resorption processes at the bone implant interface, which consequently often leads to implant loosening [3]. Although bearing materials have been introduced to reduce the generation of wear particles, osteolysis is still prevalent and is considered a major long-term complication of AJR.

Besides changing the base material of the implant, pharmaceuticals have also been investigated as a method for reducing osteolysis [4]. Bisphosphonates are well-known drugs for treating osteoporosis, but have also been shown to be effective at suppressing osteolysis [5, 6]. Similarly, statins, which are lipid-lowering agents,

\*Correspondence: [changhung0812@gmail.com](mailto:changhung0812@gmail.com)

<sup>2</sup> Department of Medical Research, MacKay Memorial Hospital, New Taipei City, Taiwan

Full list of author information is available at the end of the article



have been reported to reduce particle-induced osteolysis in a murine calvarial model [5]. However, such drugs can also introduce considerable serious side effects, such as atypical femoral bone fracture and osteonecrosis of the jaw [7–9].

Animal models are often used for investigating mechanisms that can lead to particle-induced osteolysis and for evaluating suitable treatment methods [5, 10–14]. A previous study by our institute investigated whether strontium ranelate (SR) [11], a drug for osteoporosis, could be effectively used to combat osteolysis. After gavage-feeding mice for up to 4 weeks, the results showed a significant increase in bone mineral density (BMD), bone volume/tissue volume (BV/TV) and trabecular thickness (Tb.Th), and a significant reduction in osteoclast numbers [11]. However, long-term use of SR may increase the risk of cardiovascular disease [15–17].

As an alternative to pharmaceuticals, Chinese herbal medicines are considered a more natural solution with fewer side effects [18]. The International Organization for Standardization has also assembled a technical committee (ISO/TC 249) to standardize medical fields derived from Chinese medicine. With the increasing popularity, more resources and attention have been given to the potential benefits for treating disease [19–21].

Wedelolactone (WDL) is a coumestan compound extracted from the *Wedelia Chinensis* plant [22]. WDL is considered a traditional Chinese herbal medicine with strong anti-inflammatory properties [23–27]. Recent studies have shown that WDL can promote hair growth [28] and is hepatoprotective [29, 30], neuroprotective [31] and anti-carcinogenic [32, 33]. However, few studies have reported on the potential of WDL for treating diseases of the skeletal system. Based on the ability of WDL to reduce inflammation [23–27] and inhibit osteoclastogenesis [34, 35], it is hypothesized that WDL could also play a role in reducing the risk of particle-induced osteolysis. This study used a well-established murine calvarial osteolysis model to investigate whether WDL administered orally could reduce the severity of osteolysis.

## Methods

### Establishing the calvarial particle-induced osteolysis model


The protocol for this experiment was approved by the Institutional Animal Care and Use Committee at the institute where the study was performed. Thirty-two 6-week-old C57BL/6 J female mice were supplied by BioLASCO (Taipei, Taiwan), an AAALAC certified biotechnology company. The animals were kept in a room at 24°C, 50% humidity, and with a 12 h light/dark cycle (light from AM 7:00 to PM 7:00). The animals were

randomly separated into four groups: (1) sham group ( $n=8$ ) (underwent surgery only) (2) vehicle group ( $n=8$ ) (implanted with PS particles) [36–38], (3) WDL 4w ( $n=8$ , implanted with PS particles and treated with WDL for 4 weeks. 7 animals remained at the end of the experiment) and (4) WDL 8w ( $n=8$ , treatment for 8 weeks). The vehicle group and WDL-treated groups were injected with 1 mg PS particles/100  $\mu$ l HA [10, 25]. The polystyrene particles (Polystyrene Latex Spheres, 610–38) were purchased from TED PELLA, Inc. (CA, USA). Three hundred particles were randomly selected and SEM was used to measure the particle size and aspect ratio (Fig. 1a). The particles were found to be  $1.03 \pm 0.04 \mu$ m and  $0.99 \pm 0.03$ , respectively (Fig. 1b, c). The particles were also confirmed to have an endotoxin level below 0.25 EU/mL using a Limulus Amoebocyte Lysate assay kit (ToxinSensor™ gel clot endotoxin assay kit, GenScript, NJ, USA) and then suspended in hyaluronic acid.

To inject the particles, the mice were anesthetized with 100 mg/kg of Zoletil 50 and 10 mg/kg Rompun by intraperitoneal injection. A  $0.5 \times 0.5$  cm area of the middle calvaria was exposed by sagittal incision. After removing the periosteum intact, the particle suspension was spread over the area and the incision was closed with sutures. After 2 week post-surgery, the WDL 4w and 8w groups were gavage-fed with Wedelolactone (Y0001599, European Pharmacopoeia Reference Standard, Sigma-Aldrich, USA) at a dose of 4 mg/kg/day for 5 days/week [21]. WDL was dissolved by DMSO as stock solution and diluted with phosphate-buffered saline (PBS) into 10 volumes to attain a dosage of 4 mg/kg. The vehicle group was gavage-fed the vehicle solution (10% DMSO in PBS). The sham group, vehicle group and WDL 4w group were then sacrificed after 4 weeks of feeding either the vehicle or WDL, and the WDL 8w group was sacrificed after 8 weeks of feeding WDL.

### Micro-CT imaging analysis

The calvarias were fixed in 10% buffered formalin for 24 h, and then transferred to 70% ethanol for 24 h. The specimens were scanned with the micro-CT system Skyscan 1076 (Bruker micro-CT, Kontich, Belgium) at a resolution of  $2048 \times 2048$ . Three-dimensional images were reconstructed in Skyscan with a voxel size of 9  $\mu$ m [10, 25]. A spherical volume of interest (VOI) with a diameter of 5 mm was then defined with the bregma as the center. Within this VOI, the bone mineral density (BMD, mg/cc), the ratio of bone volume to tissue volume (BV/TV, %) and trabecular thickness (Tb.Th) were recorded for each group.



### Histological analysis

The calvarias were decalcified in 10% ethylenediaminetetraacetic acid (EDTA) for 2 weeks, and then embedded in paraffin. Each section with a thickness of 5  $\mu\text{m}$  were taken in the sagittal plane centered over the particle-treated area. The sections were then stained with hematoxylin and eosin (H&E stain) to observe the morphology of cellular inflammatory responses from the connective tissue. A tartrate-resistant acid phosphatase (TRAP) stain was performed using a commercial TRAP kit (#386A, Sigma-Aldrich). The number of osteoclasts was determined by counting the number of TRAP-positive multinucleated cells by two coauthors to eliminate intra- and inter-observer error.

### Statistical analysis


The data was analyzed using one-way analysis of variance (ANOVA) to show the difference between groups. Multiple comparisons were adjusted with a Bonferroni post hoc test. Results were reported as mean  $\pm$  standard

deviation (SD). Any  $p$  value less than 0.05 was considered significantly different.

## Results

### Micro-CT imaging analysis

A visual analysis of the three dimensional (3D) reconstructed micro-CT images showed clear differences between the sham group, vehicle group, and WDL groups (Fig. 2a). The images showed typical osteolysis with pores in the sham group, but both the size and number of pores decreased in WDL-treated groups. The presence of PS particles significantly decreased the BMD in the vehicle group by 7.8% when compared to the sham group ( $0.74 \pm 0.03$  for vehicle group and  $0.801 \pm 0.03$  for sham group,  $p < 0.01$ ). Both WDL groups showed an increase in BMD (Fig. 2b), with the 8w group showing a significant increase of 5.1% in comparison to the vehicle group ( $0.78 \pm 0.01$  for 8w group and  $0.74 \pm 0.03$  for vehicle group,  $p < 0.05$ ). The BV/TV in the vehicle group decreased by 4.1% in comparison to the sham group



**Fig. 2** Reconstructed image of the VOI with the bregma at the center. The VOI is defined with a diameter of 5 mm (a). Micro-CT image of bone formation in a particle-induced osteolysis model measured at 4 and 8 weeks after feeding WDL (b) (\* $p < 0.05$ ; \*\* $p < 0.01$ , as determined using ANOVA testing)

( $21.23 \pm 1.43$  in vehicle group versus  $22.1 \pm 1.54$  in sham group) but increased in the WDL 4w group ( $22.96 \pm 2.07$ ) and WDL 8w group ( $22.44 \pm 2.78$ ). However, there were no significant differences in BV/TV and Tb.Th between the WDL 4w and WDL 8w groups (Fig. 2b).


### Histomorphometric analysis

Histological analysis with H&E staining was used to evaluate the inflammatory response. Pseudomembrane proliferation occurred in the vehicle and WDL groups. The morphology of the cells in the periosteum was observed to change to a circle-shaped contour, while the cells resembled a flat contour in the sham group. Multinucleated giant cells were found in the surrounding periosteum (Fig. 3). TRAP staining was used to highlight

polymer particles in the periosteal cells and multinucleated giant cells (Fig. 4a, b).

### Osteoclasts around the bone perimeter

TRAP stain was used to highlight osteoclasts around the calvaria to calculate the osteoclast numbers in each group. The results showed the osteoclast numbers in the vehicle group increased significantly in comparison to the sham group ( $43.7 \pm 10.1$  in vehicle group versus  $18.2 \pm 14.5$  in sham group,  $p < 0.05$ ), demonstrating that the polymer particles likely induced osteolysis. Furthermore, there was a significant reduction in osteoclast numbers in the WDL 4w group in comparison to the vehicle group ( $21.1 \pm 9.8$  in WDL 4w group versus  $43.7 \pm 10.1$  in vehicle group,  $p < 0.05$ ). However,



**Fig. 3** Hematoxylin and eosin (H&E) staining of periosteum in mice calvarial section. Multinucleated giant cells were observed in the groups injected with PS particles (Magnification:  $\times 40$ ; scale bar: 100  $\mu\text{m}$ )

osteoclast number in WDL 8w group ( $30.6 \pm 4.0$ ) was no significantly different to the WDL 4w group (Fig. 5b).


## Discussion

Osteolysis is one of the major long-term complications affecting patients who undergo AJR. Chinese herbal medicine is generally considered to be milder than pharmaceutical treatments and does not produce strong adverse effects. This study aimed to investigate the potential use of the Chinese herbal medicine wedelolactone for reducing the incidence of particle-induced osteolysis.

Bisphosphonates (BPs) are commonly used to treat conditions of metabolic bone loss, such as osteoporosis [7], but have also been shown to inhibit particle-induced osteolysis [39]. However, the long-term use of BPs and other pharmaceuticals often results in serious adverse effects, such as an increased risk of osteonecrosis of the jaw, atypical femur fractures, atrial fibrillation, and esophageal cancer [6]. Statins, a drug usually used to lower blood cholesterol levels and reduce the risk of symptoms related to atherosclerosis, targets the

mevalonate pathway of osteoclasts, which affect the same inhibition mechanism as bisphosphonates. Statins have been shown to markedly reduce the severity of particle-induced osteolysis in a murine calvarial model [5]. However, as with BPs, the use of statins can present a number of side effects when used long-term, such as rhabdomyolysis, cognitive loss, neuropathy, hepatic dysfunction, and sexual dysfunction [8].


There is increasing interest in alternative methods such as traditional Chinese medicine for treating diseases as clinicians look to reduce long-term complications associated with conventional medicine. It has reported that postmenopausal Chinese women with greater fruit intake have a significantly higher BMD than comparable women with a lower fruit intake [40]. Flavonoids, found in a wide diversity of food derived from fruit, have been recognized as potential dietary components to promote bone health [41, 42]. Nam et al. also indicated that traditional mixed extracts of medicinal herbs can effectively inhibit the expression of inflammatory mediators in gouty arthritis on monosodium urate (MSU) crystals-induced



**Fig. 4** TRAP staining indicated that PS particles exist in the periosteal cells and multinucleated giant cells in mice calvarial tissue (a) (Magnification  $\times 40$  Scale bar: 100  $\mu\text{m}$ ). Different shapes of PS particles and melanin granules observed in TRAP staining. Arrow, PS particle; Arrow head, melanin granule (b) (Magnification  $\times 100$  Scale bar: 20  $\mu\text{m}$ )

gouty inflammation, demonstrating its potential for treating gouty arthritis [43]. Phytoestrogens, which are natural compounds that act to maintain healthy bones,

have been shown to protect against postmenopausal bone loss [20]. This protective mechanism has been demonstrated with flavones [44, 45], flavanones [46,



**Fig. 5** Typical samples from micro-CT with purple staining showing TRAP-positive osteoclasts (a) (Magnification: x 40; Scale bar: 100 μm). Average number of TRAP-positive cells from each group are presented as the mean ± SD (b) (\* $p < 0.05$ )

47], flavonols [48], coumestans [49], and triterpenoids [46, 50]. Some phytoestrogens also have the ability to reduce osteolysis by blocking some modules in the RANKL signaling pathway, and subsequently reducing the release of cytokines [44, 45, 47, 51].

To our best knowledge, no studies to date have investigated whether WDL can reduce the risk of particle-induced osteolysis using an *in-vivo* murine calvarial model. The concentration of WDL used in this study was adopted from Tsai et al. [21] who showed

that a low oral dose of WDL (4 mg/kg) administered for 4 weeks significantly suppressed the growth of prostate cancer cells. The results of our study showed that the BMD was significantly greater in the WDL 8w group ( $0.784 \pm 0.014$  mg/cc) than in the vehicle group ( $0.744 \pm 0.032$  mg/cc) (Fig. 2b). However, there was no significant difference between the WDL 4w and 8w groups in terms of the mean values of BV/TV and Tb.Th. On the other hand, the osteoclast numbers were significantly lower in the WDL 4w group ( $21.1 \pm 9.8$ ) and WDL 8w group ( $30.6 \pm 4.0$ ) than the vehicle group ( $44.1 \pm 9.8$ ) (Fig. 5).

Multinucleated giant cells are typically generated after implantation of medical devices, artificial joint or biomaterials, taking the form of foreign body giant cells. The formation of these giant cells is the end-stage of the inflammation or wound healing response [52]. Osteoclasts are specialized multinuclear giant cells derived from monocyte/macrophage lineage cells [53]. In this study, multinucleated giant cells were observed in the groups injected with polymer particles (Fig. 3). In addition, TRAP staining indicated that polymer particles were present in the periosteal cells and multinucleated giant cells in the mouse calvarial tissue (Fig. 4a, b).

Wear particles have been found in surrounding tissue after implantation of many different materials used in TJR, including ultra-high molecular weight polyethylene (UHMWPE) [54], poly(methyl methacrylate) (PMMA) [54], ceramics [54], metallic CoCrMo [55] and titanium alloy [55, 56]. Some studies used titanium particles to create particle-induced osteolysis animal models [44, 46, 53, 57, 58]. However, when used in joint replacements, titanium or its alloys generate fewer wear particles than polymers because of the relatively low mechanical strength of polymeric materials. In the authors' previous animal studies, UHMWPE wear debris-induced animal models were developed to analyze the *in vivo* biological response to highly cross-linked and vitamin E-stabilized polyethylene [10] and to evaluate the potential role of strontium ranelate-contained medicine to treat osteolysis [11]. However, considerable time was spent preparing enough wear debris for these animal models, while polystyrene (PS) particles are readily available and are widely used both commercially and for biomedical research [59, 60]. PS particles have been used in many animal studies to create particle-induced osteolysis animal models [36–38]. Furthermore, given that the size distribution and shape of PS particles are easier to control, this study used PS particles for the animal model.

Previous studies treated murine calvarial osteolysis with bioactive compounds for 10–14 days after implantation of foreign particles [12, 44, 45, 47, 50, 61]. For instance, icariin, a bioactive flavonoid, has been

proven to inhibit postmenopausal osteoporosis. Shao et al. gavage-fed mice with icariin at doses of 0.1 mg/g and 0.3 mg/g for 14 days to examine the effects on osteolysis in a particle-induced murine calvarial model. The results showed an increase in BMD and BV/TV over the control model, and the number of TRAP positive cells decreased [12]. Similarly, ursolic acid is an abundant triterpenoid present in over one hundred species of plants. It has been reported that ursolic acid isolated from loquat leaves can reduce bone loss in OVX mice [46]. Jiang et al. treated mice with 10 mg/kg and 40 mg/kg doses of ursolic acid administered through intraperitoneal injections for 14 days and found that ursolic acid protects against wear particle-induced osteolysis by suppressing osteoclast formation and function [50]. The treatment period in this current study, 4 weeks and 8 weeks, was longer than the referenced studies which only treated for a short-term of 2 weeks. No adverse effects were observed in this study after treating the mice for 8 weeks with WDL.

Bone remodeling is a dynamic equilibrium with molecular mechanisms, such as RANK/RANKL/OPG [62], NF- $\kappa$ B [63], and Wnt/BMP (bone morphogenic protein) [57, 64] signaling pathways playing critical roles in osteolysis. Although the trigger mechanisms for osteolysis are not yet fully understood, it is known that one of the mechanisms is the receptor activation of NF- $\kappa$ B ligand (RANKL) and osteoprotegerin (OPG) secreted from osteoblasts and osteogenic stromal cells, both of which act to maintain a balance between bone generation and resorption [62]. RANKL is required for the differentiation of osteoclast precursors into mature osteoclasts [58]. As the ratio of RANKL/OPG increases, the osteoclast precursors are easier influenced by RANKL signaling through the downstream activation of NF- $\kappa$ B/c-fos/NFATc1, subsequently causing the precursors to differentiate into mature osteoclasts. On the other hand, macrophages also plays a key role in wear particle-induced osteolysis [63, 65–68]. Cytokines (TNF- $\alpha$  and IL-1 $\beta$ , etc.) and other mediators of pro-inflammation from activated macrophages can regulate or stimulate other tissue-resident macrophages to promote osteoclastogenesis [66]. These cytokines also regulate JNK and the p38/ERK signaling pathway to induce NFATc1, one of the downstream factors in the RANKL signaling pathway, which can lead to osteolysis.

Studies have shown that some compounds from Chinese herbal medicines can treat particle-induced osteolysis by inhibiting the modules in the NF- $\kappa$ B signaling pathway, the main mechanism in the regulation of osteolysis, to effect the balance of osteoclasts and osteoblasts [12, 45, 50, 61, 68]. WDL is known for its ability to block the phosphorylation of I $\kappa$ B $\alpha$ , which acts to regulate the transcription of NF- $\kappa$ B mediated



genes, inhibiting LPS-induced pro-inflammation [69]. Annie et al. demonstrated how an extract from *Wedelia chinensis* attenuated OVX-induced bone loss in mice [70]. WDL extracted from *Ecliptae herba* has been shown to inhibit osteoclastogenesis of RAW 264.7 cells treated with RANKL [35], and prevent OVX-induced bone loss by inhibiting osteoclast activity and enhancing osteoblast activity [34]. Furthermore, it has been confirmed that WDL can regulate the RANKL-related NF- $\kappa$ B/c-fos/NFATc1 pathway to suppress osteoclastogenesis [26, 71], and also regulate the Wnt/ $\beta$ -catenin signaling pathway to induce osteoblastogenesis [71, 72]. The authors concluded that oral WDL could improve bone formation and inhibit resorption by affecting the balance of osteoclasts and osteoblasts. However, the mechanism leading to the inhibition of osteolysis by WDL still needs to be determined.

Some limitations of this study should be mentioned. First, murine calvarial models allow for a low-cost study with relatively quick results, but the models use a flat bone instead of a long bone and the particles are injected on the cortical bone surface rather than into cancellous bone. Second, as detailed above, the WDL dose used in this study was adopted from other related publications. However, the most effective dose for treating osteolysis in vivo has yet to be determined and requires further study. Third, osteoclast numbers were counted through qualitative analysis, not quantitative analysis. When injected onto the calvaria, the particles randomly precipitated and then a section was chosen for histological staining. This sampling approach may not represent true osteoclast numbers. Further studies are recommended to investigate inflammatory makers. Accepting the above limitations, this animal study identified the potential role of wedelolactone for treating particle-induced osteolysis.

## Conclusions

This study indicated that wedelolactone (WDL), a Chinese herbal medicine, could help to maintain bone quality. Oral WDL was shown to suppress osteoclast numbers and maintain the level of BMD over time in a particle-induced osteolysis murine calvarial model. Moving forward, WDL could be potentially developed as a functional food for lowering the risk of particle-induced osteolysis after total joint replacement.

## Abbreviations

TJR: Total Joint Replacement; WDL: Wedelolactone; BMD: Bone Mineral Density; BV/TV: Bone Volume/Tissue Volume; Tb.Th: Trabecular Thickness; ISO: International Organization for Standardization; RANKL: Receptor activator of nuclear factor kappa-B ligand; OPG: Osteoprotegerin; OVX: Ovariectomy.

## Acknowledgements

The authors are pleased to acknowledge the financial support of Mackay Memorial Hospital (MMH). We acknowledge experimental support from the Electron Microscope Laboratory of MMH. We also thank the Taiwan Animal Consortium (MOST 107-2319-B-001-002) Taiwan Mouse Clinic and Tron Medicalbio Corp for technical support in the micro-CT experiments. The authors particularly thank Colin J. McClean for his kind assistance in language editing and proofreading of this manuscript.

## Authors' contributions

Y-CL: conceptualization and methodology. T-KC: methodology and data description. T-CL: animal study, data analysis, and writing. S-TY: investigation and data analysis. C-HH: supervision, review and editing. C-HH: supervision, conceptualization, writing, review and editing. All authors read and approved the final manuscript.

## Funding

The financial support of Mackay Memorial Hospital (MMH-106-145) and the Taiwan Animal Consortium Taiwan Mouse Clinic (MOST 107-2319-B-001-002) for the micro-CT experiments.

## Availability of data and materials

The data sets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

## Declarations

### Ethics approval and consent to participate

(1) This material has not been published in whole or in part elsewhere; (2) the manuscript is not currently being considered for publication in another journal; (3) all authors have been personally and actively involved in substantive work leading to the manuscript, and will hold themselves jointly and individually responsible for its content. (4) The animal protocol use in this study was approved by the Institutional Animal Care and Use Committee, where the study was performed.

### Consent for publication

Not applicable.

### Competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### Author details

<sup>1</sup>Department of Medicine, MacKay Medical College, New Taipei City, Taiwan. <sup>2</sup>Department of Medical Research, MacKay Memorial Hospital, New Taipei City, Taiwan. <sup>3</sup>Department of Orthopaedic Surgery, MacKay Memorial Hospital, Taipei, Taiwan. <sup>4</sup>Department of Chemical Engineering and Biotechnology, National Taipei University of Technology, Taipei, Taiwan. <sup>5</sup>Department of Orthopaedic Surgery, Changhua Christian Hospital, Changhua, Taiwan. <sup>6</sup>School of Dentistry, National Yang Ming Chiao Tung University, Taipei, Taiwan. <sup>7</sup>Institute of Geriatric Welfare Technology and Science, MacKay Medical College, New Taipei City, Taiwan.

Received: 6 May 2022 Accepted: 22 June 2022

Published online: 28 June 2022

## References

1. Sonntag R, Reinders J, Kretzer JP. What's next? Alternative materials for articulation in total joint replacement. *Acta Biomater.* 2012;8(7):2434–41. <https://doi.org/10.1016/j.actbio.2012.03.029>.
2. Colston J, Atkins B. Bone and joint infection. *Clin Med (Lond).* 2018;18(2):150–4. <https://doi.org/10.7861/clinmedicine.18-2-150>.
3. Ingham E, Fisher J. The role of macrophages in osteolysis of total joint replacement. *Biomaterials.* 2005;26(11):1271–86.

4. Ren K, Dusad A, Zhang Y, Wang D. Therapeutic intervention for wear debris-induced aseptic implant loosening. *Acta Pharm Sin B*. 2013;3(2):76–85. <https://doi.org/10.1016/j.apsb.2013.02.005>.
5. von Knoch F, Heckelei A, Wedemeyer C, Saxler G, Hilken G, Henschke F, et al. The effect of simvastatin on polyethylene particle-induced osteolysis. *Biomaterials*. 2005;26(17):3549–55. <https://doi.org/10.1016/j.biomaterials.2004.09.043>.
6. McClung M, Harris ST, Miller PD, Bauer DC, Davison KS, Dian L, et al. Bisphosphonate therapy for osteoporosis: benefits, risks, and drug holiday. *Am J Med*. 2013;126(1):13–20. <https://doi.org/10.1016/j.amjmed.2012.06.023>.
7. Yamazaki T, Yamori M, Yamamoto K, Saito K, Asai K, Sumi E, et al. Risk of osteomyelitis of the jaw induced by oral bisphosphonates in patients taking medications for osteoporosis: a hospital-based cohort study in Japan. *Bone*. 2012;51(5):882–7. <https://doi.org/10.1016/j.bone.2012.08.115>.
8. Golomb BA, Evans MA. Statin adverse effects: a review of the literature and evidence for a mitochondrial mechanism. *Am J Cardiovasc Drugs*. 2008;8(6):373–418. <https://doi.org/10.2165/0129784-200808060-00004>.
9. Bellosta S, Corsini A. Statin drug interactions and related adverse reactions. *Expert Opin Drug Saf*. 2012;11(6):933–46. <https://doi.org/10.1080/14740338.2018.1394455>.
10. Huang CH, Lu YC, Chang TK, Hsiao IL, Su YC, Yeh ST, et al. In vivo biological response to highly cross-linked and vitamin e-doped polyethylene—a particle-Induced osteolysis animal study. *J Biomed Mater Res B Appl Biomater*. 2016;104(3):561–7. <https://doi.org/10.1002/jbm.b.33426>.
11. Lu YC, Chang TK, Yeh ST, Fang HW, Lin CY, Hsu LI, et al. The potential role of strontium ranelate in treating particle-induced osteolysis. *Acta Biomater*. 2015;20:147–54. <https://doi.org/10.1016/j.actbio.2015.03.034>.
12. Shao H, Shen J, Wang M, Cui J, Wang Y, Zhu S, et al. Icarin protects against titanium particle-induced osteolysis and inflammatory response in a mouse calvarial model. *Biomaterials*. 2015;60:92–9. <https://doi.org/10.1016/j.biomaterials.2015.04.048>.
13. Millett PJ, Allen MJ, Bostrom MP. Effects of alendronate on particle-induced osteolysis in a rat model. *J Bone Joint Surg Am*. 2002;84-a(2):236–49. <https://doi.org/10.2106/00004623-200202000-00011>.
14. Du Z, Zhu Z, Wang Y. The degree of peri-implant osteolysis induced by PEEK, CoCrMo, and HXLPE wear particles: a study based on a porous Ti6Al4V implant in a rabbit model. *J Orthop Surg Res*. 2018;13(1):23. <https://doi.org/10.1186/s13018-018-0736-y>.
15. Reginster JY, Neuprez A, Dardenne N, Beaudart C, Emonts P, Bruyere O. Efficacy and safety of currently marketed anti-osteoporosis medications. *Best Pract Res Clin Endoc Metab*. 2014;28(6):809–34. <https://doi.org/10.1016/j.beem.2014.09.003>.
16. Abrahamsen B, Grove EL, Vestergaard P. Nationwide registry-based analysis of cardiovascular risk factors and adverse outcomes in patients treated with strontium ranelate. *Osteoporos Int*. 2014;25(2):757–62. <https://doi.org/10.1007/s00198-013-2469-4>.
17. Cooper C, Fox KM, Borer JS. Ischaemic cardiac events and use of strontium ranelate in postmenopausal osteoporosis: a nested case-control study in the CPRD. *Osteoporos Int*. 2014;25(2):737–45. <https://doi.org/10.1007/s00198-013-2582-4>.
18. Lam TP. Strengths and weaknesses of traditional Chinese medicine and Western medicine in the eyes of some Hong Kong Chinese. *J Epidemiol Community Health*. 2001;55(10):762–5. <https://doi.org/10.1136/jech.55.10.762>.
19. Liu YQ, Wang YX, Shi NN, Han XJ, Lu AP. Current situation of International Organization for Standardization/Technical Committee 249 international standards of traditional Chinese medicine. *Chin J Integr Med*. 2017;23(5):376–80. <https://doi.org/10.1007/s11655-015-2439-0>.
20. Leung PC, Siu WS. Herbal treatment for osteoporosis: a current review. *J Tradit Complement Med*. 2013;3(2):82–7. <https://doi.org/10.4103/2225-4110.110407>.
21. Tsai CH, Lin FM, Yang YC, Lee MT, Cha TL, Wu GJ, et al. Herbal extract of *Wedelia chinensis* attenuates androgen receptor activity and orthotopic growth of prostate cancer in nude mice. *Clin Cancer Res*. 2009;15(17):5435–44. <https://doi.org/10.1158/1078-0432.CCR-09-0298>.
22. Sarveswaran S, Gautam SC, Ghosh J. Wedelolactone, a medicinal plant-derived coumestan, induces caspase-dependent apoptosis in prostate cancer cells via downregulation of PKC $\epsilon$  without inhibiting Akt. *Int J Oncol*. 2012;41(6):2191–9. <https://doi.org/10.3892/ijo.2012.1664>.
23. Huang YT, Wen CC, Chen YH, Huang WC, Huang LT, Lin WC, et al. Dietary uptake of *Wedelia chinensis* extract attenuates dextran sulfate sodium-induced colitis in mice. *PLoS ONE*. 2013;8(5):e64152-e. <https://doi.org/10.1371/journal.pone.0064152>.
24. Lin WC, Wen CC, Chen YH, Hsiao PW, Liao JW, Peng CI, et al. Integrative approach to analyze biodiversity and anti-inflammatory bioactivity of *Wedelia* medicinal plants. *PLoS ONE*. 2015;10(6):e0129067-e. <https://doi.org/10.1371/journal.pone.0129067>.
25. Ding S, Hou X, Yuan J, Tan X, Chen J, Yang N, et al. Wedelolactone protects human bronchial epithelial cell injury against cigarette smoke extract-induced oxidant stress and inflammation responses through Nrf2 pathway. *Int Immunopharmacol*. 2015;29(2):648–55. <https://doi.org/10.1016/j.intimp.2015.09.015>.
26. Yuan F, Chen J, Sun PP, Guan S, Xu J. Wedelolactone inhibits LPS-induced pro-inflammation via NF- $\kappa$ B pathway in RAW 264.7 cells. *J Biomed Sci*. 2013;20:84. <https://doi.org/10.1186/1423-0127-20-84>.
27. Kobori M, Yang Z, Gong D, Heissmeyer V, Zhu H, Jung YK, et al. Wedelolactone suppresses LPS-induced caspase-11 expression by directly inhibiting the IKK complex. *Cell Death Differ*. 2004;11(1):123–30. <https://doi.org/10.1038/sj.cdd.4401325>.
28. Roy RK, Thakur M, Dixit VK. Hair growth promoting activity of *Eclipta alba* in male albino rats. *Arch Dermatol Res*. 2008;300(7):357–64. <https://doi.org/10.1007/s00403-008-0860-3>.
29. Lu Y, Hu D, Ma S, Zhao X, Wang S, Wei G, et al. Protective effect of wedelolactone against CCl4-induced acute liver injury in mice. *Int Immunopharmacol*. 2016;34:44–52. <https://doi.org/10.1016/j.intimp.2016.02.003>.
30. Luo Q, Ding J, Zhu L, Chen F, Xu L. Hepatoprotective Effect of Wedelolactone against Concanavalin A-Induced Liver Injury in Mice. *Ame J Chin Med*. 2018;46(4):819–33. <https://doi.org/10.1142/S0192415X1850043X>.
31. Maya S, Prakash T, Goli D. Evaluation of neuroprotective effects of wedelolactone and gallic acid on aluminium-induced neurodegeneration: relevance to sporadic amyotrophic lateral sclerosis. *Eur J Pharmacol*. 2018;835:41–51. <https://doi.org/10.1016/j.ejphar.2018.07.058>.
32. Peng YG, Zhang L. Wedelolactone suppresses cell proliferation and migration through AKT and AMPK signaling in melanoma. *J Dermatol Treat*. 2019;30(4):389–95. <https://doi.org/10.1080/09546634.2018.1527996>.
33. Nehybova T, Smarda J, Daniel L, Stiborek M, Kanicky V, Spasojevic I, et al. Wedelolactone Acts as Proteasome Inhibitor in Breast Cancer Cells. *Int J Mol Sci*. 2017;18(4):729. <https://doi.org/10.3390/ijms18040729>.
34. Liu YQ, Hong ZL, Zhan LB, Chu HY, Zhang XZ, Li GH. Wedelolactone enhances osteoblastogenesis by regulating Wnt/ $\beta$ -catenin signaling pathway but suppresses osteoclastogenesis by NF- $\kappa$ B/c-fos/NFATc1 pathway. *Sci Rep*. 2016;6:32260. <https://doi.org/10.1038/srep32260>.
35. Liu YQ, Zhan LB, Liu T, Cheng MC, Liu XY, Xiao HB. Inhibitory effect of *Ecliptae herba* extract and its component wedelolactone on pre-osteoclastic proliferation and differentiation. *J Ethnopharmacol*. 2014;157:206–11. <https://doi.org/10.1016/j.jep.2014.09.033>.
36. Ma T, Ortiz SG, Huang Z, Ren P, Smith RL, Goodman SB. In vivo murine model of continuous intramedullary infusion of particles—a preliminary study. *J Biomed Mater Res Part B Appl Biomater*. 2009;88(1):250–3. <https://doi.org/10.1002/jbm.b.31175>.
37. Ortiz SG, Ma T, Regula D, Smith RL, Goodman SB. Continuous intramedullary polymer particle infusion using a murine femoral explant model. *J Biomed Mater Res B Appl Biomater*. 2008;87(2):440–6. <https://doi.org/10.1002/jbm.b.31122>.
38. Chang TK, Lu YC, Yeh ST, Lin TC, Huang CH, Huang CH. In vitro and in vivo biological responses to graphene and graphene oxide: a murine calvarial animal study. *Int J Nanomed*. 2020;15:647–59. <https://doi.org/10.2147/IJN.S231885>.
39. Purdue PE, Koulouvaris P, Potter HG, Nestor BJ, Sculco TP. The cellular and molecular biology of periprosthetic osteolysis. *Clin Orthop Relat Res*. 2007;454:251–61. <https://doi.org/10.1097/01.blo.0000238813.95035.1b>.
40. Chen YM, Ho SC, Woo JL. Greater fruit and vegetable intake is associated with increased bone mass among postmenopausal Chinese women. *Br J Nutr*. 2006;96(4):745–51.

41. Weaver CM, Alekel DL, Ward WE, Ronis MJ. Flavonoid intake and bone health. *J Nutr Gerontol Geriatr*. 2012;31(3):239–53. <https://doi.org/10.1080/21551197.2012.698220>.
42. Welch AA, Hardcastle AC. The effects of flavonoids on bone. *Curr Osteoporos Rep*. 2014;12(2):205–10. <https://doi.org/10.1007/s11914-014-0212-5>. [https://doi.org/10.1016/S1875-5364\(17\)30084-5](https://doi.org/10.1016/S1875-5364(17)30084-5).
43. Nam JS, Jagga S, Sharma AR, Lee JH, Park JB, Jung JS, et al. Anti-inflammatory effects of traditional mixed extract of medicinal herbs (MEMH) on monosodium urate crystal-induced gouty arthritis. *Chin J Nat Med*. 2017;15(8):561–75. [https://doi.org/10.1016/S1875-5364\(17\)30084-5](https://doi.org/10.1016/S1875-5364(17)30084-5).
44. Shin DK, Kim MH, Lee SH, Kim TH, Kim SY. Inhibitory effects of luteolin on titanium particle-induced osteolysis in a mouse model. *Acta Biomater*. 2012;8(9):3524–31. <https://doi.org/10.1016/j.actbio.2012.05.002>.
45. Zhao S, Sun Y, Li X, Wang J, Yan L, Zhang Z, et al. Scutellarin inhibits RANKL-mediated osteoclastogenesis and titanium particle-induced osteolysis via suppression of NF- $\kappa$ B and MAPK signaling pathway. *Int Immunopharmacol*. 2016;40:458–65. <https://doi.org/10.1016/j.intimp.2016.09.031>.
46. Wang W, Wu C, Tian B, Liu X, Zhai Z, Qu X, et al. The inhibition of RANKL-induced osteoclastogenesis through the suppression of p38 signaling pathway by naringenin and attenuation of titanium-particle-induced osteolysis. *Int J Mol Sci*. 2014;15(12):21913–34. <https://doi.org/10.3390/ijms151221913>.
47. Yu X, Zhao X, Wu T, Zhou Z, Gao Y, Wang X, et al. Inhibiting wear particles-induced osteolysis with naringin. *Int Orthop*. 2013;37(1):137–43. <https://doi.org/10.1007/s00264-012-1668-5>.
48. Wang QS, Wang GF, Lu YR, Cui YL, Li H, Li RX, et al. The Combination of icariin and constrained dynamic loading stimulation attenuates bone loss in ovariectomy-induced osteoporotic mice. *J Orthop Res*. 2018;36(5):1415–24. <https://doi.org/10.1002/jor.23777>.
49. Zhai Y, Li Y, Wang Y, Cui J, Feng K, Kong X, et al. Psoralidin, a prenylated coumestan, as a novel anti-osteoporosis candidate to enhance bone formation of osteoblasts and decrease bone resorption of osteoclasts. *Eur J Pharmacol*. 2017;801:62–71. <https://doi.org/10.1016/j.ejphar.2017.03.001>.
50. Jiang C, Xiao F, Gu X, Zhai Z, Liu X, Wang W, et al. Inhibitory effects of ursolic acid on osteoclastogenesis and titanium particle-induced osteolysis are mediated primarily via suppression of NF- $\kappa$ B signaling. *Biochimie*. 2015;111:107–18. <https://doi.org/10.1016/j.biochi.2015.02.002>.
51. Yang C, Liu W, Zhang X, Zeng B, Qian Y. Naringin increases osteoprotegerin expression in fibroblasts from periprosthetic membrane by the Wnt/ $\beta$ -catenin signaling pathway. *J Orthop Surg Res*. 2020;15(1):600. <https://doi.org/10.1186/s13018-020-02145-z>.
52. Anderson JM, Rodriguez A, Chang DT. Foreign body reaction to biomaterials. *Semin Immunol*. 2008;20(2):86–100. <https://doi.org/10.1016/j.smim.2007.11.004>.
53. Katsuyama E, Miyamoto H, Kobayashi T, Sato Y, Hao W, Kanagawa H, et al. Interleukin-1 receptor-associated kinase-4 (IRAK4) promotes inflammatory osteolysis by activating osteoclasts and inhibiting formation of foreign body giant cells. *J Biol Chem*. 2015;290(2):716–26. <https://doi.org/10.1074/jbc.M114.568360>.
54. Gibon E, Córdova LA, Lu L, Lin TH, Yao Z, Hamadouche M, et al. The biological response to orthopedic implants for joint replacement. II: polyethylene, ceramics, PMMA, and the foreign body reaction. *J Biomed Mater Res B Appl Biomater*. 2017;105(6):1685–91. <https://doi.org/10.1002/jbm.b.33676>.
55. Gibon E, Amanatullah DF, Loi F, Pajarinen J, Nabeshima A, Yao Z, et al. The biological response to orthopaedic implants for joint replacement: part I: metals. *J Biomed Mater Res B Appl Biomater*. 2017;105(7):2162–73. <https://doi.org/10.1002/jbm.b.33734>.
56. Zhang L, Haddouti EM, Welle K, Burger C, Wirtz DC, Schildberg FA, et al. The effects of biomaterial implant wear debris on osteoblasts. *Front Cell Dev Biol*. 2020;8:352. <https://doi.org/10.3389/fcell.2020.00352>.
57. Nam JS, Sharma AR, Jagga S, Lee DH, Sharma G, Nguyen LT, et al. Suppression of osteogenic activity by regulation of WNT and BMP signaling during titanium particle induced osteolysis. *J Biomed Mater Res A*. 2017;105(3):912–26. <https://doi.org/10.1002/jbm.a.36004>.
58. Teitelbaum SL, Ross FP. Genetic regulation of osteoclast development and function. *Nat Rev Genet*. 2003;4(8):638–49. <https://doi.org/10.1038/nrg1122>.
59. Qiu J, Camargo PHC, Jeong U, Xia Y. Synthesis, transformation, and utilization of monodispersed colloidal spheres. *Acc Chem Res*. 2019;52(12):3475–87. <https://doi.org/10.1021/acs.accounts.9b00490>.
60. Li L, Sun S, Tan L, Wang Y, Wang L, Zhang Z, et al. Polystyrene nanoparticles reduced ROS and inhibited ferroptosis by triggering lysosome stress and TFEB nucleus translocation in a size-dependent manner. *Nano Lett*. 2019;19(11):7781–92. <https://doi.org/10.1021/acs.nanolett.9b02795>.
61. Jiao Z, Xu W, Zheng J, Shen P, Qin A, Zhang S, et al. Kaempferide prevents titanium particle induced osteolysis by suppressing JNK activation during osteoclast formation. *Sci Rep*. 2017;7(1):16665. <https://doi.org/10.1038/s41598-017-16853-w>.
62. Boyce BF, Xing L. Biology of RANK, RANKL, and osteoprotegerin. *Arthritis Res Ther*. 2007;9(Suppl 1):S1. <https://doi.org/10.1186/ar2165>.
63. Goodman SB, Gibon E, Pajarinen J, Lin TH, Keeney M, Ren PG, et al. Novel biological strategies for treatment of wear particle-induced periprosthetic osteolysis of orthopaedic implants for joint replacement. *J R Soc Interface*. 2014;11(93):20130962. <https://doi.org/10.1098/rsif.2013.0962>.
64. Jagga S, Sharma AR, Lee YH, Nam JS, Lee SS. Sclerostin-mediated impaired osteogenesis by fibroblast-like synoviocytes in the particle-induced osteolysis model. *Front Mol Biosci*. 2021;8:666295. <https://doi.org/10.3389/fmolb.2021.666295>.
65. Lei P, Dai Z, Zhang YS, Liu H, Niu W, Li K, et al. Macrophage inhibits the osteogenesis of fibroblasts in ultrahigh molecular weight polyethylene (UHMWPE) wear particle-induced osteolysis. *J Orthop Surg Res*. 2019;14(1):80. <https://doi.org/10.1186/s13018-019-1119-8>.
66. Qiu J, Peng P, Xin M, Wen Z, Chen Z, Lin S, et al. ZBTB20-mediated titanium particle-induced peri-implant osteolysis by promoting macrophage inflammatory responses. *Biomater Sci*. 2020;8(11):3147–63. <https://doi.org/10.1039/d0bm00147c>.
67. Lin S, Wen Z, Li S, Chen Z, Li C, Ouyang Z, et al. LncRNA Neat1 promotes the macrophage inflammatory response and acts as a therapeutic target in titanium particle-induced osteolysis. *Acta Biomater*. 2022;142:345–60. <https://doi.org/10.1016/j.actbio.2022.02.007>.
68. Guangtao F, Zhenkang W, Zhantao D, Mengyuan L, Qingtian L, Yuanchen M, et al. Icarin alleviates wear particle-induced periprosthetic osteolysis via down-regulation of the estrogen receptor  $\alpha$ -mediated NF- $\kappa$ B signaling pathway in macrophages. *Front Pharmacol*. 2021;12:746391. <https://doi.org/10.3389/fphar.2021.746391>.
69. Shen P, Yang X, Jiang J, Wang X, Liang T, He L. Wedelolactone from *Eclipta alba* inhibits lipopolysaccharide-enhanced cell proliferation of human renal mesangial cells via NF- $\kappa$ B signaling pathway. *Am J Transl Res*. 2017;9(5):2132–42.
70. Annie S, Prabhu RG, Malini S. Activity of *Wedelia calendulacea* Less in post-menopausal osteoporosis. *Phytomedicine*. 2006;13(1–2):43–8. <https://doi.org/10.1016/j.phymed.2004.01.011>.
71. Deng X, Liang LN, Zhu D, Zheng LP, Yu JH, Meng XL, et al. Wedelolactone inhibits osteoclastogenesis but enhances osteoblastogenesis through altering different semaphorins production. *Int Immunopharmacol*. 2018;60:41–9. <https://doi.org/10.1016/j.intimp.2018.04.037>.
72. Zhu D, Deng X, Han XF, Sun XX, Pan TW, Zheng LP, et al. Wedelolactone enhances osteoblastogenesis through ERK- and JNK-mediated BMP2 expression and Smad1/5/8 phosphorylation. *Molecules*. 2018;23(3):561. <https://doi.org/10.3390/molecules23030561>.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.