

3-(4-Bromophenyl)-4-[2-(4-nitrophenyl)-hydrazinyl]furan-2(5H)-one

Zhu-Ping Xiao,* Li-Cheng Yi, Jia-Liang Li, Bo Zhang and Mei-Ling Liao

College of Chemistry and Chemical Engineering, Jishou University, Jishou 416000, People's Republic of China
Correspondence e-mail: xiaozhuping2005@163.com


Received 20 October 2011; accepted 23 October 2011

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.036; wR factor = 0.096; data-to-parameter ratio = 13.5.

In the title compound, $\text{C}_{16}\text{H}_{12}\text{BrN}_3\text{O}_4$, the furan-2(5H)-one ring forms a dihedral angle of $33.19(9)^\circ$ with the 4-bromobenzene unit and is nearly perpendicular to the 4-nitrobenzene segment, making a dihedral angle of $89.93(10)^\circ$. In the crystal, $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules, generating an infinite chain along [010]. The chains are linked into a three-dimensional network by $\text{C}-\text{H}\cdots\text{O}$, $\text{C}-\text{H}\cdots\pi$ and $\pi-\pi$ contacts [centroid–centroid separation = $3.805(2)\text{ \AA}$].

Related literature

For background to 3-arylfuran-2(5H)-ones as antibacterial agents, see: Xiao *et al.* (2011a,b,c). For further details of $\text{C}-\text{H}\cdots\pi$ interactions, see: Castillo *et al.* (2009); Li *et al.* (2007); Trilleras *et al.* (2009).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{12}\text{BrN}_3\text{O}_4$

$M_r = 390.20$

Orthorhombic, $Pbca$

$a = 14.4725(11)\text{ \AA}$

$b = 6.7744(5)\text{ \AA}$

$c = 31.310(2)\text{ \AA}$

$V = 3069.8(4)\text{ \AA}^3$

$Z = 8$

Mo $K\alpha$ radiation

$\mu = 2.71\text{ mm}^{-1}$

$T = 296\text{ K}$

$0.30 \times 0.20 \times 0.20\text{ mm}$

Data collection

Bruker SMART APEX CCD

diffractometer

Absorption correction: multi-scan (*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.498$, $T_{\max} = 0.614$

16108 measured reflections

3022 independent reflections

2039 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.046$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$

$wR(F^2) = 0.096$

$S = 1.02$

3022 reflections

224 parameters

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\max} = 0.58\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.63\text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg3$ is the centroid of the C1–C6 ring.

| $D-\text{H}\cdots A$ | $D-\text{H}$ | $\text{H}\cdots A$ | $D\cdots A$ | $D-\text{H}\cdots A$ |
|-----------------------------------|--------------|--------------------|-------------|----------------------|
| N1—H1A \cdots O1 ⁱ | 0.80 (4) | 2.13 (4) | 2.913 (3) | 163 (4) |
| N2—H2A \cdots O2 ⁱ | 0.91 (3) | 2.50 (3) | 2.979 (3) | 113 (2) |
| C9—H9B \cdots O3 ⁱⁱ | 0.97 | 2.45 | 3.380 (4) | 161 |
| C2—H2 \cdots Cg3 ⁱⁱⁱ | 0.93 | 2.86 | 3.676 (3) | 147 |

Symmetry codes: (i) $x, y - 1, z$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$; (iii) $-x + \frac{3}{2}, y + \frac{1}{2}, z$.

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The work was financed by a project supported by the Hunan Provincial Natural Science Foundation of China (grant No. 11J3113), the Key Laboratory of Hunan Forest Products and Chemical Industry Engineering of Hunan Province (grant No. JDZ201102) and by an aid program for the Science and Technology Innovative Research Team (Chemicals of Forestry Resources and Development of Forest Products) in Higher Educational Institutions of Hunan Province.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6467).

References

- Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Castillo, J. C., Abonia, R., Cobo, J. & Glidewell, C. (2009). *Acta Cryst. C65*, o423–o430.
- Li, X., Zhang, J., Liu, Y., Pan, M., Zheng, S., Kang, B. & Su, C. (2007). *Inorg. Chim. Acta*, **360**, 2990–2996.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Trilleras, J., Quiroga, J., Cobo, J., Hursthouse, M. B. & Glidewell, C. (2009). *Acta Cryst. C65*, o134–o139.
- Xiao, Z.-P., He, X.-B., Peng, Z.-Y., Xiong, T.-J., Peng, J., Chen, L.-H. & Zhu, H.-L. (2011a). *Bioorg. Med. Chem.* **19**, 1571–1579.
- Xiao, Z.-P., Ma, T.-W., Liao, M.-L., Feng, Y.-T., Peng, X.-C., Li, J.-L., Li, Z.-P., Wu, Y., Luo, Q., Deng, Y. & Zhu, H.-L. (2011c). *Eur. J. Med. Chem.* **46**, 4904–4914.
- Xiao, Z.-P., Ouyang, H., Wang, X.-D., Lv, P.-C., Huang, Z.-J., Yu, S.-R., Yi, T.-F., Yang, Y.-L. & Zhu, H.-L. (2011b). *Bioorg. Med. Chem.* **19**, 3884–3891.

supplementary materials

Acta Cryst. (2011). E67, o3086 [doi:10.1107/S1600536811043947]

3-(4-Bromophenyl)-4-[2-(4-nitrophenyl)hydrazinyl]furan-2(5H)-one

Z.-P. Xiao, L.-C. Yi, J.-L. Li, B. Zhang and M.-L. Liao

Comment

Furan-2(5H)-one framework is a part of many natural and synthetic compounds, which possess useful biological activities including anti-inflammatory and antitumor activity. Recently, we demonstrated that 3-aryl furan-2(5H)-ones as antibacterial agent, some of which are potent tyrosyl t-RNA synthase inhibitors (Xiao *et al.*, 2011a, 2011b and 2011c). Herein, we reported the crystal structure of the title compound (**I**) (Fig. 1).

The bond length of C7—C10 is 1.350 (4) Å and was assigned as a double bond, and the title compound was therefore identified as a furan-2(5H)-one (scheme 1). The torsion angle of N2—N1—C10—C7 is 174.8 (3) °, indicating that N1 may adopt sp^2 hybridization. Therefore the *p* orbital of N1 is conjugated with the π molecular orbital of C7—C10 double bond, which shortens the N1—C10 bond from 1.48 Å to 1.341 (4) Å. However, the torsion angle of N1—N2—C11—C16 is 21.9 (4), indicating that N2 is unlikely to have sp^2 hybridization. In fact, the angle of C11—N2—H2A, C11—N2—N1 and N1—N2—H2A are 114.83 (184), 110.30 (216) and 118.11 (24), respectively. These angles are in the range of 108 to 120 °, indicating that N2 may show a type of hybridization between sp^2 and sp^3 . This resulted in a decrease of the overlap between the nitrogen (N2) and the benzene π -orbital and elongates the N2—C11 bond (1.393 (4) Å) in comparison with N1—C10.

In the extended structure of **I**, a line of molecules is generated along the *b* axis through N1—H1A···O1 and N2—H2A···O2 hydrogen bonds characterized by a graph-set motif of R_2^2 (7) (Fig. 2). Utilizing the oxygen in the nitro group as acceptor, C—H···O hydrogen bonds link pairs of the resulted lines into centrosymmetric dimers, which are generated by edge fused graph-set motifs of R_3^3 (23) (Fig. 2).

Adjacent dimeric lines are linked together *via* C2—H2··· π contacts, forming an infinite two-dimensional layer parallel to the plane (001). The H··· π length of the typical C—H··· π hydrogen bond is in the range of 2.70 to 3.10 Å (Trilleras *et al.*, 2009; Castillo, *et al.*, 2009; Li, *et al.*, 2007). C2—H2··· π in compound **I** is thus considered as a moderate contact with H···Cg length of 2.86 Å, where Cg is the centroid of C1 to C6 (Fig. 3). The resulted layers lie parallel to the plane (001), which are further linked to form its final three-dimensional network through π — π interactions with center-center length of 3.805 (2) Å (Fig. 4).

Experimental

3-(4-Bromophenyl)-4-hydroxyfuran-2(5H)-one (0.51 g, 2 mmol) was added to a mixture of 4-nitrophenylhydrazine (0.37 g, 2.4 mmol) and *p*-toluene sulphonic acid (13.6 mg, 0.08 mmol). The resulted mixture was heated to 375 K for 30 min. Fifteen ml of toluene was then added and refluxed for 7 h. After toluene was removed under reduced pressure, the residue was purified by column chromatography on silica gel, eluting with EtOAc/petroleum ether (*v/v* = 2/1), which was partially evaporated to give colorless blocks of (**I**).

supplementary materials

Refinement

The H atoms bonded to N1 and N2 were located in difference Fourier maps, and all other H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms with C—H = 0.93 Å for aromatic H atoms and 0.97 Å for CH₂ type H atoms, respectively. $U_{\text{iso}}(\text{H})$ values were set at 1.2 times $U_{\text{eq}}(\text{C})$ for both aromatic C and the CH₂ group.

Figures




Fig. 1. Molecular structure of the title compound, showing displacement ellipsoids drawn at the 30% probability level.




Fig. 2. A dimeric line is formed through intermolecular N—H···O and C—H···O hydrogen bonds. For the sake of clarity, the H atoms have been omitted except that involving in hydrogen bonds.




Fig. 3. A two-dimensional layer is generated through intermolecular C—H···π contacts. For the sake of clarity, the H atoms have been omitted except that involving in hydrogen bonds.




Fig. 4. A three-dimensional network is finally formed through π—π interactions. For the sake of clarity, the H atoms have been omitted except that involving in hydrogen bonds.

3-(4-Bromophenyl)-4-[2-(4-nitrophenyl)hydrazinyl]furan-2(5H)-one

Crystal data

C₁₆H₁₂BrN₃O₄
 $M_r = 390.20$
Orthorhombic, *Pbca*
Hall symbol: -P 2ac 2ab
 $a = 14.4725$ (11) Å
 $b = 6.7744$ (5) Å
 $c = 31.310$ (2) Å
 $V = 3069.8$ (4) Å³
 $Z = 8$

$F(000) = 1568$
 $D_x = 1.689$ Mg m⁻³
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 1957 reflections
 $\theta = 2.3\text{--}24.6^\circ$
 $\mu = 2.71$ mm⁻¹
 $T = 296$ K
Block, colorless
 $0.30 \times 0.20 \times 0.20$ mm

Data collection

Bruker SMART APEX CCD
diffractometer

3022 independent reflections

| | |
|--|---|
| Radiation source: fine-focus sealed tube graphite | 2039 reflections with $I > 2\sigma(I)$ |
| φ and ω scan | $R_{\text{int}} = 0.046$ |
| Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996) | $\theta_{\text{max}} = 26.0^\circ, \theta_{\text{min}} = 1.9^\circ$ |
| $T_{\text{min}} = 0.498, T_{\text{max}} = 0.614$ | $h = -17 \rightarrow 17$ |
| 16108 measured reflections | $k = -8 \rightarrow 7$ |
| | $l = -38 \rightarrow 35$ |

Refinement

| | |
|---------------------------------|---|
| Refinement on F^2 | Primary atom site location: structure-invariant direct methods |
| Least-squares matrix: full | Secondary atom site location: difference Fourier map |
| $R[F^2 > 2\sigma(F^2)] = 0.036$ | Hydrogen site location: inferred from neighbouring sites |
| $wR(F^2) = 0.096$ | H atoms treated by a mixture of independent and constrained refinement |
| $S = 1.02$ | $w = 1/[\sigma^2(F_o^2) + (0.0432P)^2 + 1.6464P]$ where $P = (F_o^2 + 2F_c^2)/3$ |
| 3022 reflections | $(\Delta/\sigma)_{\text{max}} = 0.001$ |
| 224 parameters | $\Delta\rho_{\text{max}} = 0.58 \text{ e \AA}^{-3}$ |
| 0 restraints | $\Delta\rho_{\text{min}} = -0.63 \text{ e \AA}^{-3}$ |

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | x | y | z | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|--------------|-------------|---------------|----------------------------------|
| Br1 | 0.61747 (3) | 0.27111 (6) | 0.394041 (10) | 0.05931 (16) |
| C1 | 0.62850 (19) | 0.4835 (4) | 0.53598 (9) | 0.0311 (6) |
| C2 | 0.6776 (2) | 0.5813 (5) | 0.50422 (10) | 0.0429 (8) |
| H2 | 0.7131 | 0.6909 | 0.5114 | 0.051* |
| C3 | 0.6746 (2) | 0.5187 (5) | 0.46235 (10) | 0.0452 (8) |
| H3A | 0.7078 | 0.5851 | 0.4414 | 0.054* |
| C4 | 0.6220 (2) | 0.3580 (5) | 0.45199 (9) | 0.0390 (7) |
| C5 | 0.5723 (2) | 0.2560 (4) | 0.48239 (9) | 0.0407 (7) |
| H5 | 0.5371 | 0.1465 | 0.4749 | 0.049* |
| C6 | 0.5761 (2) | 0.3203 (4) | 0.52416 (9) | 0.0361 (7) |

supplementary materials

| | | | | |
|-----|--------------|-------------|--------------|-------------|
| H6 | 0.5427 | 0.2529 | 0.5449 | 0.043* |
| C7 | 0.63070 (19) | 0.5525 (4) | 0.58069 (9) | 0.0316 (6) |
| C8 | 0.6358 (2) | 0.7596 (4) | 0.59231 (10) | 0.0389 (7) |
| C9 | 0.6233 (2) | 0.5875 (4) | 0.65466 (9) | 0.0407 (7) |
| H9A | 0.5653 | 0.5791 | 0.6701 | 0.049* |
| H9B | 0.6736 | 0.5581 | 0.6741 | 0.049* |
| C10 | 0.62402 (19) | 0.4491 (4) | 0.61743 (9) | 0.0329 (7) |
| C11 | 0.5150 (2) | 0.1371 (4) | 0.67875 (9) | 0.0326 (7) |
| C12 | 0.5037 (2) | 0.0240 (4) | 0.71579 (9) | 0.0368 (7) |
| H12 | 0.5552 | -0.0264 | 0.7298 | 0.044* |
| C13 | 0.4173 (2) | -0.0127 (4) | 0.73158 (9) | 0.0383 (7) |
| H13 | 0.4098 | -0.0871 | 0.7563 | 0.046* |
| C14 | 0.3413 (2) | 0.0617 (4) | 0.71039 (9) | 0.0352 (7) |
| C15 | 0.3506 (2) | 0.1675 (5) | 0.67330 (9) | 0.0399 (7) |
| H15 | 0.2986 | 0.2133 | 0.6589 | 0.048* |
| C16 | 0.4374 (2) | 0.2055 (4) | 0.65750 (9) | 0.0396 (7) |
| H16 | 0.4441 | 0.2774 | 0.6324 | 0.048* |
| H2A | 0.649 (2) | 0.101 (5) | 0.6741 (11) | 0.057 (11)* |
| N1 | 0.61816 (19) | 0.2544 (4) | 0.62466 (9) | 0.0414 (6) |
| N2 | 0.60423 (18) | 0.1894 (4) | 0.66655 (8) | 0.0368 (6) |
| N3 | 0.24976 (19) | 0.0297 (4) | 0.72818 (8) | 0.0421 (6) |
| O1 | 0.63888 (16) | 0.9067 (3) | 0.57072 (7) | 0.0497 (6) |
| O2 | 0.63469 (16) | 0.7778 (3) | 0.63613 (7) | 0.0464 (6) |
| O3 | 0.24263 (16) | -0.0574 (3) | 0.76267 (7) | 0.0525 (6) |
| O4 | 0.18281 (16) | 0.0904 (4) | 0.70812 (8) | 0.0560 (6) |
| H1A | 0.627 (2) | 0.175 (6) | 0.6061 (11) | 0.056* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|-------------|-------------|-------------|--------------|---------------|---------------|
| Br1 | 0.0609 (3) | 0.0811 (3) | 0.0360 (2) | 0.0105 (2) | -0.00079 (16) | -0.01532 (18) |
| C1 | 0.0306 (15) | 0.0285 (15) | 0.0342 (15) | 0.0027 (13) | 0.0037 (12) | 0.0014 (12) |
| C2 | 0.0419 (18) | 0.0435 (19) | 0.0432 (18) | -0.0119 (15) | 0.0059 (15) | -0.0043 (15) |
| C3 | 0.046 (2) | 0.052 (2) | 0.0378 (17) | -0.0066 (17) | 0.0116 (15) | 0.0034 (16) |
| C4 | 0.0411 (18) | 0.0452 (19) | 0.0308 (16) | 0.0082 (15) | -0.0015 (14) | -0.0065 (14) |
| C5 | 0.0483 (19) | 0.0339 (17) | 0.0399 (17) | -0.0005 (15) | -0.0026 (14) | -0.0024 (15) |
| C6 | 0.0403 (18) | 0.0313 (16) | 0.0367 (16) | -0.0026 (14) | 0.0049 (13) | 0.0014 (13) |
| C7 | 0.0353 (16) | 0.0270 (15) | 0.0324 (15) | -0.0005 (12) | 0.0044 (13) | -0.0015 (12) |
| C8 | 0.0439 (18) | 0.0331 (18) | 0.0398 (17) | -0.0002 (14) | 0.0019 (13) | -0.0024 (15) |
| C9 | 0.056 (2) | 0.0332 (17) | 0.0332 (16) | -0.0017 (15) | 0.0021 (14) | -0.0028 (13) |
| C10 | 0.0310 (15) | 0.0302 (16) | 0.0373 (16) | -0.0018 (12) | 0.0035 (13) | -0.0052 (12) |
| C11 | 0.0422 (17) | 0.0234 (14) | 0.0321 (15) | 0.0007 (13) | -0.0022 (13) | -0.0043 (12) |
| C12 | 0.0414 (18) | 0.0335 (17) | 0.0356 (16) | 0.0032 (13) | -0.0064 (14) | 0.0046 (14) |
| C13 | 0.0475 (19) | 0.0344 (17) | 0.0331 (16) | -0.0005 (14) | 0.0004 (14) | 0.0046 (13) |
| C14 | 0.0371 (17) | 0.0340 (17) | 0.0346 (16) | -0.0010 (13) | 0.0005 (13) | -0.0043 (13) |
| C15 | 0.0419 (18) | 0.0434 (18) | 0.0343 (16) | 0.0065 (14) | -0.0102 (14) | -0.0022 (14) |
| C16 | 0.053 (2) | 0.0349 (17) | 0.0310 (15) | 0.0034 (15) | -0.0035 (14) | 0.0033 (13) |
| N1 | 0.0598 (17) | 0.0287 (15) | 0.0357 (14) | -0.0025 (13) | 0.0105 (13) | -0.0037 (11) |

| | | | | | | |
|----|-------------|-------------|-------------|--------------|--------------|--------------|
| N2 | 0.0429 (16) | 0.0332 (14) | 0.0343 (14) | 0.0002 (12) | 0.0031 (12) | 0.0036 (12) |
| N3 | 0.0442 (16) | 0.0362 (15) | 0.0458 (15) | 0.0023 (12) | -0.0006 (14) | -0.0066 (13) |
| O1 | 0.0742 (17) | 0.0283 (12) | 0.0466 (13) | -0.0027 (11) | 0.0062 (12) | 0.0009 (10) |
| O2 | 0.0715 (16) | 0.0289 (12) | 0.0389 (12) | -0.0033 (11) | 0.0000 (11) | -0.0082 (9) |
| O3 | 0.0493 (14) | 0.0603 (15) | 0.0478 (14) | -0.0014 (12) | 0.0069 (11) | 0.0025 (12) |
| O4 | 0.0396 (13) | 0.0637 (16) | 0.0648 (15) | 0.0070 (12) | -0.0068 (12) | -0.0035 (13) |

Geometric parameters (Å, °)

| | | | |
|-----------|-----------|-------------|-----------|
| Br1—C4 | 1.909 (3) | C9—H9B | 0.9700 |
| C1—C2 | 1.390 (4) | C10—N1 | 1.341 (4) |
| C1—C6 | 1.391 (4) | C11—C16 | 1.385 (4) |
| C1—C7 | 1.476 (4) | C11—N2 | 1.393 (4) |
| C2—C3 | 1.379 (4) | C11—C12 | 1.399 (4) |
| C2—H2 | 0.9300 | C12—C13 | 1.368 (4) |
| C3—C4 | 1.368 (4) | C12—H12 | 0.9300 |
| C3—H3A | 0.9300 | C13—C14 | 1.380 (4) |
| C4—C5 | 1.378 (4) | C13—H13 | 0.9300 |
| C5—C6 | 1.380 (4) | C14—C15 | 1.371 (4) |
| C5—H5 | 0.9300 | C14—N3 | 1.453 (4) |
| C6—H6 | 0.9300 | C15—C16 | 1.374 (4) |
| C7—C10 | 1.350 (4) | C15—H15 | 0.9300 |
| C7—C8 | 1.451 (4) | C16—H16 | 0.9300 |
| C8—O1 | 1.205 (3) | N1—N2 | 1.398 (4) |
| C8—O2 | 1.378 (4) | N1—H1A | 0.80 (4) |
| C9—O2 | 1.423 (4) | N2—H2A | 0.91 (3) |
| C9—C10 | 1.496 (4) | N3—O4 | 1.226 (3) |
| C9—H9A | 0.9700 | N3—O3 | 1.235 (3) |
| C2—C1—C6 | 117.9 (3) | N1—C10—C9 | 119.0 (3) |
| C2—C1—C7 | 121.1 (3) | C7—C10—C9 | 109.8 (2) |
| C6—C1—C7 | 121.1 (3) | C16—C11—N2 | 122.3 (3) |
| C3—C2—C1 | 121.2 (3) | C16—C11—C12 | 119.1 (3) |
| C3—C2—H2 | 119.4 | N2—C11—C12 | 118.3 (3) |
| C1—C2—H2 | 119.4 | C13—C12—C11 | 120.4 (3) |
| C4—C3—C2 | 119.2 (3) | C13—C12—H12 | 119.8 |
| C4—C3—H3A | 120.4 | C11—C12—H12 | 119.8 |
| C2—C3—H3A | 120.4 | C12—C13—C14 | 119.2 (3) |
| C3—C4—C5 | 121.7 (3) | C12—C13—H13 | 120.4 |
| C3—C4—Br1 | 119.3 (2) | C14—C13—H13 | 120.4 |
| C5—C4—Br1 | 118.9 (2) | C15—C14—C13 | 121.3 (3) |
| C4—C5—C6 | 118.4 (3) | C15—C14—N3 | 119.5 (3) |
| C4—C5—H5 | 120.8 | C13—C14—N3 | 119.2 (3) |
| C6—C5—H5 | 120.8 | C14—C15—C16 | 119.5 (3) |
| C5—C6—C1 | 121.7 (3) | C14—C15—H15 | 120.2 |
| C5—C6—H6 | 119.2 | C16—C15—H15 | 120.2 |
| C1—C6—H6 | 119.2 | C15—C16—C11 | 120.4 (3) |
| C10—C7—C8 | 107.0 (3) | C15—C16—H16 | 119.8 |
| C10—C7—C1 | 129.9 (3) | C11—C16—H16 | 119.8 |
| C8—C7—C1 | 123.0 (3) | C10—N1—N2 | 118.5 (3) |

supplementary materials

| | | | |
|--------------|------------|-----------------|------------|
| O1—C8—O2 | 119.0 (3) | C10—N1—H1A | 122 (3) |
| O1—C8—C7 | 131.4 (3) | N2—N1—H1A | 119 (3) |
| O2—C8—C7 | 109.6 (3) | C11—N2—N1 | 118.1 (2) |
| O2—C9—C10 | 104.4 (2) | C11—N2—H2A | 115 (2) |
| O2—C9—H9A | 110.9 | N1—N2—H2A | 110 (2) |
| C10—C9—H9A | 110.9 | O4—N3—O3 | 122.8 (3) |
| O2—C9—H9B | 110.9 | O4—N3—C14 | 118.3 (3) |
| C10—C9—H9B | 110.9 | O3—N3—C14 | 118.9 (3) |
| H9A—C9—H9B | 108.9 | C8—O2—C9 | 109.0 (2) |
| N1—C10—C7 | 131.2 (3) | | |
| C6—C1—C2—C3 | -0.1 (5) | O2—C9—C10—C7 | -2.7 (3) |
| C7—C1—C2—C3 | -179.1 (3) | C16—C11—C12—C13 | -2.2 (4) |
| C1—C2—C3—C4 | 0.2 (5) | N2—C11—C12—C13 | 172.9 (3) |
| C2—C3—C4—C5 | -0.3 (5) | C11—C12—C13—C14 | 0.5 (4) |
| C2—C3—C4—Br1 | 179.9 (2) | C12—C13—C14—C15 | 1.6 (5) |
| C3—C4—C5—C6 | 0.3 (5) | C12—C13—C14—N3 | -177.4 (3) |
| Br1—C4—C5—C6 | -179.9 (2) | C13—C14—C15—C16 | -1.9 (4) |
| C4—C5—C6—C1 | -0.1 (5) | N3—C14—C15—C16 | 177.1 (3) |
| C2—C1—C6—C5 | 0.0 (4) | C14—C15—C16—C11 | 0.1 (4) |
| C7—C1—C6—C5 | 179.0 (3) | N2—C11—C16—C15 | -173.0 (3) |
| C2—C1—C7—C10 | -148.8 (3) | C12—C11—C16—C15 | 1.9 (4) |
| C6—C1—C7—C10 | 32.2 (5) | C7—C10—N1—N2 | -174.8 (3) |
| C2—C1—C7—C8 | 34.9 (4) | C9—C10—N1—N2 | 5.0 (4) |
| C6—C1—C7—C8 | -144.0 (3) | C16—C11—N2—N1 | -21.9 (4) |
| C10—C7—C8—O1 | -176.9 (3) | C12—C11—N2—N1 | 163.1 (2) |
| C1—C7—C8—O1 | 0.1 (5) | C10—N1—N2—C11 | 99.7 (3) |
| C10—C7—C8—O2 | 1.7 (3) | C15—C14—N3—O4 | 3.7 (4) |
| C1—C7—C8—O2 | 178.7 (3) | C13—C14—N3—O4 | -177.3 (3) |
| C8—C7—C10—N1 | -179.4 (3) | C15—C14—N3—O3 | -176.5 (3) |
| C1—C7—C10—N1 | 3.8 (5) | C13—C14—N3—O3 | 2.6 (4) |
| C8—C7—C10—C9 | 0.7 (3) | O1—C8—O2—C9 | 175.3 (3) |
| C1—C7—C10—C9 | -176.0 (3) | C7—C8—O2—C9 | -3.5 (3) |
| O2—C9—C10—N1 | 177.4 (3) | C10—C9—O2—C8 | 3.7 (3) |


Hydrogen-bond geometry (\AA , $^\circ$)

Cg3 is the centroid of the C1–C6 ring.

| $D\cdots H$ | $D\cdots A$ | $H\cdots A$ | $D\cdots A$ | $D\cdots H\cdots A$ |
|----------------------------|-------------|-------------|-------------|---------------------|
| N1—H1A···O1 ⁱ | 0.80 (4) | 2.13 (4) | 2.913 (3) | 163 (4) |
| N2—H2A···O2 ⁱ | 0.91 (3) | 2.50 (3) | 2.979 (3) | 113 (2) |
| C9—H9B···O3 ⁱⁱ | 0.97 | 2.45 | 3.380 (4) | 161 |
| C2—H2···Cg3 ⁱⁱⁱ | 0.93 | 2.86 | 3.676 (3) | 147 |

Symmetry codes: (i) $x, y-1, z$; (ii) $-x+1, y+1/2, -z+3/2$; (iii) $-x+3/2, y+1/2, z$.

Fig. 1



supplementary materials

Fig. 2





Fig. 3



supplementary materials

Fig. 4

