

PRIMARY RESEARCH

Open Access

Induction of H2AX phosphorylation in tumor cells by gossypol acetic acid is mediated by phosphatidylinositol 3-kinase (PI3K) family

Zhong Guo[†], Jin Zhao^{*†}, Lei Song, Jian-Xiu Ma, Chen-Jing Wang, Shu-Yan Pei, Chao Jiang and Shang-Biao Li

Abstract

Background: H2AX is phosphorylated (γ H2AX) by members of the phosphatidylinositol 3-kinase (PI3K) family, including Ataxia telangiectasia-mutated (ATM), ATM- and Rad3-related (ATR) and DNA-PK in response to DNA damage. Our study shows that gossypol acetic acid (GAA) alone can induce γ H2AX in Human mucoepidermoid carcinoma cell line (MEC-1) in vitro. Thus, we further examined the possible mechanisms of GAA to induce γ H2AX in tumor cells.

Materials and methods: The PI3K inhibitors caffeine and wortmannin were used in an effort to identify the kinase(s) responsible for GAA-induced γ H2AX in MEC-1 cells. DNA dependent protein kinase (DNA-PK) - proficient and -deficient cells, human glioma cell lines M059K and M059J, were also used to evaluate the kinases responsible for GAA induced H2AX phosphorylation. γ H2AX expression was detected by immunofluorescent microscopy. Flow cytometry assay was used to assay γ H2AX and cell cycle.

Results: GAA induced H2AX phosphorylation in a cell cycle-dependent manner and a significant G0/G1 phase arrest in MEC-1 cells was shown. Caffeine and wortmannin significantly inhibited GAA-induced H2AX phosphorylation in MEC-1 cells. GAA induced H2AX phosphorylation in M059K, but not in M059J. Taken together, these data suggested that GAA treatment alone could induce H2AX phosphorylation in a cell cycle dependent manner in MEC-1 and M059K, but not in M059J cells. A significant G0/G1 phase arrest was shown in MEC-1.

Conclusions: The member of PI3K family, DNA-PK, ATM and ATR are involved in the H2AX phosphorylation of MEC-1 cells.

Keywords: Gossypol acetic acid, H2AX phosphorylation, DNA-PK, ATM, ATR

Introduction

Gossypol is a naturally occurring polyphenolic pigment present in cottonseed and in cotton plant byproducts, such as cottonseed oil and cottonseed meal flour. This small molecule, which is well tolerated, exhibits anti-proliferative as well as antimetastatic effects and is also described as a specific inhibitor of the anti-apoptotic proteins Bcl-XL and Bcl-2, thereby inducing apoptosis [1]. Gossypol was also found to increase the radiosensitivity of PC3 cells, contributing to an enhanced rate of apoptosis [2].

Among the many forms of DNA damage, DNA double strand breaks (DSB) are considered the most serious

threat to the cell. When not repaired or misrepaired, a single DSB can be lethal. If they do not kill the cell, these breaks can result in mutations, chromosomal rearrangements and compromised genomic integrity [3]. Recently, the relationship between DSB and the histone variant H2AX has attracted much attention. It has been shown that upon DSB generation, the H2AX protein is phosphorylated (termed γ H2AX) at serine residue 139 and forms localized "foci" at DSB sites. These foci can then recruit numerous other repair or checkpoint proteins to the damaged sites, including BRCA1, 53BP1, Werner Syndrome protein (WRN), and others [4]. Due to its close relationship with DSB, γ H2AX foci formation has been suggested by our laboratory and others as a sensitive method to detect DNA damage [5,6].

* Correspondence: gz6768@163.com

[†]Equal contributors

Medical College of Northwest University for Nationalities, Lanzhou 730030, PR China

The kinases responsible for H2AX phosphorylation have also been studied in detail. It is now clear that members of the phosphatidylinositol 3-kinase family (PI3K), including ATM (ataxia telangiectasia mutated), ATR (ATM and Rad3-related), and DNA-PK (DNA-dependent protein kinase), are involved in the phosphorylation of H2AX, although their roles may differ under different genotoxic stress or in different cell types [7]. For this reason, various kinase inhibitors have been used to differentiate between the involvements of the individual PI3K family members in specific situations. Caffeine is one such inhibitor, and it has been shown that ATM is inhibited by caffeine with an IC₅₀ of 0.2 mmol/l, ATR is inhibited with an IC₅₀ of 1.1 mmol/l, while DNA-PK is relatively resistant to caffeine with an IC₅₀ of greater than 10 mmol/l [8]. Therefore, caffeine is widely used to inhibit ATM and ATR activation. On the other hand, the microbial product wortmannin is a relatively potent inhibitor of DNA-PK (IC₅₀:16 μmol/l) and ATM (IC₅₀:150 μmol/l) activities, whereas ATR activity is less sensitive to this drug (IC₅₀:1.8 μmol/l) [9].

Recently, gossypol was reported to interfere with inositol phosphate metabolism by inhibiting the activity of inositol-1,4,5-triphosphate 3-kinase (IP3K) isoforms, which are essential for the formation of higher phosphorylated inositols. The level of one isoform of this kinase (IP3K-A) is enhanced in many human tumor cell lines and appears to be associated with a malignant phenotype. One important product of the IP3K dependent metabolism is InsP6. This inositol phosphate is known to be involved in chromatin remodeling and DSB repair processes such as Non-homologous end-joining (NHEJ), as it was shown to interact directly with the Ku70/Ku80 heterodimer, leading to a stimulation of NHEJ [10,11]. These data suggest that a disturbance of the inositol phosphate metabolism by gossypol might also have an effect on DSB repair. Therefore, it was of interest to examine gossypol-triggered induction of γH2AX in tumor cells.

In our previous study, DBS and γH2AX foci have been shown by neutral comet and immunostaining assay in GAA induced MEC-1 cells [12]. Thus, we further examined the possible mechanisms of GAA to induce γH2AX in different tumor cells. By using caffeine and wortmannin, as well as DNA dependent protein kinase (DNA-PK) –proficient and -deficient cells, Human glioma cell line M059K and M059J, the kinases responsible for H2AX phosphorylation induced by GAA were investigated. As reported here, GAA treatment alone could induce H2AX phosphorylation in a cell cycle-dependent manner in MEC-1 and M059K, but not in M059J cells. Also G0/G1 phase arrest is shown MEC-1 cells. The member of PI3K family, DNA-PK, ATM and ATR are involved in the H2AX phosphorylation of MEC-1 cells.

Materials and methods

Chemicals and antibodies

GAA, caffeine, wortmannin, 4,6-diamidino-2-phenylindole (DAPI), Propidium Iodide (PI) and dimethylsulfoxide (DMSO) were purchased from Sigma (St Louis, MO). Gossypol and wortmannin were dissolved in DMSO and stored at –80°C. Caffeine was dissolved in PBS and stored at –20°C. Anti-γH2AX monoclonal antibody was purchased from Upstate Technology (Lake Placid, NY). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG (H + L) were purchased from Jackson ImmunoResearch (West Grove, PA).

Cell culture and treatment

Human mucoepidermoid carcinoma MEC-1 cell was purchased from School of Stomatology Fourth Military Medical University (Xi'an, Shanxi, PRC). DNA dependent protein kinase (DNA-PK) –proficient and -deficient cells, the two human malignant glioma cell lines M059K and M059J, isolated and established by Allalunis-Turner et al. [13], were kept in liquid nitrogen at the Institution of Modern Physics of the Chinese Academy of Science. The cells were maintained as monolayer cultures in Dulbecco's Modified Eagle's Medium (DMEM, Invitrogen, Scotland, UK) for MEC-1 and DMEM/F12 (1:1) medium supplemented (10% newborn calf serum, 100 U/ml penicillin, 125 μg/ml streptomycin, and 0.03% glutamine) (Life Technology, Paisley, UK) for M059J and M059K cells. For GAA treatment, various concentrations of GAA were added directly to the appropriate culture plates. Caffeine (8 μmol/l) or wortmannin (20 μmol/l) were added 1 h before GAA treatment to inhibit the function of ATM, ATR and DNA-PK.

Immunofluorescence microscopy for γH2AX

Immunofluorescent microscopy was conducted according to previously reported procedures with modifications [5]. Briefly, 2×10^4 cells were seeded into 35 mm dishes containing a glass cover slip in each well. After irradiation, slides were air-dried, and fixed for 0.5 h in 2% paraformaldehyde in TBS. Cells were rinsed in TBS, placed in –20°C methanol for 1 min, rinsed, then placed for 20 min in TBS plus 1% bovine serum albumin and 0.2% Tween-20 (TTN) and finally incubated for 2 h with diluted antiphosphohistone H2AX (Ser-139) mAb (Upstate, Lake Placid, NY) diluted 1:500 in TTN. Slides were washed and incubated with FITC-conjugated anti-mouse goat F (ab')² fragment (DAKO, Carpinteria, CA) diluted 1:200 in TTN for 1 h at room temperature. Slides were rinsed and then immersed in 0.05 mg/ml DAPI for 15 min, rinsed and mounted with cover slips using 10 μl Fluorogard (Bio-Rad) as the antifade mounting medium, and sealed. To prevent bias in selection of cells that display foci, over 800 randomly selected cells were counted.

Cells with three or more foci of any size were classified as positive. All experiments were repeated in triplicate.

Flow cytometry assay for γ H2AX and cell cycle

Flow cytometry analysis was conducted as described earlier [14,15]. After the various treatments, cells were fixed with cold 70% methanol and kept at -20°C for up to 2 weeks until further analysis. Cells were centrifuged and rinsed with PBS, blocked with PST (4% fetal bovine serum in PBS) for 15 min at room temperature and rinsed with PBS. Cells were incubated with anti- γ H2AX monoclonal antibody at a 300-fold dilution for 2 h at room temperature, rinsed with PBS, incubated with Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG (H + L) at a 100-fold dilution for another 1 h at room temperature and rinsed again in PBS. Cells were further incubated for 0.5 h at room temperature with 50 $\mu\text{g/ml}$ PI. Filter the sample through a cell strainer (40 μm pore size) and collect single cancer cells samples were analyzed using a flow cytometer (Becton-Dickinson, Bedford, MA, USA). Cell cycle analysis was conducted as described by Amrein et al. [16].

To examine the relationship between GAA-induced γ H2AX and cell cycle, the changes in γ H2AX immunofluorescence intensity (H2AX IF) were calculated in each phase of the cycle by gating G1, S and G2M cells based on differences in DNA content. The means of γ H2AX positive ratios for G1, S and G2/M populations of cells in the DMSO control groups were subtracted from the respective means of the GAA-treated cells. After this subtraction, the GAA-induced changes in positive γ H2AX ratio were obtained. Data is presented as the mean of the γ H2AX positive ratio of each cell cycle compartment. All experiments were performed three times.

Statistical analysis

SPSS version 13.0 software (SPSS Inc., Chicago, Illinois, USA) was used for the statistical analysis. Data are expressed as mean \pm standard deviation (SD). A two-tailed Student's *t*-test was performed to assess the differences between two groups. Statistical inferences were based on two-sided tests at a significance level of $p < 0.05$.

Results

GAA induces H2AX phosphorylation in a cell cycle-dependent manner in MEC-1 cells


Typical flow cytometry histograms of GAA-induced phosphorylation of H2AX and cell cycle distribution in MEC-1 cells are shown in Figure 1A and D. The percentage of γ H2AX positive cells showed little change in all phases when cells were incubated with 2.5 and 5 $\mu\text{mol/l}$ GAA for 24 h, 10 $\mu\text{mol/l}$ GAA had no effect on S and G2/M phase cells. The percentage of γ H2AX positive cells had

a significant increase at higher doses (20 and 40 $\mu\text{mol/l}$ GAA). G0/G1 phase cells were most sensitive to DNA damage expressing approximately 13.9% and 26.9% higher levels of γ H2AX than S and 14.4% and 31.6% higher levels of γ H2AX than G2/M phase cells after being treated with 20 and 40 $\mu\text{mol/l}$ GAA respectively for 24 h. It appeared a linear time-related increase in expression γ H2AX, especially when the incubation time increased to 24 and 48 h. It is quite evident that compared to S and G2/M phase cells, more G0/G1 phase cells expressed increased levels of γ H2AX following GAA treatment with the increase of incubation time (Figure 1B and C).

The dependency of the cell cycle of the MEC-1 cells to GAA exposure is presented in Figure 1E and F. There was a significant G0/G1 phase arrest. After 24 h incubation, 10, 20 and 40 $\mu\text{mol/l}$ GAA induced 59.7%, 63.2% and 70.2% cells in G0/G1 and 20 $\mu\text{mol/l}$ GAA treatment for 12, 24 and 48 h, there were 61.6%, 63.0% and 76.7% cells in G0/G1.

Wortmannin and caffeine inhibited GAA-induced γ H2AX in MEC-1 cells

Since both flow cytometry and immunofluorescent assays [12] proved GAA could induce the phosphorylation of H2AX in MEC-1 cells, we then were interested in determining which specific PI3K might be responsible for this process. Caffeine, one PI3K inhibitor, should block the action of ATM and ATR, and wortmannin, which should block the action of DNA-PK and ATM. Therefore, MEC-1 cells were preincubated for 1 h with 8 $\mu\text{mol/l}$ caffeine and 20 $\mu\text{mol/l}$ wortmannin, a concentration that should inhibit the action of ATM, ATR and DNA-PK. Subsequently the cells underwent GAA treatment [17]. Shown in Figure 2 are representative immunofluorescent images of MEC-1 cells treated with GAA alone, with or without caffeine and wortmannin preincubation. The incubation of cells with caffeine and wortmannin alone led to an increase in the level of γ H2AX (30.9%, 37.5% and 37.2% in untreated, caffeine and wortmannin treated MEC-1 cells). The increased percentages of γ H2AX positive cells were calculated by the γ H2AX positive cells percentages of GAA and caffeine (wortmannin) treated groups subtracted from the γ H2AX positive cell percentages of GAA treatment alone group. The detailed dose and time response analyses are shown in Figure 3, which illustrates that both caffeine and wortmannin decreased the level of γ H2AX in a dose-dependent manner and the inhibitory action of wortmannin was more obvious than caffeine. Caffeine and wortmannin did not decrease the level of γ H2AX until 24 to 48 h treatment. Together, these data demonstrate that the member of PI3K family, DNA-PK, ATM and ATR are involved in the H2AX phosphorylation in MEC-1 cells.



Immunofluorescent assay showed GAA-induced γ H2AX in M059K and M059J cells

To further distinguish the kinases responsible for GAA-induced H2AX phosphorylation, DNA dependent protein kinase (DNA-PK) – proficient and -deficient cells, M059K and M059J, were treated with GAA (from 2.5 to 20 μ mol/l) at different time points. Figure 2 shows representative immunofluorescent images of M059K and M059J cells treated with 20 μ mol/l GAA for 24 h.

Figure 4 depicts the percentage of γ H2AX positive cells, clearly demonstrating that GAA could indeed induce γ H2AX in a dose- and time-dependent manner in M059K

cells, however, for M059J cells, the levels of γ H2AX did not increased with the GAA dose, the increase only could be seen at 20 μ mol/l GAA incubation for 24 and 48 h. These data further demonstrate that DNA-PK is involved in GAA-induced H2AX phosphorylation in M059K and M059J cells.

GAA induces γ H2AX phosphorylation in a cell cycle-dependent manner in M059K and M059J cells

Typical flow cytometry histograms of GAA-induced phosphorylation of H2AX and cell cycle distribution in M059K and M059J are shown in Figure 5A,B,E and F. The




Figure 2 Digitized images of γ H2AX foci in MEC-1 cells, M059K and M059J cells. MEC-1 cells were pretreated with 8 mmol/l caffeine and 20 nmol/l wortmannin for 1 h and then were exposed to 20 μ mol/l GAA for 24 h. M059K and M059J cells were treated with 20 μ mol/l GAA for 24 h. DNA was stained with DAPI and γ H2AX was detected using an Alexa 488- conjugated secondary antibody after staining using a monoclonal anti- γ H2AX antibody (Scale bar, 15 μ m).

dose–response for M059K and M059J cells was similar with the results of the immunofluorescent assay. For M059K cells, the percentage of γ H2AX positive cells there was little change in all phases when cells were incubated with lower doses of GAA, however, the percentage of γ H2AX positive cells had a significant increase at higher doses. After 24 h incubation of 20 and 40 μ mol/l GAA, G0/G1 phase cells express approximately 4.9% and 17.1% higher levels of γ H2AX than S and 6.0% and 20.7% higher levels of γ H2AX than G2/M phase cells. S phase cells express approximately 1.1% and 3.6% higher levels of γ H2AX than G2/M phase cells after 20 and 40 μ mol/l GAA treated for 24 h. For M059J cells, the percentage of γ H2AX positive cells showed little change in all phases even when the GAA doses reached to 40 μ mol/l (Figure 5C and D).

The dependency of the cell cycle of M059K and M059J cells to GAA exposure are presented in Figure 5G and H. S phase cells are more sensitive than other cell cycle-phases cells. After 24 h incubation, 20 and 40 μ mol/l GAA induced 24.0% and 31.8% M059K cells in S, 10, 20 and 40 μ mol/l GAA induced 30.2%, 30.8% and 36.5% cells in S for M059J cells.

Discussion

A wealth of published evidence supports the conclusion that H2AX is specifically phosphorylated in response to

the occurrence of DSB caused by a variety of genotoxic agents [18,19]. Moreover, at least with ionizing radiation, there appears to be a direct linear correlation between the number of γ H2AX foci and the number of DSB, which suggests that each γ H2AX focus may represent an individual break [20]. Therefore, when results were reported that gossypol might also have an effect on DSB, it was of interest to investigate this issue in more detail.

As reported here, the result of flow cytometry assay shows that GAA treatment at doses ranging from 2.5–40 μ mol/l can induce γ H2AX in a dose and time-dependent manner in MEC-1 cells. This was in accordance with the results of immunofluorescence assays we had observed [12]. Interestingly, the degree of H2AX phosphorylation across the cell cycle-phase after GAA exposure was different and a significant G0/G1 arrest was visible. G0/G1 phase cells are much more sensitive and expressed higher levels of γ H2AX compared to other cell cycle-phase cells following GAA treatment in MEC-1 cells.

What is the possible significance of these data? One reason maybe that G0/G1 phase cells have an increased vulnerability to oxidant damage during DNA replication. These data can also be explained by the possibility that successful repair of DSB depends on timing in the cell cycle [21]. At last, if proliferating cells exposed to GAA experience similar levels of DSB during each phase of




Figure 3 The dose and time responses of caffeine and wortmannin inhibition of the GAA-induced H2AX phosphorylation in MEC-1 cells. MEC-1 cells were pretreated with 8 mmol/l caffeine and 20 nmol/l wortmannin for 1 h and then were exposed to different doses of GAA. γ H2AX foci formation was observed by immunofluorescent microscopy. **(A)** Dose responses of GAA-induced H2AX phosphorylation (GAA treated for 24 h); **(B)** Time responses of 20 μ mol/l GAA-induced H2AX phosphorylation. (*: $P < 0.05$, **: $P < 0.01$ vs GAA treated alone).

the cell cycle but dissimilar repair rates, they may be particularly susceptible to accumulating deleterious DNA defects during that specific phase.

The observation from the study of PI3K inhibitors is that caffeine and wortmannin inhibited GAA-induced γ H2AX in a dose but not time-dependent manner in MEC-1 cells. This observation was not surprising, given the reported action of caffeine and wortmannin, inhibition of ATM, ATR and DNA-PK enzymes that are thought to catalyze γ H2AX formation [22,23]. We wondered why for caffeine, the inhibition of GAA-induced γ H2AX was not so obvious as wortmannin. Cortez has reported that

caffeine does not inhibit ATM and ATR in HCT116 cells. The reason for that, as shown by Cortez, is that caffeine can promote the phosphorylation and activation of ATM under some circumstances [24]. Some other possible explanations may also exist. For example, caffeine may delay the normal dephosphorylation pattern [25] and inhibit homology directed repair [26]. Also, if only a portion of ATR or ATM molecules that are required to prevent DNA damage are accessible to caffeine then the remaining, uninhibited molecules could become superactivated [24]. Finally, it is also possible that ATM or ATR did not phosphorylate H2AX at all in HeLa cells. In




Figure 4 The dose and time responses of GAA induced H2AX phosphorylation in M059K and M059J cells. γ H2AX foci formation was observed by immunofluorescent microscopy. The dose responses of GAA-induced H2AX phosphorylation in M059K and M059J cells **(A)** (GAA treated for 24 h); The time responses of 20 μ mol/l GAA-induced H2AX phosphorylation in M059K and M059J cells **(B)**. (*: $P < 0.05$, **: $P < 0.01$ vs Control).




Figure 5 (See legend on next page.)

(See figure on previous page.)

Figure 5 GAA induced H2AX phosphorylation in a cell cycle-dependent manner in M059K and M059J cells. (A) (B) Bivariate(γ H2AX IF vs DNA content) distributions of control and GAA-treated M059K and M059J cells (GAA treated for 24 h). (C) (D) The percentage of γ H2AX positive cells in G1, S and G2/M cells after M059K and M059J cells were treated with different doses GAA for 24 h. (E) (F) Cell cycle distribution of M059K and M059J cells 24 h following GAA treatment. (G) (H) Cell cycle distribution of M059K and M059J cells after treatment with different doses of GAA for 24 h. (ACEG: M059K; BDFH: M059J. a: Control. b: 2.5 μ mol/l GAA. c: 5 μ mol/l GAA. d: 10 μ mol/l GAA. e: 20 μ mol/l GAA. f: 40 μ mol/l), n = 3, *: P < 0.05, **: P < 0.01 vs Control.

all, our data suggest that DNA-PK, ATM and ATR are involved in the H2AX phosphorylation of MEC-1 cells.

Since it is clear that DNA-PK participated in the process of phosphorylation of H2AX [27], DNA dependent protein kinase (DNA-PK)- proficient and -deficient cells, human glioma cell line M059K and M059J, were used to observe the effect of GAA-induced H2AX phosphorylation. Interestingly, differences were observed between M059K and M059J cells in terms of the degree of H2AX phosphorylation. In M059K cells, it was found GAA-induced γ H2AX in a dose and time-dependent manner, by contrast, GAA almost could not induce the formation of H2AX in M059J, indicating that DNA-PK is actually responsible for GAA-induced H2AX phosphorylation. Perhaps more interesting was the cell cycle-phase differences on GAA-induced γ H2AX phosphorylation in different cells. In M059K cells, S phase cells were much more sensitive and expressed higher levels of H2AX compared to other cell cycle-phase cells following GAA treatment. For M059J cells, the expression of H2AX was not different across the cell cycle phases, but S phase cells was also more sensitive than other cell cycle-phases cells. These results were in accordance with others who have reported that after exposure to different DNA double-strand break-inducing agents, different G2/M or S-phase arrests in M059J and M059K cells are seen [27,28].

In summary, in the present study, we have shown that GAA treatment alone can induce H2AX phosphorylation in the cell lines examined, and members of the PI3K family, DNA-PK, ATM and ATR are also involved in the H2AX phosphorylation in MEC-1 cells. In addition, the degree of H2AX phosphorylation across the cell cycle and cell cycle arrest were different for examined cells. The underlying mechanism is worth further investigation.

Abbreviations

GAA: Gossypol acetic acid; DMEM: Dulbecco's modified Eagle's medium; PI: Propidium Iodide; DMSO: Dimethyl sulfoxide; γ H2AX: H2AX phosphorylation; ATM: Ataxia telangiectasia-mutated; ATR: ATM and Rad3-related; DNA-PK: DNA dependent protein kinase; DSB: DNA double strand breaks; NHEJ: Non-homologous end-joining; PI3K: Phosphatidylinositol 3-kinase.

Competing interests

All authors have full access to all of the data in the study and have full responsibility for the decision to submit for publication. The authors declare that they have no competing interests.

Authors' contributions

ZG was project manager, JZ was project manager and wrote most of the article. LS and JM was principal investigator of this Flow cytometry assay, CW and SP was principal investigator of this Immunofluorescence microscopy, CJ and SL was principal investigator of this Cell culture and treatment. All authors read and approved the final manuscript.

Acknowledgements

This work was supported in part by grants from the National Natural Science Foundation of China (Nos.31060127 and 81260442), Gansu Provincial Natural Science Foundation (No.1010RJZA078 and No.1208RJZA246), the Project Sponsored by the Scientific Research Foundation for the Returned Overseas Chinese Scholars, State Education Ministry(2011)1568 and The State Ethnic Affairs Commission Research Project (No.12XB11).

Received: 26 June 2014 Accepted: 4 December 2014

Published online: 16 December 2014

References

1. Lin J, Wu Y, Yang D, Zhao Y: Induction of apoptosis and antitumor effects of a small molecule inhibitor of Bcl-2 and Bcl-xl, gossypol acetate, in multiple myeloma in vitro and in vivo. *Oncol Rep* 2013, **30**:731-738.
2. Akagunduz O, Karaca B, Atmaca H, Uzunoglu S, Karabulut B, Sanli UA, Haydaroglu A, Uslu R: Radiosensitization of hormone-refractory prostate cancer cells by gossypol treatment. *J Buon* 2010, **15**:763-767.
3. Shrivastav M, De Haro LP, Nickoloff JA: Regulation of DNA double-strand break repair pathway choice. *Cell Res* 2008, **18**:134-147.
4. Sharma A, Singh K, Almasan A: Histone H2AX phosphorylation: a marker for DNA damage. *Methods Mol Biol* 2012, **920**:613-626.
5. Zhao J, Guo Z, Zhang H, Wang ZH, Song L, Ma JX, Pei SY, Wang CJ: The potential value of the neutral comet assay and γ H2AX foci assay in assessing the radiosensitivity of carbon beam in human tumor cell lines. *Radiol Oncol* 2013, **47**:247-257.
6. Banuelos CA, Banáth JP, MacPhail SH, Zhao J, Reitsema T, Olive PL: Radiosensitization by the histone deacetylase inhibitor PCI-24781. *Clin Cancer Res* 2007, **15**:6816-6826.
7. Koike M, Mashino M, Sugawara J, Koike A: Histone H2AX phosphorylation independent of ATM after X-irradiation in mouse liver and kidney in situ. *J Radiat Res* 2008, **49**:445-449.
8. Daniel R, Marusich E, Argyris E, Zhao RY, Skalka AM, Pomerantz RJ: Caffeine inhibits human immunodeficiency virus type 1 transduction of nondividing cells. *J Virol* 2005, **79**:2058-2065.
9. Collis SJ, DeWeese TL, Jeggo PA, Parker AR: The life and death of DNA-PK. *Oncogene* 2005, **24**:949-961.
10. Mayr GW, Windhorst S, Hillemeier K: Antiproliferative plant and synthetic polyphenolics are specific inhibitors of vertebrate inositol-1,4,5-trisphosphate 3-kinases and inositol polyphosphate multikinase. *J Biol Chem* 2005, **280**:13229-13240.
11. Byrum J, Jordan S, Safrany ST, Rodgers W: Visualization of inositol phosphate-dependent mobility of Ku: depletion of the DNA-PK cofactor InsP6 inhibits Ku mobility. *Nucleic Acids Res* 2004, **32**:2776-2784.
12. Guo Z, Zhao J, Xue TM, Ma JX, Wang CJ, Huang SS: Gossypol acetic acid induces DNA double-strand breaks in human mucoepidermoid carcinoma cell MEC-1. *Sheng Li Xue Bao* 2011, **63**:164-170.
13. Allalunis-Turner MJ, Barron GM, Day RS 3rd, Dobler KD, Mirzayans R: Isolation of two cell lines from a human malignant glioma specimen differing in sensitivity to radiation and chemotherapeutic drugs. *Radiat Res* 1993, **134**:349-354.

14. Albino AP, Huang X, Jorgensen ED, Gietl D, Traganos F, Darzynkiewicz Z: Induction of DNA double-strand breaks in A549 and normal human pulmonary epithelial cells by cigarette smoke is mediated by free radicals. *Int J Oncol* 2006, **28**:1491–1505.
15. Schmid TE, Dollinger G, Beisker W, Hable V, Greubel C, Auer S, Mittag A, Tarnok A, Friedl AA, Molls M, Röper B: Differences in the kinetics of gamma-H2AX fluorescence decay after exposure to low and high LET radiation. *Int J Radiat Biol* 2010, **86**:682–691.
16. Loignon M, Amrein L, Dunn M, Aloyz R: XRCC3 depletion induces spontaneous DNA breaks and p53-dependent cell death. *Cell Cycle* 2007, **6**:606–611.
17. Yan C, Lu J, Zhang G, Gan T, Zeng Q, Shao Z, Duerksen-Hughes PJ, Yang J: Benzo(a)pyrene induces complex H2AX phosphorylation patterns by multiple kinases including ATM, ATR, and DNA-PK. *Toxicol In Vitro* 2011, **25**:91–99.
18. Garcia-Canton C, Anadón A, Meredith C: γ H2AX as a novel endpoint to detect DNA damage: applications for the assessment of the in vitro genotoxicity of cigarette smoke. *Toxicol In Vitro* 2012, **26**:1075–1086.
19. Chua ML, Rothkamm K: Biomarkers of radiation exposure: can they predict normal tissue radiosensitivity? *Clin Oncol (R Coll Radiol)* 2013, **25**:610–616.
20. Klokov D, MacPhail SM, Banáth JP, Byrne JP, Olive PL: Phosphorylated histone H2AX in relation to cell survival in tumor cells and xenografts exposed to single and fractionated doses of X-rays. *Radiother Oncol* 2006, **80**:223–229.
21. Rothkamm K, Löbrich M: Misrepair of radiation-induced DNA double-strand breaks and its relevance for tumorigenesis and cancer treatment (review). *Int J Oncol* 2002, **21**:433–440.
22. Blasina A, Price BD, Turenne GA, McGowan CH: Caffeine inhibits the checkpoint kinase ATM. *Curr Biol* 1999, **9**:1135–1138.
23. Sarkaria JN, Busby EC, Tibbetts RS, Roos P, Taya Y, Karnitz LM, Abraham RT: Inhibition of ATM and ATR kinase activities by the radiosensitizing agent, caffeine. *Cancer Res* 1999, **59**:4375–4382.
24. Cortez D: Caffeine inhibits checkpoint responses without inhibiting the ataxia-telangiectasia-mutated (ATM) and ATM- and Rad3-related (ATR) protein kinases. *J Biol Chem* 2003, **278**:37139–37145.
25. Wang X, Wang H, Iliakis G, Wang Y: Caffeine-induced radiosensitization is independent of nonhomologous end joining of DNA double-strand breaks. *Radiat Res* 2003, **159**:426–432.
26. Wang H, Wang X, Iliakis G, Wang Y: Caffeine could not efficiently sensitize homologous recombination repair-deficient cells to ionizing radiation-induced killing. *Radiat Res* 2003, **159**:420–425.
27. Anderson JA, Harper JV, Cucinotta FA, O'Neill P: Participation of DNA-PKcs in DSB repair after exposure to high- and low-LET radiation. *Radiat Res* 2010, **174**:195–205.
28. Holgersson A, Heiden T, Castro J, Edgren MR, Lewensohn R, Meijer AE: Different G2/M accumulation in M059J and M059K cells after exposure to DNA double-strand break-inducing agents. *Int J Radiat Oncol Biol Phys* 2005, **61**:915–921.

doi:10.1186/s12935-014-0141-5

Cite this article as: Guo et al.: Induction of H2AX phosphorylation in tumor cells by gossypol acetic acid is mediated by phosphatidylinositol 3-kinase (PI3K) family. *Cancer Cell International* 2014 **14**:141.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

