

# RNA Toxicity and Perturbation of rRNA Processing in Spinocerebellar Ataxia Type 2

Pan P. Li, PhD,<sup>1\*</sup> Roumita Moulick, PhD,<sup>2</sup> Hongxuan Feng, PhD,<sup>1</sup> Xin Sun, PhD,<sup>1</sup> Nicolas Arbez, PhD,<sup>1</sup> Jing Jin, MD, PhD,<sup>1</sup> Leonard O. Marque, BS,<sup>1</sup> Erin Hedglen, MA,<sup>1</sup> H.Y. Edwin Chan, PhD,<sup>3</sup> Christopher A. Ross, MD, PhD,<sup>1,4,5</sup> Stefan M. Pulst, MD,<sup>6</sup> Russell L. Margolis, MD,<sup>1,4</sup> Sarah Woodson, PhD,<sup>2</sup> and Dobrila D. Rudnicki, PhD<sup>1</sup>

<sup>1</sup>Department of Psychiatry and Behavioral Sciences, Division of Neurobiology, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

<sup>2</sup>T.C. Jenkins Department of Biophysics, Johns Hopkins University, Baltimore, Maryland, USA

<sup>3</sup>Biochemistry Program, School of Life Sciences, The Chinese University of Hong Kong, Hong Kong, China

<sup>4</sup>Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

<sup>5</sup>Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

<sup>6</sup>Department of Neurology, University of Utah, Salt Lake City, Utah, USA

**ABSTRACT: Background:** Spinocerebellar ataxia type 2 (SCA2) is a neurodegenerative disease caused by expansion of a CAG repeat in *Ataxin-2* (*ATXN2*) gene. The mutant *ATXN2* protein with a polyglutamine tract is known to be toxic and contributes to the SCA2 pathogenesis.

**Objective:** Here, we tested the hypothesis that the mutant *ATXN2* transcript with an expanded CAG repeat (*expATXN2*) is also toxic and contributes to SCA2 pathogenesis.

**Methods:** The toxic effect of *expATXN2* transcripts on SK-N-MC neuroblastoma cells and primary mouse cortical neurons was evaluated by caspase 3/7 activity and nuclear condensation assay, respectively. RNA immunoprecipitation assay was performed to identify RNA binding proteins (RBPs) that bind to *expATXN2* RNA. Quantitative PCR was used to examine if ribosomal RNA (rRNA) processing is disrupted in SCA2 and Huntington's disease (HD) human brain tissue.

**Results:** *expATXN2* RNA induces neuronal cell death, and aberrantly interacts with RBPs involved in RNA metabolism. One of the RBPs, transducin  $\beta$ -like protein 3 (TBL3), involved in rRNA processing, binds to both *expATXN2* and expanded *huntingtin* (*expHTT*) RNA *in vitro*. rRNA processing is disrupted in both SCA2 and HD human brain tissue.

**Conclusion:** These findings provide the first evidence of a contributory role of *expATXN2* transcripts in SCA2 pathogenesis, and further support the role of *expHTT* transcripts in HD pathogenesis. The disruption of rRNA processing, mediated by aberrant interaction of RBPs with *expATXN2* and *expHTT* transcripts, suggest a point of convergence in the pathogenesis of repeat expansion diseases with potential therapeutic implications. © 2021 The Authors. *Movement Disorders* published by Wiley Periodicals LLC on behalf of International Parkinson and Movement Disorder Society

**Key Words:** RNA toxicity; RNA binding protein; *ATXN2*

Spinocerebellar ataxia type 2 (SCA2) is an autosomal dominant disorder caused by a CAG repeat expansion

in the first exon of the *Ataxin-2* (*ATXN2*) gene located on chromosome 12q24.<sup>1</sup> The repeat is in-frame to

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

\***Correspondence to:** Dr. Pan P. Li, Department of Psychiatry and Behavioral Sciences, Division of Neurobiology, Johns Hopkins University School of Medicine, 600 N. Wolfe St., Baltimore, MD 21287, USA; E-mail: ple5@jhmi.edu

Roumita Moulick and Hongxuan Feng contributed equally to this paper.

**Relevant conflicts of interest/financial disclosures:** Nothing to report.

**Funding agencies:** This work was supported by the National Institutes of Health grants NS064138 (to D.D.R.), NS112796 (to P.P.L.), NS112687 (to P.P.L.), NS099397 (to S.W.) and NS033123 (to S.M.P.), and the ABCD Charitable Trust (to R.L.M.).

**Received:** 5 April 2021; **Revised:** 3 June 2021; **Accepted:** 12 July 2021

**Published online 14 August 2021 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mds.28729**

encode polyglutamine (polyQ). The signs and symptoms of SCA2 include progressive deterioration in balance and coordination, neuropathies, nystagmus and slow saccadic eye movements, slurred speech, and cognitive impairment.<sup>2-5</sup> SCA2 is the second most common form of autosomal dominant ataxia, with a prevalence of 1–2 cases/10<sup>5</sup> inhabitants, varying somewhat by ethnicity and geographic location.<sup>2,3,6-8</sup> The highest prevalence of the SCA2 mutation occurs in Cuba (6.57 cases/10<sup>5</sup> inhabitants)<sup>9</sup> and is likely a consequence of a founder effect.<sup>10</sup> SCA2 neuropathology is characterized by a significant loss of cerebellar Purkinje neurons, a less prominent loss of cerebellar granule cells<sup>11</sup>; marked neuronal loss in the inferior olive, pontocerebellar nuclei, and substantia nigra; degeneration of the thalamus and pons, and thinning of the cerebellar cortex without changes in neuronal density.<sup>11-14</sup> The normal *ATXN2* allele contains 15 to 32 CAG triplets, whereas the disease allele typically has 33 to 64 triplets.<sup>15</sup> The most common disease allele has 37 triplets, and neonatal onset SCA2 cases with over 200 CAG repeats have been reported.<sup>16</sup> Similar to other CAG repeat diseases, the repeat length in SCA2 is inversely correlated to age of onset.<sup>17,18</sup> Recently, intermediate CAG expansion in *ATXN2* has been associated with a higher risk for amyotrophic lateral sclerosis (ALS).<sup>19</sup> Current evidence indicates that neurotoxicity of *ATXN2* protein, which is involved in multiple cellular pathways, including messenger RNA (mRNA) maturation, translation, and endocytosis, is central to SCA2 pathogenesis.<sup>20,21</sup> This is supported by data from several SCA2 cell and mouse models expressing mutant *ATXN2* protein.<sup>22-24</sup>

However, multiple laboratories, including ours, have demonstrated an important neurotoxic role for mutant RNA transcripts in CAG/CTG repeat expansion diseases, including myotonic dystrophy type 1 (DM1),<sup>25,26</sup> Huntington's disease (HD),<sup>27,28</sup> Huntington's disease-like 2 (HDL2),<sup>29</sup> SCA3,<sup>30-32</sup> and SCA8.<sup>33</sup> RNA-triggered pathogenic processes are thought to be, at least in part, mediated by aberrant interaction between expanded repeat-containing RNA transcripts and RNA-binding proteins (RBPs).<sup>34-36</sup> The basic hypothesis is that expanded CAG/CUG repeats in transcripts form hairpin structures that sequester multiple RBPs and hence, prevent the RBPs from performing their normal function in cells.<sup>35</sup> To add to the pathomechanistic complexity of CAG/CUG repeat diseases, antisense transcripts that span the CAG/CUG repeat regions are also expressed at the DM1 (CAG direction),<sup>37</sup> HDL2 (CAG direction),<sup>38,39</sup> SCA7 (CUG direction),<sup>40</sup> SCA8 (CUG direction),<sup>33</sup> and HD (CUG direction)<sup>41</sup> loci. We have recently described a transcript expressed antisense to *ATXN2* at the SCA2 locus<sup>42</sup> and provided evidence that this antisense *ATXN2* (*ATXN2-AS*) transcript contributes to SCA2 pathogenesis and potentially to

ALS associated with an intermediate repeat expansion at the *ATXN2* locus.<sup>19</sup>

We hypothesized that, in addition to mutant *ATXN2* protein and mutant *ATXN2-AS* transcript,<sup>42</sup> mutant sense *ATXN2* RNA also contributes to SCA2 pathogenesis.

As predicted by this hypothesis, the data presented here demonstrate that sense *expATXN2* transcripts are neurotoxic in cell models in the absence of expression of mutant *ATXN2* protein, aberrantly interact with RBPs that are involved in ribosomal RNA (rRNA) processing, and lead to disruption of rRNA processing. We demonstrate a similar disruption of rRNA processing in HD patient brain tissue. Similar to findings in other repeat expansion diseases, SCA2 is, therefore, the fifth neurodegenerative CAG/CTG repeat expansion disease in which pathogenesis is likely a consequence of a combination of expression of mutant protein and bi-directionally expressed mutant RNA.


## Materials and Methods

A description of the materials and methods is provided in the Supporting Data.

## Results

### The Non-Translatable *expATXN2* Transcript Is Neurotoxic

To confirm the toxicity of *expATXN2* transcripts, we cloned full-length (FL) *ATXN2* complementary DNA (cDNA) with 22, 58, or 104 CAG triplets into the 3' untranslated (UTR) region of *Renilla luciferase* (*Rluc*) cDNA, thereby allowing expression of *expATXN2* RNA transcripts, but preventing ATG-initiated translation of the RNA into FL *ATXN2* protein (Fig. 1A). Primers F1/R1 and F1/R2, spanning the junction of the fusion *Rluc-ATXN2* construct, were used to detect mRNA transcript that contains CAG repeat in SK-N-MC cells overexpressing *Rluc-ATXN2-(CAG)*n** (Supplementary Fig. S1). Anti-*ATXN2* Western blotting suggested no detectable expression of FL *ATXN2-Q58* or *ATXN2-Q104* protein in SK-N-MC cells overexpressing *Rluc-ATXN2-(CAG)58* or *Rluc-ATXN2-(CAG)104*, respectively (Supplementary Fig. S2). Further, no evidence of expression of an expanded polyglutamine tract was detected by Western blotting using the expanded polyglutamine-specific antibody 1C2,<sup>43</sup> confirming that the 3' UTR cloning approach indeed eliminated detectable ATG-initiated translation of the FL *ATXN2* (Fig. 1B). As an additional test of the 3' UTR approach, expression from EGFP ORF was eliminated when placed at the 3' UTR of *Rluc* ORF (Supplementary Fig. S3). Caspase 3/7 activity assay showed that overexpression of *Rluc-ATXN2-(CAG)58*



**FIG. 1.** Non-translatable full length (FL) *expATXN2* transcript is neurotoxic to SK-N-MC cells. **(A)** Schematic presentation of the non-translatable FL *ATXN2* cell model. FL *ATXN2* cDNA was cloned into 3' UTR region of *Renilla luciferase* (*Rluc*) cDNA to prevent its translation. Primer locations were indicated. **(B)** *Rluc-ATXN2-(CAG)*n** constructs do not express canonically translated polyglutamine (polyQ), as confirmed by immunoblotting with polyQ-specific 1C2 antibody.  $\beta$ -Actin was used as a loading control. A representative blot was shown. **(C)** At 72 hours after overexpression, both *Rluc-ATXN2-(CAG)58* and *Rluc-ATXN2-(CAG)104* transcripts are toxic to neuronal-like SK-N-MC cells, as determined by caspase 3/7 activity assay. The caspase 3/7 activity in *Rluc-ATXN2-(CAG)22* transfected SK-N-MC cells was normalized to 100. Data were expressed as mean  $\pm$  SEM from 4 independent samples per condition ( $n = 4$ ); \*\* $P < 0.01$  by Kruskal-Wallis test and Dunn's multiple comparison test. **(D)** Comparable expression levels of exogenous *Rluc-ATXN2-(CAG)*n** transcripts were confirmed by qPCR. *ACTB* transcript was used as an internal control. Locations of qPCR primers for *Rluc* were indicated in **(A)**. The *Rluc/ACTB* ratio in *Rluc-ATXN2-(CAG)22* transfected SK-N-MC cells was normalized to 1. Data were expressed as mean  $\pm$  SEM from 4 independent samples per condition ( $n = 4$ ); Kruskal-Wallis test. **(E)** At 48 hours after overexpression, *Rluc-ATXN2-(CAG)104* is toxic to primary mouse cortical neurons, as determined by a nuclear condensation assay. **(F)** *expATXN2* transcript toxicity depends on the repeat's ability to form toxic hairpin structures. *ATXN2-(CAG)104*, but not the interrupted *ATXN2-(CAG/CAA)105*, is toxic to primary mouse cortical neurons, as indicated by a nuclear condensation assay. Data are expressed as mean  $\pm$  SEM from 4–8 independent samples per condition ( $n = 4$ –8). In each sample, 4000 neurons per condition were analyzed. \* $P < 0.05$ , \*\* $P < 0.01$  by Kruskal-Wallis test and Dunn's multiple comparison test.

or *Rluc-ATXN2-(CAG)104* was significantly more toxic than *Rluc-ATXN2-(CAG)22* in SK-N-MC cells (Fig. 1C). Comparable expression levels of overexpressed transcripts in SK-N-MC cells were confirmed by quantitative polymerase chain reaction (qPCR) using *Rluc* primers (Fig. 1D). However, hairpin-forming expanded CAG repeats can also be translated in the absence of ATG start codon through the mechanism of repeat-associated non-ATG translation (RAN translation).<sup>44</sup> To exclude the possibility that RAN translation of protein fragments with expanded amino acid tracts leads to neurotoxicity in our SK-N-MC model system, we cloned an *ATXN2* fragment containing a CAG repeat expansion, multiple upstream stop codons, and a

total of  $\sim 150$  bp of *ATXN2* sequence flanking the repeat (thereby excluding all ATGs) into a vector with tags for each of the three open reading frames (Supplementary Fig. S4A). It was previously shown that HEK293 cells support the RAN translation of *expATXN2* transcript,<sup>45</sup> however, there were no detectable protein fragments from any of the three reading frames when overexpressed in SK-N-MC cells (Supplementary Fig. S4 B–E), indicating that SK-N-MC cells do not support the RAN translation of *expATXN2* transcripts, consistent with our previous results indicating that RAN translation is cell line-specific.<sup>28</sup> Our results, therefore, confirm that expression of FL *expATXN2* transcript is sufficient to trigger neurotoxicity in SK-N-MC

cells, even when the transcripts are not translated into proteins. Consistent with these observations in neuroblastoma cells, overexpression of *Rluc-ATXN2-(CAG)104* triggers neurotoxicity in primary mouse cortical neurons, as measured by nuclear condensation assay (Fig. 1E). *Rluc-ATXN2-(CAG)58* was not toxic in this assay, perhaps reflective of the short time frame of the experiment (nuclear condensation is a later stage event, whereas caspase 3/7 activation occurs at an early stage in cell death), differences in the levels of transcript expression in primary neurons compared to neuroblastoma cell lines, or different sensitivity to transcript-induced toxicity in primary neurons and SK-N-MC cells.

It has been suggested that the CAG repeats form stable hairpin structures,<sup>35</sup> whereas CAA interruptions either break hairpin regularity or induce the formation of branched structures.<sup>46</sup> To examine whether preventing the formation of hairpin structures in *expATXN2* ameliorate its neurotoxicity, we replaced the pure CAG repeat region in the *ATXN2-(CAG)104* with a fragment of heavily interrupted CAG/CAA triplets to obtain the *ATXN2-(CAG/CAA)105* construct. Inserting interruptions abolished *expATXN2* toxicity in primary mouse cortical neurons (Fig. 1F), suggesting that the secondary hairpin structure adopted by the pure CAG repeat may be critical for neurotoxicity.

### Full-Length *expATXN2* Transcripts Form RNA Foci in SCA2 Cell and Mouse Models and in One Human SCA2 Brain

Repeat-containing mutant transcripts form RNA foci in all CUG/CAG diseases, in which RNA neurotoxicity has been demonstrated to contribute to pathogenesis.<sup>47-49</sup> We sought to detect similar foci in SCA2 models and human brain by fluorescence in situ hybridization (FISH) using a 20-mer 2-O-methyl-CUG riboprobe that binds to the CAG repeat region on the transcript.<sup>28</sup> CAG RNA foci were absent in SK-N-MC neuroblastoma cells that overexpress *GFP* alone (Fig. 2A) or a FL *ATXN2* construct modified to have only one CAG triplet (*GFP-ATXN2Q1*, Fig. 2B) and were only rarely detected in cells overexpressing FL normal *ATXN2* (*nATXN2*) transcripts with 22 triplets (*GFP-ATXN2Q22*, Fig. 2C). Foci were much more abundant in cells overexpressing full-length (FL) expanded *ATXN2* (*expATXN2*) transcripts with 58 or 104 CAG triplets (*GFP-ATXN2Q58* or *GFP-ATXN2Q104*, Fig. 2D,F), as quantified in Figure 2E. The foci are resistant to deoxyribonuclease (DNase) treatment and are degraded by Ribonuclease (RNase) treatment (Fig. 2G,H). This set of experiments demonstrates that *expATXN2* transcripts form RNA foci, with a similar number of foci formed in cells transfected with *GFP-ATXN2Q58* and *GFP-ATXN2Q104*. Furthermore, although not detected in wild-type (WT) (Fig. 2I,J)


mice, *ATXN2* RNA foci are present in cerebellar Purkinje neurons of SCA2 transgenic mice (Fig. 2K,L), which express FL *ATXN2* with 127 CAG triplets specifically in Purkinje neurons,<sup>22</sup> whereas Purkinje cells were identified based on the size and morphology of Hoechst stained nuclei. Finally, of the five human postmortem brains available for this study, *ATXN2* RNA foci were detected in cerebellar Purkinje cells in one brain (H1 case, Table 1) that had 38 triplets for the mutant allele (Fig. 2O,P) but not in the control human brains (Fig. 2M,N.) RNA foci may be only a hallmark for RNA toxicity. Whether RNA foci are toxic or not remains to be further determined.

### *expATXN2* Transcripts Aberrantly Interact with RBPs

Next, we examined whether the neurotoxicity of *expATXN2* transcript is mediated by aberrant *expATXN2* RNA-RBP interactions. We performed an in vitro biotinylated *ATXN2* RNA pull-down assay (Fig. 3A) and identified by mass spectrometry (MS) a total of 57 RBPs that preferentially bind to the *expATXN2*, compared to the *nATXN2* transcript. Go analysis of functional annotation<sup>50</sup> and STRING analysis<sup>51</sup> of the *expATXN2* RBPs are shown in Supplementary Figure S6. The list of *expATXN2* RBPs is shown in Supplementary Table S1. Of the 57 *expATXN2* RBPs, 40 are localized in the nucleus, with 20 of them in the nucleolus, suggesting that aberrant *expATXN2*-RBP interactions may predominantly occur in the nucleus. Interestingly, among the 20 nucleolar RBPs, 7 of them contain WD40 repeat domains, of which five (PWP1, TBL3, WDR3, WDR36, and UTP18) (Supplementary Table S1 and Supplementary Fig. S6B) are components of the small subunit (SSU) processome for rRNA processing. We, therefore, became interested in the SSU processome components that were identified as *expATXN2* RBPs. Of the five SSU components<sup>52,53</sup> that are potential *expATXN2* interactors, we selected transducin  $\beta$ -like protein 3 (TBL3) for further analysis because we were interested in RNA-mediated disease mechanisms shared by both SCA2 and HD. By the same method, TBL3 appeared to interact with the expanded *Huntingtin* (*expHtt*) transcript as well (Fig. 3B,C) and has a relatively greater number of peptide hits and percentage of protein coverage compared with other SSU components identified by MS (Supplementary Table S1), although the number of peptide hits does not always imply stronger interaction.<sup>54</sup>

### TBL3 Binds to Expanded CAG Repeats *In Vitro*

We performed additional RNA pull down experiments and Western blots to confirm that TBL3 interacts with *expATXN2* *in vitro* (Fig. 3B). TBL3 was not detected when RNA pull down was performed using beads that are not loaded with transcripts (beads only,



**FIG. 2.** *expATXN2* transcripts form RNA foci. (A–H) Exogenous *expATXN2* transcripts form nuclear CAG RNA foci in SK-N-MC neuroblastoma cells. *GFP-ATXN2-(CAG)58* or *104* (D and F) transcripts formed frequent foci, whereas RNA foci were occasionally detected in *GFP-ATXN2-(CAG)22* expressing cells (C) and were absent in cells expressing GFP (A) or *GFP-ATXN2-(CAG)1* (B). (E) The percentage of cells with foci. Data were expressed as mean  $\pm$  SEM from 3 independent samples per condition ( $n = 3$ ); \* $P < 0.05$  by Kruskal–Wallis test and Dunn’s multiple comparison test, compared to *GFP-ATXN2-(CAG)22*. (G,H) *GFP-ATXN2-(CAG)104* RNA foci were resistant to DNase treatment (G) and degraded by RNase treatment (H). (I–L) *expATXN2* transcript forms RNA foci in the cerebellar Purkinje cells of SCA2 transgenic (Tg) mice in which the expression of FL *ATXN2-Q127* cDNA is driven by a Purkinje cell specific *Pcp2* promoter<sup>24</sup> (K,L). RNA foci were not detected in wildtype control mice (I,J). (M–P) *expATXN2* transcript forms RNA foci in cerebellar Purkinje cells of a human SCA2 brain (O,P), but not in human control cerebella (M,N). Scale bar, 5  $\mu$ m. Arrows point to RNA foci and asterisks indicate the Purkinje cells. [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

Fig. 3B) or when beads loaded with exogenous GFP transcripts (Supplementary Fig. S5) were used, indicating that the interaction between TBL3 and *expATXN2* transcript is specific. To test whether the interaction is disease-specific, we also included *expHTT* transcripts associated with HD, the most prevalent and most studied CAG repeat disease.<sup>27,28,41,55</sup> Studies from multiple laboratories, including ours, support the idea that RNA neurotoxicity contributes to HD.<sup>27–29,48</sup> We confirmed that TBL3 interacts *in vitro* with expanded CAG repeats flanked with either *ATXN2*- or *HTT*-specific

sequence (Fig. 3B), but not with expanded CUG repeats flanked with either antisense *ATXN2* (*ATXN2-AS*<sup>42</sup>), antisense *HTT* (*HTT-AS*; expressed on HD locus),<sup>41</sup> or junctophilin-3 (*JPH3*) flanking sequence<sup>29</sup> (Fig. 3B.) To further confirm that the interaction between TBL3 and *expATXN2* and *expHTT* was dependent on the CAG repeat, we pre-incubated *expATXN2* and *expHTT* transcripts with (CTG) 8C Morpholino (MO), which we have previously established hybridizes to CAG repeat expansions.<sup>56</sup> The pretreatment with (CTG)8C prevented TBL3 from binding to either transcript *in*


**TABLE 1** Human control and patient brain information

Disease	Case ID	ATXN2 alleles	HTT alleles	Age of death	Age of onset	Gender	PMI (hr)
Control	201	22/22	20/25	62	N/A	M	14
	219	22/22	17/18	35	N/A	F	8
	249	22/22	18/18	49	N/A	M	12
SCA2	K1	22/37	17/18	72	40	F	6
	H1	22/38	15/17	74	60	F	19
	K2	22/41	14/17	49	26	M	24
	K3	22/41	18/23	55	35	F	24
	M1	22/44	17/22	43	30	M	23
HD	791	22/22	36/43	56	38	M	10
	172	22/25	18/45	63	N/A	M	8.5
	372	22/22	15/47	62	34	M	18
	903	22/22	18/45	59	38	M	7

*in vitro* and provided further evidence that both *expATXN2*-TBL3 and *expHTT*-TBL3 interactions are dependent on the presence of an expanded CAG repeat (Fig. 3C). Taken together, these data indicate that

TBL3 binds to expanded CAG repeats independent of flanking sequence.

To investigate whether TBL3 binds to expanded CAG repeats independently of other cellular proteins,



**FIG. 3.** TBL3 aberrantly interacts with *expATXN2* and *expHTT* transcripts. **(A)** Schematic illustration of the biotinylated RNA pull-down procedure. **(B)** TBL3 interacts with biotinylated *expATXN2* (with 58 or 104 CAG repeats) and *expHTT* (with 56 or 80 CAG repeats) RNAs in an *in vitro* biotinylated RNA pull-down assay. The interaction of TBL3 is specific for expanded CAG repeats, because no binding between TBL3 and CUG repeats in *JPH3*-(CUG)55, *ATXN2-AS*-(CUG)104 or *HTT-AS*-(CUG)80 was observed. **(C)** Interaction between TBL3 and *expATXN2* or *expHTT* is dependent on the CAG repeat region. Incubation with (CTG)<sub>8</sub>C, but not control Morpholino (MO), abolished the binding of TBL3 to *ATXN2*-(CAG)104 or *HTT*-(CAG)80 transcript. SK-N-MC cell lysate was used as a positive control. n = 3 independent experiments and representative blots were shown. **(D)** Nitrocellulose filter-binding analysis of MBP-NTD-TBL3 binding to *ATXN2*-(CAG)22, 108, *ATXN2*-(CAG/CAA)105, and *ATXN2-AS*-(CUG)110 RNAs. The circles represent the mean fraction RNA bound to MBP-NTD-TBL3 in the absence or presence of competitor tRNA (w tRNA), respectively. Schematic presentation of the transcripts with various repeats and lengths of flanking regions are shown. Mean and SEM are shown; n ≥ 3 independent trials. The fits through the data are from non-linear regression analysis of the binding curves to a Scatchard plot. In the absence of competitor tRNA, the K<sub>D</sub> obtained for *ATXN2*-(CAG)22, *ATXN2*-(CAG)108, *ATXN2*-(CAG/CAA)105, and *ATXN2-AS*-(CUG)110 were 350, 420 nM, 2.1 μM, and 650 nM, respectively. [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

we purified the TBL3 N-terminal RNA binding domain as a fusion with maltose binding protein (MBP-NTD-TBL3) and measured its binding with *expATXN2* transcripts using an *in vitro* nitrocellulose filter binding assay.<sup>57</sup> The isolated TBL3 NTD associated with ATXN2 CAG RNA, with  $K_D = 350$  nM and 420 nM for ATXN2-(CAG)22 and ATXN2-(CAG)108, respectively (Fig. 3D). Although overall binding was weak, the yeast homolog of TBL3, Utp13, binds pre-rRNA as a tetramer with other UtpB complex proteins. Therefore, the weak affinity of the isolated MBP-NTD-TBL3 for *expATXN2* may be because of the absence of its normal binding partners. The *in vitro* binding reactions saturated ~20%–30% of refolded ATXN2 RNA, suggesting that a fraction of the ATXN2 RNA is unable to refold into a conformation that is competent to bind TBL3.

Despite its weak affinity for RNA, MBP-NTD-TBL3 bound ATXN2 CAG repeats more strongly than control RNAs, including the ATXN2-AS-(CUG)110 transcript ( $K_D = 650$  nM) and a CAG repeat containing CAA interruptions, ATXN2-(CAG/CAA)105 ( $K_D = 2.1$   $\mu$ M). The total saturation of the CAG repeat containing CAA interruptions was greater than that of the continuous CAG repeats, which we attribute to better refolding of the interrupted repeat-RNA. Nevertheless, tighter binding to the continuous CAG repeats raised the possibility that TBL3 recognizes the hairpin structure of CAG repeat RNA. To test this idea, the filter binding assays were also carried out in the presence of a competitor yeast transfer RNA (tRNA), which is expected to be structured under our assay conditions. The tRNA competitor abolished the interaction between MBP-NTD-TBL3 and ATXN2 or ATXN2-AS transcripts (Fig. 3D), consistent with the idea that TBL3 binding depends on the structures of ATXN2 CAG repeats.


### The Effect of TBL3 Reduction on 45S pre-rRNA Level and Processing

Depletion of UTP13, the yeast homolog of TBL3, increases the steady-state level of unprocessed 35S pre-rRNA in yeast.<sup>58</sup> We hypothesized that, although the interaction between *expATXN2* and TBL3 may not be direct and likely involves other proteins, sequestration of TBL3 in a complex that interacts with *expATXN2* may disrupt its normal function and affect rRNA maturation. We, therefore, predicted that knockdown of TBL3 in cells, mimicking its sequestration, would increase the level of unprocessed 45S pre-rRNA, the human counterpart of yeast 35S pre-rRNA. Three individual small interfering RNAs (siRNAs) were used to knock down TBL3 in HEK293T cells to minimize the possibility of alternative mechanisms of TBL3 reductions through off-target effects. Each siRNA reduced TBL3 protein level by 50%–80% at 72 hours post transfection (Fig. 4A,B). Next, we examined

45S pre-rRNA levels by qPCR using primers against the 5' external transcribed spacer<sup>59</sup> as indicated in Figure 4C. Knockdown of TBL3 in HEK293T cells using each siRNA increased steady state 45S pre-rRNA levels (normalized to *ACTB*, Fig. 4D), consistent with a previous study using stable short hairpin RNA (shRNA) transfection.<sup>60</sup> As previously reviewed,<sup>61,62</sup> a complex sequence of cleavage steps is required to release the mature RNAs (18S, 5.8S, and 28S) from the precursor 45S pre-rRNA. qPCR using primers against 18S rRNA would detect the mature 18S rRNA, unprocessed 45S pre-rRNA, as well as any intermediate rRNAs containing 18S sequence. Similarly, qPCR using primers against 28S rRNA would detect the mature 28S rRNA, the 45S pre-rRNA precursor, as well as intermediate rRNAs that contain the 28S sequence (Fig. 4C). Therefore, we used the ratios of 18S rRNA/45S pre-rRNA and 28S rRNA/45S rRNA measured by qPCR as readouts for 18S rRNA maturation and 28S rRNA maturation, respectively. Depletion of UTP13 in yeast has been previously shown to decrease 18S rRNA maturation.<sup>63,64</sup> Consistently, we found that knockdown of TBL3 in HEK293T cells decreased the ratio of 18S rRNA to 45S pre-rRNA (Fig. 4E), indicating that TBL3 may play a role in 18S rRNA maturation. In addition, 28S rRNA maturation was also decreased after TBL3 knockdown (Fig. 4F). We attempted to determine if overexpression of TBL3 has the opposite effect, however, forced expression of TBL3 by itself triggered toxicity and mislocalized the protein into nuclear aggregates (data not shown). Similarly, MBNL1, an RBP previously shown to interact with expanded CAG/CUG (*expCAG/CUG*) transcript also appeared to be toxic when overexpressed or knocked down,<sup>28</sup> suggesting that expression of certain RBPs must be tightly controlled to maintain their normal function.

### 45S pre-rRNA Level and Processing Is Altered in SCA2 and HD Postmortem Tissue

Finally, we examined the expression of 45S pre-rRNA in human postmortem SCA2 and HD cerebella. qPCR amplification suggested that there was a slight, although not statistically significant, increase of 45S pre-rRNA level (normalized to *ACTB*) in both SCA2 and HD cerebella, compared with control (Fig. 4G). The qPCR results suggested that there was a decrease in both 18S rRNA maturation and 28S rRNA maturation in SCA2 and HD cerebella, compared with the controls (Fig. 4H, I), consistent with the trend observed with the knockdown of TBL3 (Fig. 4D–F). Only the decrease in HD samples, but not in SCA2 samples, reached statistical significance, under the caveat that the relatively lower statistical power in SCA2 samples may not allow the detection of small changes. Taken together, the data support the idea that aberrant RNA-RBP interactions may affect the steady-state level and the maturation of 45S pre-rRNA in both SCA2 and HD.



**FIG. 4.** rRNA processing is affected in TBL3 knockdown cells, as well as in SCA2 and HD brains. **(A,B)** siRNAs against TBL3 (siTBL3), but not control siRNAs (siCtrl), efficiently knocked down TBL3 protein expression by 50%–80% in HEK293T cells after 72 hours. Representative blots were shown. TBL3/ $\beta$ -Actin protein expression in siCtrl-treated cells was normalized to 100. Data were expressed as mean  $\pm$  SEM from 3 independent samples per condition ( $n = 3$ ). \* $P < 0.05$  and \*\* $P < 0.01$  by Kruskal–Wallis test and Dunn’s multiple comparison test. **(C)** Locations for qPCR primers for the detection of 45S pre-rRNA, 18S rRNA and 28S rRNA. **(D–F)** TBL3 knockdown increased 45S pre-rRNA/ACTB ratio but decreased 18S rRNA/45S pre-rRNA ratio, compared to siCtrl-treated cells. 45S pre-rRNA/ACTB, 18S rRNA/45S pre-rRNA, and 28S rRNA/45S pre-rRNA ratios in siCtrl-treated cells were normalized to 1, respectively. Data were expressed as mean  $\pm$  SEM from 3 independent samples per condition ( $n = 3$ ) for **(D–F)**. \* $P < 0.05$  and \*\* $P < 0.01$  by Kruskal–Wallis test and Dunn’s multiple comparison test. **(G–I)** 45S pre-rRNA/ACTB, 18S rRNA/45S pre-rRNA, and 28S rRNA/45S pre-rRNA ratios in human control, HD, and SCA2 postmortem cerebella. 45S pre-rRNA/ACTB, 18S rRNA/45S pre-rRNA, and 28S rRNA/45S pre-rRNA ratios in Ctl group were normalized to 1, respectively. Data were expressed as mean  $\pm$  SEM from  $n = 3$  (Ctl), 4 (HD) and 5 (SCA2) individual patient cerebella samples. \* $P < 0.05$  by Kruskal–Wallis test and Dunn’s multiple comparison test.

## Discussion

We have previously shown that antisense *ATXN2-AS* transcripts contribute to SCA2 pathogenesis.<sup>42</sup> Here, we provide the first evidence that sense *expATXN2* transcripts is involved in SCA2 pathogenesis. First, in establishing its potential pathogenicity, we show that untranslatable FL *ATXN2* transcript is neurotoxic (Fig. 1). This model is not suitable to test the contribution of sense *ATXN2* transcript relative to the toxicity

of *ATXN2* protein, or antisense *ATXN2-AS* RNA,<sup>42</sup> because neither *ATXN2* protein nor *ATXN2-AS* RNA is present in this model. In the future, genome editing approaches can be used to establish human SCA2 induced pluripotent stem cells (iPSC) cell models that specifically model protein versus the RNA-triggered mechanism of pathogenesis. SCA2 iPSCs can also be subjected to transcriptome, proteome, and RNA interactome analysis to identify additional pathways that are involved in RNA-mediated aspects of SCA2



pathogenesis. Differentiation into neuronal types of greater and lesser selective vulnerability in SCA2 (eg, Purkinje cells, cortical excitatory neurons, etc.) could be used to determine cell-type vulnerability to RNA and protein mediated neurotoxicity. Next, we show that the *expATXN2* transcripts aggregate into nuclear RNA foci in SCA2 cell and transgenic mouse models, as well as in human SCA2 postmortem brain tissue. However, of five human SCA2 postmortem brains available for this study, *expATXN2* RNA foci were only detected in case H1 that had the 22/38 *ATXN2* CAG repeat lengths and the latest disease on-set (Table 1). Interestingly, we have recently characterized a transcript that is expressed in the direction antisense to *ATXN2* (*ATXN2-AS*) and contains an expanded CUG repeat.<sup>42</sup> CUG RNA foci containing this transcript were detected in SCA2 cases K3 and M1 (Table 1).<sup>42</sup> Given that SCA2 is associated with a relatively short repeat expansion, detection of foci may require a more sensitive assay. There are a number of alternative explanations for the absence of foci in the other SCA2 brains: (1) CAG RNA foci are highly toxic or appear in cells marked for early death such that Purkinje cells with foci may not have been present by the time of death; (2) CAG RNA foci are protective and associated with late onset disease and perhaps slower disease progression; (3) detectable foci were lost consequent to the process of brain collection or storage; (4) RNA foci are a byproduct of neurotoxic processes and have a neutral role in neurotoxicity; and (5) RNA foci are an epiphenomenon, present in some SCA2 brains because of an unknown genetic or environmental factor and with no relevance to disease. Recent work describing a transgenic BAC mouse model expressing expanded *C9orf72* (*expC9orf72*) and exhibiting widespread RNA foci, but lacking behavioral abnormalities and neurodegeneration, even at advanced ages, suggests that RNA foci are not sufficient to trigger toxicity in ALS.<sup>65</sup> A transgenic mouse model expressing non-translatable FL *ATXN2*, which could be tracked in live cells in real time,<sup>66</sup> might help determine the relevance of CAG RNA foci to disease pathogenesis.

Our data strongly suggest that the neurotoxicity of *expATXN2* transcript involves aberrant *expATXN2*-RBP interactions that perturb rRNA maturation. We initially focused on TBL3, a component of the SSU processome required for rRNA processing. Although our RNA pull-down assay indicates that TBL3 interacts with *expATXN2* RNA, this assay cannot be used to prove a direct interaction. Filter-binding assays showed that recombinant TBL3 NTD can weakly interact with *expATXN2* RNA, and preferentially binds the structures of the CAG repeats. (Fig. 3D). Other components of the SSU processome likely stabilize the interaction of TBL3 with the *expATXN2* RNA in the cell. The yeast homolog of TBL3, Utp13, recognizes double-stranded regions of the pre-rRNA as a heterotetramer with other Utp

proteins. Indeed, mass spectrometry analysis of *expATXN2* interactors did identify other proteins from the SSU processome in our isolated complexes (Supplementary Table S1). One interesting possibility is that the multi-dentate recognition of structured RNA by TBL3 and its binding partners, which is a normal feature of their function in pre-rRNA processing, also contributes to the toxic aggregation of CAG repeat RNAs. Future experiments will be needed to determine which proteins are most important for neuronal toxicity.

Our results indicate that a subset of RBPs bind to both expanded *ATXN2* and *HTT* transcripts. This is not surprising because it is well established that transcripts containing expanded CAG repeats form similar secondary structures *in vitro*<sup>35,67,68</sup> and, hence, at least some of the downstream effects are likely to be shared between different CAG repeat diseases. Although SCA2 primarily affects cerebellum, HD is primarily characterized by atrophy of striatum and cerebral cortex.<sup>69</sup> However, recent evidence indicates that cerebellum is also affected in HD<sup>70,71</sup> and, in fact, appears to degenerate early<sup>72</sup> and independently from the striatal atrophy.<sup>72</sup> This suggests that similar mechanism of pathogenesis may contribute to cerebellar pathology in both SCA2 and HD. Whether and to which degree mutant RNA-triggered mechanisms contribute to this pathology remain to be further determined.

Interestingly, it was previously reported that expanded *Ataxin-3* (*ATXN3*) transcripts involved in spinocerebellar ataxia type 3 (SCA3) interact with nucleolin. In SCA3, this aberrant nucleolin-*ATXN3* interaction decreases 45S pre-rRNA levels in cell and *Drosophila* models of SCA3.<sup>31</sup> Moreover, aberrant interaction between the *expC9orf72* transcripts and nucleolin may contribute to the decreased maturation of 28S, 18S, and 5.8S rRNAs from the precursor 45S pre-rRNA in ALS patients associated with CCCC GG hexamer expansion in *C9orf72* gene.<sup>59</sup> Ribosomes are ribonucleoprotein complexes that consist of rRNA and ribosomal protein components. Perturbation of rRNA processing disrupts ribosome biogenesis, which in turn compromises protein translation and causes neuronal dysfunction. As a consequence of rRNA processing failure, ribosomal proteins are not assembled into ribosome ribonucleoprotein subunits. These unincorporated ribosomal proteins accumulate in the neuronal cytosol and prevent proteasomal degradation of p53, which induces nucleolar stress and neuronal cell death.<sup>31</sup> Protein translation and nucleolar stress are therefore two possible mechanisms for further investigation. It is quite possible that a therapeutic agent that prevents aberrant RNA-RBP interactions between toxic hairpin-forming transcripts and RBPs may be at least partially effective across multiple diseases. Alternatively, similar therapies may target shared pathogenic pathways downstream of the toxic transcripts.

In summary, we provide the first evidence that the *ATXN2* transcript with an expanded repeat may contribute to SCA2 pathogenesis with similar properties to transcript-mediated toxicity in HD. The *ATXN2* transcript with an expanded CAG repeat itself, or its protein interactors, may provide valuable therapeutic targets in the future. ■

**Acknowledgments:** We acknowledge support for the statistical analysis from the National Center for Research Resources and the National Center for Advancing Translational Sciences (NCATS) of the National Institutes of Health through grant 1UL1TR001079. We thank the Johns Hopkins University School of Medicine Mass Spectrometry and Proteomics Facility for assistance with Mass Spectrometry analysis and data interpretation. We thank Dr. Laura Ranum for the kind gift of the A8(\*KKQ<sub>EXP</sub>)-3Tf1 construct. We thank Dr. Arnulf Koeppen, for providing frozen brain samples of three SCA2 patients. We thank Dr. Olga Pletnikova for providing a frozen brain sample of one SCA2 patient. We thank Dr. Shanshan Zhu for technical assistance regarding confocal imaging. We thank Kathryn A. Carson, ScM, for the advice on statistical analysis.

### Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request. ■

### References

- Pulst, S. M. in *GeneReviews*(R) (ed Adam, M. P. et al.) University of Washington, Seattle. *GeneReviews* is a registered trademark of the University of Washington, Seattle. All rights reserved, Seattle (WA). 1993.
- Geschwind DH, Perlman S, Figueroa CP, Treiman LJ, Pulst SM. The prevalence and wide clinical spectrum of the spinocerebellar ataxia type 2 trinucleotide repeat in patients with autosomal dominant cerebellar ataxia. *Am J Hum Genet* 1997;60:842–850.
- Cancel G, Dürr A, Didierjean O, et al. Molecular and clinical correlations in spinocerebellar ataxia 2: a study of 32 families. *Hum Mol Genet* 1997;6:709–715.
- Rottnek M, Riggio S, Byne W, Sano M, Margolis RL, Walker RH. Schizophrenia in a patient with spinocerebellar ataxia 2: coincidence of two disorders or a neurodegenerative disease presenting with psychosis? *Am J Psychiatry* 2008;165:964–967.
- Chen KH, Lin CH, Wu RM. Psychotic-affective symptoms and multiple system atrophy expand phenotypes of spinocerebellar ataxia type 2. *BMJ Case Rep* 2012;2012. <https://doi.org/10.1136/bcr.2011.5061>
- Riess O, Laccone FA, Gispert S, et al. SCA2 trinucleotide expansion in German SCA patients. *Neurogenetics* 1997;1:59–64.
- Lorenzetti D, Bohlega S, Zoghbi HY. The expansion of the CAG repeat in ataxin-2 is a frequent cause of autosomal dominant spinocerebellar ataxia. *Neurology* 1997;49:1009–1013.
- Moseley ML, Benzow KA, Schut LJ, et al. Incidence of dominant spinocerebellar and Friedreich triplet repeats among 361 ataxia families. *Neurology* 1998;51:1666–1671.
- Cruz-Marino T, Vázquez-Mojena Y, Velázquez-Pérez L, et al. SCA2 predictive testing in Cuba: challenging concepts and protocol evolution. *J Community Genet* 2015;6:265–273.
- Pérez LV, Cruz GS, Falcón NS, et al. Molecular epidemiology of spinocerebellar ataxias in Cuba: insights into SCA2 founder effect in Holguín. *Neurosci Lett* 2009;454:157–160.
- Orozco G, Estrada R, Perry TL, et al. Dominantly inherited olivopontocerebellar atrophy from eastern Cuba. Clinical, neuropathological, and biochemical findings. *J Neurol Sci* 1989;93:37–50.
- Rub U, del Turco D, del Tredici K, et al. Thalamic involvement in a spinocerebellar ataxia type 2 (SCA2) and a spinocerebellar ataxia type 3 (SCA3) patient, and its clinical relevance. *Brain* 2003;126:2257–2272.
- Rub U, Farrag K, Seidel K, et al. Involvement of the cholinergic basal forebrain nuclei in spinocerebellar ataxia type 2 (SCA2). *Neuropathol Appl Neurobiol* 2013;39:634–643.
- Rub U, Turco DD, Bürk K, et al. Extended pathoanatomical studies point to a consistent affection of the thalamus in spinocerebellar ataxia type 2. *Neuropathol Appl Neurobiol* 2005;31:127–140.
- Rub U, Schöls L, Paulson H, et al. Clinical features, neurogenetics and neuropathology of the polyglutamine spinocerebellar ataxias type 1, 2, 3, 6 and 7. *Prog Neurobiol* 2013;104:38–66.
- Moretti P, Blazo M, Garcia L, et al. Spinocerebellar ataxia type 2 (SCA2) presenting with ophthalmoplegia and developmental delay in infancy. *Am J Med Genet A* 2004;124A:392–396.
- Giunti P, Sabbadini G, Sweeney MG, et al. The role of the SCA2 trinucleotide repeat expansion in 89 autosomal dominant cerebellar ataxia families. Frequency, clinical and genetic correlates. *Brain* 1998;121(Pt 3):459–467.
- Pulst SM, Nechiporuk A, Nechiporuk T, et al. Moderate expansion of a normally biallelic trinucleotide repeat in spinocerebellar ataxia type 2. *Nat Genet* 1996;14:269–276.
- Elden AC, Kim HJ, Hart MP, et al. Ataxin-2 intermediate-length polyglutamine expansions are associated with increased risk for ALS. *Nature* 2010;466:1069–1075.
- Huynh DP, Yang HT, Vakharia H, Nguyen D, Pulst SM. Expansion of the polyQ repeat in ataxin-2 alters its Golgi localization, disrupts the Golgi complex and causes cell death. *Hum Mol Genet* 2003;12:1485–1496.
- Magana JJ, Velazquez-Perez L, Cisneros B. Spinocerebellar ataxia type 2: clinical presentation, molecular mechanisms, and therapeutic perspectives. *Mol Neurobiol* 2013;47:90–104.
- Hansen ST, Meera P, Otis TS, Pulst SM. Changes in Purkinje cell firing and gene expression precede behavioral pathology in a mouse model of SCA2. *Hum Mol Genet* 2013;22:271–283.
- Huynh DP, Figueroa K, Hoang N, Pulst SM. Nuclear localization or inclusion body formation of ataxin-2 are not necessary for SCA2 pathogenesis in mouse or human. *Nat Genet* 2000;26:44–50.
- Dansithong W, Paul S, Figueroa KP, et al. Ataxin-2 regulates RGS8 translation in a new BAC-SCA2 transgenic mouse model. *PLoS Genet* 2015;11:e1005182.
- Thornton CA. Myotonic dystrophy. *Neurol Clin* 2014;32:705–719.
- Petersson OJ, Aagaard L, Jensen TG, Damgaard CK. Molecular mechanisms in DM1 - a focus on foci. *Nucleic Acids Res* 2015;43:2433–2441.
- Banez-Coronel M, Porta S, Kagerbauer B, et al. A pathogenic mechanism in Huntington's disease involves small CAG-repeated RNAs with neurotoxic activity. *PLoS Genet* 2012;8:e1002481.
- Sun X, Li PP, Zhu S, et al. Nuclear retention of full-length HTT RNA is mediated by splicing factors MBNL1 and U2AF65. *Sci Rep* 2015;5:12521.
- Rudnicki DD, Holmes SE, Lin MW, et al. Huntington's disease—like 2 is associated with CUG repeat-containing RNA foci. *Ann Neurol* 2007;61:272–282.
- Tsoi H, Lau CK, Lau KF, Chan HY. Perturbation of U2AF65/NXF1-mediated RNA nuclear export enhances RNA toxicity in polyQ diseases. *Hum Mol Genet* 2011;20:3787–3797.
- Tsoi H, Lau TC, Tsang SY, Lau KF, Chan HY. CAG expansion induces nucleolar stress in polyglutamine diseases. *Proc Natl Acad Sci U S A* 2012;109:13428–13433.
- Li LB, Yu Z, Teng X, Bonini NM. RNA toxicity is a component of ataxin-3 degeneration in drosophila. *Nature* 2008;453:1107–1111.
- Moseley ML, Zu T, Ikeda Y, et al. Bidirectional expression of CUG and CAG expansion transcripts and intranuclear polyglutamine inclusions in spinocerebellar ataxia type 8. *Nat Genet* 2006;38:758–769.
- Echeverria GV, Cooper TA. RNA-binding proteins in microsatellite expansion disorders: mediators of RNA toxicity. *Brain Res* 2012;1462:100–111.
- Krzyzosiak WJ, Sobczak K, Wojciechowska M, Fiszler A, Mykowska A, Kozłowski P. Triplet repeat RNA structure and its role as pathogenic agent and therapeutic target. *Nucleic Acids Res* 2012;40:11–26.

36. Goodwin M, Swanson MS. RNA-binding protein misregulation in microsatellite expansion disorders. *Adv Exp Med Biol* 2014;825:353–388.
37. Cho DH, Thienes CP, Mahoney SE, Analau E, Filippova GN, Tapscott SJ. Antisense transcription and heterochromatin at the DMI CTG repeats are constrained by CTCF. *Mol Cell* 2005;20:483–489.
38. Wilburn B, Rudnicki DD, Zhao J, et al. An antisense CAG repeat transcript at JPH3 locus mediates expanded polyglutamine protein toxicity in Huntington's disease-like 2 mice. *Neuron* 2011;70:427–440.
39. Seixas AI, Holmes SE, Takeshima H, et al. Loss of junctophilin-3 contributes to Huntington disease-like 2 pathogenesis. *Ann Neurol* 2012;71:245–257.
40. Sopher BL, Ladd PD, Pineda VV, et al. CTCF regulates ataxin-7 expression through promotion of a convergently transcribed, antisense noncoding RNA. *Neuron* 2011;70:1071–1084.
41. Chung DW, Rudnicki DD, Yu L, Margolis RL. A natural antisense transcript at the Huntington's disease repeat locus regulates HTT expression. *Hum Mol Genet* 2011;20:3467–3477.
42. Li PP, Sun X, Xia G, et al. ATXN2-AS, a gene antisense to ATXN2, is associated with spinocerebellar ataxia type 2 and amyotrophic lateral sclerosis. *Ann Neurol* 2016;80:600–615.
43. Trotter Y, Devys D, Imbert G, et al. Cellular localization of the Huntington's disease protein and discrimination of the normal and mutated form. *Nat Genet* 1995;10:104–110.
44. Zu T, Gibbens B, Doty NS, et al. Non-ATG-initiated translation directed by microsatellite expansions. *Proc Natl Acad Sci U S A* 2011;108:260–265.
45. Scoles DR, Ho MH, Dansithong W, et al. Repeat associated non-AUG translation (RAN translation) dependent on sequence downstream of the ATXN2 CAG repeat. *PLoS One* 2015;10:e0128769.
46. Sobczak K, Krzyzosiak WJ. CAG repeats containing CAA interruptions form branched hairpin structures in spinocerebellar ataxia type 2 transcripts. *J Biol Chem* 2005;280:3898–3910.
47. Rudnicki DD, Margolis RL, Pearson CE, Krzyzosiak WJ. Diced triplets expose neurons to RISC. *PLoS Genet* 2012;8:e1002545.
48. Wojciechowska M, Krzyzosiak WJ. CAG repeat RNA as an auxiliary toxic agent in polyglutamine disorders. *RNA Biol* 2011;8:565–571.
49. Urbanek MO, Krzyzosiak WJ. RNA FISH for detecting expanded repeats in human diseases. *Methods* 2016;98:115–123.
50. Zhou Y, Zhou B, Pache L, et al. Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nat Commun* 2019;10:1523–1526.
51. Szklarczyk D, Gable AL, Lyon D, et al. STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res* 2019;47:D607–D613.
52. Wada K, Sato M, Araki N, et al. Dynamics of WD-repeat containing proteins in SSU processome components. *Biochem Cell Biol* 2014;92:191–199.
53. Hutchinson SA, Tooke-Locke E, Wang J, Tsai S, Katz T, Trede NS. Tbl3 regulates cell cycle length during zebrafish development. *Dev Biol* 2012;368:261–272.
54. Zhang Y, Xu T, Shan B, et al. ProteinInferencer: confident protein identification and multiple experiment comparison for large scale proteomics projects. *J Proteomics* 2015;129:25–32.
55. Ross CA, Tabrizi SJ. Huntington's disease: from molecular pathogenesis to clinical treatment. *Lancet Neurol* 2011;10:83–98.
56. Sun X, Marque LO, Cordner Z, et al. Phosphorodiamidate morpholino oligomers suppress mutant huntingtin expression and attenuate neurotoxicity. *Hum Mol Genet* 2014;23:6302–6317.
57. Yuan Y, Compton SA, Sobczak K, et al. Muscleblind-like 1 interacts with RNA hairpins in splicing target and pathogenic RNAs. *Nucleic Acids Res* 2007;35:5474–5486.
58. Hughes JM, Ares M. Depletion of U3 small nucleolar RNA inhibits cleavage in the 5' external transcribed spacer of yeast pre-ribosomal RNA and impairs formation of 18S ribosomal RNA. *EMBO J* 1991;10:4231–4239.
59. Haeusler AR, Donnelly CJ, Periz G, et al. C9orf72 nucleotide repeat structures initiate molecular cascades of disease. *Nature* 2014;507:195–200.
60. Wang J, Tsai S. Tbl3 encodes a WD40 nucleolar protein with regulatory roles in ribosome biogenesis. *World J Hematol* 2014;3:93–104.
61. Henras AK, Plisson-Chastang C, O'Donohue MF, Chakraborty A, Gleizes PE. An overview of pre-ribosomal RNA processing in eukaryotes. *Wiley Interdiscip Rev RNA* 2015;6:225–242.
62. Aubert M, O'Donohue MF, Lebaron S, Gleizes PE. Pre-ribosomal RNA processing in human cells: from mechanisms to congenital diseases. *Biomolecules* 2018;8:123. <https://doi.org/10.3390/biom8040123>
63. Gallagher JE, Dunbar DA, Granneman S, et al. RNA polymerase I transcription and pre-rRNA processing are linked by specific SSU processome components. *Genes Dev* 2004;18:2506–2517.
64. Albert B, Kos-Braun IC, Henras AK, et al. A ribosome assembly stress response regulates transcription to maintain proteome homeostasis. *Elife* 2019;8:e45002. <https://doi.org/10.7554/eLife.45002>
65. O'Rourke JG, Bogdanik L, Muhammad AK, et al. C9orf72 BAC transgenic mice display typical pathologic features of ALS/FTD. *Neuron* 2015;88:892–901.
66. Lionnet T, Czaplinski K, Darzacq X, et al. A transgenic mouse for in vivo detection of endogenous labeled mRNA. *Nat Methods* 2011;8:165–170.
67. Kiliszek A, Kierzek R, Krzyzosiak WJ, Rypniewski W. Atomic resolution structure of CAG RNA repeats: structural insights and implications for the trinucleotide repeat expansion diseases. *Nucleic Acids Res* 2010;38:8370–8376.
68. Tawani A, Kumar A. Structural insights reveal the dynamics of the repeating r(CAG) transcript found in Huntington's disease (HD) and spinocerebellar ataxias (SCAs). *PLoS One* 2015;10:e0131788.
69. Ross CA, Aylward EH, Wild EJ, et al. Huntington disease: natural history, biomarkers and prospects for therapeutics. *Nat Rev Neurol* 2014;10:204–216.
70. Rees EM, Farmer R, Cole JH, et al. Cerebellar abnormalities in Huntington's disease: a role in motor and psychiatric impairment? *Mov Disord* 2014;29:1648–1654.
71. Klionsky DJ, Abdel-Aziz AK, Abdelfatah S, et al. Guidelines for the use and interpretation of assays for monitoring autophagy. *Autophagy* 2012;8:445–544.
72. Rub U, Hoche F, Brunt ER, et al. Degeneration of the cerebellum in Huntington's disease (HD): possible relevance for the clinical picture and potential gateway to pathological mechanisms of the disease process. *Brain Pathol* 2013;23:165–177.

## Supporting Data

Additional Supporting Information may be found in the online version of this article at the publisher's web-site.