

Research Article

Baseline Triglyceride Level Affected the Efficacy of Vildagliptin in Treating Type 2 Diabetes: A Post Hoc Analysis of the VISION Study

Lingli Zhou , Xiaoling Cai , Yingying Luo, Fang Zhang, and Linong Ji 

Peking University People's Hospital, 11 Xizhimen Street, Xicheng District, Beijing 100044, China

Correspondence should be addressed to Linong Ji; prof_jilinong@aliyun.com

Received 25 February 2019; Accepted 5 July 2019; Published 8 August 2019

Academic Editor: Rosa Fernandes

Copyright © 2019 Lingli Zhou et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Identifying factors that may impact vildagliptin's efficacy could contribute to individualized treatment for patients with type 2 diabetes. In the current study, we aimed to assess the correlation between patient baseline triglyceride (TG) and efficacy of vildagliptin in Chinese patients with type 2 diabetes in a post hoc analysis of the VISION study. TG-based subgroup analysis was performed to evaluate baseline TG's impact on the decrease of glycated hemoglobin (HbA_{1c}) in patients receiving vildagliptin plus low-dose metformin (VLDM) vs. high-dose metformin (HDM). Additionally, multivariate linear regression was performed to assess the association between baseline TG and HbA_{1c} reduction at weeks 12 and 24 for patients receiving VLDM vs. HDM. For patients receiving VLDM, baseline TG ≤ 2.03 mmol/L was associated with significantly greater HbA_{1c} reduction vs. TG > 2.03 mmol/L at week 12, but not at week 24. Additionally, multivariate linear regression analysis revealed a significant independent association and an association short of statistical significance between patient baseline TG and the HbA_{1c}-reducing efficacy of VLDM at weeks 12 ($P < 0.001$) and 24 ($P = 0.082$), respectively, while such association was absent for HDM. Collectively, baseline TG was an independent predictive factor for the efficacy of a dipeptidyl peptidase-IV in treating type 2 diabetes during its initial use.

1. Introduction

Vildagliptin is a dipeptidyl peptidase-IV (DPP-4) inhibitor approved for the treatment of type 2 diabetes mellitus (T2DM). By inhibiting DPP-4, vildagliptin prevents the degradation of glucagon-like peptide-1 (GLP-1) and glucose-dependent insulinotropic polypeptide (GIP) and increases the level of biological active, intact plasma GLP-1, and GIP-1; as a result, it could restore or improve pancreatic α - and β -cell sensitivity to glucose and suppress glucagon release [1]. Clinical studies found that vildagliptin in combination with metformin resulted in better glycemic control than high-dose metformin alone [2–5]. Despite a better glucose control achieved by vildagliptin plus low-dose metformin (VLDM) than high-dose metformin (HDM) in our previous VISION study, it showed 47.1% of patients have not yet satisfied the recommended target glycated hemoglobin (HbA_{1c}) level of $\leq 6.5\%$ after a six-month treatment by

VLDM [5]. Identifying factors that impact the efficacy of vildagliptin could contribute to individualized treatment for T2DM.

It has been reported that nonesterified fatty acid (NEFA) treatment decreased GLP-1 receptor (GLP-1R) expression in rodent insulinoma cell lines and isolated islet and led to impaired GLP-1 signaling. Further, lowering of plasma TGs by fibrates increased the efficacy of the DPP-4 inhibitor des-fluoro-sitagliptin in *db/db* mice [6]. Interestingly, Duca et al. found that high-energy (HE)/high-fat (HF) feeding led to significant downregulation of GLP-1R expression in the vagal nodose ganglia of obesity-prone (OP) but not obesity-resistant (OR) rats and that the combination of HE/HF feeding and the OP phenotype led to reduced endogenous GLP-1 and GLP-1R activation [7]. These findings of animal studies suggested the possibility that patient baseline TG levels could impact the efficacy of incretin-based therapies. A recent study found that markers of higher insulin

resistance, including triglyceride, are consistently associated with reduced glycemic response to DPP-4 inhibitors in T2DM patients with DPP-4 inhibitor monotherapy [8]. However, it is still unknown whether the inverse relationship between triglyceride and glucose control is solely contributed to DPP-4 inhibitor treatment or due to drugs targeting on insulin resistance. Therefore, the current post hoc analysis of the VISION study was undertaken to determine whether a patient baseline TG level was associated with the HbA_{1c}-reducing effect of VLDM and HDM.

2. Methods

2.1. Design. Details of the design and method of the 24-week, phase 4, multicenter, randomized, open-label, prospective, parallel-group VISION study have been described previously [5]. Briefly, patients with T2DM inadequately controlled with metformin 1000 mg daily were divided 1:1:1:1 into four prespecified subgroups based on age and body mass index (BMI). Patients in each subgroup were randomized 5:1 to receive either vildagliptin (50 mg twice daily) plus metformin [500 mg twice daily; vildagliptin and low-dose metformin (VLDM) group] or metformin uptitration [1000 mg twice daily; high-dose metformin (HDM) group]. The primary endpoint was change in glycated hemoglobin (HbA_{1c}) from baseline at week 24. This clinical trial was designed, conducted, and reported in accordance with Good Clinical Practice (GCP) and the Declaration of Helsinki. The study protocol has also been approved by the Independent Ethics Committee (IEC) or Institutional Review Board (IRB) of each participating study center.

2.2. Post Hoc Analysis Procedures. The analysis of the VISION data was conducted on the full analysis set (FAS) population (patients who received at least one dose of study drug and had at least one primary or secondary efficacy evaluation after baseline) with last observation carried forward (LOCF). The VLDM and HDM groups were further grouped into 3 tertiles/subgroups based on their baseline TG levels: “TG < 1.28 mmol/L” (1st tertile), “1.28 mmol/L ≤ TG ≤ 2.03 mmol/L” (2nd tertile), and “TG > 2.03 mmol/L” (3rd tertile). HbA_{1c} changes from baseline (Δ HbA_{1c}) at weeks 12 and 24 served as endpoints that were assessed in all subgroups. Correlations of baseline TG and HbA_{1c} reduction at weeks 12 and 24 were also analyzed.

2.3. Statistical Analysis. The mean ± standard deviation (SD) was used to describe Δ HbA_{1c} at weeks 12 and 24. Since total cholesterol and TG did not follow a normal distribution, the values were log transferred (e.g., log (baseline total cholesterol) and log (baseline TG)). Baseline characteristics were compared among the three subgroups by the one-way analysis of variance (ANOVA). Both univariate and multivariate analyses were performed for the intersubgroup comparison of Δ HbA_{1c}. The univariate analysis used the TG subgroups as the variable and baseline HbA_{1c} as the covariate. The multivariate analysis used the TG subgroups as the variable; covariates included baseline HbA_{1c}, BMI, systolic blood pres-

sure (SBP), total cholesterol (TC), ALT, creatinine, and duration of T2DM, whether the patients took lipid-lowering medication and whether the patients took $\alpha\beta$ -blockers or β -blockers. Patients' compliance with the treatment was also used as one of the covariates.

Correlations between the patient baseline TG level and Δ HbA_{1c} at weeks 12 and 24 for the VLDM and HDM groups were further assessed using multivariate linear regression analysis (an analysis of covariance (ANCOVA)), wherein Δ HbA_{1c} at weeks 12 and 24 was a dependent variable. Independent variables included patient baseline demographics and clinical characteristics such as age, gender, baseline alanine transaminase (ALT), 120 min postprandial blood glucose (PPG), fast plasma glucose (FPG), serum creatinine, HbA_{1c}, BMI, duration of type 2 diabetes, log (baseline total cholesterol), log (baseline TG), systolic blood pressure (SBP), and diastolic blood pressure (DBP), whether the patients took lipid-lowering medication and whether the patients took $\alpha\beta$ -blockers or β -blockers. Patients' compliance with the treatment was also included as one of the independent variables.

The statistical significance was accepted with a *P* value < 0.05. Statistical analyses were performed using SAS version 9.2.

3. Results

3.1. Characteristics of Participants. Data from the FAS population that included 2,501 patients in the VLDM arm and 484 patients in the HDM arm were analyzed. Patients in the VLDM and HDM arms were grouped into 3 subgroups according to their baseline TG: “TG < 1.28 mmol/L” (1st tertile; 826 and 172 patients in VLDM and HDM, respectively), “1.28 mmol/L ≤ TG ≤ 2.03 mmol/L” (2nd tertile; 855 and 156 patients in VLDM and HDM, respectively), and “TG > 2.03 mmol/L” (3rd tertile; 819 and 155 patients in VLDM and HDM, respectively), and their baseline demographics and clinical characteristics are summarized in Table 1. For patients in both the VLDM and the HDM arms, there were baseline differences among the 3 TG-based subgroups (Table 1).

3.2. Impact of Baseline TG on HbA_{1c} Change from Baseline and Predictive Value of Baseline TG. For the VLDM group, at week 12, patients in the 1st, 2nd, and 3rd tertiles had a mean Δ HbA_{1c} of -0.55% ± 0.816%, -0.52% ± 0.754%, and -0.48% ± 0.789%, respectively, and at week 24, they had a mean Δ HbA_{1c} of -0.57% ± 0.905%, -0.55% ± 0.858%, and -0.51% ± 0.882%, respectively. Univariate analysis revealed significant differences in HbA_{1c} reduction across the 3 subgroups at weeks 12 and 24 (*P* = 0.007 and *P* = 0.025, respectively) (Table 2). Particularly, univariate analysis showed that patients in both the 1st and 2nd tertiles had significantly greater HbA_{1c} reduction than patients in the 3rd tertile at weeks 12 and 24 (*P* < 0.05 for both) (Table 2). Multivariate analysis reveals the same significant differences in HbA_{1c} reduction across the 3 subgroups at week 12 (*P* = 0.008) and that patients in both the 1st and 2nd tertiles had significantly greater HbA_{1c} reduction than patients in the 3rd tertile at week 12 (*P* < 0.05); however, at week 24, multivariate analysis failed to reveal any

TABLE 1: Patients' baseline demographics and clinical characteristics (FAS) according to the 3 TG-based tertiles.

	Vildagliptin+metformin (VLDM)				Metformin (HDM)			
	1 st tertile <i>n</i> = 826	2 nd tertile <i>n</i> = 855	3 rd tertile <i>n</i> = 919	<i>P</i> value	1 st tertile <i>n</i> = 172	2 nd tertile <i>n</i> = 156	3 rd tertile <i>n</i> = 155	<i>P</i> value
Age (years)	58 (11)	57 (11)	55 (11)	<0.0001	58 (10)	56 (12)	55 (11)	0.0555
Male, <i>n</i> (%)	462 (55.9)	419 (49.0)	479 (58.5)	<0.0001	79 (45.9)	73 (46.8)	87 (56.1)	0.1314
Weight (kg)	67.2 (11.1)	68.9 (10.8)	71.7 (11.4)	<0.0001	66.4 (10.4)	68.7 (11.4)	71.2 (12.2)	0.0007
BMI (kg/m ²)	24.4 (3.1)	25.1 (7.9)	25.7 (3.1)	<0.0001	24.6 (3.0)	25.1 (3.4)	25.6 (3.3)	0.0119
Duration of DM (years)	4.6 (4.5)	4.1 (4.2)	4.1 (3.9)	0.0369	4.0 (4.3)	3.8 (4.1)	4.4 (4.4)	0.4291
HbA _{1c} (%)	7.2 (0.9)	7.2 (0.9)	7.3 (0.9)	0.0358	7.2 (0.8)	7.2 (0.8)	7.3 (0.8)	0.5528
FPG (mmol/L)	7.4 (1.9)	7.5 (1.7)	8.0 (2.0)	<0.0001	7.3 (1.7)	7.5 (1.6)	7.8 (1.9)	0.0225
120 min PPG (mmol/L)	11.8 (3.6)	12.6 (3.3)	13.3 (3.8)	0.013	11.0 (3.5)	12.8 (3.5)	11.6 (3.8)	0.1754
TG (mmol/L)	0.95 (0.23)	1.63 (0.21)	3.44 (2.11)	<0.0001	0.97 (0.21)	1.62 (0.22)	3.36 (2.12)	<0.0001
Total cholesterol (mmol/L)	4.37 (0.87)	4.78 (0.83)	5.04 (0.93)	<0.0001	4.42 (0.90)	4.88 (0.89)	5.03 (0.98)	<0.0001
SBP (mmHg)	125 (11)	126 (11)	126 (11)	0.1743	125 (12)	126 (12)	127 (12)	0.4498
DBP (mmHg)	77 (8)	77 (7)	78 (8)	0.0030	77 (7)	77 (8)	79 (7)	0.0366
ALT (U/L)	22 (13)	26 (14)	29 (15)	<0.0001	21 (10)	25 (13)	28 (14)	<0.0001
Plasma creatinine (μmol/L)	64 (15)	65 (16)	66 (15)	0.0286	64 (16)	64 (15)	66 (17)	0.4972

All data except for the gender were in mean (standard deviation (SD)). FAS: full analysis set; VLDM: vildagliptin (50 mg bid) plus metformin (500 mg bid); HDM: metformin uptitration (1000 mg bid); 1st tertile: TG < 1.28 mmol/L; 2nd tertile: 1.28 mmol/L ≤ TG ≤ 2.03 mmol/L; 3rd tertile: TG > 2.03 mmol/L; TG: triglyceride; BMI: body mass index; FPG: fasting plasma glucose; 120 min PPG: 120-minute postprandial blood glucose; SBP: systolic blood pressure; DBP: diastolic blood pressure; ALT: alanine aminotransferase.

TABLE 2: ΔHbA_{1c} at weeks 12 and 24 for patients of the 3 TG-based tertiles in the VLDM and HDM groups (FAS LOCF).

ΔHbA _{1c}	VLDM				HDM			
	1 st tertile	2 nd tertile	3 rd tertile	<i>P</i>	1 st tertile	2 nd tertile	3 rd tertile	<i>P</i>
Week 12								
Sample size (<i>n</i>)	799	837	798	—	168	154	150	—
Mean ± SD (%)	-0.55 ± 0.82	-0.52 ± 0.75	-0.48 ± 0.79	—	-0.33 ± 1.03	-0.25 ± 0.88	-0.22 ± 0.79	—
<i>Univariate analysis</i>								
LSMean (SE) (%)	-0.56 ± 0.03	-0.53 ± 0.02	-0.45 ± 0.03* ^c	0.007	-0.35 ± 0.07	-0.25 ± 0.07	-0.20 ± 0.07	0.332
<i>Multivariate analysis</i>								
LSMean (SE) (%)	-0.57 ± 0.03	-0.52 ± 0.02	-0.45 ± 0.03* ^c	0.008	-0.34 ± 0.07	-0.25 ± 0.07	-9	0.503
Week 24								
Sample size (<i>n</i>)	808	847	804	—	169	154	148	—
Mean ± SD (%)	-0.57 ± 0.91	-0.55 ± 0.86	-0.51 ± 0.88	—	-0.49 ± 0.85	-0.39 ± 0.82	-0.31 ± 0.82	—
<i>Univariate analysis</i>								
LSMean (SE) (%)	-0.58 ± 0.03	-0.56 ± 0.03	-0.48 ± 0.03* ^c	0.025	-0.51 ± 0.06	-0.38 ± 0.06	-0.30 ± 0.07	0.064
<i>Multivariate analysis</i>								
LSMean (SE) (%)	-0.57 ± 0.03	-0.56 ± 0.03	-0.49 ± 0.03	0.113	-0.49 ± 0.06	-0.38 ± 0.06	-0.32 ± 0.07	0.184

*3rd tertile vs. 1st tertile: *P* < 0.05; ^c3rd tertile vs. 2nd tertile: *P* < 0.05. 1st tertile: TG < 1.28 mmol/L; 2nd tertile: 1.28 mmol/L ≤ TG ≤ 2.03 mmol/L; 3rd tertile: TG > 2.03 mmol/L; ΔHbA_{1c}: glycated hemoglobin (HbA_{1c}) change from baseline; VLDM: vildagliptin (50 mg bid) plus metformin (500 mg bid); HDM: metformin uptitration (1000 mg bid); FAS: full analysis set; LOCF: last observation carried forward; LSmean: least square mean. The univariate analysis used the TG subgroups as the variable and baseline HbA_{1c} as the covariate.

significant difference in HbA_{1c} reduction among the 3 TG-based groups (*P* = 0.113) (Table 2).

For patients receiving HDM, no significant difference in HbA_{1c} reduction was found among the 3 TG-based subgroups at both week 12 and week 24 (*P* > 0.05) (Table 2).

Multivariate linear regression analysis further revealed significant association and an association short of statistical significance between Log (baseline TG) and ΔHbA_{1c} at week 12 (regression coefficient: 0.1083 [95% CI: 0.0537, 0.1629]; *P* < 0.001) and week 24 (regression coefficient: 0.0551 [95%

TABLE 3: Relationship between baseline characteristics and ΔHbA_{1c} in the VLDM and HDM groups by multivariate linear regression analysis (FAS LOCF).

	VLDM		HDM	
	B (95% CI)	P value	B (95% CI)	P value
Week 12				
Log (baseline TG)	0.1083 (0.0537, 0.1629)	<0.001	0.0800 (-0.0821, 0.2422)	0.333
Plasma creatinine	-0.0035 (-0.0053, -0.0017)	<0.001	-0.0052 (-0.0104, -0.0000)	0.052
Duration of DM	0.0231 (0.0163, 0.0298)	<0.001	0.0083 (-0.0112, 0.0279)	0.403
Baseline HbA_{1c}	-0.4127 (-0.4455, -0.3799)	<0.001	-0.2582 (-0.3620, -0.1544)	<0.001
Week 24				
Log (baseline TG)	0.0551 (-0.0071, 0.1173)	0.082	0.0761 (-0.0691, 0.2212)	0.304
Plasma creatinine	-0.0036 (-0.0057, -0.0016)	<0.001	-0.0014 (-0.0061, 0.0032)	0.541
Duration of DM	0.0234 (0.0157, 0.0311)	<0.001	0.0314 (0.0139, 0.0490)	<0.001
Baseline HbA_{1c}	-0.4290 (-0.4663, -0.3917)	<0.001	-0.3325 (-0.4248, -0.2402)	<0.001

TG: triglyceride; FAS: full analysis set; LOCF: last observation carried forward; VLDM: vildagliptin (50 mg bid) plus metformin (500 mg bid); HDM: metformin uptitration (1000 mg bid); 95% CI: 95% confidence interval; HbA_{1c} : glycated hemoglobin; TC: total cholesterol.

CI: -0.0071, 0.1173]; $P = 0.082$), respectively, for patients in the VLDM group (Table 3), but not for patients in the HDM group.

4. Discussion

This post hoc analysis of the VISION study revealed that patient baseline TG was an independent predictive factor for the HbA_{1c} -lowering efficacy of VLDM at week 12, although such predictive effect diminished as the VLDM treatment continued and became insignificant at week 24. Since such association was lacking for HDM, it could be deduced that it was vildagliptin's efficacy that was affected by patients' baseline TG. Our study reported a statistically significant association between the patient baseline TG level and a DPP-4 inhibitor's efficacy in glycemic control during its early use. This finding is just agreed with previous animal and human studies [2–4, 8]. Our finding was further supported by the recently published study (Tanabe et al.) reporting that hypertriglyceridemia was an independent predictor for the efficacy of the GLP-1 analog liraglutide [9]. Our analysis further expanded their findings by TG-based subgroup analysis to examine the association between patient baseline TG and treatment effects of vildagliptin.

Importantly, we found this relationship did not exist in patients treated with metformin alone. The underlying mechanisms that the differences responded between DPP-4 inhibitor and metformin are still unknown. The differences of the pharmacological function between two drugs may partially explain the divergence. Unlike vildagliptin that prevents GLP-1 and GIP-1 degradation and increases the level of biological active, the efficacy of metformin depends on its capacity to reduce hepatic glucose production and improve metabolic parameters [10]. Duca et al. suggested that high-energy/high-fat feeding combined with an obesity-prone phenotype led to reduced endogenous GLP-1 and GLPR activation [7], suggesting that a high circulation TG level may decrease endogenous GLP-1R activation, therefore affecting the treatment efficacy of the DPP-4 inhibitor.

That lowering of the TG level by fibrates which improved the efficacy of the DPP-4 inhibitor in a *db/db* mouse model verified this hypothesis [6]. A further intervention clinical study should be performed.

We speculated that for those patients with $\text{TG} > 2.03$ mmol/L, adding a lipid-lowering medication to the vildagliptin or other DPP-4 inhibitor-containing antidiabetic regimen may improve treatment efficacy at least during their early treatment period. The current American Diabetes Association guideline on standards in medical care in diabetes recommends lifestyle change and dietary for diabetic patients with hypertriglyceridemia, evaluation for the second cause, and considering medical therapy for patients with $\text{TG} > 5.7$ mmol/L and immediate pharmacological therapy only for patients with severe hypertriglyceridemia ($\text{TG} > 11.4$ mmol/L) to reduce the risk of pancreatitis [11]. However, our results suggested that for a patient taking vildagliptin or possibly other DPP-4 inhibitors with $\text{TG} > 2.03$ mmol/L, lipid-lowering pharmacological intervention during the early use of these DPP-4 inhibitors might be beneficial. More studies are needed to explore this position and also to validate the clinical relevance of our results in view of the fact that the difference in ΔHbA_{1c} between the 1st and the 3rd TG-based tertiles was small although the fact that both our multivariate analysis based on the 3 TG subgroups and our multivariate linear regression analysis produced consistent results suggested the robustness of our findings.

Our analysis had certain limitations. Our sample size for HDM was relatively small, and there is the possibility of insufficient statistic power to detect any significant association between baseline TG and HDM's efficacy especially in view of our results that both VLDM and HDM led to the greatest HbA_{1c} reduction in the 1st TG-based tertile and the smallest reduction in the 3 tertile (Table 2). Another limitation was the relatively short treatment duration of the VISION study; therefore, we cannot ascertain the impact of baseline TG on the HbA_{1c} -reducing effect of vildagliptin beyond the 24-week treatment period. A prespecified,

properly randomized, large-scale, longer-term study is currently being planned. Additionally, the present study demonstrated that the impact of baseline TG on treatment effects of vildagliptin decreased as the vildagliptin use continued. Possible reason(s) for such change over time is currently unclear. Nevertheless, our preliminary finding provided a new evidence of a precision approach for the DPP-4 inhibitor in treating T2DM.

In summary, our results suggested that the patient baseline TG level affected and was an independent predictive factor for the efficacy in glycemic control of vildagliptin, not metformin, for patients with T2DM.

In conclusion, our analysis is the first to report that the patient baseline TG level was an independent predictive factor for the glycemic control effect of a DPP-4 inhibitor, but not metformin, in patients with T2DM during its initial use.

Data Availability

The previously reported VISION data were used to support this study and are available at DOI 10.1111/dom.12667. These datasets are available from the corresponding author upon request.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgments

We would like to thank all the investigators, study teams, and patients for their participation in this study. Particularly, we would like to thank all the members of the VISION Study Group for their participation and generous support. Statistical analyses were provided by Hejing Hong, Applied mathematics Master of Fountain Medical Development Ltd.

References

- [1] J. J. Neumiller, L. Wood, and R. K. Campbell, "Dipeptidyl peptidase-4 inhibitors for the treatment of type 2 diabetes mellitus," *Pharmacotherapy*, vol. 30, no. 5, pp. 463–484, 2010.
- [2] E. Bosi, F. Dotta, Y. Jia, and M. Goodman, "Vildagliptin plus metformin combination therapy provides superior glycaemic control to individual monotherapy in treatment-naive patients with type 2 diabetes mellitus," *Diabetes, Obesity and Metabolism*, vol. 11, no. 5, pp. 506–515, 2009.
- [3] G. Derosa, P. D. Ragonesi, A. Carbone et al., "RETRACTED: evaluation of the positive effects on insulin-resistance and β -cell measurements of vildagliptin in addition to metformin in type 2 diabetic patients," *Pharmacological Research*, vol. 73, pp. 20–26, 2013.
- [4] C. Filozof, S. Schwartz, and J. E. Foley, "Effect of vildagliptin as add-on therapy to a low-dose metformin," *World Journal of Diabetes*, vol. 1, no. 1, pp. 19–26, 2010.
- [5] L. N. Ji, C. Y. Pan, J. M. Lu et al., "Efficacy and safety of combination therapy with vildagliptin and metformin versus metformin uptitration in Chinese patients with type 2 diabetes inadequately controlled with metformin monotherapy: a randomized, open-label, prospective study (VISION)," *Diabetes, Obesity & Metabolism*, vol. 18, no. 8, pp. 775–782, 2016.
- [6] Z. F. Kang, Y. Deng, Y. Zhou et al., "Pharmacological reduction of NEFA restores the efficacy of incretin-based therapies through GLP-1 receptor signalling in the beta cell in mouse models of diabetes," *Diabetologia*, vol. 56, no. 2, pp. 423–433, 2013.
- [7] F. A. Duca, Y. Sakar, and M. Covasa, "Combination of obesity and high-fat feeding diminishes sensitivity to GLP-1R agonist exendin-4," *Diabetes*, vol. 62, no. 7, pp. 2410–2415, 2013.
- [8] J. M. Dennis, B. M. Shields, A. V. Hill et al., "Precision medicine in type 2 diabetes: clinical markers of insulin resistance are associated with altered short- and long-term glycemic response to DPP-4 inhibitor therapy," *Diabetes Care*, vol. 41, no. 4, pp. 705–712, 2018.
- [9] A. Tanabe, H. Kaneto, S. Kamei et al., "Clinical effects of liraglutide are possibly influenced by hypertriglyceridemia and remaining pancreatic β -cell function in subjects with type 2 diabetes mellitus," *Journal of Diabetes and its Complications*, vol. 30, no. 6, pp. 1201–1203, 2016.
- [10] E. Guarino, L. Nigi, A. Patti, C. Fondelli, and F. Dotta, "Combination therapy with metformin plus vildagliptin in type 2 diabetes mellitus," *Expert Opinion on Pharmacotherapy*, vol. 13, no. 9, pp. 1377–1384, 2012.
- [11] American Diabetes Association, "Cardiovascular disease and risk management: standards of medical care in diabetes—2019," *Diabetes Care*, vol. 42, Supplement 1, pp. S103–S123, 2019.