

RESEARCH**Open Access**

A randomized, controlled trial to evaluate the effect of an anti-interleukin-9 monoclonal antibody in adults with uncontrolled asthma

Chad K Oh^{1,3}, Richard Leigh², Kimmie K McLaurin¹, Keunpyo Kim¹, Micki Hultquist^{1*} and Nestor A Molfino^{1,3}

Abstract

Background: Preclinical studies suggest that interleukin-9 may be a central mediator in the development and maintenance of airway inflammation in asthma. The aim of this study was therefore to evaluate the effects of MEDI-528, an anti-interleukin-9 monoclonal antibody, in adults with confirmed uncontrolled moderate-to-severe asthma.

Methods: In this prospective double-blind, multicenter, parallel-group study, 329 subjects were randomized (1:1:1:1) to subcutaneous placebo or MEDI-528 (30, 100, 300 mg) every 2 weeks for 24 weeks, in addition to their usual asthma medications. The primary endpoint was change in mean Asthma Control Questionnaire-6 (ACQ-6) score at week 13. Secondary endpoints included weighted asthma exacerbation rates and pre-bronchodilator forced expiratory volume in 1 second (FEV₁) at weeks 13 and 25, as well as Asthma Quality of Life Questionnaire scores at weeks 12 and 25 and the safety of MEDI-528 throughout the study period. The primary endpoint was analyzed using analysis of covariance.

Results: The study population (n = 327) was predominantly female (69%) with a mean age of 43 years (range 18–65). The mean (SD) baseline ACQ-6 score for placebo (n = 82) and combined MEDI-528 (n = 245) was 2.8 (0.7) and 2.8 (0.8); FEV₁ % predicted was 70.7% (15.9) and 71.5% (16.7). Mean (SD) change from baseline to week 13 in ACQ-6 scores for placebo vs combined MEDI-528 groups was -1.2 (1.0) vs -1.2 (1.1) (p = 0.86). Asthma exacerbation rates (95% CI) at week 25 for placebo vs MEDI-528 were 0.58 (0.36–0.88) vs 0.49 (0.37–0.64) exacerbations/subject/year (p = 0.52). No significant improvements in FEV₁ % predicted were observed between the placebo and MEDI-528 groups. Adverse events were comparable for placebo (82.9%) and MEDI-528 groups (30 mg, 76.5%; 100 mg, 81.9%; 300 mg, 85.2%). The most frequent were asthma (placebo vs MEDI-528, 30.5% vs 33.5%), upper respiratory tract infection (14.6% vs 17.1%), and headache (9.8% vs 9.8%).

Conclusions: The addition of MEDI-528 to existing asthma controller medications was not associated with any improvement in ACQ-6 scores, asthma exacerbation rates, or FEV₁ values, nor was it associated with any major safety concerns.

Trial registration: ClinicalTrials.gov: NCT00968669.

Keywords: Asthma, Anti-IL-9 monoclonal antibody, MEDI-528

Background

Asthma is a chronic inflammatory airway disease characterized by variable airflow limitation, airway hyperresponsiveness, mucus hypersecretion, and structural changes to the airways that include proliferation of smooth muscle [1]. Interleukin (IL)-9, a multifunctional cytokine produced

by type 2 T helper cells (Th2), lymphocytes, and mast cells, is proposed to be a central mediator in the pathogenesis of asthma. Interest in IL-9 was first triggered by genetic linkage studies [2] and supported by the finding that expression of IL-9 and its associated receptor is higher in the airways of subjects with asthma compared with healthy controls [3,4]. Further evidence has come from *in vitro* studies showing that IL-9 enhances the growth and/or activity of a variety of cell types and pro-

* Correspondence: hultqustm@medimmune.com

¹MedImmune, One MedImmune Way, Gaithersburg, MD, USA

Full list of author information is available at the end of the article

inflammatory and pro-fibrotic mediators that are implicated in the pathogenesis of asthma [5]. IL-9 induced the release of Th2-associated chemokines in cultured human airway smooth muscle cells [6] and enhanced the stem cell factor-dependent growth of human mast cell progenitors, particularly those from children with asthma [7]. Another study reported that IL-9 upregulated mucin expression in human lung cells, suggesting that it is involved in the control of mucus secretion [8].

Preclinical studies in animal models of asthma support a contributing role for an IL-9 mast cell axis in the immunopathology of asthma [5]. In one study, anti-IL-9 antibody treatment had a protective effect against airway remodeling in mouse models of airway inflammation, together with a concomitant reduction in the number and activation of mature mast cells [9]. Furthermore, impaired lung function related to airway remodeling was reversed by IL-9 neutralization. Taken together, evidence from these and other experimental studies indicated that targeting IL-9 may offer a novel approach to the treatment of asthma.

MEDI-528 is a humanized immunoglobulin G1 monoclonal antibody that binds to IL-9 [10], and hence reduces the activity of a variety of cell types implicated in asthma pathogenesis [5]. Preliminary clinical studies in healthy adults and subjects with mild or mild-to-moderate asthma have demonstrated that MEDI-528 administered subcutaneously has linear pharmacokinetics and an acceptable safety profile with no reports of serious adverse events (SAEs) [11,12]. One study in subjects with mild asthma showed that MEDI-528 had no effect on pulmonary function or the use of rescue medication, although there were positive trends for improvement in asthma symptom scores and for a reduction in the number of asthma exacerbations [11]. Another study in a small number of subjects with mild-to-moderate asthma suggested that MEDI-528 may decrease sensitivity to exercise-induced bronchoconstriction that is dependent on mast cell degranulation [11]. We now report the results from a larger study (ClinicalTrials.gov, NCT00968669) that was designed to further investigate the potential clinical benefits of MEDI-528 in treating subjects with asthma. The primary objective of this study was to evaluate the effect of multiple dose subcutaneous administration of MEDI-528 on symptom control in adults with uncontrolled, moderate-to-severe, persistent asthma.

Methods

Subjects

Eligible subjects were aged 18–65 years with body mass index 18–35 kg/m² and a clinical diagnosis of asthma, confirmed by pre-bronchodilator forced expiratory volume in 1 second (FEV₁) ≥ 40% predicted and post-bronchodilator FEV₁ reversibility ≥ 12% and ≥ 200 mL.


Subjects were required to have poor asthma symptom control [1], as demonstrated by an Asthma Control Questionnaire-6 (ACQ-6) score ≥ 1.5 [13], daytime symptoms on ≥ 2 days/week, night-time awakening ≥ 1 night/week, rescue medication use on ≥ 2 days/week, and ≥ 1 asthma exacerbation in the past year. Subjects were either currently taking medium to high-dose inhaled corticosteroids (ICS) or were eligible to take them based on Expert Panel Report 3 guidelines, and were started on medium to high-dose ICS at the start of the run-in phase of the study [1]. Key exclusion criteria were the presence of other lung diseases, severe depression or history of suicidal behavior, tobacco smoking (≥ 10 pack-years), recent illness or infection, or clinically significant electrocardiogram (ECG) abnormalities. Treatment with any other biologic agent or immunosuppressive medication (apart from ≤ 10 mg/day oral prednisone or equivalent, or inhaled or topical corticosteroids) was not allowed during the month prior to and throughout the study. Subjects were permitted to receive a short burst of systemic corticosteroid therapy or be hospitalized to control an asthma exacerbation.

All subjects provided written informed consent before participating in the study. The protocol was approved by an Independent Ethics Committee at each study center, and the study was conducted in accordance with International Conference on Harmonisation Guidance for Good Clinical Practice and the Declaration of Helsinki.

Study design and randomization

This phase 2b, randomized, double-blind, placebo-controlled, parallel group study was conducted at 53 sites in North America, Central America, South America, and Asia between October 2009 and November 2011. The study consisted of a 4-week screening period, a 13-week steroid stable treatment period, an 11-week steroid reduction treatment period, and a 22-week follow-up period (Figure 1). The ICS dosage was optimized and stabilized during the screening period, maintained unchanged throughout the steroid-stable treatment period, and was reduced in subjects with clinically stable asthma during the steroid-reduction treatment period. Subjects were seen at the clinic twice during the screening period (weeks -4 and -2), at two-weekly intervals during the treatment period (weeks 0 to 24) with an additional visit at week 13, and then at four-weekly intervals during the post-treatment follow-up period (weeks 25 to 46).

Subjects were randomized (1:1:1:1) using an interactive voice response system to receive placebo or one of three doses of MEDI-528 (30, 100, or 300 mg) subcutaneously every 2 weeks for 24 weeks (13 doses), in addition to their usual asthma medications. Because MEDI-528 and placebo are visually distinct, study medication was administered at the clinic by unblinded personnel who were



not otherwise involved in the study in order to maintain blinding.

Randomization was stratified by asthma status (atopic or non-atopic) and ICS dose (medium or high). High-dose ICS was defined as 1 puff twice a day of fluticasone/salmeterol (500 µg/50 µg) and medium dose ICS was defined as 1 puff twice a day fluticasone/sal meterol (250 µg/50 µg) or 2 puffs twice a day budesonide/formoterol (160 µg/4.5 µg). Only subjects who were taking medium- to high-dose ICS at screening were enrolled. Subjects who were uncontrolled while receiving medium-dose ICS were enrolled to the medium-dose or high-dose ICS group at the study physician's discretion depending on their level of asthma control, and subjects receiving high-dose ICS were enrolled to the high-ICS group. Randomization of subjects with non-atopic asthma was restricted to ensure that a minimum of 160 subjects with atopic asthma were included in the study.

Primary and secondary outcomes

The primary efficacy outcome was change from baseline in mean ACQ-6 score at week 13 among individual MEDI-528 treatment groups and placebo. Secondary outcomes included change from baseline in mean ACQ-6 score at week 25, asthma exacerbation rates (week 25), pre-bronchodilator FEV₁ (weeks 13 and 25), health-related quality of life (Asthma Quality of Life Questionnaire (Standardized version) [AQLQ(S)] scores; weeks 12 and 25), and the safety of MEDI-528 throughout the study period.

Assessments

Clinical activity

The ACQ-6 was completed at home on a weekly basis, using an electronic patient-reported outcome device. The occurrence of any asthma exacerbations during the study was recorded by the study physician at each clinic visit. Asthma exacerbation was defined as a progressive

increase in asthma symptoms (cough, wheeze, chest tightness, and/or shortness of breath) with a reduction from baseline or best previous measurement of ≥ 20% in FEV₁ or peak expiratory flow (PEF) rate that did not resolve after the initiation of rescue medication and resulted in a prescription for, or administration of, systemic corticosteroid burst therapy. An exacerbation was considered to be resolved when the subject's asthma symptoms had diminished, and PEF or FEV₁ returned to > 80% of baseline for ≥ 7 days after completion of systemic corticosteroid burst therapy. Pre-bronchodilator spirometry was conducted according to the American Thoracic Society/European Respiratory Society guidelines [14]. All spirometry assessments for each subject were performed at the same time of day (± 1 hour) and in the same manner. Between three and eight forced expiratory efforts were made at each session and the maximum FEV₁ of the two best efforts was recorded.

Health-related quality of life

The AQLQ(S) was completed every 4 weeks at home using the electronic patient-reported outcome device.

Safety profile

Details of any adverse events (AEs) and SAEs were recorded. AEs were defined as any untoward medical occurrence that occurred after initiation of study medication. Blood and urine samples were collected at screening, prior to the first dose, at 4-weekly intervals during the treatment period, and 8-weekly intervals during the follow-up period for routine clinical laboratory evaluations. Vital signs (blood pressure, heart rate, temperature, and respiratory rate) were monitored at every clinic visit. A computerized 12-lead ECG was recorded. Blood samples were collected for determination of the presence of anti-drug antibodies (ADAs) using an enzyme-linked immunosorbent or electrochemiluminescent assay [12].

Statistical methods

Sample size calculations were based on an assumed change from baseline to week 13 in mean ACQ-6 score of -0.4 to -0.5 for placebo and -0.9 to -1.1 for the MEDI-528 groups, with a common standard deviation (SD) of 0.8 to 1.0 [15,16]. Sixty subjects in each treatment group would allow detection of a statistically significant difference between treatment groups with 58–99% power based on a 2-sided type I error of 0.05 and 0.10. Assuming a drop-out rate of 20%, it was planned to randomize 80 subjects into each treatment group.

The intent-to-treat (ITT) population was defined as all randomized subjects who received either placebo or MEDI-528. The safety population included all subjects who received either placebo or MEDI-528 and had safety data. Statistical analysis of clinical activity was based on a comparison of placebo with the individual MEDI-528 treatment groups for the ITT population.

Change from baseline (week 0) in mean ACQ-6 score at weeks 13 (primary endpoint) and 25 was analyzed using an analysis of covariance model. The model included the treatment group, atopic asthma status, geographic region, baseline eosinophil level, and mean ACQ score at baseline. Any missing ACQ-6 scores at weeks 13 and 25 were imputed using the last observation carried forward method. A sensitivity analysis was performed after excluding subjects with missing ACQ-6 scores at weeks 13 and 25. The proportion of subjects with an improvement in mean ACQ-6 score of ≥ 0.5 at weeks 13 and 25 was compared between placebo and MEDI-528 using Fisher's exact test. Subgroup analyses were conducted on the primary efficacy outcome by atopic asthma status (atopic asthma, non-atopic asthma) and baseline use of

ICS (medium dose, high dose). A subgroup analysis was conducted on the primary efficacy outcome by peripheral blood eosinophil count $< 0.3 \times 10^3/\mu\text{L}$ or $\geq 0.3 \times 10^3/\mu\text{L}$.

Weighted asthma exacerbation rate and the 95% confidence interval (CI) were calculated as the total number of exacerbations/total duration of person-year follow-up. The weighted asthma exacerbation rate at week 25 was analyzed using a pair-wise Poisson regression. Change from baseline in pre-bronchodilator FEV₁ at week 25 was compared between placebo and MEDI-528 using a pair-wise t-test. The proportion of subjects with an improvement in overall AQLQ(S) score of ≥ 0.5 at weeks 12 and 25 was compared between placebo and MEDI-528 using Fisher's exact test. Safety variables were analyzed descriptively based on the safety population.

Results

Subjects

Of the 329 randomized subjects, 327 received either placebo or MEDI-528 and were included in the ITT and safety populations. Overall, 64/82 (78.0%) of the placebo-treated and 207/245 (84.5%) of the MEDI-528 treated subjects completed the study. The most common reason for discontinuation was withdrawal of consent (Figure 2).

Demographic characteristics were generally balanced between the groups (Table 1). The study population was predominantly female (69%) with a mean age of 43.0 years (range 18–65). Most subjects (84.4%) had atopic asthma and 50.3% had evidence of a peripheral blood eosinophilia (eosinophil count $\geq 0.3 \times 10^3/\mu\text{L}$). Mean baseline ACQ-6 scores (2.8 for both placebo and combined MEDI-528 groups) indicated uncontrolled




Figure 2 Subject disposition.

Table 1 Demographic and baseline asthma characteristics (ITT Population)

Variable	MEDI-528					<i>P</i> value ^a
	Placebo (n = 82)	30 mg (n = 81)	100 mg (n = 83)	300 mg (n = 81)	Combined (n = 245)	
Age, years, mean (SD)	43.6 (11.6)	41.8 (11.1)	45.1 (11.6)	41.5 (12.3)	42.8 (11.7)	0.16
Male, n (%)	29 (35.4)	26 (32.1)	21 (25.3)	26 (32.1)	73 (29.8)	0.56
Race, n (%)						0.62
White	42 (51.2)	34 (42.0)	33 (39.8)	40 (49.4)	107 (43.7)	
Asian	18 (22.0)	17 (21.0)	25 (30.1)	18 (22.2)	60 (24.5)	
American Indian or Alaskan Native	5 (6.1)	8 (9.9)	8 (9.6)	4 (4.9)	20 (8.2)	
Black or African American	2 (2.4)	3 (3.7)	4 (4.8)	5 (6.2)	12 (4.9)	
Mixed race	1 (1.2)	0 (0.0)	0 (0.0)	2 (2.5)	2 (0.8)	
Other	14 (17.1)	19 (23.5)	13 (15.7)	12 (14.8)	44 (18.0)	
BMI, kg/m ² , mean (SD)	27.6 (4.3)	27.1 (4.4)	26.9 (4.3)	27.2 (4.4)	27.1 (4.3)	0.79
FEV ₁ (L)	2.21 (0.74)	2.22 (0.71)	2.05 (0.62)	2.27 (0.66) ^b	2.18 (0.67) ^c	0.19
FEV ₁ , % predicted	70.7 (15.9)	71.8 (17.6)	70.6 (16.6)	72.2 (15.9) ^b	71.5 (16.7) ^c	0.90
FVC, % predicted	83.9 (14.6)	86.1 (16.7)	84.8 (14.3)	84.7 (14.8)	85.2 (15.2)	0.83
FEV ₁ /FVC	68.0 (10.4)	67.6 (10.8)	67.0 (11.3)	69.4 (10.8) ^b	68.0 (11.0) ^c	0.57
FEV ₁ reversibility, %, mean (SD)	25.2 (13.8) ^b	23.9 (10.2) ^d	25.6 (15.3) ^d	21.8 (11.1) ^e	23.8 (12.5) ^f	0.25
Atopic asthma, n (%)	70 (85.4)	69 (85.2)	69 (83.1)	68 (84.0)	206 (84.1)	0.98
Number of exacerbations requiring oral steroids in past year, n (%) ^g						0.33
0	2 (2.4)	5 (6.2)	1 (1.2)	4 (4.9)	10 (4.1)	
1	45 (54.9)	47 (58.0)	42 (51.2)	39 (48.1)	128 (52.5)	
2	16 (19.5)	12 (14.8)	16 (19.5)	12 (14.8)	40 (16.4)	
3	5 (6.1)	5 (6.2)	10 (12.2)	15 (18.5)	30 (12.3)	
4+	14 (17.1)	12 (14.8)	13 (15.7)	11 (13.6)	36 (14.7)	
ICS use, n (%)						0.81
Medium dose	43 (52.4)	38 (46.9)	38 (45.8)	41 (50.6)	117 (47.8)	
High dose	39 (47.6)	43 (53.1)	45 (54.2)	40 (49.4)	128 (52.2)	
Rescue medication use, puffs per day, mean (SD)	4.3 (4.0)	4.9 (5.6)	4.5 (4.5)	5.9 (7.3)	5.1 (5.9)	0.25
Mean ACQ-6 score, mean (SD)	2.8 (0.7)	2.7 (0.9)	2.8 (0.7)	2.9 (0.9)	2.8 (0.8)	0.62
Overall AQLQ(S) score, mean (SD)	3.6 (0.8) ^d	3.8 (1.1) ^h	3.8 (0.9) ⁱ	3.8 (0.9) ^h	3.8 (0.9) ^j	0.60
Eosinophil count $\geq 0.3 \times 10^3/\mu\text{L}$, n (%)	44 (53.7)	38 (47.5) ^b	43 (51.8)	39 (48.1)	120 (49.2) ^c	0.84
Eosinophil count ($10^3/\mu\text{L}$)						0.47
Mean	0.40	0.34 ^b	0.40	0.35	0.36 ^c	
Median	0.31	0.28 ^b	0.37	0.29	0.29 ^c	
Basophil count ($10^3/\mu\text{L}$)						0.05
Mean	0.03	0.03 ^b	0.03	0.03	0.03 ^c	
Median	0.03	0.03 ^b	0.02	0.03	0.02 ^c	

Abbreviations: ACQ-6 Asthma Control Questionnaire, AQLQ Asthma Quality of Life Questionnaire, BMI Body mass index, FEV₁ Forced expiratory volume in 1 second, FVC Forced vital capacity, ICS Inhaled corticosteroids, ITT Intent-to-treat.

^aP value for differences among groups (placebo and MEDI-528 groups) by analysis of variance for continuous variables and the chi-square test for categorical variables.

^bn = 80; ^cn = 244; ^dn=79; ^en=78; ^fn=236; ^gOne value is missing in the MEDI-528 100 mg group; ^hn = 76; ⁱn = 81; ^jn = 233.

asthma, although 4 subjects (placebo, n = 1; 30 mg MEDI-528, n = 3) had a mean ACQ-6 score < 1.5 suggesting that their asthma was more controlled. Almost all subjects had experienced ≥ 1 exacerbation requiring oral steroids in the past year (placebo 80/82, 97.6%; MEDI-528 234/244, 95.9%). At baseline, an average of 2.06 exacerbations/

subject required oral steroids during the previous year. Subjects in the placebo group appeared to have less severe asthma than those in the MEDI-528 groups, according to the proportion taking high-dose ICS (47.6% vs 52.2%) and rescue medication use (mean number of puffs per day 4.3 vs 5.1), although there were no statistically significant

differences between placebo and any of the three MEDI-528 groups.

Clinical activity

Mean ACQ-6 score

The primary outcome measure analyzed using an analysis of covariance model, change from baseline to week 13 in mean ACQ scores, was not significantly different between the placebo and the individual MEDI-528 groups ($p = 0.80$). Improvements in mean ACQ-6 score occurred in all groups at weeks 13 and 25, with little difference between placebo and the individual MEDI-528 treatment groups (Figure 3). Clinically relevant improvements of ≥ 0.5 in mean ACQ-6 score occurred in similar proportions of placebo and combined MEDI-528 subjects at week 13 (64/82 [78.0%] vs 184/245 [75.1%], $p = 0.66$) and week 25 (59/82 [72.0%] vs 194/245 [79.2%], $p = 0.22$). In the individual MEDI-528 groups, a clinically relevant improvement was seen in the following proportions of subjects at week 13: 30 mg, 54/81 (66.7%); 100 mg, 66/83 (79.5%); and 300 mg, 64/81 (79.0%); and at week 25: 30 mg, 61/81 (75.3%); 100 mg, 67/83 (80.7%); and 300 mg, 66/81 (81.5%).

Differences between placebo and the combined MEDI-528 groups were not significantly different when analyzed by atopic or non-atopic asthma, baseline medium- or high-dose ICS use, or peripheral blood eosinophil count < 0.3 or $\geq 0.3 \times 10^3/\mu\text{L}$ (Table 2). In addition, there was no difference between the steroid-stable and steroid reduction phases with regard to the effect of ICS and systemic steroids (data not shown).

Asthma exacerbation rate

At week 25, the weighted asthma exacerbation rate (exacerbations/subject/year, 95% CI) for the placebo group was 0.58 (0.36 – 0.88) compared with 0.49 (0.37 – 0.64) in the combined MEDI-528 groups ($p = 0.52$). Results were similar for the individual MEDI-528 dose groups (Figure 4); weighted asthma exacerbation rates (95% CI) at week 25 were 0.64 (0.40 – 0.96) ($p = 0.76$) for 30 mg, 0.28 (0.14 – 0.50) ($p = 0.05$) for 100 mg, and 0.58 (0.36 – 0.88) ($p = 1.00$) for 300 mg.

Pre-bronchodilator FEV₁

Mean increase from baseline in pre-bronchodilator FEV₁ was similar between the placebo and MEDI-528 groups at weeks 13 and 25 (Figure 5). The FEV₁ % predicted mean (SD) change from baseline at week 25 for placebo vs the combined MEDI-528 groups was 1.44% (11.08) vs 1.12% (9.81) ($p = 0.82$); mean (SD) change in FEV₁ absolute values were 0.04 L (0.37) vs 0.03 L (0.30) ($p = 0.82$). In the individual MEDI-528 dose groups, mean (SD) change from baseline at week 25 was -0.01 L (0.32) ($p = 0.41$) for 30 mg, 0.06 L (0.30) ($p = 0.85$) for 100 mg, and 0.05 L (0.29) ($p = 0.97$) for 300 mg.

Health-related quality of life

A clinically relevant improvement of ≥ 0.5 in overall AQ LQ(S) score was seen in similar proportions of placebo and MEDI-528 subjects at week 12 (53/69, 76.8% vs 144/207, 69.6%; $p = 0.21$) and week 25 (50/62, 80.6% vs 136/182, 74.7%; $p = 0.57$).




Table 2 ACQ score change from baseline to week 13 (ITT population)

Variable	Placebo		Combined MEDI-528		P Value ^b
	n	Mean (SD)	n	Mean (SD)	
All subjects ^a	82	-1.23 (0.95)	245	-1.22 (1.07)	0.86
Asthma type ^a					
Atopic	70	-1.28 (0.97)	206	-1.19 (1.04)	0.55
Non-atopic	12	-0.94 (0.82)	39	-1.42 (1.18)	0.41
ICS use ^a					
Medium dose	43	-1.21 (0.97)	117	-1.17 (1.02)	0.55
High dose	39	-1.26 (0.94)	128	-1.28 (1.11)	0.77
Eosinophil count ^a					
≥ 0.3 × 10 ³ /μL	44	-1.39 (0.95)	120	-1.17 (1.01)	0.14
< 0.3 × 10 ³ /μL	38	-1.04 (0.92)	124	-1.26 (1.09)	0.22
All subjects without imputation	67	-1.30 (0.94)	208	-1.28 (1.10)	0.75

Abbreviations: ACQ Asthma Control Questionnaire, ICS Inhaled corticosteroid.

^aMissing mean ACQ score at Week 13 was imputed using the last observation carried forward method.

^bP-values are calculated using an analysis of covariance with treatment and baseline value as covariates.

Safety

The proportion of subjects who reported at least one AE was similar between the placebo (68/82, 82.9%) and MEDI-528 (30 mg: 62/81, 76.5%; 100 mg: 68/83, 81.9%; 300 mg: 69/81, 85.2%) groups. The most frequent AEs in the combined MEDI-528 groups were asthma, upper respiratory tract infection, and headache (Table 3). Most AEs were mild or moderate in severity. One subject in the MEDI-528 100 mg group had a mild hypersensitivity reaction on the day of dosing that was considered possibly related to study medication and resolved the following day.

SAEs were reported by 4 (4.9%) and 15 (6.1%) subjects in the placebo and combined MEDI-528 groups, respectively (Table 4). Two SAEs (pneumonia and asthma) in one subject who received 30 mg MEDI-528 were considered to be possibly related to study medication. Six subjects discontinued the study due to AEs: one placebo (rash); two 30 mg MEDI-528 (ischemic stroke, headache); one 100 mg MEDI-528 (musculoskeletal pain and pain in extremity); and two 300 mg MEDI-528 (headache, bladder transitional cell carcinoma). Three of the AEs that resulted in discontinuation were considered likely to be related




Figure 4 Weighted rate of asthma exacerbation (95% CI) at week 25 (ITT population). Abbreviation: ITT, Intent-to-treat.




Figure 5 Mean (SE) change from baseline in pre-bronchodilator FEV₁ at week s 13 and 25 (ITT population). Abbreviations: FEV₁, Forced expiratory volume in 1 second; ITT, Intent-to-treat; SE, Standard error.

to study medication (100 mg, musculoskeletal pain and pain in extremity; 300 mg, headache).

No clinically relevant changes occurred in clinical laboratory values, vital signs, or ECGs.

ADAs occurred in 4/82 subjects (4.9%) in the placebo group and 22/245 subjects (9.0%) in the MEDI-528 groups (30 mg: 15/81, 18.5%; 100 mg: 4/83, 4.8%; 300 mg: 3/81, 3.7%). ADAs were detected prior to day 0 in 1 subject in the placebo group and 14 subjects in the MEDI-528 group.

Discussion

This is the largest double-blind, placebo-controlled, dose-ranging study performed to date to determine whether an anti-IL-9 monoclonal antibody has any clinical benefits in subjects with poorly controlled, moderate-to-severe asthma. The results indicate that adding MEDI-528 to existing controller medications was not associated with any major safety concerns, but failed to achieve a clinically important effect on mean ACQ-6 scores at 13 weeks or on asthma exacerbation rates, lung function, or asthma-related quality of life at the pre-specified time points. However, this study was powered to detect a change in the primary study endpoint of mean ACQ-6 score from baseline to week 13, thus statistical results for the secondary endpoints must be viewed in this context.

Data from preclinical studies provide strong evidence that an IL-9 mast cell axis regulates airway inflammation, mucus production, airway hyperresponsiveness,

and subepithelial fibrosis, with increased IL-9 expression in the airways in humans with asthma [5]. For example, both Th9 and Th17 are involved in asthma pathogenesis and produce IL-9 [17]. Kim et al. demonstrated in Balb/c mice that cellular infiltration related to chronic airway inflammation and remodeling can be reduced by an anti-IL-9 antibody. In addition, the number of Th9 cells, Th17 cells, mast cells, eosinophils, and neutrophils in the airway were reduced; the synthesis and secretion of cytokines was inhibited; and IgE synthesis in B cells was reduced [18]. These reported observations support our hypothesis that targeting IL-9 via an anti-IL-9 mAb would be an effective treatment for patients with poorly controlled asthma. Despite convincing *in vitro* and animal model data, there have been very few studies in humans examining IL-9 involvement in asthma.

In a previous small study in subjects with mild asthma, fewer subjects experienced ≥ 1 asthma exacerbation in the combined MEDI-528 group (1/27) compared with the placebo group (2/9; $p = 0.15$). The subject in the MEDI-528 group who experienced an exacerbation received the lowest dose (0.3 mg/kg), which indicated that it was not due to a dose effect. These results suggested a potential improvement in asthma exacerbation rates in subjects with mild asthma [11]. However, in the current study, no significant differences in the asthma exacerbation rate occurred between placebo and MEDI-528 in adults with uncontrolled moderate-to-severe persistent asthma.

Table 3 Adverse events that occurred in ≥ 2% of the combined MEDI-528 group (safety population)

Adverse event, n (%) of subjects	Placebo (n = 82)	MEDI-528			Combined (n = 245)
		30 mg (n = 81)	100 mg (n = 83)	300 mg (n = 81)	
Asthma	25 (30.5)	30 (37.0)	21 (25.3)	31 (38.3)	82 (33.5)
Upper respiratory tract infection	12 (14.6)	6 (7.4)	16 (19.3)	20 (24.7)	42 (17.1)
Headache	8 (9.8)	13 (16.0)	5 (6.0)	6 (7.4)	24 (9.8)
Nasopharyngitis	12 (14.6)	11 (13.6)	5 (6.0)	7 (8.6)	23 (9.4)
Sinusitis	4 (4.9)	10 (12.3)	4 (4.8)	6 (7.4)	20 (8.2)
Influenza	6 (7.3)	10 (12.3)	5 (6.0)	2 (2.5)	17 (6.9)
Urinary tract infection	6 (7.3)	7 (8.6)	7 (8.4)	2 (2.5)	16 (6.5)
Pharyngitis	2 (2.4)	4 (4.9)	8 (9.6)	4 (4.9)	16 (6.5)
Bronchitis	8 (9.8)	4 (4.9)	2 (2.4)	5 (6.2)	11 (4.5)
Injection site erythema	2 (2.4)	1 (1.2)	4 (4.8)	5 (6.2)	10 (4.1)
Allergic rhinitis	7 (8.5)	5 (6.2)	1 (1.2)	3 (3.7)	9 (3.7)
Gastroenteritis	3 (3.7)	2 (2.5)	5 (6.0)	2 (2.5)	9 (3.7)
Diarrhea	2 (2.4)	3 (3.7)	3 (3.6)	3 (3.7)	9 (3.7)
Rhinitis	5 (6.1)	3 (3.7)	3 (3.6)	2 (2.5)	8 (3.3)
Back pain	2 (2.4)	3 (3.7)	3 (3.6)	2 (2.5)	8 (3.3)
Pneumonia	0 (0.0)	5 (6.2)	0 (0.0)	3 (3.7)	8 (3.3)
Dizziness	5 (6.1)	2 (2.5)	5 (6.0)	0 (0.0)	7 (2.9)
Injection site pain	5 (6.1)	1 (1.2)	4 (4.8)	2 (2.5)	7 (2.9)
Rash	3 (3.7)	2 (2.5)	1 (1.2)	4 (4.9)	7 (2.9)
Acute sinusitis	2 (2.4)	3 (3.7)	2 (2.4)	2 (2.5)	7 (2.9)
Hypertension	2 (2.4)	4 (4.9)	3 (3.6)	0 (0.0)	7 (2.9)
Pain in extremity	2 (2.4)	1 (1.2)	3 (3.6)	3 (3.7)	7 (2.9)
Cough	0 (0.0)	2 (2.5)	2 (2.4)	3 (3.7)	7 (2.9)
Oropharyngeal pain	0 (0.0)	2 (2.5)	3 (3.6)	2 (2.5)	7 (2.9)
Pyrexia	0 (0.0)	0 (0.0)	4 (4.8)	3 (3.7)	7 (2.9)
Migraine	1 (1.2)	4 (4.9)	1 (1.2)	1 (1.2)	6 (2.4)
Pruritus	1 (1.2)	1 (1.2)	2 (2.4)	3 (3.7)	6 (2.4)
Toothache	1 (1.2)	2 (2.5)	4 (4.8)	0 (0.0)	6 (2.4)
Nasal congestion	0 (0.0)	2 (2.5)	3 (3.6)	1 (1.2)	6 (2.4)
Pharyngotonsillitis	0 (0.0)	5 (6.2)	0 (0.0)	1 (1.2)	6 (2.4)
Bacterial upper respiratory tract infection	0 (0.0)	2 (2.5)	2 (2.4)	2 (2.5)	6 (2.4)
Asthenia	2 (2.4)	3 (3.7)	1 (1.2)	1 (1.2)	5 (2.0)
Malaise	1 (1.2)	0 (0.0)	3 (3.6)	2 (2.5)	5 (2.0)
Dyspepsia	0 (0.0)	3 (3.7)	1 (1.2)	1 (1.2)	5 (2.0)
Musculoskeletal pain	0 (0.0)	1 (1.2)	4 (4.8)	0 (0.0)	5 (2.0)
Bacterial pharyngitis	0 (0.0)	2 (2.5)	1 (1.2)	2 (2.5)	5 (2.0)

Likewise, the mean increase from baseline in pre-bronchodilator FEV₁ was similar between the MEDI-528 and placebo groups, which is consistent with the earlier study where FEV₁ was essentially unchanged and short-acting beta agonist use was comparable among groups [11]. Studies with other monoclonal antibodies, including omalizumab [19] and mepolizumab [20], also reported modest, if any, improvements in FEV₁.

Although the study did not meet its primary endpoint, all groups, including placebo, demonstrated improvements in mean ACQ-6 score at 13 weeks. The placebo response has been well described in asthma studies and various explanations have been proposed. For example, Wise et al. reported that the placebo effect on asthma symptoms could be augmented by an optimistic message from the physician that enhanced the subject's expectation of

Table 4 Serious adverse events by treatment group (safety population)

Serious adverse event, n (%) of subjects	Placebo (n = 82)	MEDI-528			
		30 mg (n = 81)	100 mg (n = 83)	300 mg (n = 81)	Combined (n = 245)
Asthma	0 (0.0)	6 (7.4)	1 (1.2)	2 (2.5)	9 (3.7)
Pneumonia	0 (0.0)	3 (3.7)	0 (0.0)	0 (0.0)	3 (1.2)
Food allergy	0 (0.0)	1 (1.2)	0 (0.0)	0 (0.0)	1 (0.4)
Bronchitis	0 (0.0)	1 (1.2)	0 (0.0)	0 (0.0)	1 (0.4)
Lobar pneumonia	0 (0.0)	0 (0.0)	1 (1.2)	0 (0.0)	1 (0.4)
Limb traumatic amputation	0 (0.0)	1 (1.2)	0 (0.0)	0 (0.0)	1 (0.4)
Thoracic vertebral fracture	0 (0.0)	0 (0.0)	1 (1.2)	0 (0.0)	1 (0.4)
Muscle spasms	0 (0.0)	0 (0.0)	1 (1.2)	0 (0.0)	1 (0.4)
Uterine leiomyoma	0 (0.0)	0 (0.0)	1 (1.2)	0 (0.0)	1 (0.4)
Complicated migraine	0 (0.0)	1 (1.2)	0 (0.0)	0 (0.0)	1 (0.4)
Ischemic stroke	0 (0.0)	1 (1.2)	0 (0.0)	0 (0.0)	1 (0.4)
Vaginal hemorrhage	0 (0.0)	0 (0.0)	1 (1.2)	0 (0.0)	1 (0.4)
Pancreatitis	1 (1.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Cholelithiasis	1 (1.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Hydrocholecystis	1 (1.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Appendicitis	1 (1.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Nephrolithiasis	1 (1.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)

benefit from the drug [21]. Similarly, a large placebo response was observed in two pivotal studies of omalizumab and was explained by improved compliance with ICS treatment and/or intensive medical input [22].

It is becoming increasingly apparent that asthma is a heterogeneous disease, and that identification of potential subgroups or individual subject characteristics is likely to be key in delivering optimal response with therapeutics such as MEDI-528 that target specific immunological mechanisms. Subgroup analyses of the mean ACQ-6 score results provided no evidence that MEDI-528 may be effective in subjects with atopic asthma, peripheral blood eosinophilia, or in those taking moderate or high doses of ICS at entry to the study. Interpretation of these results should take into account the small numbers of subjects in some subgroups.

The heterogeneous nature of asthma makes it difficult for a targeted therapy with monoclonal antibodies such as MEDI-528 to show significant beneficial effects in a non-selected asthma population. Identification of biological markers may be of great help in determining more accurate asthma diagnosis and severity, predicting treatment response, or monitoring of disease control. Corren et al. showed that lebrikizumab increased FEV₁ and reduced exacerbations in the high Th2 subgroup vs placebo [23]. Unfortunately, a pharmacodynamic marker for the IL-9 pathway has not been identified and IL-9 levels couldn't be measured in blood or sputum in this study. Furthermore, subgroup analyses of the primary endpoint

in pre-specified subpopulations, stratified by atopic status or medium- vs high-dose ICS at screening, and other pre-specified subpopulations based on FEV₁, ACQ score, reversibility, and blood eosinophilia at randomization did not identify a subpopulation with improved efficacy over placebo.

Overall, the safety profile of MEDI-528 was similar to earlier clinical studies [11,12] and no new safety concerns were identified. ADAs were detected in the placebo and all three MEDI-528 treatment groups. None of these events were considered to be serious, but it is unclear why ADAs were detected in the placebo group or in some subjects prior to administration of MEDI-528. Our ADA assays may nonspecifically detect cross-reacting antigen or other epitopes; however, it is not uncommon for subjects to develop antibodies against a biologic agent [24].

Conclusions

This study was designed to further evaluate the clinical effects of MEDI-528, an anti-IL-9 antibody, in light of the inconclusive results obtained in previous small studies. In this large study, we clearly demonstrated that adding MEDI-528 to existing controller medications was not associated with an improvement in ACQ-6 score, asthma exacerbation rate, FEV₁, health-related quality of life, or any major safety concerns.

Abbreviations

ACQ-6: Asthma control questionnaire-6; ADA: Anti-drug antibody; AE: Adverse event; AQLQ(S): Asthma quality of life questionnaire (Standardized version); CI: Confidence interval; ECG: Electrocardiogram;

FEV₁: Forced expiratory volume in 1 second; ICS: Inhaled corticosteroids; IL: Interleukin; ITT: Intent-to-treat; PEF: Peak expiratory flow; SAE: Serious adverse event; SD: Standard deviation; Th2: T helper cells.

Competing interests

Chad Oh and Nestor Molfino were employed by MedImmune during the conduct of the study and owned stock in AstraZeneca. Micki Hultquist, Kimmie McLaurin, and Keunpyo Kim are employees of MedImmune and own stock in AstraZeneca. Richard Leigh has received research funding from MedImmune, payable to the University of Calgary, for conducting this and other clinical studies over the past 3 years.

Authors' contributions

CKO participated in design and coordination of the study. RL was a principal investigator for the study. KKM participated in the design of the study and interpretation of results. KK participated in the design of the study and performed the statistical analysis. MH participated in study design and interpretation of results. NAM conceived the study and participated in its design and coordination. All authors have been involved in drafting the manuscript or revising it critically for important intellectual content, read, and approved the final manuscript.

Acknowledgements

The authors thank all subjects who consented to participate in this study, the study coordinators who facilitated the conduct of the study, and the following principal investigators.

Argentina: F Colodenco, J Maspero, L Nannini, A Stok, A Tolcachier, A Yanez. **Brazil:** E Fiss, C Fritscher, J Hetzel, E Pizzichini, R Stelmach. **Canada:** J Hebert, M Fitzgerald, R Olivenstein, I Mayers, A Cheema, W Yang. **Colombia:** A Casas, D Severiche, F Serrano, R Contreras, A Orozco. **Costa Rica:** T Mitchell. **Panama:** L Noriega. **Peru:** A Matsuno, C Villaran, J Portugal, O Cubas. **Philippines:** T Aquino, T Atienza, R Samoro. **Taiwan:** C-C Wang. **USA:** A Nayak, H Kaiser, J Winder, T Casale, J Corren, R Nathan, M White, D Dvorin, E Kerwin, J Matz, I Melamed, J Diaz, B Collins, B Chippis, C Chappel, L Mansfield, J Zwetchkenbaum, E Meltzer, S Pollard, N Kao.

We acknowledge Jennifer Stewart, MSc, (QXV Communications, Macclesfield, UK) for medical writing support, which was funded by MedImmune; Diana Swanson, PhD, (MedImmune) for document management and medical writing support, and Lourdes Briz (MedImmune) for editorial support. The study sponsor, MedImmune, was involved in the study design; collection, analysis, and interpretation of data; provided funding for writing support; and approved the decision to submit the manuscript for publication.

Author details

¹MedImmune, One MedImmune Way, Gaithersburg, MD, USA. ²Department of Medicine, University of Calgary, Calgary, Alberta, Canada. ³Currently at Kalobios Pharmaceuticals, Inc, South San Francisco, CA, USA.

Received: 15 May 2013 Accepted: 5 September 2013

Published: 19 September 2013

References

- National Heart, Lung, and Blood Institute: *National Asthma Education and Prevention Program: Guidelines for the diagnosis and management of asthma: expert panel report 3*, NIH Publication No. 07-4051. Bethesda, MD: National Institutes of Health; 2007.
- Nicolaides NC, Holroyd KJ, Ewart SL, Eleff SM, Kiser MB, Dragwa CR, Sullivan CD, Grasso L, Zhang LY, Messler CJ, Zhou T, Kleeberger SR, Buettow KH, Levitt RC: **Interleukin 9: a candidate gene for asthma.** *Proc Natl Acad Sci USA* 1997, **94**:13175–13180.
- Longphore M, Li D, Gallup M, Drori E, Ordonez CL, Redman T, Wenzel S, Bice DE, Fahy JV, Basbaum C: **Allergen-induced IL-9 directly stimulates mucin transcription in respiratory epithelial cells.** *J Clin Invest* 1999, **104**:1375–1382.
- Shimbara A, Christodouloupolous P, Soussi-Gounni A, Olivenstein R, Nakamura Y, Levitt RC, Nicolaides NC, Holroyd KJ, Tsicopoulos A, Lafitte JJ, Wallaert B, Hamid QA: **IL-9 and its receptor in allergic and nonallergic lung disease: increased expression in asthma.** *J Allergy Clin Immunol* 2000, **105**:108–115.
- Oh CK, Raible D, Geba GP, Molfino NA: **Biology of the interleukin-9 pathway and its therapeutic potential for the treatment of asthma.** *Inflamm Allergy Drug Targets* 2011, **10**:180–186.
- Gounni AS, Hamid Q, Rahman SM, Hoeck J, Yang J, Shan L: **IL-9-mediated induction of eotaxin1/CCL11 in human airway smooth muscle cells.** *J Immunol* 2004, **173**:2771–2779.
- Matsuzawa S, Sakashita K, Kinoshita T, Ito S, Yamashita T, Koike K: **IL-9 enhances the growth of human mast cell progenitors under stimulation with stem cell factor.** *J Immunol* 2003, **170**:3461–3467.
- Louahed J, Toda M, Jen J, Hamid Q, Renaud JC, Levitt RC, Nicolaides NC: **Interleukin-9 upregulates mucus expression in the airways.** *Am J Respir Cell Mol Biol* 2000, **22**:649–656.
- Kearley J, Erjefalt JS, Andersson C, Benjamin E, Jones CP, Robichaud A, Pegorier S, Brewah Y, Burwell TJ, Bjerner L, Kiener PA, Kolbeck R, Lloyd CM, Coyle AJ, Humbles AA: **IL-9 governs allergen-induced mast cell numbers in the lung and chronic remodeling of the airways.** *Am J Respir Crit Care Med* 2011, **183**:865–875.
- Humbles AA, Reed JL, Parker J, Kiener PA, Molfino NA, Kolbeck R, Coyle AJ: **Monoclonal antibody therapy directed against interleukin-9: MEDI-528.** In *New drugs and targets for asthma and COPD*. Edited by Hansel TT, Barnes PJ. Basel: Karger; 2010:137–140.
- Parker JM, Oh CK, LaForce C, Miller SD, Pearlman DS, Le C, Robbie GJ, White WI, White B, Molfino NA: **Safety profile and clinical activity of multiple subcutaneous doses of MEDI-528, a humanized anti-interleukin-9 monoclonal antibody, in two randomized phase 2a studies in subjects with asthma.** *BMC Pulm Med* 2011, **11**:14.
- White B, Leon F, White W, Robbie G: **Two first-in-human, open-label, phase I dose-escalation safety trials of MEDI-528, a monoclonal antibody against interleukin-9, in healthy adult volunteers.** *Clin Ther* 2009, **31**:728–740.
- Juniper EF, O'Byrne PM, Guyatt GH, Ferrie PJ, King DR: **Development and validation of a questionnaire to measure asthma control.** *Eur Respir J* 1999, **14**:902–907.
- Miller MR, Hankinson J, Busasco V, Burgos F, Casaburi R, Coates A, Crapo R, Enright P, van der Grinten CP, Gustafsson P, Jensen R, Johnson DC, MacIntyre N, McKay R, Navajas D, Pedersen OF, Pellegrino R, Viegi G, Wanger J: **Standardisation of spirometry.** *Eur Respir J* 2005, **26**:319–338.
- Giraud V, Rogeaux Y, Dusser D: **Asthma control following initial inhaled corticosteroid monotherapy in mild to moderate asthma: a 4- to 8-week observational study.** *Respiration* 2006, **73**:617–622.
- Molimard M, Martinat Y, Rogeaux Y, Moyse D, Pello JY, Giraud V: **Improvement of asthma control with beclomethasone extrafine aerosol compared to fluticasone and budesonide.** *Respir Med* 2005, **99**:770–778.
- Stassen M, Schmitz E, Bopp T: **From interleukin-9 to T helper 9 cells.** *Ann N Y Acad Sci* 2012, **1247**:56–68.
- Kim MS, Cho KA, Cho YJ, Woo SY: **Effects of interleukin-9 blockade on chronic airway inflammation in murine asthma models.** *Allergy Asthma Immunol Res* 2013, **5**:197–206.
- Rodrigo GJ, Neffen H, Catro-Rodriguez JA: **Efficacy and safety of subcutaneous omalizumab vs placebo as add-on therapy to corticosteroids for children and adults with asthma.** *Chest* 2011, **139**:28–35.
- Pavord ID, Korn S, Howarth P, Bleeker ER, Buhl R, Keene ON, Ortega H, Chaney P: **Mepolizumab for severe eosinophilic asthma (DREAM): a multicentre, double-blind, placebo-controlled trial.** *Lancet* 2012, **380**:651–659.
- Wise RA, Bartlett SJ, Brown ED, Castro M, Cohen R, Holbrook JT, Irvin CG, Rand CS, Sockrider MM, Sugar EA: **Randomized trial of the effect of drug presentation on asthma outcomes: the American Lung Association Asthma Clinical Research Centers.** *J Allergy Clin Immunol* 2009, **124**:436–438.
- Bousquet J, Wenzel S, Holgate S, Lumry W, Freeman P, Fox H: **Predicting response to omalizumab, an anti-IgE antibody, in patients with allergic asthma.** *Chest* 2004, **125**:1378–1386.
- Corren J, Lemanske RF, Hanania NA, Korenblat PE, Parsey MV, Arron JR, Harris JM, Scheerens H, Wu LC, Su Z, Mosesova S, Eisner MD, Bohen SP, Matthews JG: **Lebrikizumab treatment in adults with asthma.** *N Engl J Med* 2011, **365**:1088–1098.
- Jones TD, Crompton LJ, Carr FJ, Baker MP: **Deimmunization of monoclonal antibodies.** *Methods Mol Biol* 2009, **525**:405–423.

doi:10.1186/1465-9921-14-93

Cite this article as: Oh et al.: A randomized, controlled trial to evaluate the effect of an anti-interleukin-9 monoclonal antibody in adults with uncontrolled asthma. *Respiratory Research* 2013 **14**:93.