

POSTER PRESENTATION

Open Access

Cryptococcosis in a patient with HIV infection

Vasile Benea^{1*}, Simona Roxana Georgescu¹, Mircea Tampa¹, Mihaela-Anca Benea-Mălin¹, Diana Leahu¹, Cristina Răileanu¹, Șerban Benea²

From The 9th Edition of the Scientific Days of the National Institute for Infectious Diseases Prof Dr Matei Bals Bucharest, Romania. 23-25 October 2013

Background

Cryptococcus neoformans is encapsulated yeast with worldwide distribution, which may cause a self-limited pulmonary infection or disseminate (especially to the meninges, but sometimes to the skin, bones, viscera, or other sites). The infection is acquired by inhalation of contaminated soil.

Cryptococcosis is a defining opportunistic infection for AIDS; also at increased risk for infection are patients with Hodgkin's or other lymphomas, sarcoidosis, or those receiving long-term corticosteroid therapy. Progressive disseminated cryptococcosis also sometimes affects men over 40 years that aren't obviously immunocompromised.

Most cryptococcal infections have a self-limited, subacute or chronic course; in AIDS patients cryptococcal infection may present with acute, severe pneumonia. The brain is the most common organ infected by the hematogenous route; cryptococcal meningitis is an important complication of AIDS. Disseminated cutaneous involvement occurs also by hematogenous dissemination; it causes pustular, papular, nodular, or ulcerated lesions, sometimes resembling acne, molluscum contagiosum, or basal cell carcinoma. The initial treatment is based on amphotericin B and flucytosine; oral fluconazole is needed for chronic suppressive therapy, especially in AIDS patients.

Case report

We present a 32 years old man who was referred for a disseminated polymorph eruption with papular, nodular and ulcerated lesions. The serology for HIV infection was positive and the biopsy identified encapsulated yeasts. Because the patient had fever, headache, confusion, agitation and blurred vision, he was transferred in a neurology clinic.

Authors' details

¹Department of Dermatology, Clinical Hospital of Infectious and Tropical Diseases "Dr. Victor Babeș", Bucharest, Romania. ²National Institute for Infectious Diseases "Prof. Dr. Matei Bals", Bucharest, Romania.

Published: 16 December 2013

doi:10.1186/1471-2334-13-S1-P109

Cite this article as: Benea et al.: Cryptococcosis in a patient with HIV infection. *BMC Infectious Diseases* 2013 13(Suppl 1):P109.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: beneav@yahoo.com

¹Department of Dermatology, Clinical Hospital of Infectious and Tropical Diseases "Dr. Victor Babeș", Bucharest, Romania

Full list of author information is available at the end of the article