

SCIENTIFIC REPORTS



OPEN

Ultraviolet-B Radiation (UV-B) Relieves Chilling-Light-Induced PSI Photoinhibition And Accelerates The Recovery Of CO₂ Assimilation In Cucumber (*Cucumis sativus* L.) Leaves

Received: 14 May 2016
Accepted: 13 September 2016
Published: 30 September 2016

Zi-Shan Zhang^{1,2,3}, Li-Qiao Jin^{1,3}, Yu-Ting Li^{1,3}, Mikko Tikkanen⁴, Qing-Ming Li^{1,2}, Xi-Zhen Ai^{1,2} & Hui-Yuan Gao^{1,3}

Ultraviolet-B radiation (UV-B) is generally considered to negatively impact the photosynthetic apparatus and plant growth. UV-B damages PSII but does not directly influence PSI. However, PSI and PSII successively drive photosynthetic electron transfer, therefore, the interaction between these systems is unavoidable. So we speculated that UV-B could indirectly affect PSI under chilling-light conditions. To test this hypothesis, the cucumber leaves were illuminated by UV-B prior or during the chilling-light treatment, and the leaves were then transferred to 25 °C and low-light conditions for recovery. The results showed that UV-B decreased the electron transfer to PSI by inactivating the oxygen-evolving complex (OEC), thereby protecting PSI from chilling-light-induced photoinhibition. This effect advantages the recoveries of PSI and CO₂ assimilation after chilling-light stress, therefore should minimize the yield loss caused by chilling-light stress. Because sunlight consists of both UV-B and visible light, we suggest that UV-B-induced OEC inactivation is critical for chilling-light-induced PSI photoinhibition in field. Moreover, additional UV-B irradiation is an effective strategy to relieve PSI photoinhibition and yield loss in protected cultivation during winter. This study also demonstrates that minimizing the photoinhibition of PSI rather than that of PSII is essential for the chilling-light tolerance of the plant photosynthetic apparatus.

Ultraviolet-B radiation (UV-B) is a common abiotic stress that restrains leaf expansion, Rubisco activity and the stomatal opening to reduce carbon assimilation and vegetative growth^{1,2}. UV-B also damages DNA and proteins, particularly the photosynthetic pigment-protein complex in the thylakoid membrane³. The filtration of UV-B markedly increases the biomass of plants, but additional UV-B impairs the growth of plants^{2,4}. Consequently, UV-B is generally considered to negatively impact plants^{2,4}, although some research has shown that UV-B can suppress insect predation and pathogen infection in plants^{5–7}.

However, this opinion considered only the Calvin cycle and photosystem II (PSII), while neglecting photosystem I (PSI) and the interaction between PSII and PSI. Although UV-B does not directly influence PSI^{8,9}, PSI and PSII successively drive electron transfer in C3 plants; therefore, the interaction between them is unavoidable¹⁰. PSI is more stable than PSII under most abiotic stresses, such as high-light and high-temperature conditions^{9,11}. However, when chilling-sensitive plants experience chilling-light stress^{12–15}, or under the sequence of saturation pulses^{16–18}, PSI becomes the primary photoinhibition site. Unlike PSII, which can quickly recovers

¹State Key Lab of Crop Biology, Tai'an, Shandong Province, China. ²College of Horticulture Science and Engineering, Shandong Agricultural University, Tai'an, 271018, China. ³College of Life Sciences, Shandong Agricultural University, Tai'an, 271018, China. ⁴Plant Physiology and Molecular Biology, Department of Biochemistry and Food Chemistry, University of Turku, FIN-20014 Turku, Finland. Correspondence and requests for materials should be addressed to X.-Z.A. (email: axz@sdau.edu.cn) or H.-Y.G. (email: gaohy@sdau.edu.cn)




Figure 1. The influence of UV-B to PSI and PSII photoinhibition under chilling-light condition. The activity of the PSI complex and the maximum quantum yield of PSII (Fv/Fm) in leaves exposed to chilling-light and UV-B conditions for the indicated times. In plots (a,c), the leaves were exposed to approximately $6 \mu\text{mol m}^{-2} \text{s}^{-1}$ UV-B (+UV) or darkness (–UV) at 25°C for 2 h (shaded part), and the leaves were then transferred to chilling-light conditions (6°C , $200 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 6 h. In plots (b,d), the leaves were exposed to chilling-light conditions (6°C , $200 \mu\text{mol m}^{-2} \text{s}^{-1}$) in the presence (+UV) and absence (–UV) of approximately $2 \mu\text{mol m}^{-2} \text{s}^{-1}$ UV-B for 6 h. The means \pm SE, $n = 12$. Different letters indicate significant differences between +UV and –UV treated leaves at $P < 0.05$.

from damage^{18–20}, the recovery of the damaged PSI is quite slow and may be irreversible, even under optimum conditions^{13,20}.

The electron transfer from PSII has been shown to be necessary for PSI photoinhibition^{13–15,21}. Specifically, blocking the electron transfer from PSII to PSI with DCMU prevents the PSI photoinhibition induced by chilling-light conditions or fluctuating light^{13,22}. During the recovery from chilling-light stress, the limited electron transport from PSII to PSI protects PSI from further photoinhibition and accelerates the recovery of PSI¹³.

Recent studies have identified several effective mechanisms that control the electron transport from PSII to PSI in C3 leaves^{23,24}. The phosphorylation of PSII light-harvesting antenna (LHCII) balances the excitation of PSI and PSII and maintains the low reducing state of intersystem electron transfer chain under low-light conditions, which avoids excess electron transfer to the PSI acceptor side when the light intensity suddenly increases²⁵. In addition, the protection of PSI from photodamage upon a shift from low light to high light requires a PGR5-dependent cyclic electron flow around PSI (CEF) that controls the speed of intersystem electron transfer via the Cyt b6f complex in the high light phase^{22,26–28}. The ΔpH -dependent regulation of the Cyt b6f complex can slow the electron transfer from the plastoquinone (PQ) pool to plastocyanin (PC), avoiding excess electron transfer to PSI under high-light conditions²⁹. Tikkanen *et al.*³⁰ reported that the inhibition of de novo D1 synthesis protects PSI from photodamage in a *pgr5* mutant, which indicates that the PSII photoinhibition-repair cycle is an active regulatory component of the photosynthetic electron transfer to PSI. Although the sustained non-photochemical quenching (NPQ) that limits the light energy capture of PSII helps to maintain PSI activity in overwinter plants during harsh winters^{31–34}, the NPQ adversely affects PSI in other species. Recent research showed that the deletion of PsbS protein exacerbates PSII photoinhibition but alleviates PSI photoinhibition under high-light conditions in *Arabidopsis thaliana*³⁵ and *Oryza sativa* L.³⁶.

UV-B does not directly influence PSI but significantly inhibits PSII³⁷. PSI and PSII successively drive photosynthetic electron transfer, therefore, the interaction between these systems is unavoidable. So we speculated that UV-B could indirectly affect PSI under chilling-light conditions. Here, we provide evidence to show that UV-B controls the electron transfer to PSI by restraining the activity of the oxygen-evolving complex (OEC), which ultimately protects PSI from photoinhibition under chilling-light stress. After chilling-light stress, the recovery of PSI activity and carbon assimilation capacity were accelerated in UV-B exposed leaves, which should attenuate the loss of yield and growth of cucumbers caused by chilling-light stress.

Results

The influence of UV-B on PSI and PSII activities. Exposure to UV-B at normal temperatures did not markedly affect the activity of the PSI complex but steeply decreased the maximum quantum yield of PSII (Fv/Fm) (Fig. 1a,c). During the subsequent chilling-light treatments, the Fv/Fm was consistently lower in leaves pre-exposed to UV-B than in leaves that had not been pre-exposed to UV-B. Interestingly, UV-B pretreatment markedly




Figure 2. Immunoblot analysis of thylakoid membranes proteins extracted from leaves. Line 1, the leaves were exposed to darkness at 25 °C for 2 h; line 2, the leaves were exposed to approximately $6 \mu\text{mol m}^{-2} \text{s}^{-1}$ UV-B at 25 °C for 2 h; line 3, the leaves were exposed to darkness at 25 °C for 2 h and the leaves were then transferred to chilling-light conditions ($6 \text{ }^\circ\text{C}$, $200 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 6 h; line 4, the leaves were exposed to approximately $6 \mu\text{mol m}^{-2} \text{s}^{-1}$ UV-B at 25 °C for 2 h and the leaves were then transferred to chilling-light conditions ($6 \text{ }^\circ\text{C}$, $200 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 6 h; line 5, before treatment; line 6, the leaves were exposed to chilling-light conditions for 6 h; line 7, the leaves were exposed to chilling-light conditions in the presence of approximately $2 \mu\text{mol m}^{-2} \text{s}^{-1}$ UV-B for 6 h; Thylakoid membranes ($10 \mu\text{g}$ of chlorophyll) were separated by SDS-PAGE, electroblotted, and probed using specific antibodies against PsbO, PsbA, and PsaA.




Figure 3. The PSI (DCPIP-MV) and PSII (H₂O-BQ) electron transport activities of isolated thylakoid membranes in leaves exposed to chilling-light treatment and UV-B. In plots (a,c), the leaves were exposed to approximately $6 \mu\text{mol m}^{-2} \text{s}^{-1}$ UV-B (+UV) or darkness (–UV) at 25 °C for 2 h, and the leaves were then transferred to chilling-light conditions ($6 \text{ }^\circ\text{C}$, $200 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 6 h. In plots (b,d), the leaves were exposed to chilling-light ($6 \text{ }^\circ\text{C}$, $200 \mu\text{mol m}^{-2} \text{s}^{-1}$) in the presence (+UV) or absence (–UV) of approximately $2 \mu\text{mol m}^{-2} \text{s}^{-1}$ UV-B for 6 h. The means \pm SE, $n = 12$. Different letters indicate significant differences between +UV and –UV treated leaves at $P < 0.05$.

alleviated the decrease in the activity of the PSI complex during the chilling-light treatment (Fig. 1a,c). In another experiment, the leaves were irradiated with weaker UV-B during chilling-light treatment; PSI photoinhibition was also alleviated in this experiment, and the PSII photoinhibition was more severe (Fig. 1b,d).

The above results were verified by the following immunoblot analysis. Treatment with UV-B before or during the chilling-light treatment decreased the amount of PSII protein (PsbO and PsbA) but prevented the decrease in PSI protein (PsaA) during chilling-light treatment (Fig. 2).

To study the effect of UV-B to the capacity of electron transfer from PSII to PSI, we next analysed the electron transport rate in the thylakoid membrane. UV-B markedly decreased the electron transfer capacity of PSII to the acceptor side (H₂O to BQ). During chilling-light treatment, the capacity of electron transfer from donor side to acceptor side of PSI (DCPIP to MV) was protected from decreases by UV-B irradiation or UV-B pre-irradiation (Fig. 3).




Figure 4. The maximum fluorescence (F_m or $F_{m_{DCMU}}$; **(a,b)**) and maximum quantum yield of PSII (F_v/F_m or $F_v/F_{m_{DCMU}}$; **(c,d)**) in leaves exposed to chilling-light and UV-B conditions for the indicated times. In plots **(a,c)**, the leaves were exposed to approximately $6 \mu\text{mol m}^{-2} \text{s}^{-1}$ UV-B (+UV) or darkness (–UV) at 25°C for 2 h (shaded part), and the leaves were then transferred to chilling-light conditions (6°C , $200 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 6 h. In plots **(b,d)**, the leaves were exposed to chilling-light conditions (6°C , $200 \mu\text{mol m}^{-2} \text{s}^{-1}$) in the presence (+UV) or absence (–UV) of approximately $2 \mu\text{mol m}^{-2} \text{s}^{-1}$ UV-B for 6 h. After chilling-light and UV-B treatment, the leaves were infiltrated with DCMU or water in the dark for 1 h. The original F_m or the F_m measured in the presence of DCMU ($F_{m_{DCMU}}$) was used to calculate F_v/F_m (or $F_v/F_{m_{DCMU}}$). The means \pm SE, $n = 12$. Different letters indicate significant differences between +UV and –UV treated leaves at $P < 0.05$.

The above results clearly demonstrated that UV-B protected PSI from chilling-light-induced photoinhibition by inactivating PSII and decreasing the electron transfer from PSII to PSI.

The influence of UV-B on the donor and acceptor sides of PSII. The immunoblot analysis (Fig. 2) showed that UV-B damaged both the core protein of oxygen-evolving complex (OEC; PsbO) and Q_B -banding protein (PsbA), which may suppress the electron support at the PSII donor side and electron transfer from Q_A to Q_B at PSII acceptor side, respectively. Although the decrease in F_v/F_m has been used to reflect PSII photoinhibition, both the donor side inactivation and acceptor side inactivation have been proven to decrease the F_v/F_m ^{13,38–40}. Thus, to compare the inactivation of the donor and acceptor sides of PSII after UV-B exposure, leaves exposed to chilling-light (–UV leaves) or chilling-light-UV-B (+UV leaves) conditions for 2 h were infiltrated with DCMU or water in the dark, and the maximum fluorescence (F_m) was then measured with a saturated flash (Fig. 4).

DCMU did not influence the F_m in –UV leaves, which indicates that a flash of light can completely reduce Q_A^- , irrespective of the presence of DCMU (Fig. 4). However, DCMU markedly enhanced the F_m in +UV leaves (Fig. 4), which indicates that the Q_A cannot be completely reduced by a flash in +UV leaves without the presence of DCMU. This effect is attributed to that the electron transfer from OEC to Q_A was too weak to overcome the electron transfer beyond Q_A . To avoid the underestimation of F_m and F_v/F_m due to the inactivation of OEC, the F_m measured in the presence of DCMU ($F_{m_{DCMU}}$) was used instead of the original F_m to calculate F_v/F_m ($F_v/F_{m_{DCMU}}$). The $F_v/F_{m_{DCMU}}$ was higher than original F_v/F_m in +UV leaves, but the $F_v/F_{m_{DCMU}}$ was similar in +UV and –UV leaves (Fig. 4c,d). These results proved that, in UV-exposed leaves, the inactivation of OEC, rather than the inhibition of electron transfer from Q_A to Q_B , dominated the electron transfer from PSII to PSI and PSI photoprotection.

The influence of UV-B on the recovery of the photosynthetic apparatus. The eventual effect of abiotic stress on the photosynthesis and growth of plants is not only a result of the extent of the damage to the photosynthetic apparatus but also depends on the capacity for recovery after the damage has taken place^{13,41}. Thus, leaves exposed to chilling-light (–UV leaves) or chilling-light-UV-B (+UV leaves) conditions for 6 h were transferred to normal temperature and low-light conditions (25°C and $15 \mu\text{mol m}^{-2} \text{s}^{-1}$ PAR) for recovery. The activity of the PSI complex in +UV leaves recovered to approximately 87% of the initial activity within 36 h (Fig. 5a). In sharp contrast to +UV leaves, the activity of PSI complex in –UV leaves did not recover in the first 36 h of the recovery process; it recovered to less 50% of the initial activity after 72 h (Fig. 5a). Next, the photosynthetic rate (P_n) during the stress and recovery process was also investigated. Six hours of chilling-light stress




Figure 5. UV-B during chilling-light treatment accelerated the following recovery of the activity of the PSI complex and photosynthetic rate. The activity of the PSI complex (a) and photosynthetic rate (Pn; (b)) in leaves exposed to chilling-light conditions in the presence (+UV) or absence (–UV) of approximately $2 \mu\text{mol m}^{-2} \text{s}^{-1}$ UV-B for 6 h (shaded part) and subsequent repair process at $25 \text{ }^\circ\text{C}$ under low-light conditions ($15 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 72 h. The means \pm SE, $n = 12$ (a) or 6 (b). Different letters indicate significant differences between +UV and –UV treated leaves at $P < 0.05$.

decreased the Pn to negative values, irrespective of the presence of UV-B (Fig. 5b). However, during the subsequent recovery process, the recovery of Pn was much faster in +UV leaves than in –UV leaves (Fig. 5b).

Discussion

The PSI photoinhibition depends on the balance between the electron transfer to the acceptor side of PSI and the consumption of electrons by the Calvin cycle and other metabolic processes. Chilling stress causes massive electron transfer to O_2 due to the severe inhibition of Calvin cycle^{21,42}. Chilling inactivated the thylakoid-bound ascorbate peroxidases and superoxide dismutase, which caused the accumulation of ROS^{42,43}. ROS invariably damage the neighbouring iron-sulphur cluster and PsbA/B protein. In chilling-sensitive species, the Calvin cycle and ROS-scavenging enzymes are more sensitive to cold than in other species^{10,21}. Thus, limiting electron transfer to PSI is critical to avoiding PSI photoinhibition caused by chilling-light stress in chilling-sensitive species.

Previous studies reported that the mechanisms contributing to the control of electron transport to PSI involve LHCII phosphorylation²⁵, the ΔpH -²⁹, CEF-dependent control of Cyt b6^{22,26}, and the dynamic regulation of D1 protein turnover³⁰. The LHCII phosphorylation is only operational under fluctuating light, but this study examined only constant light conditions. Although D1 protein degradation (Figs 1,2 and 3), PGR5-dependent CEF¹³, and ΔpH dependent NPQ^{44–46} have been shown to be enhanced in cucumber leaves under chilling-light stress, however, PSI photoinhibition was not prevented (Figs 1,2 and 3). These factors indicate that the above mechanisms were not sufficiently strong to limit electron flow to PSI and thereby protect PSI in cucumber leaves under chilling-light stress. Here, we proved that UV-B inactivate OEC and thereby limited the photosynthetic electron flow to PSI (Fig. 6). Due to sunlight contains both UV-B and PAR, we suggest that the UV-B-induced PSII inactivation is critical for PSI photoprotection, irrespective of dynamic or constant light conditions. Because the OEC activity cannot immediately recover, the UV-B-induced OEC inactivation could protect PSI in the long term.

Tikkanen *et al.*³⁰ reported that pre-illumination in the presence of lincomycin protected PSI from photoinhibition during subsequent high-light treatment. The presence of Lin and pre-illumination resulted in the degradation of D1 protein, which inhibited the electron transfer at the acceptor side of PSII. In contrast, this study proved that in UV-exposed leaves, the inactivation of OEC, rather than the inhibition of electron transfer from Q_A to Q_B ,




Figure 6. Scheme of UV-B protects PSI from photoinhibition. OEC, oxygen-evolving complex; Fd, ferredoxins.

caused the increased electron transfer from PSII to PSI (Figs 2 and 4). Consequently, the dynamic regulation of D1 protein turnover and the sensitivity of OEC to UV-B are distinct mechanisms for the control of electron transfer to PSI. It has been reported that after the cucumber leaves were pretreated by dark-chilling, which inactivating the OEC, the PSI was protected from photoinhibition under following chilling-light condition⁴¹.

The entire photosynthetic apparatus, including the dark and light reactions, was inhibited by chilling-light stress, but the activities of ATPase, Calvin-cycle enzymes and PSII were restored during the recovery process^{47–49}. PSI is the limiting factor of the recovery of Pn after chilling-light stress^{13,20}. Consequently, the prevention of PSI photoinhibition and the acceleration of recovery in PSI activity are essential for the recovery of Pn after chilling-light stress.

Although UV-B damaged PSII (Figs 1,2 and 3), suppressed the opening of stoma and inhibited the Calvin-cycle^{1,2}, this study showed that these negative effects minimally influenced the photosynthesis rate (Fig. 5). However, the photoprotective effect of UV-B on PSI improved the recovery of Pn. Irreversible decreases in Pn result in a loss of yield during and after chilling-light stress, and additional UV-B irradiation can effectively prevent this loss to protect cultivation during winter. In contrast, the other mechanisms that control electron transport to PSI, such as LHCII phosphorylation and the Δ pH- and PGR5-dependent control of Cyt b6f, are complex, and their regulatory pathways are poorly understood. In addition, these mechanisms are multifunctional, and artificially modifying them may result in metabolic disturbances in cells. Consequently, these mechanisms cannot currently be used to improve the cold tolerance of PSI and Pn, whereas UV-B can be used to this end, as demonstrated in this study.

Conclusion

Previously, UV-B was considered to negatively impact the photosynthesis and growth of plants. However, this study demonstrated that UV-B decreased the electron transfer to PSI by inactivating the OEC, thereby protecting PSI from chilling-light-induced photoinhibition. This effect advantages the recovery of Pn after chilling-light




Figure 7. Spectral transmissions of the two spectral filters used in this study. Mylar film excluding most UV-B or Aclar film characterized by high UV-B transmittance. The variation between multiple samples of identical films was negligible, and the SEs are too small to present.

stress and minimized yield losses caused by chilling-light stress. This study also demonstrated that minimizing the photoinhibition of PSI rather than that of PSII is essential for the cold-tolerance of the photosynthetic apparatus. In addition, in contrast to sunlight, the artificial light source used in photosynthetic research does not contain UV-B, which may increase PSI photoinhibition. Thus, a light spectrum close to that of solar radiation spectrum should be considered in future studies.

Methods

Plant materials and growth conditions. Cucumber (*Cucumis sativus* L. cv. Jinyou 3) plants were grown in pots (7 cm in diameter, 10 cm in height) filled with rich soil. This rich soil supplied sufficient nutrients to plants, and the plants were supplied with sufficient water to grow. The plants were placed in a growth chamber at 25/22 °C and $150 \mu\text{mol m}^{-2} \text{s}^{-1}$ with a 14 h/10 h photoperiod. The youngest fully developed leaves of approximately 4-week-old plants were used in the experiments.

Treatments. The abaxial sides of attached leaves of the cucumber plants were placed on the surface of water at 6 °C. The temperature of the water was controlled by an RTE-211 water circulator (Thermo, USA). The leaves were illuminated by photosynthetically active radiation (PAR) and UV-B, which were obtained from a red and blue (8:1) light-emitting diode (Senpro, China) and a PL-S 9W/01/2P UV-B light source (Philips, Netherlands), respectively. The leaves were separated with PAR and UV-B light source by a 0.13 mm polyester plastic film (Mylar D; DuPont, USA) that excluded most UV-B or a 0.08-mm Aclar film (22A; Allied Signal, USA), which is characterized by a high transmittance of UV-B. The films were placed on 5 cm above the leaves. The spectral transmittance of the filters was measured using a spectrophotometer UV-1601 (Shimadzu, Japan; as shown in Fig. 7). The transmittance of PAR (400–700 nm) is high in both films (Fig. 7). The intensities of PAR and UV-B on the surface of leaves were adjusted by altering the distance between the surface of the leaf and light source. A 3414F LightScout UV meter (Spectrum, USA) or a 3415F LightScout quantum meter (Spectrum, USA) were used to measure the intensity of UV-B or PAR radiation.

Two series UV-B treatments were performed in this study. Series 1 is the UV-B pre-treatment experiment, in which the leaves were exposed to approximately $6 \mu\text{mol m}^{-2} \text{s}^{-1}$ UV-B or darkness at a normal temperature (25 °C) for 2 h. The leaves were then transferred to chilling-light conditions ($200 \mu\text{mol m}^{-2} \text{s}^{-1}$ PAR and 6 °C) for 6 h. Series 2 refers to the UV-B and PAR co-processing experiment, in which the leaves were exposed to chilling-light conditions ($200 \mu\text{mol m}^{-2} \text{s}^{-1}$ PAR and 6 °C) in the presence or absence of approximately $2 \mu\text{mol m}^{-2} \text{s}^{-1}$ UV-B for 6 h. The intensity of UV-B used in series 2 was lower than it in series 1 due to the treatment period in series 2 was much longer than it in series 1.

For inhibitor treatment, the leaves were submerged in $70 \mu\text{M}$ 3-(3,4-Dichlorophenyl)-1,1-dimethylurea (DCMU) or water for 1 h in the dark and then placed in air for 20 min before further experiments.

Measurements of the chlorophyll a fluorescence transient (OJIP) and 820-nm transmission.

The Chl a fluorescence transient, and the 820-nm transmission were synchronously measured using an integral Multifunctional Plant Efficiency Analyser (M-PEA; Hansatech, UK) with dark-adapted leaves at 25 °C. A 2 s saturating red light pulse of $5000 \mu\text{mol m}^{-2} \text{s}^{-1}$ produced by an array of four light-emitting diodes (peak 650 nm) was used in the measurements. The maximum quantum yield of PSII (Fv/Fm) was calculated with the following equation: $Fv/Fm = 1 - (Fo/Fm)$. The initial slope of the oxidation phase indicates the capability of P700 to be oxidized, which was used to reflect the activity of the PSI complex^{50–54}.

Isolation of Thylakoid Membranes. Five grams of leaf discs were homogenized on ice in 400 mM sucrose, 50 mM HEPES-KOH (pH 7.8), 10 mM NaCl, and 2 mM MgCl_2 and filtered through two layers of cheesecloth. The resulting filtrates were centrifuged at 5000 g for 10 min at 4 °C, and the thylakoid pellets were resuspended in the homogenization buffer. Chlorophyll content was determined according to the method of⁵⁵.

SDS-PAGE and Immunoblot analysis. Thylakoid membranes (10 µg chlorophyll) were loaded on and separated by a 12% (w/w) SDS-PAGE gel. Proteins from the gel were subsequently blotted onto nitrocellulose by standard methods. The blot was blocked with 5% (w/w) skimmed milk and then incubated for 2 h with the primary antibody. Subsequently, the blot was incubated with horseradish peroxidase-conjugated anti-rabbit IgG antibody (Solarbio, Beijing, China) for 2 h. The Supersignal West Pico substrate (Thermo Fisher Scientific, Shanghai, China) was applied to detect the immunoreaction. The chemiluminescence was recorded on blots using a Tanon-5500 cooled CCD camera (Tanon, Shanghai, China). The primary anti-PsbO, anti-PsbA, anti-PsaA antibodies against synthetic polypeptides containing amino acid residues CQPSDTDLGAKVPKD, APPVDIDGIREPVC, CRDYDPTTRYNDLLD, respectively, were purchased from Genscript company (Nanjing, China).

Photosynthetic electron transport measurements. Electron transport was measured using a Clark-type oxygen electrode under 1000 µmol m⁻² s⁻¹ white light⁵⁶. PSI electron transport was assayed in a 3 mL reaction mixture composed of 40 mM Tricine, pH 7.5, 10 mM Suc, 167 µM MV, 0.1 mM DCPIP, 1 mM sodium ascorbate, 10 mM NH₄Cl, 10 µM DCMU, 5 mM sodium azide, and thylakoids corresponding to 15 µg of chlorophyll. Similarly, PSII electron transport was assayed in a 3 mL reaction mixture composed of 50 mM Tricine, pH 7.5, 20 mM NaCl, 5 mM MgCl₂, 100 mM Suc, 1 mM phenyl-p-benzoquinone, and thylakoids corresponding to 15 µg of chlorophyll.

Photosynthetic gas exchange measurements. The net photosynthetic rate (P_n) was measured using a CIRAS-3 portable photosynthesis system (PP Systems, USA) at 25 °C, 800 µmol m⁻² s⁻¹ light (90% red light and 10% blue light), 400 µmol mol⁻¹ CO₂ and 60–65% relative humidity. Leaf temperature, light intensity, CO₂ concentration and relative humidity were controlled using the automatic control device of the CIRAS-3 portable photosynthetic system. After chilling-light treatment, the P_n were measured 2 h after the leaves were transferred from cool water to normal temperature.

References

- Jansen, M. A., Gaba, V. & Greenberg, B. M. Higher plants and UV-B radiation: balancing damage, repair and acclimation. *Trends Plant Sci.* **3**, 131–135 (1998).
- Comont, D., Winters, A., Gomez, L. D., McQueen-Mason, S. J. & Gwynn-Jones, D. Latitudinal variation in ambient UV-B radiation is an important determinant of *Lolium perenne* forage production, quality, and digestibility. *J. Exp. Bot.* **64**, 2193–2204 (2013).
- Kataria, S., Jajoo, A. & Guruprasad, K. N. Impact of increasing Ultraviolet-B (UV-B) radiation on photosynthetic processes. *J. Photochem. Photobiol. B* **137**, 55–66 (2014).
- Zhao, D., Reddy, K. R., Kakani, V. G., Read, J. J. & Sullivan, J. H. Growth and physiological responses of cotton (*Gossypium hirsutum* L.) to elevated carbon dioxide and ultraviolet-B radiation under control environmental conditions. *Plant Cell Environ.* **26**, 771–782 (2003).
- Ballaré, C. L., Mazza, C. A., Austin, A. T. & Pierik, R. Canopy light and plant health. *Plant Physiol.* **160**, 145–155 (2012).
- Mazza, C. A., Giménez, P. I., Kantolic, A. G. & Ballaré, C. L. Beneficial effects of solar UV-B radiation on soybean yield mediated by reduced insect herbivory under field conditions. *Physiol. Plantarum* **147**, 307–315 (2013).
- Đinh, S. T., Galis, I. & Baldwin, I. T. UVB radiation and 17-hydroxygeranylinalool diterpene glycosides provide durable resistance against mirid (*Tupiocoris notatus*) attack in field-grown *Nicotiana attenuata* plants. *Plant Cell Environ.* **36**, 590–606 (2013).
- Jones, L. W. & Kok, B. Photoinhibition of chloroplast reactions. II. Multiple effects. *Plant Physiol.* **41**, 1044–1049 (1966).
- Teramura, A. H. & Ziska, L. H. [Ultraviolet-B radiation and photosynthesis] *Photosynthesis and the Environment* [Baker, N. R. (eds)] [435–450] (Kluwer Academic Publishers, Dordrecht, 1996).
- Sonoike, K. [Photoinhibition and protection of photosystem I] *The light-driven plastocyanin: ferredoxin oxidoreductase*. [Golbeck, J. H. (eds)] [657–668] (Springer, Dordrecht, 2006).
- Scheller, H. V. & Haldrup, A. Photoinhibition of Photosystem I. *Planta* **221**, 5–8 (2005).
- Terashima, I., Funayama, S. & Sonoike, K. The site of photoinhibition in leaves of *Cucumis sativus* L. at low temperatures is photosystem I, not photosystem II. *Planta* **193**, 300–306 (1994).
- Zhang, Z. S. *et al.* Characterization of PSI recovery after chilling-induced photoinhibition in cucumber (*Cucumis sativus* L.) leaves. *Planta* **234**, 883–889 (2011).
- Zhang, Z. S. *et al.* Water status related root-to-shoot communication regulates the chilling tolerance of shoot in cucumber (*Cucumis sativus* L.) plants. *Sci. Rep.* **5**, 13094 (2015).
- Zhang, Z. S. *et al.* The higher sensitivity of PSI to ROS results in lower chilling-light tolerance of photosystems in young leaves of cucumber. *J. Photochem. Photobiol. B* **137**, 127–134 (2014).
- Zivcak, M., Brestic, M., Kunderlikova, K., Sytar, O. & Allakhverdiev, S. I. Repetitive light pulse-induced photoinhibition of photosystem I severely affects CO₂ assimilation and photoprotection in wheat leaves. *Photosynth. Res.* **126**, 449–463 (2015).
- Zivcak, M., Brestic, M., Kunderlikova, K., Olsovska, K. & Allakhverdiev, S. I. Effect of photosystem I inactivation on chlorophyll a fluorescence induction in wheat leaves: Does activity of photosystem I play any role in OJIP rise? *J. Photochem. Photobiol. B* **152**, 318–324 (2015).
- Melis, A. Photosystem-II damage and repair cycle in chloroplasts: what modulates the rate of photodamage *in vivo*?. *Trends Plant Sci.* **4**, 130–135 (1999).
- Vass, I. Molecular mechanisms of photodamage in the Photosystem II complex. *BBA-Bioenergetics* **1817**, 209–217 (2012).
- Kudoh, H. & Sonoike, K. Irreversible damage to photosystem I by chilling in the light: cause of the degradation of chlorophyll after returning to normal growth temperature. *Planta* **215**, 541–548 (2002).
- Sonoike, K. Photoinhibition of photosystem I. *Physiol. Plantarum* **142**, 56–64 (2011).
- Suorsa, M. *et al.* PROTON GRADIENT REGULATION5 is essential for proper acclimation of Arabidopsis photosystem I to naturally and artificially fluctuating light conditions. *Plant Cell* **24**, 2934–2948 (2012).
- Suorsa, M. *et al.* PGR5 ensures photosynthetic control to safeguard photosystem I under fluctuating light conditions. *Plant Signal. Behav.* **8**, e22741 (2013).
- Allahverdiyeva, Y., Suorsa, M., Tikkanen, M. & Aro, E. M. Photoprotection of photosystems in fluctuating light intensities. *J. Exp. Bot.* **66**, 2427–2436 (2015).
- Grieco, M., Tikkanen, M., Paakkariinen, V., Kangasjärvi, S. & Aro, E. M. Steady-state phosphorylation of light-harvesting complex II proteins preserves photosystem I under fluctuating white light. *Plant Physiol.* **160**, 1896–1910 (2012).
- Kono, M., Noguchi, K. & Terashima, I. Roles of the cyclic electron flow around PSI (CEF-PSI) and O₂-dependent alternative pathways in regulation of the photosynthetic electron flow in short-term fluctuating light in *Arabidopsis thaliana*. *Plant Cell Physiol.* **55**, 990–1004 (2014).

27. Huang, W., Yang, Y. J., Hu H., Cao, K. F. & Zhang, S. B. Sustained diurnal stimulation of cyclic electron flow in two tropical tree species *Erythrophleum guineense* and *Khaya ivorensis*. *Front. Plant Sci.* **7**, 1068 (2016).
28. Takagi, D., Hashiguchi, M. & Sejima, T. Photorespiration provides the chance of cyclic electron flow to operate for the redox-regulation of P700 in photosynthetic electron transport system of sunflower leaves. *Photosynth. Res.* in press (2016).
29. Joliot, P. & Johnson, G. N. Regulation of cyclic and linear electron flow in higher plants. *P. Natl. Acad. Sci. USA* **108**, 13317–13322 (2011).
30. Tikkanen, M., Mekala, N. R. & Aro, E. M. Photosystem II photoinhibition-repair cycle protects Photosystem I from irreversible damage. *BBA-Bioenergetics* **1837**, 210–215 (2014).
31. Ivanov, A. G. *et al.* Photosynthetic electron transport adjustments in overwintering Scots pine (*Pinus sylvestris* L.). *Planta* **213**, 575–585 (2001).
32. Ivanov, A. G. *et al.* Characterization of the photosynthetic apparatus in cortical bark chlorenchyma of Scots pine. *Planta* **223**, 1165–1177 (2006).
33. Öquist, G. & Huner, N. P. Photosynthesis of overwintering evergreen plants. *Annu. Rev. Plant Biol.* **54**, 329–355 (2003).
34. Ruban, A. V. Nonphotochemical chlorophyll fluorescence quenching: mechanism and effectiveness in protecting plants from photodamage. *Plant Physiol.* **170**, 1903–1916 (2016).
35. Roach, T. & Krieger-Liszka, A. The role of the PsbS protein in the protection of photosystems I and II against high light in *Arabidopsis thaliana*. *BBA-Bioenergetics* **1817**, 2158–2165 (2012).
36. Zulfugarov, I. S., Tovuu, A. & Lee, C. H. Acceleration of cyclic electron flow in rice plants (*Oryza sativa* L.) deficient in the PsbS protein of Photosystem II. *Plant Physiol. Bioch.* **84**, 233–239 (2014).
37. Takahashi, S. *et al.* The solar action spectrum of photosystem II damage. *Plant Physiol.* **153**, 988–993 (2010).
38. Murakami, R. *et al.* Characterization of an *Arabidopsis thaliana* mutant with impaired psbO, one of two genes encoding extrinsic 33-kDa proteins in photosystem II. *FEBS Lett.* **523**, 138–142 (2002).
39. Dwyer, S. A. *et al.* Antisense reductions in the PsbO protein of photosystem II leads to decreased quantum yield but similar maximal photosynthetic rates. *J. Exp. Bot.* **63**, 4781–4795 (2012).
40. Fan, X. *et al.* Photoinhibition-like damage to the photosynthetic apparatus in plant leaves induced by submergence treatment in the dark. *Plos One* **9**, 2 (2014).
41. Zhang, S. & Scheller, H. V. Photoinhibition of photosystem I at chilling temperature and subsequent recovery in *Arabidopsis thaliana*. *Plant Cell Physiol.* **45**, 1595–1602 (2004).
42. Choi, S. *et al.* Chloroplast Cu/Zn-superoxide dismutase is a highly sensitive site in cucumber leaves chilled in the light. *Planta* **216**, 315–324 (2002).
43. Terashima, I. *et al.* The cause of PSI photoinhibition at low temperatures in leaves of *Cucumis sativus*, a chilling-sensitive plant. *Physiol. Plantarum* **103**, 295–303 (1998).
44. Li, X. G., Meng, Q. W., Jiang, G. Q. & Zou, Q. The susceptibility of cucumber and sweet pepper to chilling under low irradiance is related to energy dissipation and water-water cycle. *Photosynthetica* **41**, 259–265 (2003).
45. Zhou, Y. H. *et al.* Role of thermal dissipation in the photoprotection in cucumber plants after exposure to a chill stress. *Photosynthetica* **44**, 262–267 (2006).
46. Kudoh, H. & Sonoike, K. Dark-chilling pretreatment protects PSI from light-chilling damage. *J. Photosci.* **9**, 59–62 (2002).
47. Terashima, I., Kashino, Y. & Katoh, S. Exposure of leaves of *Cucumis sativus* L. to low temperatures in the light causes uncoupling of thylakoids I. Studies with isolated thylakoids. *Plant Cell Physiol.* **32**, 1267–1274 (1991).
48. Terashima, I., Sonoike, K., Kawazu, T. & Katoh, S. Exposure of leaves of *Cucumis sativus* L. to low temperatures in the light causes uncoupling of thylakoids II. Non-destructive measurements with intact leaves. *Plant Cell Physiol.* **32**, 1275–1283 (1991).
49. Wise, R. R. Chilling-enhanced photooxidation: the production, action and study of reactive oxygen species produced during chilling in the light. *Photosynth. Res.* **45**, 79–97 (1995).
50. Strasser, R. J., Tsimilli-Michael, M., Qiang, S. & Goltsev, V. Simultaneous *in vivo* recording of prompt and delayed fluorescence and 820-nm reflection changes during drying and after rehydration of the resurrection plant *Haberlea rhodopensis*. *BBA-Bioenergetics* **1797**, 1313–1326 (2010).
51. Gao, J., Li, P., Ma, F. & Goltsev, V. Photosynthetic performance during leaf expansion in *malus micromalus* probed by chlorophyll a fluorescence and modulated 820 nm reflection. *J. Photoch. Photobiol. B* **137**, 144–150 (2014).
52. Oukarroum, A., Goltsev, V. & Strasser, R. J. Temperature effects on pea plants probed by simultaneous measurements of the kinetics of prompt fluorescence, delayed fluorescence and modulated 820 nm reflection. *Plos One* **8**, e59433 (2013).
53. Yang, C., Zhang, Z., Gao, H., Liu, M. & Fan, X. Mechanisms by which the infection of *Sclerotinia sclerotiorum* (lib) de bary affects the photosynthetic performance in tobacco leaves. *BMC Plant Biol.* **14**, 240 (2014).
54. Yan, K. *et al.* Dissection of Photosynthetic Electron Transport Process in Sweet Sorghum under Heat Stress. *Plos One* **8**, e62100 (2013).
55. Porra, R. J. The chequered history of the development and use of simultaneous equations for the accurate determination of chlorophyll a and b. *Photosynth. Res.* **73**, 149–156 (2002).
56. Tjus, S. E., Möller, B. L. & Scheller, H. V. Photosystem I is an early target of photoinhibition in barley illuminated at chilling temperatures. *Plant Physiol.* **116**, 755–764 (1998).

Acknowledgements

This work was supported by the China Postdoctoral Science Foundation (2015M572068), National Natural Science Foundation of China (31370276, 31471918), the State Key Basic Research and Development Plan of China (2015CB150105) and the National Plan of Key Basic Research in China (2009CB119000).

Author Contributions

Z.-S.Z., X.-Z.A. and H.-Y.G. designed the research; Z.-S.Z., L.-Q.J. and Y.-T.L. performed research; Z.-S.Z. and Q.-M.L. analyzed data; Z.-S.Z. and H.-Y.G. wrote the article; Q.-M.L. and M.T. read and approved the final manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Zhang, Z.-S. *et al.* Ultraviolet-B Radiation (UV-B) Relieves Chilling-Light-Induced PSI Photoinhibition And Accelerates The Recovery Of CO₂ Assimilation In Cucumber (*Cucumis sativus* L.) Leaves. *Sci. Rep.* **6**, 34455; doi: 10.1038/srep34455 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016