

Research Article

Analysis and Simulation of Fractional Order Smoking Epidemic Model

Aqeel Ahmad,¹ Muhammad Farman ,² Abdul Ghafar ,¹ Mustafa Inc ,^{3,4,5}
Mohammad Ozair Ahmad,² and Ndolane Sene ⁶

¹Department of Mathematics, Ghazi University, DG Khan, Pakistan

²Department of Mathematics and Statistics, University of Lahore, Lahore 54590, Pakistan

³Department of Computer Engineering, Biruni University, Istanbul, Turkey

⁴Department of Mathematics, Science Faculty, Firat University, Elazig, Turkey

⁵Department of Medical Research, China Medical University, Taichung, Taiwan

⁶Laboratoire Lmdan, Departement De Mathematiques De Decision, Faculté des Sciences Economiques et Gestion, Université Cheikh Anta Diop De Dakar, BP 5683 Dakar Fann, Senegal

Correspondence should be addressed to Mustafa Inc; minc@firat.edu.tr and Ndolane Sene; ndolanesene@yahoo.fr

Received 23 September 2021; Revised 8 December 2021; Accepted 12 January 2022; Published 20 May 2022

Academic Editor: Asep Kuswandi Supriatna

Copyright © 2022 Aqeel Ahmad et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In recent years, there are many new definitions that were proposed related to fractional derivatives, and with the help of these definitions, mathematical models were established to overcome the various real-life problems. The true purpose of the current work is to develop and analyze Atangana-Baleanu (AB) with Mittag-Leffler kernel and Atangana-Toufik method (ATM) of fractional derivative model for the Smoking epidemic. Qualitative analysis has been made to verify the steady state. Stability analysis has been made using self-mapping and Banach space as well as fractional system is analyzed locally and globally by using first derivative of Lyapunov. Also derive a unique solution for fractional-order model which is a new approach for such type of biological models. A few numerical simulations are done by using the given method of fractional order to explain and support the theoretical results.

1. Introduction

Mathematics was firstly used in biology in the twelfth century when Fibonacci used his popular Fibonacci series to explain a growing population. Daniel Bernoulli used mathematics to describe the effect of small pox. The term biological mathematics was primarily used by Johannes Reinke in 1901. It is aimed at the mathematical image and modeling of biological processes. It is also used to recognize phenomena in the living organism. Bio math has made major progress during the last few decades, and this progress will continue in upcoming decades. Math has played a large role in natural science but now it also will be more useful in biology. We should teach basic concepts of bio math at early stages. The basic steps in mathematical biology are few. The initial step is to explain the biological process and raise a

question about the basics. The second step is to build up a mathematical model that represents the underlying biological process. The 3rd step is to apply the method and concepts of math to obtain predictions about the model. The last step is to check whether this prediction answers the raised questions. After this, anyone can further explore the biological question by using mathematical models [1].

Now in the modern world, tobacco smoke is the most inhaled substance. Tobacco is made by mixing its agricultural form with many substances. The smoke is inhaled through the lungs. The most dangerous epidemic in the world is the smoking epidemic. Due to smoking 50% of its users died. Every year, about 60 million people die due to smoking. During the last few decades, there has been a huge boost in deaths. The death rate will rise thrice annually in 2030, almost seventy percent is in developing countries.

According to WHO, 10 million people will die due to smoking. As compared to other diseases, the death ratio is higher than all. The person who uses tobacco dies 14 years earlier than someone who does not smoke [2]. Tobacco smoking is the major reason for cancer and another disease. About 70% of people died due to tobacco-related diseases in developing countries [3]. Three million people died due to smoking yearly.

Nowadays, smoking is the most dangerous habit. The heart attack ratio is 70 percent more than a nonsmoker. There are 900 million men smokers and 200 women smokers in the world. After every 6 seconds, there is a death due to smoking. Smoking is a major cause of lung and heart attack in the world. Due to smoking, the chances of other diseases like heart attack, stroke, especially lung cancer, throat, mouth, esophagus, and pancreas are increased. Tobacco causes many tissue-related diseases. Tobacco smoke is a mixture of several toxic gases. It includes 98 of which are linked with an increased risk of cardiovascular disease, 69 of which are known to be carcinogenic. Daily, a smoker takes 1 or 2 milligrams of nicotine per cigarette. Thus, if a person smokes 5 cigarettes, it takes at least 5-10 milligrams of nicotine. The effect of smoking is not limited to the person who but also adverse effects for other people. It causes 22% of death annually.

The chemical composition of tobacco varies according to the environment. These leaves are mixed with many chemicals. Tobacco smoke contains a large number of different chemicals such as benzopyrene, NNK, aldehydes, carbon monoxide, hydrogen cyanide, phenol, nicotine, and harmful alkoids. The radioactive element polonium 210 is also occurring in tobacco. There are almost 4 thousand noxious substances in smoke which is the main reason for cancer [4, 5]. The chemical composition of smoke depends on puff frequency and other materials. Nicotine is the main issue for disturbing the nervous system, rise in heartbeat, raising blood pressure, and shrinking the small blood vessels which are the main basis of wrinkles. The amount of oxygen decreased in the lungs due to carbon monoxide (CO). The natural lungs cleaner that is minuscule hairs is destroyed by hydrogen cyanide. Lead, nickel arsenic, and cadmium are also present in smoke. Some pesticides like DDT are also found in smoke. The major reason for skin and lung cancer is a toxic chemical that is present in smoke. 10 million deaths will occur in the 20th century and 1 billion in the 21st century due to smoking [6]. Cigarette smoking affects human fertility badly [7].

The generalization of classical calculus is called fractional calculus which is concerned with the operation of integration and differentiation of fractional order. In the 19th century, fractional calculus mathematicians introduced fractional differential equations, fractional dynamics, and fractional geometry. Fractional calculus is used in almost every field of science. It is used to model physical as well as engineering processes. In many cases, standard mathematical models of integer order do not work properly. Due to this reason, fractional calculus made a major contribution to the field of mechanics, chemistry, biology, and image processing. By using fractional calculus, several physical prob-

lems are solved. By using integer-order derivatives, the system shows many problems such as history and nonlocal effects. Primarily, all the studies were dependent on Caputo fractional-order and Reimann Liouville fractional (RLF) derivatives. Nowadays, it has been highlighted that these derivatives have the issue, and the issue is they have a singular kernel. That is the reason so many new definitions were presented in the studies [8–16]. These new definitions were very impactful because they have nonsingular kernels which are according to their needs. Caputo fractional derivatives [17], the Caputo-Fabrizio derivative [11], and AB [18] fractional derivative have differed from each other only because Caputo is defined by a power law, Fabrizio defined by using exponential decay law, and AB defined by ML law. Tateishi et al. describe the role of fractional time operator derivative in a study of anomalous diffusion [12]. With the help of analytical techniques, Bulut et al. deliberate the role of differential equations of arbitrary order [19]. The key concepts of fractional differential equations and their application are explained by Kilbas et al. [20]. Atangana and Koca examined the Keller-Segel model about a fractional derivative having a nonsingular kernel [21]. Fractional logistic maps are newly introduced by Huang et al. [22]. Zaman studied the qualitative response of the dynamics of giving up smoking [23]. The giving up smoking model linked with Caputo fractional derivative is a probe by Singh et al. [24].

Numerous studies identified sociodemographic, environmental, and behavioral risk factors such as age, sex, occupations, indoor air pollutions, smoking, and alcohol consumption [25–27] as being associated with the development of TB in humans. In [28], a simple model for the effect of tobacco smoking in the in-host dynamics of HIV is formulated with the aim of studying how tobacco smoking affects HIV in-host dynamics. Zoonotic tuberculosis (zTB) knowledge, prevention, and control practices via a survey in Bangladesh considering impact of smoking are also in [29].

In this work, we get the approximate solutions of the fractional smoking model by using the Atangana-Toufik method.

2. Basic Concepts of Fractional Operators

Definition 1. For a function $g(t) \in W_2^1(0, 1)$, $b > a$ and $\sigma \in [0, 1]$, the definition of AB derivative in the Caputo sense is given by

$${}_{0}^{ABC}D_t^\sigma g(t) = \frac{AB(\sigma)}{1-\sigma} \int_0^t \frac{d}{d\tau} g(\tau) M_\sigma \cdot \left[-\frac{\sigma}{1-\sigma} (t-\tau)^\sigma \right] d\tau, \quad n-1 < \sigma < n, \quad (1)$$

where

$$AB(\sigma) = 1 - \sigma + \frac{\sigma}{\Gamma(\sigma)}. \quad (2)$$

By using ST for (1), we obtain

$$\text{ST} [{}_{0}^{ABC}D_t^\sigma g(t)](s) = \frac{q(\sigma)}{1-\sigma} \left\{ \sigma \Gamma(\sigma+1) M_\sigma \left(-\frac{1}{1-\sigma} V^\sigma \right) \right\} \times [ST(g(t)) - g(0)]. \quad (3)$$

Definition 2. The Laplace transform (LT) of the Caputo fractional derivative of a function $g(t)$ of order $\sigma > 0$ is defined as

$$L [{}_{0}^C D_t^\sigma g(t)] = s^\sigma g(s) - \sum_{\sigma=0}^{n-1} g^{(\sigma)}(0) s^{\sigma-1}. \quad (4)$$

Definition 3. The LT of the function $t^{\sigma_1-1} E_{\sigma,\sigma_1}(\pm \mu t^\sigma)$ is defined as

$$L [t^{\sigma_1-1} E_{\sigma,\sigma_1}(\pm \mu t^\sigma)] = \frac{s^{\sigma-\sigma_1}}{s^\sigma \mp \mu}, \quad (5)$$

where E_{σ,σ_1} is the two-parameter ML function with $\sigma, \sigma_1 > 0$. Further, the ML function satisfies the following equation [17].

$$E_{\sigma,\sigma_1}(f) = f E_{\sigma,\sigma+\sigma_1}(f) + \frac{1}{\Gamma(\sigma_1)}. \quad (6)$$

Definition 4. Suppose that $g(t)$ is continuous on an open interval (a, b) , then the fractal-fractional integral of $g(t)$ of order σ having ML type kernel and given by

$$\begin{aligned} {}^{FFM} J_{0,t}^{\sigma,\sigma_1}(g(t)) &= \frac{\sigma \sigma_1}{AB(\sigma) \Gamma(\sigma_1)} \int_0^t s^{\sigma_1-1} g(s) (t-s)^\sigma ds \\ &+ \frac{\sigma_1 (1-\sigma) t^{\sigma_1-1} g(t)}{AB(\sigma)}. \end{aligned} \quad (7)$$

3. Model Formulation

We will study the giving up smoking model for overall population at time t . We separate the population into 5 groups, potential smokers $P(t)$, occasional smokers $L(t)$, heavy smokers $S(t)$, temporary quitters $Q(t)$, and smokers who quit permanently $R(t)$ specified by $T(t) = P(t) + L(t) + S(t) + Q(t) + R(t)$. Due to smoking, the chances of other diseases like heart attack, stroke, especially lung cancer, throat, mouth, esophagus, and pancreas are increased [24]. The model is developed as follows

$$\frac{dP}{dt} = a(1-P) - bPS, \quad (8)$$

$$\frac{dL}{dt} = -aL + bPL - cLS, \quad (9)$$

$$\frac{dS}{dt} = -(a+d)S + cLS + fQ, \quad (10)$$

$$\frac{dQ}{dt} = -(a+f)Q + d(1-e)S, \quad (11)$$

$$\frac{dR}{dt} = -aR + edS. \quad (12)$$

With the initial conditions

$$P(0) = \delta_1, \quad (13)$$

$$L(0) = \delta_2, \quad (14)$$

$$S(0) = \delta_3, \quad (15)$$

$$Q(0) = \delta_4, \quad (16)$$

$$R(0) = \delta_5. \quad (17)$$

The rate of change between potential smoker and occasional smokers is represented by b , a represents the rate of natural death, the rate of occasional smoker and temporary smokers by c , the rate of change between quitters and smoker is shown by f , the rate of giving up smoking is shown by d , fraction of temporary giving up smoker is represented by $(1-e)$ (at the rate of d), e shows the remaining fraction of smokers who give up smoking forever (at a rate d).

4. Qualitative Analysis

By substituting the values of parameters in given system of differential equations and the rate of change with respect to time is zero, we get

$$a(1-P) - bPS = 0, \quad (18)$$

$$-aL + bPS - cLS = 0, \quad (19)$$

$$-(a+d)S + cLS + fQ = 0, \quad (20)$$

$$-(a+f)Q + d(1-e)S = 0, \quad (21)$$

$$-aR + edS = 0. \quad (22)$$

By simplifying the above equations, we get disease-free equilibrium, denoted by E_0 , i.e., $E_0 = (1, 0, 0, 0, 0)$.

Endemic equilibrium is found in terms of one of the infected compartment, denoted by E_1 , i.e., $E_1 = (P^*, L^*, S^*, Q^*, R^*)$ where $P^* = a/a + bS^*$, $L^* = ab/(a + bS^*)(a + cS^*)$, $Q^* = d(1-e)S^*/a + f$, $R^* = edS^*/a$.

4.1. Stability Analysis and Reproductive Number. It is important to find the verge conditions to check the status of population, whether the disease persist or dies out. In case of disease free equilibrium point, $R_0 < 1$, which shows

that the disease will die out. In case of endemic equilibrium, $R_0 > 1$. Consider the Jacobian matrix (JM) as

$$J = \begin{bmatrix} -a - bS & 0 & -bP & 0 & 0 \\ bS & -a - cS & bP - cL & 0 & 0 \\ 0 & cS & -a - d + cL & f & 0 \\ 0 & 0 & d(1 - e) & -a - f & 0 \\ 0 & 0 & ed & 0 & -a \end{bmatrix}. \quad (23)$$

Since the JM is $J = F - V$ where

We know that $K = FV^{-1}$ and using the relation $|K - \lambda I| = 0$ solving on mathematica for the Eigenvalue λ , which represents the reproductive number R_0 , i.e.,

$$R_0 = \frac{df(1 - e)}{(a + d)(a + f)}. \quad (24)$$

Hence, $R_0 = 0.431034 < 1$, according to the given parameter values.

Theorem 5. *The disease free equilibrium E_0 is locally asymptotically stable for $R_0 < 1$, if $\text{Re}(\lambda) < 0$, otherwise, unstable.*

Proof. E_0 of the given system is locally asymptotically stable if $\text{Re}(\lambda) < 0$ where λ can be evaluated from the relation $|J_0 - \lambda I| = 0$.

By using the relation $|J_0 - \lambda I| = 0$, we get. $\text{Re}(\lambda)$ as

$$\lambda_1 = -a, \lambda_2 = \frac{1}{2} \left[-2a - d - f - \sqrt{d^2 + 2df - 4def + f^2} \right] < 0, \quad (25)$$

$$\lambda_3 = \frac{1}{2} \left[-2a - d - f + \sqrt{d^2 + 2df - 4def + f^2} \right] < 0\lambda_1. \quad (26)$$

All the Eigenvalues are negative real parts which represent that the given system is locally asymptotically stable. \square

Theorem 6. *When the reproductive number $R_0 > 1$, the endemic equilibrium points E_1 of the PLSQR model is globally asymptotically stable.*

Proof. The Lyapunov function can be written as

$$\begin{aligned} M(P^*, L^*, S^*, Q^*, R^*) &= \left(P - P^* - P^* \log \frac{P}{P^*} \right) + \left(L - L^* - L^* \log \frac{L}{L^*} \right) \\ &+ \left(S - S^* - S^* \log \frac{S}{S^*} \right) + \left(Q - Q^* - Q^* \log \frac{Q}{Q^*} \right) \\ &+ \left(R - R^* - R^* \log \frac{R}{R^*} \right). \end{aligned} \quad (27)$$

Therefore, applying the derivative respect to t on both sides yields

$$\begin{aligned} \frac{dM}{dt} &= \dot{M} = \left(\frac{P - P^*}{P} \right) \dot{P} + \left(\frac{L - L^*}{L} \right) \dot{L} + \left(\frac{S - S^*}{S} \right) \dot{S} \\ &+ \left(\frac{Q - Q^*}{Q} \right) \dot{Q} + \left(\frac{R - R^*}{R} \right) \dot{R}. \end{aligned} \quad (28)$$

Now, we can write their values for derivatives as follows

$$\begin{aligned} \frac{dM}{dt} &= \dot{M} = \left(\frac{P - P^*}{P} \right) (a(1 - P) - bPS) \\ &+ \left(\frac{L - L^*}{L} \right) (-aL + bPS - cLS) \\ &+ \left(\frac{S - S^*}{S} \right) (-(a + d)S + cLS + fQ) \\ &+ \left(\frac{Q - Q^*}{Q} \right) (-(a + f)Q + d(1 - e)S) \\ &+ \left(\frac{R - R^*}{R} \right) (-aR + edS). \end{aligned} \quad (29)$$

Putting $P = P - P^*$, $L = L - L^*$, $S = S - S^*$, $Q = Q - Q^*$, $R = R - R^*$ leads to

$$\begin{aligned} \frac{dM}{dt} &= \left(\frac{P - P^*}{P} \right) (a(1 - (P - P^*)) - b(P - P^*)(S - S^*)) \\ &+ \left(\frac{L - L^*}{L} \right) (-a(L - L^*) + b(P - P^*)(S - S^*) \\ &- c(L - L^*)(S - S^*)) + \left(\frac{S - S^*}{S} \right) (-(a + d)(S - S^*) \\ &+ c(L - L^*)(S - S^*) + f(Q - Q^*)) \\ &+ \left(\frac{Q - Q^*}{Q} \right) (-(a + f)(Q - Q^*) + d(1 - e)(S - S^*)) \\ &+ \left(\frac{R - R^*}{R} \right) (-a(R - R^*) + ed(S - S^*)). \end{aligned} \quad (30)$$

We can organize the above as follows

$$\begin{aligned} \frac{dM}{dt} &= a - \left(\frac{P^*}{P} \right) a - \frac{1}{P} (P - P^*)^2 - \frac{b}{P} (P - P^*)^2 (S) \\ &+ \frac{b}{P} (P - P^*)^2 (S^*) - \frac{a}{L} (L - L^*)^2 + bPS - bP^*S \\ &+ bP^*S^* - bPS^* - \left(\frac{L^*}{L} \right) bPS + \left(\frac{L^*}{L} \right) bP^*S \\ &- \left(\frac{L^*}{L} \right) bP^*S^* + \left(\frac{L^*}{L} \right) bPS^* - \frac{C}{L} (L - L^*)^2 (S) \\ &+ \frac{C}{L} (L - L^*)^2 (S^*) - \frac{(a + d)}{S} (S - S^*)^2 + \frac{c}{S} (L)(S - S^*)^2 \end{aligned}$$

$$\begin{aligned}
 & -\frac{c}{S}(L^*)(S-S^*)^2 + f(Q) - f(Q^*) - \left(\frac{S^*}{S}\right)fQ \\
 & + \left(\frac{S^*}{S}\right)fQ^* - \frac{(a+f)}{Q}(Q-Q^*)^2 + d(1-e)(S) \\
 & - d(1-e)(S^*) - d(1-e)(S)\left(\frac{Q^*}{Q}\right) \\
 & + d(1-e)(S^*)\left(\frac{Q^*}{Q}\right) - \frac{a}{R}(R-R^*)^2 + ed(S) - ed(S^*) \\
 & - ed(S)\left(\frac{R^*}{R}\right) + ed(S^*)\left(\frac{R^*}{R}\right).
 \end{aligned} \tag{31}$$

To avoid the complexity, the above can be written as

$$\frac{dM}{dt} = \Sigma - \Omega, \tag{32}$$

where

$$\begin{aligned}
 \Sigma = & a + \frac{b}{P}(P-P^*)^2(S^*) + bPS + bP^*S^* + \left(\frac{L^*}{L}\right)bP^*S \\
 & + \left(\frac{L^*}{L}\right)bPS^* + \frac{C}{L}(L-L^*)^2(S^*) + \frac{c}{S}(L)(S-S^*)^2 + f \\
 & + \left(\frac{S^*}{S}\right)fQ^* + d(1-e)(S) + d(1-e)(S^*)\left(\frac{Q^*}{Q}\right) \\
 & + ed(S) + ed(S^*)\left(\frac{R^*}{R}\right),
 \end{aligned} \tag{33}$$

$$\begin{aligned}
 \Omega = & \left(\frac{P^*}{P}\right)a + \frac{1}{P}(P-P^*)^2 + \frac{b}{P}(P-P^*)^2(S) + \frac{a}{L}(L-L^*)^2 \\
 & + bP^*S + bPS^* + \left(\frac{L^*}{L}\right)bPS + \left(\frac{L^*}{L}\right)bP^*S^* \\
 & + \frac{C}{L}(L-L^*)^2(S) + \frac{(a+d)}{S}(S-S^*)^2 + \frac{c}{S}(L^*)(S-S^*)^2 \\
 & + fQ^* + \left(\frac{S^*}{S}\right)fQ + \frac{(a+f)}{Q}(Q-Q^*)^2 + d(1-e)(S^*) \\
 & + d(1-e)(S)\left(\frac{Q^*}{Q}\right) + \frac{a}{R}(R-R^*)^2 + ed(S^*) \\
 & + ed(S)\left(\frac{R^*}{R}\right).
 \end{aligned} \tag{34}$$

It is concluded that if $\Sigma < \Omega$, this yields, $dM/dt < 0$, however when, $P = P^*, L = L^*, S = S^*, Q = Q^*, R = R^*$

$$0 = \Sigma - \Omega \Rightarrow \frac{dM}{dt} = 0. \tag{35}$$

We can see that the largest compact invariant set for the suggested model in

$$\left\{ (P^*, L^*, S^*, Q^*, R^*) \in \Gamma; \frac{dM}{dt} = 0 \right\}, \tag{36}$$

is the point $\{E_1\}$ the endemic equilibrium of the considered model. By the help of the Lasalles invariance concept, it follows that E_1 is globally asymptotically stable in Γ if $\Sigma < \Omega$. \square

5. Atangana-Baleanu Caputo Sense with Mittag-Leffler Kernel

By applying AB fractional derivative of order σ and $\sigma \in (0, 1]$, into ML kernel, then, the system (8) becomes

$${}_{0}^{ABC}D_t^\sigma P = a(1-P) - bPS, \tag{37}$$

$${}_{0}^{ABC}D_t^\sigma L = -aL + bPL - cLS, \tag{38}$$

$${}_{0}^{ABC}D_t^\sigma S = -(a+d)S + cLS + fQ, \tag{39}$$

$${}_{0}^{ABC}D_t^\sigma Q = -(a+f)Q + d(1-e)S, \tag{40}$$

$${}_{0}^{ABC}D_t^\sigma R = -aR + edS. \tag{41}$$

The initial conditions associated with the system (37) are

$$P(0) = \delta_1, \tag{42}$$

$$L(0) = \delta_2, \tag{43}$$

$$S(0) = \delta_3, \tag{44}$$

$$Q(0) = \delta_4, \tag{45}$$

$$R(0) = \delta_5. \tag{46}$$

We will discuss the numerical value of solution of system for different values of ρ . With the help of iterative method and the Padé approximation results are obtained.

We use the values of the parameters $a = 0.04$, $b = 0.23$, $c = 0.3$, $d = 0.2$, $e = 0.4$, and $f = 0.25$. The initial conditions are given by $P(0) = 0.60301$, $L(0) = 0.24000$, $S(0) = 0.10628$, $Q(0) = 0.03260$, and $R(0) = 0.01811$.

Taking ST on both sides of (37), we get

$$\begin{aligned}
 & \frac{q(\sigma)\sigma\Gamma(\sigma+1)}{1-\sigma}N_\sigma\left(-\frac{1}{1-\sigma}V^\sigma\right)ST\{P(t) - P(0)\} \\
 & = ST[a(1-P) - bPS],
 \end{aligned} \tag{47}$$

$$\begin{aligned}
 & \frac{q(\sigma)\sigma\Gamma(\sigma+1)}{1-\sigma}N_\sigma\left(-\frac{1}{1-\sigma}V^\sigma\right)ST\{L(t) - L(0)\} \\
 & = ST[-aL + bPL - cLS],
 \end{aligned} \tag{48}$$

$$\begin{aligned}
 & \frac{q(\sigma)\sigma\Gamma(\sigma+1)}{1-\sigma}N_\sigma\left(-\frac{1}{1-\sigma}V^\sigma\right)ST\{S(t) - S(0)\} \\
 & = ST[-(a+d)S + cLS + fQ],
 \end{aligned} \tag{49}$$

$$\begin{aligned} & \frac{q(\sigma)\sigma\Gamma(\sigma+1)}{1-\sigma} N_\sigma\left(-\frac{1}{1-\sigma}V^\sigma\right) \text{ST}\{Q(t) - Q(0)\} \\ & = \text{ST}[-(a+f)Q + d(1-e)S], \end{aligned} \quad (50)$$

$$\begin{aligned} & \frac{q(\sigma)\sigma\Gamma(\sigma+1)}{1-\sigma} N_\sigma\left(-\frac{1}{1-\sigma}V^\sigma\right) \text{ST}\{R(t) - R(0)\} \\ & = \text{ST}[-aR + edS]. \end{aligned} \quad (51)$$

Rearranging, we get

$$\begin{aligned} \text{ST}(P(t)) = P(0) + & \frac{1-\sigma}{q(\sigma)\sigma\Gamma(\sigma+1)N_\sigma(-(1/1-\sigma)V^\sigma)} \\ & \cdot \text{ST}[a(1-P) - bPS], \end{aligned} \quad (52)$$

$$\begin{aligned} \text{ST}(L(t)) = L(0) + & \frac{1-\sigma}{q(\sigma)\sigma\Gamma(\sigma+1)N_\sigma(-(1/1-\sigma)V^\sigma)} \\ & \cdot \text{ST}[-aL + bPL - cLS], \end{aligned} \quad (53)$$

$$\begin{aligned} \text{ST}(S(t)) = S(0) + & \frac{1-\sigma}{q(\sigma)\sigma\Gamma(\sigma+1)N_\sigma(-(1/1-\sigma)V^\sigma)} \\ & \cdot \text{ST}[-(a+d)S + cLS + fQ], \end{aligned} \quad (54)$$

$$\begin{aligned} \text{ST}(Q(t)) = Q(0) + & \frac{1-\sigma}{q(\sigma)\sigma\Gamma(\sigma+1)N_\sigma(-(1/1-\sigma)V^\sigma)} \\ & \cdot \text{ST}[-(a+f)Q + d(1-e)S], \end{aligned} \quad (55)$$

$$\begin{aligned} \text{ST}(R(t)) = R(0) + & \frac{1-\sigma}{q(\sigma)\sigma\Gamma(\sigma+1)N_\sigma(-(1/1-\sigma)V^\sigma)} \\ & \cdot \text{ST}[-aR + edS]. \end{aligned} \quad (56)$$

Now taking inverse ST on both sides of equation (52), we get

$$\begin{aligned} P(t) = P(0) + \text{ST}^{-1} & \left[\frac{1-\sigma}{q(\sigma)\sigma\Gamma(\sigma+1)N_\sigma(-(1/1-\sigma)V^\sigma)} \right. \\ & \left. \cdot \text{ST}\{a(1-P) - bPS\} \right], \end{aligned} \quad (57)$$

$$\begin{aligned} L(t) = L(0) + \text{ST}^{-1} & \left[\frac{1-\sigma}{q(\sigma)\sigma\Gamma(\sigma+1)N_\sigma(-(1/1-\sigma)V^\sigma)} \right. \\ & \left. \cdot \text{ST}\{-aL + bPL - cLS\} \right], \end{aligned} \quad (58)$$

$$\begin{aligned} S(t) = S(0) + \text{ST}^{-1} & \left[\frac{1-\sigma}{q(\sigma)\sigma\Gamma(\sigma+1)N_\sigma(-(1/1-\sigma)V^\sigma)} \right. \\ & \left. \cdot \text{ST}\{-(a+d)S + cLS + fQ\} \right], \end{aligned} \quad (59)$$

$$\begin{aligned} Q(t) = Q(0) + \text{ST}^{-1} & \left[\frac{1-\sigma}{q(\sigma)\sigma\Gamma(\sigma+1)N_\sigma(-(1/1-\sigma)V^\sigma)} \right. \\ & \left. \cdot \text{ST}\{-(a+f)Q + d(1-e)S\} \right], \end{aligned} \quad (60)$$

$$\begin{aligned} R(t) = R(0) + \text{ST}^{-1} & \left[\frac{1-\sigma}{q(\sigma)\sigma\Gamma(\sigma+1)N_\sigma(-(1/1-\sigma)V^\sigma)} \right. \\ & \left. \cdot \text{ST}\{-aR + edS\} \right]. \end{aligned} \quad (61)$$

We next attain the following recursive formula.

$$\begin{aligned} P_{(n+1)}(t) = P_n(0) + \text{ST}^{-1} & \left[\frac{1-\sigma}{q(\sigma)\sigma\Gamma(\sigma+1)N_\sigma(-(1/1-\sigma)V^\sigma)} \right. \\ & \left. \cdot \text{ST}\{a(1-P_n) - bP_nS_n\} \right], \end{aligned} \quad (62)$$

$$\begin{aligned} L_{(n+1)}(t) = L_n(0) + \text{ST}^{-1} & \left[\frac{1-\sigma}{q(\sigma)\sigma\Gamma(\sigma+1)N_\sigma(-(1/1-\sigma)V^\sigma)} \right. \\ & \left. \cdot \text{ST}\{-aL_n + bP_nL_n - cL_nS_n\} \right], \end{aligned} \quad (63)$$

$$\begin{aligned} S_{n+1}(t) = S_n(0) + \text{ST}^{-1} & \left[\frac{1-\sigma}{q(\sigma)\sigma\Gamma(\sigma+1)N_\sigma(-(1/1-\sigma)V^\sigma)} \right. \\ & \left. \cdot \text{ST}\{-(a+d)S_n + cL_nS_n + fQ_n\} \right], \end{aligned} \quad (64)$$

$$\begin{aligned} Q_{(n+1)}(t) = Q_n(0) + \text{ST}^{-1} & \left[\frac{1-\sigma}{q(\sigma)\sigma\Gamma(\sigma+1)N_\sigma(-(1/1-\sigma)V^\sigma)} \right. \\ & \left. \cdot \text{ST}\{-(a+f)Q_n + d(1-e)S_n\} \right], \end{aligned} \quad (65)$$

$$\begin{aligned} R_{(n+1)}(t) = R_n(0) + \text{ST}^{-1} & \left[\frac{1-\sigma}{q(\sigma)\sigma\Gamma(\sigma+1)N_\sigma(-(1/1-\sigma)V^\sigma)} \right. \\ & \left. \cdot \text{ST}\{-aR_n + edS_n\} \right]. \end{aligned} \quad (66)$$

And the solution of (62) is

$$P(t) = \lim_{n \rightarrow \infty} P_n(t), \quad (67)$$

$$L(t) = \lim_{n \rightarrow \infty} L_n(t), \quad (68)$$

$$S(t) = \lim_{n \rightarrow \infty} S_n(t), \quad (69)$$

$$Q(t) = \lim_{n \rightarrow \infty} Q_n(t), \quad (70)$$

$$R(t) = \lim_{n \rightarrow \infty} R_n(t). \quad (71)$$

Theorem 7. Let $(X, |\cdot|)$ be a Banach space and H a self-map of X satisfying

$$\|H_x - H_r\| \leq \theta \|X - H_x\| + \theta \|x - r\|, \quad (72)$$

for all $x, r \in X$, and $0 \leq \theta < 1$. Suppose that H is Picard H -stable. Suppose that system (62), we have

$$P_{n+1}(t) = P_n(0) + ST^{-1} \left\{ \frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \cdot ST[a(1 - P_n) - bP_nS_n] \right\}, \quad (73)$$

$$L_{n+1}(t) = L_n(0) + ST^{-1} \left\{ \frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \cdot ST[-aL_n + bP_nL_n - cL_nS_n] \right\}, \quad (74)$$

$$S_{n+1}(t) = S_n(0) + ST^{-1} \left\{ \frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \cdot ST[-(a + d)S_n + cL_nS_n + fQ_n] \right\}, \quad (75)$$

$$Q_{n+1}(t) = Q_n(0) + ST^{-1} \left\{ \frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \cdot ST[-(a + f)Q_n + d(1 - e)S_n] \right\}, \quad (76)$$

$$R_{n+1}(t) = R_n(0) + ST^{-1} \left\{ \frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \cdot ST[-aR_n + edS_n] \right\}, \quad (77)$$

where $1 - \sigma/q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)$ is the fractional Lagrange multiplier.

Theorem 8. Define K be a self-map is given by

$$K \left[P_{(n+1)}(t) \right] = P_n(0) + ST^{-1} \cdot \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \cdot ST\{a(1 - P_n) - bP_nS_n\} \right], \quad (78)$$

$$K \left[L_{(n+1)}(t) \right] = L_n(0) + ST^{-1} \cdot \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \cdot ST\{-aL_n + bP_nL_n - cL_nS_n\} \right], \quad (79)$$

$$K \left[S_{n+1}(t) \right] = S_n(0) + ST^{-1} \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \cdot ST\{-(a + d)S_n + cL_nS_n + fQ_n\} \right], \quad (80)$$

$$Q_{(n+1)}(t) = Q_n(0) + ST^{-1} \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \cdot ST\{-(a + f)Q_n + d(1 - e)S_n\} \right], \quad (81)$$

$$R_{(n+1)}(t) = R_n(0) + ST^{-1} \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \cdot ST\{-aR_n + edS_n\} \right], \quad (82)$$

$$K \left[S_{n+1}(t) \right] = S_n(0) + ST^{-1} \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \cdot ST\{-(a + d)S_n + cL_nS_n + fQ_n\} \right], \quad (83)$$

$$K \left[Q_{(n+1)}(t) \right] = Q_n(0) + ST^{-1} \cdot \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \cdot ST\{-(a + f)Q_n + d(1 - e)S_n\} \right], \quad (84)$$

$$K \left[R_{(n+1)}(t) \right] = R_n(0) + ST^{-1} \cdot \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \cdot ST\{-aR_n + edS_n\} \right]. \quad (85)$$

Proof. In first step, we will show that K is fixed point

$$\forall (m, n) \in N \times N, \quad (86)$$

$$\begin{aligned}
& K(P_n(t)) - K(P_m(t)) \\
&= P_n(t) - P_m(t) + ST^{-1} \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \right. \\
&\quad \cdot ST\{a(1 - P_n) - bP_nS_n\} \left. \right] - ST^{-1} \\
&\quad \cdot \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \right. \\
&\quad \cdot ST\{a(1 - P_m) - bP_mS_m\} \left. \right], \tag{87}
\end{aligned}$$

$$\begin{aligned}
& K(L_n(t)) - K(L_m(t)) \\
&= L_n(t) - L_m(t) + ST^{-1} \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \right. \\
&\quad \cdot ST\{-aL_n + bP_nL_n - cL_nS_n\} \left. \right] - ST^{-1} \\
&\quad \cdot \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} ST \right. \\
&\quad \cdot \{-aL_m + bP_mL_m - cL_mS_m\} \left. \right], \tag{88}
\end{aligned}$$

$$\begin{aligned}
& K(S_n(t)) - K(S_m(t)) \\
&= S_n(t) - S_m(t) + ST^{-1} \\
&\quad \cdot \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \right. \\
&\quad \cdot ST\{-(a + d)S_n + cL_nS_n + fQ_n\} \left. \right] - ST^{-1} \tag{89} \\
&\quad \cdot \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \right. \\
&\quad \cdot ST\{-(a + d)S_m + cL_mS_m + fQ_m\} \left. \right],
\end{aligned}$$

$$\begin{aligned}
& K(Q_n(t)) - K(Q_m(t)) \\
&= Q_n(t) - Q_m(t) + ST^{-1} \\
&\quad \cdot \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \right. \\
&\quad \cdot ST\{-(a + f)Q_n + d(1 - e)S_n\} \left. \right] \tag{90} \\
&\quad - ST^{-1} \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \right. \\
&\quad \cdot ST\{-(a + f)Q_m + d(1 - e)S_m\} \left. \right],
\end{aligned}$$

$$\begin{aligned}
& K(R_n(t)) - K(R(t)) \\
&= R_n(t) - R_m(t) + ST^{-1} \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \right. \\
&\quad \cdot ST\{-aR_n + edS_n\} \left. \right] - ST^{-1} \\
&\quad \cdot \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \right. \\
&\quad \cdot ST\{-aR_m + edS_m\} \left. \right]. \tag{91}
\end{aligned}$$

Applying the properties of the norm and also using the triangular inequality, we obtain

$$\begin{aligned}
& \|K(P_n(t)) - K(P_m(t))\| \\
&\leq \|P_n(t) - P_m(t)\| ST^{-1} \\
&\quad \cdot \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \right. \\
&\quad \cdot ST\{\|a(1 - (P_n - P_m))\| + \|b(P_nS_n - P_mS_m)\|\} \left. \right], \tag{92}
\end{aligned}$$

$$\begin{aligned}
& \|K(L_n(t)) - K(L_m(t))\| \\
&\leq \|L_n(t) - L_m(t)\| + ST^{-1} \\
&\quad \cdot \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \right. \tag{93} \\
&\quad \cdot ST\{-\|a(L_n - L_m)\| + \|b(P_nL_n - P_mL_m)\| \\
&\quad \left. + \|-c(L_nS_n - L_mS_m)\|\} \right],
\end{aligned}$$

$$\begin{aligned}
& \|K(S_n(t)) - K(S_m(t))\| \\
&\leq \|S_n(t) - S_m(t)\| + ST^{-1} \\
&\quad \cdot \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \right. \tag{94} \\
&\quad \cdot ST\{-\|(a + d)(S_n - S_m)\| + \|c(L_nS_n - L_mS_m)\| \\
&\quad \left. + \|f(Q_n - Q_m)\|\} \right],
\end{aligned}$$

$$\begin{aligned}
& \|K(Q_n(t)) - K(Q_m(t))\| \\
&\leq \|Q_n(t) - Q_m(t)\| + ST^{-1} \\
&\quad \cdot \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \right. \\
&\quad \cdot ST\{-\|(a + f)(Q_n - Q_m)\| + \|d(1 - e)(S_n - S_m)\|\} \left. \right], \tag{95}
\end{aligned}$$

$$\begin{aligned}
 & \|K(R_n(t)) - K(R_m(t))\| \\
 & \leq \|R_n(t) - R_m(t)\| + ST^{-1} \\
 & \cdot \left[\frac{1 - \sigma}{q(\sigma)\sigma\Gamma(\sigma + 1)N_\sigma(-(1/1 - \sigma)V^\sigma)} \right. \\
 & \left. \cdot ST\{\| -a(R_n - R_m)\| + \|ed(S_n - S_m)\|\} \right]. \quad (96)
 \end{aligned}$$

K fulfills the conditions associated with Theorem 6, when

$$\theta = (0, 0, 0, 0, 0), \theta \begin{cases} \|P_n(t) - P_m(t)\| \times \| -P_n(t) - P_m(t)\| + a\|(1 - (P_n - P_m))\| - b\|(P_n S_n - P_m S_m)\|, \\ \times \|L_n(t) - L_m(t)\| \times \| -L_n(t) - L_m(t)\| - a\|L_n(t) - L_m(t)\| + b\|(P_n L_n - P_m L_m)\| - c\|(L_n S_n - L_m S_m)\|, \\ \times \|S_n(t) - S_m(t)\| \times \| -S_n(t) - S_m(t)\| - (a + d)\|P_n - P_m\| + c\|(L_n S_n - L_m S_m)\| + f\|(Q_n - Q_m)\|, \\ \times \|Q_n(t) - Q_m(t)\| \times \| -Q_n(t) - Q_m(t)\| - (a + f)\|Q_n - Q_m\| + d(1 - e)\|d(1 - e)(S_n - S_m)\|, \\ \times \|R_n(t) - R_m(t)\| \times \| -R_n(t) - R_m(t)\| - a\|R_n - R_m\| + ed\|(S_n - S_m)\|. \end{cases} \quad (97)$$

And we add that K is Picard K -stable. \square

So, the following operators are considered

Theorem 9. Prove that system (27) has special solution is unique.

$$\theta(0, 0, 0, 0, 0), \theta = \begin{cases} a(1 - p) - bps, \\ -aL + bPL - cLS, \\ -(a + d)S + cLS + fQ, \\ -(a + f)Q + d(1 - e)S, \\ -aR + edS. \end{cases} \quad (99)$$

Proof. Let H be the Hilbert space defined as $H = L^2((p, q) \times (0, T))$ where

$$h : (p, q) \times (0, T) \longrightarrow \mathbb{R}, \iint gh dg dh < \infty. \quad (98)$$

We establish the inner product of

$$P((P_{11} - P_{12}, L_{21} - L_{22}, S_{31} - S_{32}, Q_{41} - Q_{42}, R_{51} - R_{52}), (V_1, V_2, V_3, V_4, V_5)). \quad (100)$$

where $(S_{11} - S_{12}, I_{21} - I_{22}, A_{31} - A_{32}, T_{41} - T_{42}, R_{51} - R_{52})$ are the special solutions of the system. By using the inner function and the norm, we have

$$\begin{aligned}
 & \{a - a(P_{11} - P_{12}) - b(P_{11} - P_{12})(S_{31} - S_{32})\} \\
 & \leq a\|V_1\| + a\|P_{11} - P_{12}\| + b\|P_{11} - P_{12}\| \\
 & \times \|S_{31} - S_{32}\| \|V_1\|, \quad (101)
 \end{aligned}$$

$$\begin{aligned}
 & \{-a(L_{21} - L_{22}) + b(P_{11} - P_{12})(L_{21} - L_{22}) \\
 & - c(L_{21} - L_{22})(S_{31} - S_{32})\} \\
 & \leq a\|L_{21} - L_{22}\| \|V_2\| + b\|P_{11} - P_{12}\| \|L_{21} - L_{22}\| \|V_2\| \\
 & + c\|L_{21} - L_{22}\| \|S_{31} - S_{32}\| \|V_2\|, \quad (102)
 \end{aligned}$$

$$\begin{aligned}
 & \{-(a + d)(S_{31} - S_{32}) + c(L_{21} - L_{22})(S_{31} - S_{32}) \\
 & + f(Q_{41} - Q_{42})\} \\
 & \leq (a + d)\|S_{31} - S_{32}\| \|V_3\| + c\|L_{21} - L_{22}\| \|S_{31} - S_{32}\| \|V_3\| \\
 & + f\|Q_{31} - Q_{32}\| \|V_3\|, \quad (103)
 \end{aligned}$$

$$\begin{aligned}
 & \{-(a + f)(Q_{41} - Q_{42}) + d(1 - e)(S_{31} - S_{32})\} \\
 & \leq (a + f)\|Q_{31} - Q_{32}\| \|V_4\| + d(1 - e)\|S_{31} - S_{32}\| \|V_4\|, \quad (104)
 \end{aligned}$$

$$\begin{aligned}
 & \{-a(R_{51} - R_{52}) + ed(S_{31} - S_{32})\} \\
 & \leq a\|R_{31} - R_{32}\| \|V_5\| + ed\|S_{31} - S_{32}\| \|V_5\|. \quad (105)
 \end{aligned}$$

Due to large number of $e_1, e_2, e_3, e_4,$ and $e_5,$ both solutions converge to the exact solution. Applying the

topological idea, we have the very small positive five parameters (χ_{e_1} , χ_{e_2} , χ_{e_3} , χ_{e_4} , and χ_{e_5}).

$$\|P - P_{11}\|, \|P - P_{12}\| \leq \frac{\chi_{e_1}}{\omega}, \quad (106)$$

$$\|L - L_{21}\|, \|L - L_{22}\| \leq \frac{\chi_{e_2}}{\varsigma}, \quad (107)$$

$$\|S - S_{31}\|, \|S - S_{32}\| \leq \frac{\chi_{e_3}}{\nu}, \quad (108)$$

$$\|Q - Q_{41}\|, \|Q - Q_{42}\| \leq \frac{\chi_{e_4}}{\kappa}, \quad (109)$$

$$\|R - R_{51}\|, \|R - R_{52}\| \leq \frac{\chi_{e_5}}{\varrho}, \quad (110)$$

$$\omega = 5(a + a\|P_{11} - P_{12}\| + b\|P_{11} - P_{12}\|\|S_{31} - S_{32}\|)\|V_1\|, \quad (111)$$

$$\varsigma = 5(a\|L_{21} - L_{22}\| + b\|P_{11} - P_{12}\|\|L_{21} - L_{22}\| + c\|L_{21} - L_{22}\|\|S_{31} - S_{32}\|)\|V_2\|, \quad (112)$$

$$\nu = 5((a + d)\|S_{31} - S_{32}\| + c\|L_{21} - L_{22}\|\|S_{31} - S_{32}\| + f\|Q_{31} - Q_{32}\|)\|V_3\|, \quad (113)$$

$$\kappa = 5((a + f)\|Q_{31} - Q_{32}\| + d(1 - e)\|S_{31} - S_{32}\|)\|V_4\|, \quad (114)$$

$$\varrho = 5(a\|R_{31} - R_{32}\| + ed\|S_{31} - S_{32}\|)\|V_5\|. \quad (115)$$

But, it is obvious that

$$(a + a\|P_{11} - P_{12}\| + b\|P_{11} - P_{12}\|\|S_{31} - S_{32}\|) \neq 0, \quad (116)$$

$$(a\|L_{21} - L_{22}\| + b\|P_{11} - P_{12}\|\|L_{21} - L_{22}\| + c\|L_{21} - L_{22}\|\|S_{31} - S_{32}\|) \neq 0, \quad (117)$$

$$((a + d)\|S_{31} - S_{32}\| + c\|L_{21} - L_{22}\|\|S_{31} - S_{32}\| + f\|Q_{31} - Q_{32}\|) \neq 0, \quad (118)$$

$$((a + f)\|Q_{31} - Q_{32}\| + d(1 - e)\|S_{31} - S_{32}\|) \neq 0, \quad (119)$$

$$(a\|R_{31} - R_{32}\| + ed\|S_{31} - S_{32}\|) \neq 0, \quad (120)$$

where $\|V_1\|, \|V_2\|, \|V_3\|, \|V_4\|, \|V_5\| \neq 0$.

Therefore, we have

$$\|P_{11} - P_{12}\| = 0, \quad (121)$$

$$\|L_{21} - L_{22}\| = 0, \quad (122)$$

$$\|S_{31} - S_{32}\| = 0, \quad (123)$$

$$\|Q_{41} - Q_{42}\| = 0, \quad (124)$$

$$\|R_{51} - R_{52}\| = 0. \quad (125)$$

Which yields that

$$P_{11} = P_{12}, \quad (126)$$

$$L_{21} = L_{22}, \quad (127)$$

$$S_{31} = S_{32}, \quad (128)$$

$$Q_{41} = Q_{42}, \quad (129)$$

$$R_{51} = R_{52}. \quad (130)$$

This completes the proof of uniqueness. \square

6. New Numerical Scheme

We define the AT proposed scheme for fractional derivative model (37) for the smoking epidemic. For this purpose, we suppose that we obtain the following results for system (37)

$$\begin{aligned} P(t) - P(0) &= \frac{(1 - \sigma)}{ABC(\sigma)} \{a - aP(t) - bP(t)S(t)\} \\ &+ \frac{\sigma}{\Gamma(\sigma) \times ABC(\sigma)} \\ &\cdot \int_0^t \{a - aP(\tau) - bP(\tau)S(\tau)\}(t - \tau)^{\sigma-1} d\tau, \end{aligned} \quad (131)$$

$$\begin{aligned} L(t) - L(0) &= \frac{(1 - \sigma)}{ABC(\sigma)} \{-aL(t) + bP(t)L(t) - cL(t)S(t)\} \\ &+ \frac{\sigma}{\Gamma(\sigma) \times ABC(\sigma)} \int_0^t \{-aL(\tau) + bP(\tau)L(\tau) \\ &- cL(\tau)S(\tau)\}(t - \tau)^{\sigma-1} d\tau, \end{aligned} \quad (132)$$

$$\begin{aligned} S(t) - S(0) &= \frac{(1 - \sigma)}{ABC(\sigma)} \{-(a + d)S(t) + cL(t)S(t) + fQ(t)\} \\ &+ \frac{\sigma}{\Gamma(\sigma) \times ABC(\sigma)} \int_0^t \{-(a + d)S(\tau) + cL(\tau)S(\tau) \\ &+ fQ(\tau)\}(t - \tau)^{\sigma-1} d\tau, \end{aligned} \quad (133)$$

$$\begin{aligned} Q(t) - Q(0) &= \frac{(1 - \sigma)}{ABC(\sigma)} \{-(a + f)Q(t) + d(1 - e)S(t)\} \\ &+ \frac{\sigma}{\Gamma(\sigma) \times ABC(\sigma)} \int_0^t \{-(a + f)Q(\tau) \\ &+ d(1 - e)S(\tau)\}(t - \tau)^{\sigma-1} d\tau, \end{aligned} \quad (134)$$

$$\begin{aligned} R(t) - R(0) &= \frac{(1 - \sigma)}{ABC(\sigma)} \{-aR(t) + edS(t)\} \\ &+ \frac{\sigma}{\Gamma(\sigma) \times ABC(\sigma)} \\ &\cdot \int_0^t \{-aR(\tau) + edS(\tau)\}(t - \tau)^{\sigma-1} d\tau. \end{aligned} \quad (135)$$

At a given point $t_{n+1}, n = 0, 1, 2, 3, \dots$, the above equation is reformulated as

$$P(t) - P(0) = \frac{(1 - \sigma)}{ABC(\sigma)} \{a - aP(t_n) - bP(t_n)S(t_n)\} + \frac{\sigma}{\Gamma(\sigma) \times ABC(\sigma)} \cdot \int_0^t \{a - aP(\tau) - bP(\tau)S(\tau)\} (t - \tau)^{\sigma-1} d\tau, \quad (136)$$

$$L(t) - L(0) = \frac{(1 - \sigma)}{ABC(\sigma)} \{-aL(t_n) + bP(t_n)L(t_n) - cL(t_n)S(t_n)\} + \frac{\sigma}{\Gamma(\sigma) \times ABC(\sigma)} \int_0^t \{-aL(\tau) + bP(\tau)L(\tau) - cL(\tau)S(\tau)\} (t - \tau)^{\sigma-1} d\tau, \quad (137)$$

$$S(t) - S(0) = \frac{(1 - \sigma)}{ABC(\sigma)} \{-(a + d)S(t_n) + cL(t_n)S(t_n) + fQ(t_n)\} + \frac{\sigma}{\Gamma(\sigma) \times ABC(\sigma)} \int_0^t \{-(a + d)S(\tau) + cL(\tau)S(\tau) + fQ(\tau)\} (t - \tau)^{\sigma-1} d\tau, \quad (138)$$

$$Q(t) - Q(0) = \frac{(1 - \sigma)}{ABC(\sigma)} \{-(a + f)Q(t_n) + d(1 - e)S(t_n)\} + \frac{\sigma}{\Gamma(\sigma) \times ABC(\sigma)} \int_0^t \{-(a + f)Q(\tau) + d(1 - e)S(\tau)\} (t - \tau)^{\sigma-1} d\tau, \quad (139)$$

$$R(t) - R(0) = \frac{(1 - \sigma)}{ABC(\sigma)} \{-aR(t_n) + edS(t_n)\} + \frac{\sigma}{\Gamma(\sigma) \times ABC(\sigma)} \int_0^t \{-aR(\tau) + edS(\tau)\} (t - \tau)^{\sigma-1} d\tau. \quad (140)$$

Also, we have

$$P(t_{n+1}) - P(0) = \frac{(1 - \sigma)}{ABC(\sigma)} \{a - aP(t_n) - bP(t_n)S(t_n)\} + \frac{\sigma}{\Gamma(\sigma) \times ABC(\sigma)} \sum_{j=0}^n \int_{t_j}^{t_{j+1}} \{a - aP(\tau) - bP(\tau)S(\tau)\} (t_{n+1} - \tau)^{\sigma-1} d\tau, \quad (141)$$

$$L(t_{n+1}) - L(0) = \frac{(1 - \sigma)}{ABC(\sigma)} \{-aL(t_n) + bP(t_n)L(t_n) - cL(t_n)S(t_n) - cL(\tau)S(\tau)\} + \frac{\sigma}{\Gamma(\sigma) \times ABC(\sigma)} \sum_{j=0}^n \int_{t_j}^{t_{j+1}} \{-aL(\tau) + bP(\tau)L(\tau) - cL(\tau)S(\tau)\} (t_{n+1} - \tau)^{\sigma-1} d\tau, \quad (142)$$

$$S(t_{n+1}) - S(0) = \frac{(1 - \sigma)}{ABC(\sigma)} \{-(a + d)S(t_n) + cL(t_n)S(t_n) + fQ(t_n)\} + \frac{\sigma}{\Gamma(\sigma) \times ABC(\sigma)} \sum_{j=0}^n \int_{t_j}^{t_{j+1}} \{-(a + d)S(\tau) + cL(\tau)S(\tau) + fQ(\tau)\} (t_{n+1} - \tau)^{\sigma-1} d\tau, \quad (143)$$

$$Q(t_{n+1}) - Q(0) = \frac{(1 - \sigma)}{ABC(\sigma)} \{-(a + f)Q(t_n) + d(1 - e)S(t_n)\} + \frac{\sigma}{\Gamma(\sigma) \times ABC(\sigma)} \sum_{j=0}^n \int_{t_j}^{t_{j+1}} \{-(a + f)Q(\tau) + d(1 - e)S(\tau)\} (t_{n+1} - \tau)^{\sigma-1} d\tau, \quad (144)$$

$$R(t_{n+1}) - R(0) = \frac{(1 - \sigma)}{ABC(\sigma)} \{-aR(t_n) + edS(t_n)\} + \frac{\sigma}{\Gamma(\sigma) \times ABC(\sigma)} \cdot \sum_{j=0}^n \int_{t_j}^{t_{j+1}} \{-aR(\tau) + edS(\tau)\} (t_{n+1} - \tau)^{\sigma-1} d\tau. \quad (145)$$

By using equation (133), we have

$$P_{n+1} = P_0 + \frac{(1 - \sigma)}{ABC(\sigma)} \{a - aP(t_n) - bP(t_n)S(t_n)\} + \frac{\sigma}{\Gamma(\sigma) \times ABC(\sigma)} \cdot \sum_{j=0}^n \left(\frac{\{a - aP_j - bP_jS_j\}}{h} \times \int_{t_j}^{t_{j+1}} (\tau - t_{j-1})(t_{n+1} - \tau)^{\sigma-1} d\tau - \frac{\{a - aP_{j-1} - bP_{j-1}S_{j-1}\}}{h} \times \int_{t_j}^{t_{j+1}} (\tau - t_j)(t_{n+1} - \tau)^{\sigma-1} d\tau \right), \quad (146)$$

$$\begin{aligned}
L_{n+1} = & L_0 + \frac{(1-\sigma)}{\text{ABC}(\sigma)} \{-aL(t_n) + bP(t_n)L(t_n) - cL(t_n)S(t_n)\} \\
& + \frac{\sigma}{\Gamma(\sigma) \times \text{ABC}(\sigma)} \sum_{j=0}^n \left(\frac{\{-aL_j + bP_jL_j - cL_jS_j\}}{h} \right. \\
& \times \int_{t_j}^{t_{j+1}} (\tau - t_{j-1})(t_{n+1} - \tau)^{\sigma-1} d\tau \\
& \left. - \frac{\{-aL_{j-1} + bP_{j-1}L_{j-1} - cL_{j-1}S_{j-1}\}}{h} \times \int_{t_j}^{t_{j+1}} (\tau - t_j)(t_{n+1} - \tau)^{\sigma-1} d\tau \right), \tag{147}
\end{aligned}$$

$$\begin{aligned}
S_{n+1} = & S_0 + \frac{(1-\sigma)}{\text{ABC}(\sigma)} \{-(a+d)S(t_n) + cL(t_n)S(t_n) + fQ(t_n)\} \\
& + \frac{\sigma}{\Gamma(\sigma) \times \text{ABC}(\sigma)} \sum_{j=0}^n \left(\frac{\{-(a+d)S_j + cL_jS_j + fQ_j\}}{h} \right. \\
& \times \int_{t_j}^{t_{j+1}} (\tau - t_{j-1})(t_{n+1} - \tau)^{\sigma-1} d\tau \\
& \left. - \frac{\{-(a+d)S_{j-1} + cL_{j-1}S_{j-1} + fQ_{j-1}\}}{h} \times \int_{t_j}^{t_{j+1}} (\tau - t_j)(t_{n+1} - \tau)^{\sigma-1} d\tau \right), \tag{148}
\end{aligned}$$

$$\begin{aligned}
Q_{n+1} = & Q_0 + \frac{(1-\sigma)}{\text{ABC}(\sigma)} \{-(a+f)Q(t_n) + d(1-e)S(t_n)\} \\
& + \frac{\sigma}{\Gamma(\sigma) \times \text{ABC}(\sigma)} \sum_{j=0}^n \left(\frac{\{-(a+f)Q_j + d(1-e)S_j\}}{h} \right. \\
& \times \int_{t_j}^{t_{j+1}} (\tau - t_{j-1})(t_{n+1} - \tau)^{\sigma-1} d\tau \\
& \left. - \frac{\{-(a+f)Q_{j-1} + d(1-e)S_{j-1}\}}{h} \times \int_{t_j}^{t_{j+1}} (\tau - t_j)(t_{n+1} - \tau)^{\sigma-1} d\tau \right), \tag{149}
\end{aligned}$$

$$\begin{aligned}
R_{n+1} = & R_0 + \frac{(1-\sigma)}{\text{ABC}(\sigma)} \{-aR(t_n) + edS(t_n)\} \\
& + \frac{\sigma}{\Gamma(\sigma) \times \text{ABC}(\sigma)} \sum_{j=0}^n \left(\frac{\{-aR_j + edS_j\}}{h} \right. \\
& \times \int_{t_j}^{t_{j+1}} (\tau - t_{j-1})(t_{n+1} - \tau)^{\sigma-1} d\tau \\
& \left. - \frac{\{-aR_{j-1} + edS_{j-1}\}}{h} \times \int_{t_j}^{t_{j+1}} (\tau - t_j)(t_{n+1} - \tau)^{\sigma-1} d\tau \right). \tag{150}
\end{aligned}$$

By using equations (133) and (144), we get

$$\begin{aligned}
P_{n+1} = & P_0 + \frac{(1-\sigma)}{\text{ABC}(\sigma)} \{a - aP(t_n) - bP(t_n)S(t_n)\} \\
& + \frac{\sigma}{\text{ABC}(\sigma)} \sum_{j=0}^n \left(\frac{h^\sigma \{a - aP_j - bP_jS_j\}}{\Gamma(\sigma+2)} \times \{p_1p_2 - p_3p_4\} \right. \\
& \left. - \frac{h^\sigma \{a - aP_{j-1} - bP_{j-1}S_{j-1}\}}{\Gamma(\sigma+2)} \times \{p_5 - p_3p_6\} \right), \tag{151}
\end{aligned}$$

$$\begin{aligned}
L_{n+1} = & L_0 + \frac{(1-\sigma)}{\text{ABC}(\sigma)} \{-aL(t_n) + bP(t_n)L(t_n) - cL(t_n)S(t_n)\} \\
& + \frac{\sigma}{\text{ABC}(\sigma)} \sum_{j=0}^n \left(\frac{h^\sigma \{-aL_j + bP_jL_j - cL_jS_j\}}{\Gamma(\sigma+2)} \times \{p_1p_2 - p_3p_4\} \right. \\
& \left. - \frac{h^\sigma \{-aL_{j-1} + bP_{j-1}L_{j-1} - cL_{j-1}S_{j-1}\}}{\Gamma(\sigma+2)} \times \{p_5 - p_3p_6\} \right), \tag{152}
\end{aligned}$$

$$\begin{aligned}
S_{n+1} = & S_0 + \frac{(1-\sigma)}{\text{ABC}(\sigma)} \{-(a+d)S(t_n) + cL(t_n)S(t_n) + fQ(t_n)\} \\
& + \frac{\sigma}{\text{ABC}(\sigma)} \sum_{j=0}^n \left(\frac{h^\sigma \{-(a+d)S_j + cL_jS_j + fQ_j\}}{\Gamma(\sigma+2)} \times \{p_1p_2 - p_3p_4\} \right. \\
& \left. - \frac{h^\sigma \{-(a+d)S_{j-1} + cL_{j-1}S_{j-1} + fQ_{j-1}\}}{\Gamma(\sigma+2)} \times \{p_5 - p_3p_6\} \right), \tag{153}
\end{aligned}$$

$$\begin{aligned}
Q_{n+1} = & Q_0 + \frac{(1-\sigma)}{\text{ABC}(\sigma)} \{-(a+f)Q(t_n) + d(1-e)S(t_n)\} \\
& + \frac{\sigma}{\text{ABC}(\sigma)} \sum_{j=0}^n \left(\frac{h^\sigma \{-(a+f)Q_j + d(1-e)S_j\}}{\Gamma(\sigma+2)} \times \{p_1p_2 - p_3p_4\} \right. \\
& \left. - \frac{h^\sigma \{-(a+f)Q_{j-1} + d(1-e)S_{j-1}\}}{\Gamma(\sigma+2)} \times \{p_5 - p_3p_6\} \right), \tag{154}
\end{aligned}$$

$$\begin{aligned}
R_{n+1} = & R_0 + \frac{(1-\sigma)}{\text{ABC}(\sigma)} \{-aR(t_n) + edS(t_n)\} \\
& + \frac{\sigma}{\text{ABC}(\sigma)} \sum_{j=0}^n \left(\frac{h^\sigma \{-aR_j + edS_j\}}{\Gamma(\sigma+2)} \times \{p_1p_2 - p_3p_4\} \right. \\
& \left. - \frac{h^\sigma \{-aR_{j-1} + edS_{j-1}\}}{\Gamma(\sigma+2)} \times \{p_5 - p_3p_6\} \right). \tag{155}
\end{aligned}$$




FIGURE 1: Simulation of $P(t)$ with ABC fractional-order scheme.




FIGURE 2: Simulation of $L(t)$ with ABC fractional-order scheme.

where

$$p_1 = (m + 1 - j)^\sigma, \tag{156}$$

$$p_2 = (m - j + 2 + \sigma), \tag{157}$$

$$p_3 = (m - j)^\sigma, \tag{158}$$

$$p_4 = (m - j + 2 + 2\sigma), \tag{159}$$

$$p_5 = (m + 1 - j)^{\sigma+1}, \tag{160}$$

$$p_6 = (m - j + 1 + \sigma). \tag{161}$$

7. Numerical Results and Discussion

A mathematical study of the nonlinear epidemic model of smoking has been presented. For checking of parameters effects on smoking dynamical model, relatively some numerical simulations according to the value of the parameters are




FIGURE 3: Simulation of $S(t)$ with ABC fractional-order scheme.




FIGURE 4: Simulation of $Q(t)$ with ABC fractional-order scheme.

accomplished to confirm the effect of the fractional derivative on the different compartments. We got mathematical consequences of the model for different fractional values with the help of the ATM. If we note the impacts of variables on the dynamics of the model of fractional order, the end-

time value of the given parameter can be observed in various numerical ways. We can observe that the results of fractional value are more accurate as compared to classical derivatives. Desired results can be achieved to analyze the epidemic that occurs due to smoking. The graphs of the approximate




FIGURE 5: Simulation of $R(t)$ with ABC fractional-order scheme.

solutions are given in Figures 1–5 against different fractional-order α . $P(t)$ and $R(t)$ start decreasing by decreasing the fractional values while $L(t)$, $S(t)$, and $Q(t)$ start increasing by decreasing fractional values which can be easily observed in Figures 1–5. When the fractional values decrease then the behavior approaches steady-state in all figures, which shows that the solution will be more effective by decreasing the fractional values.

8. Conclusion

The advanced numerical scheme of fractional differential equation has been investigated in this article for smoking model by using ATM. With the help of fixed point theory uniqueness and stability of the smoking, the model has been examined. System is analyzed qualitatively to verify the steady state position of the dynamic. Proposed system is analyzed locally; also, global stability has been made using first derivative of Lyapunov. The arbitrary derivative of fractional order has been taken in ATM with no singular kernel. Effective results are obtained for the proposed model. Also discussed some theoretical results and proved the efficiency of the proposed techniques. Numerical simulations are carried out to check the actual behavior of the dynamic using the advanced ATM. These results will be helpful to understand further analysis and to control different outbreak caused by smoking.

Data Availability

No data were used to support this study.

Conflicts of Interest

The authors state that there is no conflict of interest.

Authors' Contributions

All authors contributed equally and significantly to writing this article. All authors read and approved the final manuscript.

References

- [1] C. S. Chou and A. Friedman, "Introduction," in *Introduction to Mathematical Biology. Springer Undergraduate Texts in Mathematics and Technology*, Springer, Cham, 2016.
- [2] Centers for Disease Control and Prevention (CDC), "Annual smoking –attributable mortality, years of potential life lost, and economic cost–united state 1995–1999," *MMWR. Morbidity and Mortality Weekly Report*, vol. 51, no. 14, pp. 300–303, 2002.
- [3] A. Jemal, M. M. Center, C. Desantis, and E. M. Ward, "Global patterns of cancer incidence and mortality rates and trends," *Cancer Epidemiology, Biomarkers & Prevention*, vol. 19, pp. 1893–1907, 2010.
- [4] D. Hoffmann, I. Hoffmann, and K. El-Bayoumy, "The less harmful cigarette: a controversial issue. A tribute to Ernst L Wynder," *Chemical Research in Toxicology*, vol. 14, pp. 767–790, 2001.
- [5] A. Charlton, "Medicinal use of tobacco in history," *Journal of the Royal Society of Medicine*, vol. 97, pp. 292–296, 2004.
- [6] P. Garodia, H. Ichikawa, N. Malani, G. Sethi, and B. B. Aggarwal, "From ancient medicine to modern medicine: ayurvedic concepts of health and their role in inflammation and cancer," *Journal of the Society for Integrative Oncology*, vol. 5, no. 1, pp. 25–27, 2007.

- [7] J. Handelsman, J. Conway, M. Boylan, and R. Turtle, "Testicular function in potential sperm donors: normal ranges and the effects of smoking and varicocele," *International Journal of Andrology*, vol. 7, no. 5, pp. 369–382, 1984.
- [8] A. Atangana and B. S. T. Alkahtani, "Analysis of the Keller–Segel model with a fractional derivative without singular kernel," *Entropy*, vol. 17, no. 12, pp. 4439–4453, 2015.
- [9] A. Alsaedi, J. J. Nieto, and V. Venkatesh, "Fractional electrical circuits," *Mechanical Engineering*, vol. 7, no. 12, 2015.
- [10] M. Caputo and M. Fabrizio, "A new definition of fractional derivative without singular kernel," *Progress in Fractional Differentiation and Applications (PFDA)*, vol. 1, no. 2, pp. 1–13, 2015.
- [11] J. Losada and J. J. Nieto, "Properties of a new fractional derivative without singular kernel," *Progress in Fractional Differentiation and Applications (PFDA)*, vol. 1, no. 2, pp. 87–92, 2015.
- [12] A. A. Tateishi, H. V. Ribeiro, and E. K. Lenzi, "The role of fractional time-derivative operators on anomalous diffusion," *Frontiers of Physics*, vol. 5, p. 52, 2017.
- [13] D. Kumar, J. Singh, M. al Qurashi, and D. Baleanu, "Analysis of logistic equation pertaining to a new fractional derivative with non-singular kernel," *Advances in Mechanical Engineering*, vol. 9, no. 2, 2017.
- [14] K. M. Owolabi and A. Atangana, "Analysis and application of new fractional Adams–Bashforth scheme with Caputo–Fabrizio derivative," *Chaos, Solitons and Fractals*, vol. 105, pp. 111–119, 2017.
- [15] D. Kumar, F. Tchier, J. Singh, and D. Baleanu, "An efficient computational technique for fractal vehicular traffic flow," *Entropy*, vol. 20, p. 259, 2018.
- [16] J. Singh, D. Kumar, and D. Baleanu, "New aspects of fractional Biswas–Milovic model with Mittag–Leffler law," *Mathematical Modelling of Natural Phenomena*, vol. 14, no. 3, p. 303, 2019.
- [17] K. Diethelm, *The Analysis of Fractional Differential Equations: An Application-Oriented Exposition Using Differential Operators of Caputo Type*, Springer, Berlin, 2010.
- [18] A. Atangana, "Blind in a commutative world: simple illustrations with functions and chaotic attractors," *Chaos, Solitons and Fractals*, vol. 114, pp. 347–363, 2018.
- [19] H. Bulut, H. M. Baskonus, and F. B. M. Belgecam, "The analytic solutions of some fractional ordinary differential equation by Sumudu transform method," *Abstr. Appl. Anal.*, vol. 2013, article 203875, 6 pages, 2013.
- [20] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier, Amsterdam, 2006.
- [21] A. Atangana and I. Koca, "Chaos in a simple nonlinear system with Atangana–Baleanu derivatives with fractional order," *Chaos, Solitons and Fractals*, vol. 89, pp. 447–454, 2016.
- [22] L. L. Huang, D. Baleanu, and S. D. Wu Zeng, "A new application of the fractional logistic map," *Romanian Journal of Physics*, vol. 61, pp. 1172–1179, 2016.
- [23] G. Zaman, "Qualitative behavior of giving up smoking model," *Bulletin of the Malaysian Mathematical Sciences Society*, vol. 34, pp. 403–415, 2011.
- [24] J. Singh, D. Kumar, M. A. Qurashi, and D. Baleanu, "A new fractional model for giving up smoking dynamics," *Advances in Difference Equations*, vol. 2017, no. 1, Article ID 88, 2017.
- [25] A. E. Nava, "Risk factors associated with recent transmission of tuberculosis: systematic review and meta-analysis," *The International Journal of Tuberculosis and Lung Disease*, vol. 13, pp. 17–26, 2009.
- [26] A. K. Pokhrel, M. N. Bates, S. C. Verma, H. S. Joshi, C. T. Sreeramareddy, and K. R. Smith, "Tuberculosis and indoor biomass and kerosene use in Nepal: a case–control study," *Environmental Health Perspectives*, vol. 118, pp. 558–564, 2010.
- [27] P. Narasimhan, J. Wood, C. R. MacIntyre, and D. Mathai, "Risk factors for tuberculosis," *Pulmonary Medicine*, vol. 2013, Article ID 828939, 11 pages, 2013.
- [28] J. I. Irunde, L. S. Luboobi, and Y. Nkansah-Gyekye, "Modeling the effect of tobacco smoking on the in-host dynamics of hiv/aids," *Journal of Mathematical and Computational Science*, vol. 6, no. 3, p. 406, 2016.
- [29] S. S. Islam, T. B. Rumi, S. L. Kabir et al., "Zoonotic tuberculosis knowledge and practices among cattle handlers in selected districts of Bangladesh," *PLoS Neglected Tropical Diseases*, vol. 15, no. 4, article e0009394, 2021.