

First checklist of the fruit flies of Morocco, including new records (Diptera, Tephritidae)

Younes El Harym¹, Boutaina Belqat¹

¹ Department of Biology, Faculty of Sciences, University Abdelmalek Essaâdi, Tétouan

Corresponding author: Boutaina Belqat (belqat@gmail.com)

Academic editor: M. De Meyer | Received 22 April 2017 | Accepted 16 July 2017 | Published 26 September 2017

<http://zoobank.org/AD5C2A87-50BD-42FF-812E-9DC81EB33FBC>

Citation: El Harym Y, Belqat B (2017) First checklist of the fruit flies of Morocco, including new records (Diptera, Tephritidae). ZooKeys 702: 137–171. <https://doi.org/10.3897/zookeys.702.13368>

Abstract

The first checklist of the Tephritidae of Morocco, containing 59 species, is presented here. Out of 38 species collected during the present project, three (*Campiglossa martii* (Becker, 1908), *Tephritis divisa* (Rondani, 1871), and *Terellia* sp. near *longicauda*) present new records for North Africa, and ten (*Carpomyia incompleta* (Becker, 1903), *Chaetorellia conjuncta* (Becker, 1913), *Chetostoma curvinerve* Rondani, 1856, *Dacus frontalis* (Becker, 1922), *D. longistylus* (Wiedemann, 1830), *Dioxyna sororcula* (Wiedemann, 1830), *Ensina sonchi* (Linnaeus, 1767), *Myopites inulaedysenteriae* Blot, 1827, *M. stylatus* Fabricius, 1794, and *Tephritis vespertina* (Loew, 1844)) are new for Morocco.

Keywords

Anti Atlas, Beni Snassen, High Atlas, Middle Atlas, North Africa, Rif, Sahara

Introduction

Tephritidae is one of the largest families of the acalyprate Diptera, and more than 4300 valid species are known in the world (Norrbom 2004). Prior to this study, the Tephritid fauna of Morocco has not been the subject of focused research. Only few

studies from North Africa were devoted to the fruit flies (e.g., Tunisia, Arambourg and Soria 1961; Algeria, Hering 1937; Morocco, Algeria and Tunisia, Heršman and Dirlbek 2006). Some new species were described from Algeria (Dirlbek and Dirlbekova 1976; Hendel 1927; Hering 1938; Séguy 1934a) and Morocco (Becker and Stein 1913; Séguy 1930, 1941).

The Tephritidae of Morocco have been also published upon among some Diptera studies (Wiedemann 1824; Becker and Stein 1913; Séguy (1930, 1934b, 1941, 1949, 1953). Faunal records from all relevant publications have been registered in the Catalogue of the Tephritidae of the World (Norrbom et al. 1999). A total of 46 species have been previously recorded from Morocco. New findings increase the number of fruit flies known from Morocco to 59.

The material reported in the present article was mostly collected in Morocco between 2013–2016 in 99 field expeditions over mountainous areas (Rif, Beni Snassen (eastern Morocco), Middle Atlas, Anti Atlas) and the arid area of Morocco (Sahara). Thirty-eight species of Tephritidae were identified from this material, of which ten species are new to Morocco, and three species are new to North Africa. One of the main achievements of this project is the first checklist of Tephritidae from Morocco, containing 59 species, which is presented here.

Materials and methods

Collecting methods

A total of 924 specimens of Tephritidae were collected by sweeping nets and malaise traps, or reared from flower heads or fruits of plants, examined, and preserved in 70% ethanol. When terminalia were necessary to confirm species identity, they were prepared by boiling the abdomen in 10% KOH for 20 minutes at 95°C and preserved in glycerine. Species were recognised according to the identification keys of Hendel (1927), Freidberg and Kugler (1989), White and Elson-Harris (1992), Merz (1994), Korneyev and White (1992, 2000). The systematic classification is based on White and Elson-Harris (1992), Norrbom et al. (1999), White et al. (2000) and White and Goodger (2009).

A list of 99 sampling sites, with coordinates and altitudes, is given in Table 2, and the locations of the sites are shown in Map 1, elaborated using the logiciel GisArc (Geographic Information System, version 9.3). All the specimens are deposited in the collection of Diptera of the Department of Biology, Faculty of Sciences, University Abdelmalek Essaâdi, Tétouan.

The following checklist summarizes the species inventory presently known from Morocco, and their worldwide distribution.

Table I. Species (in alphabetical order) of Tephritidae known from North Africa.

Species	Morocco	Algeria	Tunisia	Libya	Egypt
<i>Acanthiophilus helianthi</i> (Rossi, 1794)	X*	X	X		
<i>Aciura coryli</i> Rossi, 1794	X				X
<i>Bactrocera cucurbitae</i> (Coquillett, 1899)					X
<i>Bactrocera oleae</i> Rossi, 1790	X**				X
<i>Bactrocera zonata</i> Saunders, 1842				X	
<i>Campiglossa hofferi</i> (Dirlbek & Dirlbekova, 1976)		X			
<i>Campiglossa martii</i> (Becker, 1908)	X****				
<i>Campiglossa producta</i> Loew, 1844	X*	X	X		
<i>Capitites ramulosa</i> (Loew, 1844)	X	X			X
<i>Capparimyia savastani</i> (Martelli, 1911)	X	X	X	X	X
<i>Carpomya incompleta</i> (Becker, 1903)	X***				X
<i>Carpomya pardalina</i> (Bigot, 1891)					X
<i>Ceratitis capitata</i> (Wiedemann, 1824)	X*	X	X	X	X
<i>Chaetorellia conjuncta</i> (Becker, 1913)	X***				X
<i>Chaetorellia hestia</i> (Hering, 1937)	X**	X			
<i>Chaetorellia succinea</i> Costa, 1844	X				X
<i>Chaetostomella cylindrica</i> Robineau-Desvoidy, 1830	X	X	X		
<i>Chetostoma curvinerve</i> Rondani, 1856	X***	X?	X?	X?	X?
<i>Dacus annulatus</i> (Becker, 1903)					X
<i>Dacus ciliatus</i> (Loew, 1862)					X
<i>Dacus frontalis</i> (Becker, 1922)	X***	X		X	X
<i>Dacus longistylus</i> (Wiedemann, 1830)	X***			X	X
<i>Dacus semisphaereus</i> (Becker, 1903)					X
<i>Dacus sexmaculatus</i> (Walker, 1871)					X
<i>Desmella conyzae</i> (Frauenfeld, 1857)					X
<i>Desmella rostellata</i> (Séguy, 1941)	X				
<i>Dioxyna bidentis</i> (Robineau-Desvoidy, 1830)	X?	X?	X?	X?	X?
<i>Dioxyna sororcula</i> (Wiedemann, 1830)	X***	X	X		
<i>Ensina sonchi</i> (Linnaeus, 1830)	X***	X			
<i>Euaresta bullans</i> (Wiedemann, 1830)	X*	X	X		
<i>Euarestella iphionae</i> (Efflatoun, 1924)					X
<i>Euarestella kugleri</i> (Freidberg, 1974)					X
<i>Euarestella pninae</i> (Freidberg, 1981)					X
<i>Euleia heraclei</i> (Linnaeus, 1758)	X**	X			
<i>Euleia marmorea</i> (Fabricius, 1805)	X				
<i>Goniurellia lacerata</i> (Becker, 1913)					X
<i>Goniurellia longicauda</i> (Freidberg, 1980)	X**	X	X	X	X
<i>Goniurellia persignata</i> (Freidberg, 1980)	X*				X
<i>Goniurellia spinifera</i> (Freidberg, 1980)					X
<i>Hyalotephritis planiscutellata</i> (Becker, 1903)					X
<i>Hypenidium graecum</i> (Loew, 1862)	X				
<i>Katonaia aida</i> (Hering, 1938)					X
<i>Metasphenisca gracilipes</i> (Loew, 1862)					X
<i>Metasphenisca haematopoda</i> (Bezzi, 1924)					X
<i>Metasphenisca negeviana</i> (Freidberg, 1974)					X

Species	Morocco	Algeria	Tunisia	Libya	Egypt
<i>Myopites apicatus</i> (Freidberg, 1979)	X				
<i>Myopites boghariensis</i> (Séguy, 1934)		X			
<i>Myopites cypriacus</i> Hering, 1938	X				
<i>Myopites inulaedysenteriae</i> Blot, 1827	X***	X			
<i>Myopites stylatus</i> Fabricius, 1794	X***	X	X?	X?	X?
<i>Myopites variofasciatus</i> (Becker, 1903)					X
<i>Neoceratitis efflatouni</i> (Hendel, 1931)					X
<i>Notomma mutilum</i> Bezzi, 1923					X
<i>Oedaspis daphne</i> (Séguy, 1930)	X				
<i>Oedaspis farinosa</i> (Hendel, 1927)		X			
<i>Oedaspis fissa</i> Loew, 1862		X			
<i>Oedaspis multifasciata</i> (Hering, 1937)	X	X			
<i>Oedaspis trotteriana</i> (Bezzi, 1913)	X	X		X	X
<i>Oedaspis villeneuvei</i> (Bezzi, 1913)		X		X	X
<i>Oxyaciura tibialis</i> Robineau-Desvoidy 1830	X*				
<i>Oxyna superflava</i> (Freidberg, 1974)	X				X
<i>Paradesis augur</i> (Frauenfeld, 1857)		X	X		X
<i>Paraspheniscus debskii</i> (Efflatoun, 1924)					X
<i>Schistopterum moebiusi</i> (Becker, 1903)					X
<i>Spathulina acroleuca</i> (Schiner, 1868)					X
<i>Spathulina sricula</i> (Rondani, 1856)	X*		X		
<i>Sphaeniscus filiolus</i> (Loew, 1869)	X**				X
<i>Sphenella marginata</i> (Fallen, 1814)	X*				X
<i>Tephritis bimaculata</i> (Freidberg, 1981)					X
<i>Tephritis dioscorea</i> Loew, 1856	X				
<i>Tephritis divisa</i> (Rondani, 1871)	X****				
<i>Tephritis formosa</i> (Loew, 1844)	X**				
<i>Tephritis jabeliae</i> (Freidberg, 1981)					X
<i>Tephritis leontodontis</i> De Géer, 1776	X	X?	X?	X?	X?
<i>Tephritis matricariae</i> (Loew, 1844)	X*				X
<i>Tephritis nigricauda</i> Loew 1856	X**	X			X
<i>Tephritis postica</i> (Loew, 1844)	X**	X	X		
<i>Tephritis praecox</i> (Loew, 1844)	X**	X			
<i>Tephritis pulchra</i> Loew, 1844	X	X			
<i>Tephritis simplex</i> (Loew, 1844)	X	X	X		
<i>Tephritis stictica</i> Loew, 1862	X				
<i>Tephritis spreta</i> (Loew, 1861)					X
<i>Tephritis theryi</i> (Séguy, 1930)	X				
<i>Tephritis truncata</i> (Loew, 1844)			X		
<i>Tephritis vespertina</i> (Loew, 1844)	X***		X		
<i>Tephritomyia lauta</i> (Loew, 1869)	X**		X		X
<i>Terellia colon</i> Meigen, 1826	X	?	X	?	?
<i>Terellia fuscicornis</i> (Loew, 1844)	X	X?	X?	X?	X?
<i>Terellia longicauda</i> Meigen, 1838	X				
<i>Terellia luteola</i> (Wiedemann, 1830)			X		X
<i>Terellia oasis</i> (Hering, 1938)		X			

Species	Morocco	Algeria	Tunisia	Libya	Egypt
<i>Terellia serratulae</i> (Linnaeus, 1758)	X*	X	X		
<i>Terellia</i> sp. near <i>longicauda</i>	X****				
<i>Terellia vectensis</i> Collin, 1937		X			
<i>Terellia virens</i> (Loew, 1846)	X**		X		
<i>Trupanea amoena</i> (Frauenfeld, 1857)	X**		X		
<i>Trupanea guimari</i> (Becker, 1908)	X**	X			
<i>Trupanea pseudoamoena</i> (Freidberg, 1974)					X
<i>Trupanea repleta</i> (Bezzi, 1918)					X
<i>Trupanea stellata</i> (Fuesslin, 1775)	X**		X		
<i>Urelliosoma desertorum</i> (Efflatoun, 1924)					X
<i>Urelliosoma pulcherrimum</i> (Efflatoun, 1924)		X	X		X
<i>Urophora calcitrupae</i> (White & Korneyev, 1989)					X
<i>Urophora mauritanica</i> (Macquart, 1851)	X**	X		X	
<i>Urophora pauperata</i> Zaitzev, 1945					X
<i>Urophora phaeocera</i> (Hering, 1961)					X
<i>Urophora quadrifasciata</i> (Meigen, 1826)	X	X	X		
<i>Urophora quadrifasciata algerica</i> (Hering, 1941)		X	X		
<i>Urophora solstitialis</i> Linnaeus 1758	X	X	X		

X****: new species for North Africa; X***: new species for Morocco; X**: new species for one or more of the geographic regions of Morocco; X*: species known from Morocco and collected by the authors; ?: species considered, in the literature, present in North Africa without specifying country.

Table 2. Sampling sites (in alphabetical order) harboring the species collected in Morocco with localities, geographical coordinates, and altitudes.

Province	Station	Locality	Altitude (m)	Geographical coordinates
Rif				
Al Hoceïma	1. Affluent Tarmast	Parc National d'Al Hoceïma	168	N35°10.666', W004°03.088'
	2. Oued Azila	Azila, Jbel Tidghine	1601	N34°52.028', W004°32.609'
Chefchaouen	3. Aïn Afersiw	Mezine	746	N35°05.945', W005°20.439'
	4. Aïn El Ma Bared	Bouzthate	1267	N35°00.333', W005°12.105'
	5. Aïn El Malâab	Parc National Talassemiane	1278	N35°05.509', W005°09.443'
	6. Aïn El Maounzil	Parc National Talassemiane	1106	N35°04.577', W005°10.406'
	7. Aïn Tiouila	Parc National Talassemiane	1502	N35°07.194', W005°09.978'
	8. Bab El Karne	Tamakoute	1248	N34°58.510', W005°11.838'
	9. Cascade Chrafate	Chrafate	820	N35°03.997', W005°06.434'
	10. Daya Afrate	Tanaqoub	600	N35°05.634', W005°26.028'
	11. Daya El Ânassar	Bab Berred	1183	N35°00.788', W004°57.419'
	12. Daya El Birdiyel	Ânasar	1300	N35°01.089', W004°59.477'
	13. Douar Abou Boubnar (Marabout Sidi Gile)	Parc National Talassemiane	1247	N35°10.812', W005°07.500'

Chefchaouen	14. Douar Ouslaf	Parc National Talassemtnane	625	N35°13.787', W005°11.376'
	15. Douar Tamakoute	Tamakoute	1089	N34°58.609', W005°12.743'
	16. Maison forestière	Parc National Talassemtnane	1674	N35°08.076', W005°08.262'
	17. Mizoghar	El Khizana	1005	N35°02.725', W005°12.969'
	18. Oued Abou Bnar	Parc National Talassemtnane	1260	N35°10.854', W005°07.840'
	19. Oued Derdara	Derdara	400	N35°06.484', W005°17.147'
	20. Oued El Kanar	Stehate	50	N35°17.233', W004°59.639'
	21. Oued El Kelâa	Akchour	510	N35°13.514', W005°08.553'
	22. Oued Jbara	Tanaqoub	600	N35°05.634', W005°26.028'
	23. Oued Jnane Niche	Jnane Niche	36	N35°17.019', W004°51.233'
	24. Oued Majjou	Village Majjou	799	N35°06.186', W005°10.935'
	25. Oued Mezine	Mezine	778	N35°06.104', W005°21.177'
	26. Oued Sidi Ben Sâada	Derdara	220	N35°03.921', W005°19.895'
	27. Oued Sidi Yahya Aârab	Sidi Yahya Aârab	62	N35°17.545', W004°53.503'
Larache	28. Affluent Oued Amsemlil	Jbel Bouhachem	1130	N35°15.657', W005°26.159'
	29. Daya Mtahen	Jbel Bouhachem	966	N35°16.195', W005°26.158'
	30. Daya Tazia	Route Moulay Abdessalam	721	N35°20.814', W005°33.139'
	31. Oued Tkaraâ	Jbel Bouhachem	959	N35°16.063', W005°25.829'
M'diq-Fnideq	32. Ksar Rimal	Kabila	11	N35°43.806', W005°20.509'
Tanger-Assilah	33. Daya Aïn Jdioui	Aïn Jdioui	5	N35°34.074', W005°55.499'
	34. Daya El Hajjami	Barrage 9 Avril	21	N35°31.306', W005°50.263'
	35. Oued Aïn Jdioui (Touaret)	Aïn Jdioui	2	N35°33.768', W005°55.002'
Taounate	36. Aïn Boharroch	Aïn Aïcha	234	N34°29.049', W004°40.442'
	37. Dhar Sbagh Mâasra	Khemis Hoara	396	N34°26.977', W004°28.972'
	38. El Hajria	Sahel Botaher	184	N34°29.496', W004°49.886'
	39. Koudia El Aouinate	Douar Kda	403	N34°26.170', W004°28.167'
	40. Lâazaba	Khemis Hoara	357	N34°26.820', W004°28.408'
Tétouan	41. Aïn El Âakba Larbaâ	Mokdassene	95	N35°31.328', W005°19.012'
	42. Barrage Nakhla	Zinat	240	N35°26.954', W005°24.326'
	43. Barrage Smir	Bouzaghlale (M'diq)	20	N35°41.491', W005°22.673'
	44. Daya Amsemlil	Jbel Bouhachem	1059	N35°15.596', W005°25.917'
	45. Daya Jbel Zemzem	Jbel Zemzem	216	N35°45.457', W005°22.189'
	46. Douar Dacheryène	Dacheryène	132	N35°33.863', W005°27.162'
	47. Douar Kitane	Kitane	52	N35°32.412', W005°20.393'
	48. Douar Taghbaloute	Larbaa Beni Hassen	379	N35°15.323', W005°20.887'
	49. Douar Tizga	Amsa	516	N35°26.237', W005°13.694'
	50. El Haouta	Kitane	52	N35°32.230', W005°20.166'
	51. El Malâab	Kitane	75	N35°32.348', W005°20.290'
	52. Jumb Kitane	Kitane	38	N35°32.759', W005°20.420'
	53. Oued Achekrade	Douar Aouzighen	642	N35°22.931', W005°20.364'
	54. Oued Al Mizzine	Mezine	790	N35°06.143', W005°21.272'
	55. Oued Amsa	Amsa	11	N35°31.926', W005°13.747'

Tétouan	56. Oued Boumarouil	Larbaa Beni Hassen	551	N35°18.733', W005°21.271'
	57. Oued El Hamma	Beni Ydder	200	N35°23.532', W005°30.051'
	58. Oued El Kebir	Route Vers Beni Yedder	89	N35°27.351', W005°25.796'
	59. Oued Halila	Azla	3	N35°33.137', W005°14.678'
	60. Oued Maâza (Âachira)	Kitane	15	N35°32.596', W005°20.350'
	61. Oued Maâza (Tarik El Ouasâa)	Kitane	39	N35°32.237', W005°19.891'
	62. Oued Martil	Mhannech 2	9	N35°33.693', W005°22.510'
	63. Oued Martil (Tamouda)	Tamouda	8	N35°33.627', W005°24.799'
	64. Oued Sahel	Ben Karrich	40	N35°29.238', W005°26.352'
	65. Oued Zarka	Zarka	128	N35°31.211', W005°20.477'
	66. Oued Zinat	Zinat	190	N35°28.207', W005°24.224'

Beni Snassen (Eastern Morocco)

Berkane	67. Oued Beni Ouaklane	Beni Snassen	630	N34°51.581', W002°15.180'
---------	------------------------	--------------	-----	---------------------------


Middle Atlas

Khénifra	68. Oued Oum Er Rbia	Farra	826	N32°55.358', W005°40.390'
	69. Sensla	Arougo	1111	N32°55.387', W005°34.081'

Anti Atlas

Chtouka-Aït Baha	70. Oued Assa	Assa	306	N28°36.507', W009°25.800'
	71. Avant Sidi Binzarne	Massa	10	N30°03.144', W009°39.067'
	72. Centre Sidi Ouassay	Massa	41	N30°03.387', W009°41.249'
	73. Douar Sidi Abou	Massa	55	N29°59.322', W009°35.627'
	74. Environs Massa	Massa	24	N29°59.353', W009°38.708'
	75. Oued Massa (Pont Aghbalou)	Massa	2	N30°02.007', W009°38.768'
Errachidia	76. Oued Ziz (Pont Errachidia)	Oued Ziz	1026	N31°56.253', W004°25.455'
Guelmim	77. Msidira	El Filaha	257	N28°56.865', W010°02.979'
	78. Oued Sayad	Taghjijt	568	N29°03.784', W009°26.027'
	79. Oued Tisla	Bouizakarne	613	N29°09.908', W009°43.280'
Ouarzazate	80. Douar Zaouia	Kelâa M'gouna	1238	N31°14.275', W006°08.169'
	81. Oued Ouarzazate	Ouarzazate	1135	N30°54.917', W006°54.187'
Rissani	82. Ksibat Elhdeb	Rissani	770	N31°16.494', W004°16.310'
Tata	83. Douar Tighrimt	Tighrimt	706	N29°45.742', W007°58.521'
	84. Douar Zaouiet	Akka	554	N29°26.485', W008°16.535'
	85. Imzine	Tissint	566	N29°53.415', W007°18.928'
	86. Oued Foum Zguid (Douar Ouaiftoute)	Foum Zguid	665	N30°06.786', W006°52.639'
	87. Oued Tamanarne	Foum El Hisn	480	N29°00.941', W008°53.588'
Tinghir	88. Oued Tata	Tata	277	N29°45.348', W007°58.690'
	89. Oued Tinghir	Douar Ihertane	1280	N31°31.366', W005°31.677'
Tiznit	90. Airport Sidi Ifni	Sidi Ifni	51	N29°22.654', W010°10.711'
	91. Atbane	Tiznit	191	N29°44.388', W009°43.740'
	92. Route Bab El Khemis	Tiznit	225	N29°42.647', W009°43.702'

	93. Aït Aissa O Brahim	Mhamid El Ghizlane	556	N29°48.948', W005°43.676'
	94. Douar Rggaga	Tagonite	614	N29°57.874', W005°34.532'
	95. Isdaoun	Tazzarine	852	N30°44.748', W005°32.194'
Zagora	96. Jnane Makadir	Zagora	726	N30°19.209', W005°50.498'
	97. Kasbah Asma	Amzrou	718	N30°19.015', W005°50.016'
	98. Oued Drâa (Ikhf Mezrou)	Zagora	872	N30°41.466', W006°11.405'
	99. Oued Drâa (Tahtah)	Zagora	718	N30°19.665', W005°49.813'



Map 1. Map showing all collecting sites in Morocco.

Results

Faunistic records

Family TEPHRITIDAE (Newman, 1834)

Subfamily DACINAE (Loew, 1862)

Tribe CERATITIDINI Bezzi, 1910

Genus *CAPPARIMYIA* Bezzi, 1920

Capparimyia savastani Martelli, 1911

Literature records. Morocco, Anti Atlas: Tiznit (Séguy 1953).

World distribution. Algeria, Tunisia, Libya, Egypt, Italy, Greece, Malta, Israel, Oman, France, Yemen and Pakistan (Freidberg and Kugler 1989, Norrbom et al. 1999, Donati and Belcari 2003, De Meyer and Freidberg 2005, Miranda et al. 2008, Paphristos et al. 2009).

Genus *CERATITIS* MacLeay, 1829

Ceratitidis capitata (Wiedemann, 1824)

Literature records. Morocco, Rif: Tangier (Becker and Stein 1913), High Atlas (Séguy 1930), Oujad-Ouli-Mohamed, env. Settat, Insgane (De Meyer 2000).

New records. Morocco, Rif: Douar Kitane, 4♂♂, 4♀♀, 23-IX-2014- 4♂♂, 24-IX-2014- 4♂♂, 3♀♀, 25-IX-2014- 1♂, 2♀♀, 27-IX-2014- 1♂, 3♀♀, 02-X-2014- 4♂♂, 1♀, 27-VII-2015 (reared); El Haouta, 1♂, 1♀, 20-IX-2014 (net sweeping)- 5♂♂, 11♀♀, 02-X-2014 (reared); Jumb Kitane, 4♂♂, 7♀♀, 30-VI-2015. Middle Atlas: Sensla, 1♀, 14-VII-2016 (net sweeping). Anti Atlas: Environs Massa, 1♀, 11-V-2015; Oued Massa (Pont Aghbalou), 2♂♂, 12-V-2015; Douar Sidi Abou, 23♂♂, 34♀♀, 13-V-2015; Douar Tighrimt, 1♂, 29-V-2015; Douar Zaouia, 1♂, 11-VI-2015 (net sweeping).

Host plants. A highly polyphagous species, including the infestation of fruits of *Ficus carica* L., *Prunus armeniaca* L., *Prunus persica* L., and *Pyrus communis* L. from which the recorded specimens were reared, and *Argania spinosa* L. (Fig. 1) from which our specimens were collected by sweep-netting.

World distribution. Algeria, Tunisia, Libya, Egypt, Angola, Burundi, Cameroon, Congo, Ghana, Guinea, Kenya, Madagascar, Mozambique, Nigeria, South Africa, Seychelles Is, St Helena I., Tanzania, Benin, Burkino Faso, Cape Verde Is, Ethiopia, Gabon, Ivory Coast, Liberia, Malawi, Mali, Mauritius I., Niger, Reunion I., São Tomé I., Senegal, Sudan, Togo, Uganda, Zimbabwe, South Europe, Middle East, Neotropics, West Australia, Hawaii (Norrbom et al 1999, De Meyer 2000, De Meyer et al. 2013).



Figure 1. *Argania spinosa* L. host plant of *Ceratitis capitata*.

Tribe DACINI Loew, 1862

Genus *BACTROCERA* Macquart, 1835

Subgenus *Bactrocera* Macquart, 1835

***Bactrocera oleae* (Rossi, 1790)**

Literature record. Morocco, Rif: Tangier (Becker and Stein 1913). First record for the Middle and Anti Atlas.

New records. Rif: El Haouta, 1♂, 30-I-2014 (net sweeping); Oued Maâza (Tarik El Ouasâa), 5♂♂, 1♀ 18-III-2015 (reared); Cascade Chrafate, 1♂, 28-IV-2015 (net sweeping); Koudia El Aouinate, 3♂♂, 6♀♀, 12-XII-2015 (net sweeping); Lâazaba, 2♂♂, 12-XII-2015 (net sweeping); Dhar Sbagh Mâasra, 2♂♂, 12-XII-2015 (net sweeping); El Hajria, 1♀, 13-XII-2015 (net sweeping). Middle Atlas: Sensla, 1♂, 1♀, 14-VII-2016 (net sweeping). Anti Atlas: Route Bab El Khemis, 2♂♂, 2♀♀, 14-V-2015 (net sweeping).

Host plant. Fruits of *Olea europaea* L. from which the specimens were both reared and collected by sweep-netting.

World distribution. Mediterranean region, Eritrea, Ethiopia, Kenya, Angola, Lesotho, Namibia, South Africa, Sudan, Greece, Italy, Portugal, Spain, Madeira, Canary Islands, North Africa, Israel, Lebanon, Jordan, Réunion, Saudi Arabia, Syria, Turkey,

Caucasus, Pakistan, nw. India, USA, Mexico (Séguy 1930, Munro 1984, White and Hancock 1997, Norrbom et al. 1999, Copeland et al. 2004, White 2006).

Genus *DACUS* Fabricius, 1805

Dacus frontalis (Becker, 1922)

New records. Morocco, Anti Atlas: Oued Foum Zigid (Douar Ouaitoute), 1♂, 2♀♀, 01-VI-2015 (net sweeping); Oued Drâa (Ikhf Mezrou), 1♂, 1♀, 06-VI-2015 (net sweeping); Isdaoun, 2♂♂, 3♀♀, 07-VI-2015 (net sweeping). First record for Morocco.

World distribution. Algeria, Libya, Egypt, Angola, Benin, Botswana, Cape Verde Is., Congo, Eritrea, Kenya, Lesotho, Namibia, Saudi Arabia, South Africa, Sudan, Tanzania, Yemen, Zimbabwe (White 2006, White and Goodger 2009, De Meyer et al. 2013).

Dacus longistylus (Wiedemann, 1830)

New records. Morocco, Anti Atlas: Oued Tata, 1♀, 29-V-2015 (net sweeping); Douar Tighrimt, 1♂, 29-V-2015 (net sweeping); Oued Foum Zigid (Douar Ouaitoute), 31♂♂, 12♀♀, 01-VI-2015 (net sweeping). First record for Morocco.

Host plant. *Calotropis* L. (Fig. 2) from which the specimens were collected by sweep-netting.

World distribution. Libya, Egypt, Benin, Cameroon, Chad, Eritrea, Ethiopia, Kenya, Mali, Mauritania, Niger, Nigeria, Oman, Saudi Arabia, Senegal, Somalia, Sudan, Tanzania, Uganda, Yemen (White and Hancock 1997, White 2006, White and Goodger 2009, De Meyer et al. 2013).

Subfamily TEPHRITINAE (Newman, 1834)

Tribe DITHRYCINI Hendel, 1927

Subtribe OEDASPIDINA Hering, 1947

Genus *OEDASPIS* Loew, 1862

Oedaspis daphneae (Séguy 1930)

Literature records. Morocco, Middle Atlas: El Mers (Séguy 1930, Foote 1984, Norrbom et al. 1999).

World distribution.

Comments. The species was described from Morocco and never recorded since; we did not examine the type material.




Figure 2. *Calotropis procera* L., host plant of *Dacus longistylus*.

***Oedaspis multifasciata* (Loew, 1850)**

Literature records. Morocco, Middle Atlas: Itzer (Séguy 1953).

World distribution. Spain, France, Germany, Italy, Croatia, Austria, Ukraine (Korneyev 2002).

***Oedaspis trotteriana* (Bezzi, 1913)**

Literature records. Morocco (Ribera and Blasco-Zumeta 1998).

World distribution. Algeria, Libya, Egypt, Israel (Ribera and Blasco-Zumeta 1998, Norrbom et al. 1999).

Tribe MYOPITINI Hendel, 1927

Genus *MYOPITES* Blot, 1827

***Myopites inulaedyssentericae* Blot, 1827**

New record. Morocco, Rif: Affluent Tarmast, 1♂, 26-IV-2013; Aïn Afersiw, 1♀, 11-VI-2013 (net sweeping). First record for Morocco.

World distribution. Algeria, Great Britain, Estonia, France, Germany, Italy, Poland, Russia, Spain, Switzerland, Ukraine, Balkans, central Europe (Evstigneev 2011).

Myopites stylatus Fabricius, 1794

New record. Morocco, Rif: Affluent Tarmast, 1♀, 26-VI-2013; Oued El Hamma, 1♂, 25-IV-2014; El Haouta, 1♂, 18-VI-2014 (net sweeping). First record for Morocco.

World distribution. North Africa, southern Europe, Israel, Spain, Albania, (Zimšen 1964, Freidberg and Kugler 1989, Norrbom et al 1999, Merz 2001, Korneyev 2003).

Genus *UROPHORA* Robineau-Desvoidy, 1830

Subgenus *Urophora* Robineau-Desvoidy, 1830

= *Euribia* Meigen, 1800

Urophora sp. near *congrua*

Literature records. Morocco, Anti Atlas: Taroudant (Séguy 1941).

World distribution. France, South Germany, Austria (Norrbom et al, 1999).

Comments. The presence of *Urophora congrua* Loew, 1862 in Morocco is obviously a misidentification of another *Urophora* (Personal communication with Valery Korneyev). The literature records and the world distribution given above are those of *U. congrua* and therefore do not necessarily represent the true information for the species that occurs in Morocco

Urophora mauritanica (Macquart, 1851)

Literature records. Morocco, High Atlas: Imi N'Takandout, Dar Kaid M'tougui (Séguy 1930), Ito (White and Korneyev 1989). First record for the Anti Atlas.

New records. Morocco, Anti Atlas: Airport Sidi Ifni, 1♀, 17-V-2015; Oued Tata, 5♂♂, 29-V-2015 (net sweeping); Oued Tata, 12♂♂, 8♀♀, 06-VI-2015 (reared); Douar Tighrimt, 3♂♂, 29-V-2015; Imzine, 3♂♂, 1♀, 30-V-2015; Jnane Makadir, 1♀, 03-VI-2015; Oued Ouarzazate, 2♂♂, 12-VI-2015 (net sweeping)- 25♂♂, 14♀♀, 28-VI-2015- 7♂♂, 2♀♀, 01-VII-2015 (reared).

Host plant. *Centaurea calcitrapa* L. (Fig. 3) from which the specimens were reared.

World distribution. Algeria, Libya, Albania, Azerbaijan, Bulgaria, Cyprus, Czech Rep., France, Greece, Iran, Israel, Italy, Jordan, Kazakhstan, Lebanon, Macedonia, Malta, Moldova, Russia, Spain, Tajikistan, Turkey, Turkmenistan, Ukraine, Uzbekistan



Figure 3. *Centaurea calcitrapa* L., host plant of *Urophora mauritanica*, *Chaetorellia conjuncta* and *Terellia virens*.

(White and Korneyev 1989, Korneyev and White 1996, 1999, Norrbom et al. 1999, Gharali et al. 2005, Khaghaninia and Gharajedaghi 2012).

Urophora quadrifasciata (Meigen, 1826)

Literature records. Morocco (Séguy 1930).

World distribution. South to North Africa, Europe, Kazakhstan, Iran, North America, Australia (Norrbom et al. 1999, Heřman and Dirlbek 2006).

Urophora solstitialis Linnaeus, 1758

Literature record. Morocco, High Atlas: Haute Reraya (Séguy 1930).

World distribution. Tunisia, Great Britain, Eastern Scandianvia to Kazakhstan, China, France, Italy, Switzerland, Turkey, Iran, Armenia, Moldova, Poland, Russia, Ukraine, North America, Australia, New Zealand (Korneyev and White 1999, Heřman and Dirlbek 2006, Evstigneев 2011).

Tribe Noeetini (Norrbom & Korneyev, 1999)**Genus *ENSINA* Robineau-Desvoidy, 1830*****Ensina sonchi* (Linnaeus, 1767)**

New records. Morocco, Rif: Ksar Rimal, 5♂♂, 2♀♀, 05-VI-2013- Douar Tizga, 1♂, 25-VI-2014- Oued El Kebir, 1♀, 31-V-2015. Middle Atlas: Sensla, 1♂, 14-VII-2016 (net sweeping). Anti Atlas: Oued Massa (Pont Aghbalou), 1♀, 12-V-2015 (net sweeping); Centre Sidi Ouassay, 1♂, 12-V-2015; Aïn Boharroch, 1♀, 13-XII-2015; Atbane, 1♀, 14-V-2015; Oued Tamanarne, 1♂, 26-V-2015; Oued Drâa (Tahtah), 6♂♂, 7♀♀, 03-VI-2015; Jnane Makadir, 3♂♂, 1♀, 03-VI-2015; Kasbah Asma, 16♂♂, 13♀♀, 04-VI-2015 (net sweeping)- 7♂♂, 13♀♀, 10-VI-2015 (reared); Aït Aissa O Brahim, 1♀, 05-VI-2015; Oued Ziz (Pont Errachidia), 2♂♂, 2♀♀, 09-VI-2015; Oued Ouarzazate, 4♂♂, 2♀♀, 12-VI-2015 (net sweeping). First record for Morocco.

Host plants. Flower heads of *Hypochaeris radicata* L. (Fig. 4) from which the specimens were reared.

World distribution. South to North Africa, Great Britain, Scandinavia, Andorra, Czech Rep., France, Germany, Iran, Poland, Portugal, Spain, Sweden, Switzerland,



Figure 4. *Hypochaeris* sp., host plant of *Ensina sonchi*.

Turkey, Ukraine, Saudi Arabia, Japan, Korea, India, Ethiopia, Taiwan, Philippines, Hawaii (White and Elson-Harris 1992, Merz and Dawah 2005, Heřman and Dirlbek 2006, Gharajedaghi et al. 2012).

Genus *HYPENIDIUM* Loew, 1862

Hypenidium graecum (Loew, 1862)

= *Stephanaciura bipartita* Séguay, 1930

Literature records. Morocco, Middle Atlas: Taffert (Séguay 1930, Norrbom et al. 1999).

World distribution. Bosnia, Greece, Hungary, Israel, Italy, Portugal, Spain, Ukraine (Korneyev et al. 2011).

Tribe TEPHRELLINI Hendel, 1927

Subtribe TEPHRELLINA Hendel, 1927

Genus *ACIURA* Robineau-Desvoidy, 1830

Aciura coryli (Rossi, 1794)

= *Aciura powelli* (Séguay, 1930)

Literature records. Morocco, Middle Atlas: Azrou (Séguay 1930), Korifla (Séguay 1953).

World distribution. North Africa, Spain and southern France to southern Ukraine, Canary Islands, Israel, Syria (Korneyev and Dirlbek 2000).

Genus *OXYACIURA* Hendel, 1927

Oxyaciura tibialis (Robineau-Desvoidy, 1830)

Literature records. Morocco, Rif: Tangier (Becker and Stein 1913), Sker (Séguay 1930).

New sites. Morocco, Rif: Daya El Birdiyel, 1♂, 27-VI-2013; Oued Azila, 1♂, 27-VI-2013; Maison forestière, 1♂, 17-VI-2014; Oued Maâza (Tarik El Ouasâa), 1♂, 18-IV-2015 (net sweeping).

World distribution. North Africa, Canary Islands, Cape Verde Islands, Madeira, Spain, China, Iran, Israel, Kazakhstan, Saudi Arabia, Syria, United Arab Emirates, Afghanistan, Ethiopia (Merz 1992, 2008, Korneyev and Dirlbek 2000, Merz and Dawah 2005, Gharajedaghi et al. 2012).

Genus *SPHAENISCUS* Becker, 1908***Sphaeniscus filiolus* (Loew, 1869)**

= *Spheniscomyia aegyptiaca* (Efflatoun, 1924)

Literature records. Morocco, Anti Atlas: Goulimine (Séguy 1930, 1949). First record for the Rif.

New records. Morocco, Rif: Affluent Tarmast, 1♂, 26-VI-2013; Oued Maâza (Tarik El Ouasâa), 2♂♂, 1♀, 19-VI-2014 (net sweeping).

World distribution. Egypt, Spain, Canary Islands, Cape Verde Islands, Israel, Ethiopia (Séguy 1930, 1949; Merz 1992, 2001).

Tribe TEPHRITINI Macquart, 1835**Genus *ACANTHIOPHILUS* Becker, 1908*****Acanthiophilus helianthi* (Rossi, 1794)**

Literature records. Morocco, Rif: around Tangier (Becker and Stein 1913, Séguy 1930), Middle Atlas (Séguy 1930), Anti Atlas: Goulimine, Foum el Hassan, Alnif, Tarfaia, Boufarik (Séguy 1949).

New sites. Morocco, Rif: Oued Zinat, 1♀, 27-IV-2012; Ksar Rimal, 1♀, 10-V-2013; Oued Derdara, 1♀, 24-V-2013; Affluent Tarmast, 1♂, 26-VI-2013; Daya El Birdiyel, 1♀, 27-VI-2013; Oued Azila, 1♂, 27-VI-2013; Daya Jbel Zemzem, 1♂, 23-IV-2014; Oued Boumarouil, 2♂♂, 1♀, 10-V-2014; Douar Abou Boubnar (Mababout Sidi Gile), 1♀, 18-V-2014; El Haouta, 1♂, 18-VI-2014; Oued Maâza (Tarik El Ouasâa), 1♀, 19-VI-2014- 1♀, 18-IV-2015; Douar Tizga, 1♂, 25-VI-2014; Daya Afrate, 1♀, 18-IV-2015; Oued Mezine, 1♂, 18-IV-2015; Aïn El Malâab, 1♂, 1♀, 21-IV-2015; Oued Jnane Niche, 1♀, 25-IV-2015. Middle Atlas: Oued Oum Er Rbia, 1♂, 14-VII-2016. Anti Atlas: Centre Sidi Ouassay, 1♂, 2♀♀, 12-V-2015; Avant Sidi Binzarne, 1♀, 12-V-2015; Route Bab El Khemis, 1♂, 1♀, 14-V-2015; Airport Sidi Ifni, 22♂♂, 22♀♀, 17-V-2015; Oued Tisla, 1♀, 24-V-2015; Oued Sayad, 1♀, 25-V-2015; Oued Tamanarne, 1♂, 1♀, 26-V-2015; Oued Foum Ziguid (Douar Ouaiftoute), 1♂, 1♀, 01-VI-2015; Jnane Makadir, 2♂♂, 4♀♀, 03-VI-2015; Douar Rggaga, 1♀, 05-VI-2015; Oued Tinghir, 1♂, 1♀, 11-VI-2015; Oued Ouarzazate, 12♂♂, 6♀♀, 12-VI-2015 (net sweeping).

World distribution. Algeria, Tunisia, Albania, Andorra, Austria, Balearic Is., Belgium, Great Britain, Bulgaria, Corsica, Cyprus, Czech Rep., France, Germany, Greece, Hungary, Italy, Malta, Moldova, Netherlands, Norway, Poland, Portugal, Romania, Russia, Sardinia, Sicily, Slovakia, Spain, Switzerland, Ukraine, Turkey, Central Asia, Mongolia, China, Madeira, Canary Islands, Ethiopia, Kenya, Sudan, Afghanistan, Iran,

Israel, Kazakhstan, Kyrgyzstan, Lebanon, Saudi Arabia, United Arab Emirates, Syria, India, Nepal, Pakistan, Thailand (Séguy 1930, Merz and Korneyev 2004, Heřman and Dirlbek 2006, David and Ramani 2011, Merz 2011).

Genus **CAMPIGLOSSA** Rondani, 1870

***Campiglossa martii* (Becker, 1908)**

= *Oxyna martii* Becker, 1908

New records. North Africa, Morocco, Rif: Oued El Kebir, 1♀, 31-V-2014 (net sweeping). Anti Atlas: Centre Sidi Ouassay, 1♂, 12-V-2015 (net sweeping). First record for Morocco and North Africa.

World distribution. Canary Islands (Frey 1958a), Cape Verde Islands (Frey 1958b), Spain (Merz 1992).

***Campiglossa producta* Loew, 1844**

= *Paroxyna tessellata* Hendel, 1927

Literature records. Morocco, Rif: Tangier and its surroundings, (Becker and Stein 1913, Séguy 1930), High Atlas (Séguy 1930).

New sites. Morocco, Rif: Oued Al Mizzine, 1♀, 11-VI-2013; Aïn El Malâab, 1♂, 17-V-2014; Daya El Hajjami, 1♂, 28-III-2015 (net sweeping).

World distribution. Algeria, Tunisia, Great Britain, Finland, Netherlands, Poland, Portugal, Spain, Switzerland, Madeira, Canary Islands, Turkey, Syria, Jordan, Iran, Israel, Iraq, Afghanistan, Central Asia, India, China, Thailand, Vietnam, Korea, Japan (Korneyev and Dirlbek 2000, Hancock and McGuire 2002, Korneyev 2004, Heřman and Dirlbek 2006, Mohamadzade Namin and Nozari 2011).

Genus **CAPITITES** Foote & Freidberg, 1981

***Capitites ramulosa* (Loew, 1844)**

Literature records. Morocco, Middle Atlas: Forest of Timelilt (Séguy 1941). High Atlas: Tizi n'Test, Imdress, Taroudant (Séguy 1941), Foum El Hassan, Akka, Agdz, Alnif (Séguy 1949).

World distribution. Algeria, Tunisia, Egypt, Greece, Portugal, Spain, Cyprus, Syria, Israel, Iraq, Canary Islands, Cape Verde Islands (Séguy 1930, Freidberg and Kugler 1989, Merz 1992, Norrbom et al. 1999, Korneyev and Dirlbek 2000).

Genus *DESMELLA* Munro, 1957

Desmella rostellata (Séguy, 1941)

= *Paroxyna rostellata* (Séguy, 1941)

Literature record. Morocco, Anti Atlas: Agadir (Séguy 1941).

Genus *DIOXYNA* Frey, 1945

Dioxyna sororcula (Wiedemann, 1830)

New records. Morocco, Rif: Ksar Rimal, 2♂♂, 19-V-2013- 1♂, 2♀♀, 26-V-2013; Oued Jnane Niche, 2♀♀, 14-VI-2013; Oued Halila, 1♂, 05-XI-2013; Oued Zarka, 7♂♂, 1♀, 14-XI-2013; Oued Martil, 1♂, 1♀, 13-XII-2013; Oued Amsa, 1♀, 19-XII-2013; Oued Sahel, 1♂, 05-IV-2014; Daya Jbel Zemzem, 1♂, 17-IV-2014; Oued Majjou, 2♂♂, 10-V-2014; Dhar Sbagh Mâasra, 1♀, 12-XII-2015; Douar Kitane, 15♂♂, 5♀♀, 14-XI-2013- 1♂, 2♀♀, 13-III-2014- 1♀, 17-III-2014- 2♂♂, 1♀, 10-IV-2014- 5♂♂, 13♀♀, 02-V-2014- 26♂♂, 14♀♀, 23-I-2016 (net sweeping). First record for Morocco.

World distribution. Algeria, Tunisia, South Africa, Madeira, Canary Islands, Spain, Saudi Arabia, Yemen, India, West Bengal, Nepal, Thailand, China, Korea, Japan, Taiwan, Botswana, Namibia, Zambia, Zimbabwe, Australia, New Caledonia, Western Samoa, Niue, Cook Is., French Polynesia, Fiji, Hawaii (Hancock and McGuire 2001, Norrbom and Hancock 2004, Merz and Dawah 2005, David and Ramani 2011).

Genus *EUARESTA* Loew, 1873

Euaresta bullans (Wiedemann, 1830)

Literature records. Morocco, Anti Atlas: Tiznit, Sidi-Moussa-d'Aglou (Heřman and Dirlbek 2006).

New site. Morocco, Anti Atlas: Msidira, 2♂♂, 1♀, 18-V-2015 (net sweeping).

World distribution. Algeria, Tunisia, South Africa, Argentina, Bolivia, Chile, Peru, Uruguay, California, Arizona, Bulgaria, France, Greece, Hungary, Iran, Israel, Italy, Macedonia, Moldova, Russia, Slovakia, Spain, Turkey, Ukraine, Australia (Heřman and Dirlbek 2006, Khaghaninia and Gharajedaghi 2012).

Genus *GONIURELLIA* Hendel, 1927***Goniurellia longicauda* (Freidberg, 1980)**

Literature records. Morocco, Middle Atlas: Tizi s'Tkrine, Forest of Azrou (Séguy 1930, Freidberg 1980), High Atlas: Taroudant (Séguy 1953, Freidberg 1980). Anti Atlas: Edehby Ouarzazate (Pârvu et al. 2006).

New sites. Morocco, Anti Atlas: Airport Sidi Ifni, 4♂♂, 17-V-2015; Oued Tisla, 6♂♂, 9♀♀, 24-V-2015; Oued Tamanarne, 1♂, 26-V-2015; Douar Zaouiet, 1♂, 27-V-2015; Oued Tata, 1♀, 29-V-2015; Douar Tighrimt, 1♀, 29-V-2015; Ksibat Elhdeb, 2♂♂, 07-VI-2015; Oued Ziz (Pont Errachidia), 1♀, 09-VI-2015; Oued Ouarzazate, 2♂♂, 12-VI-2015 (net sweeping).

World distribution. Algeria, Tunisia, Libya, Egypt, France, Canary Islands, Cape Verde Islands, Iran, Iraq, Israel, Saudi Arabia, Syria, Turkey, United Arab Emirates, Kenya (Freidberg 1980, Freidberg and Kugler 1989, Merz 1992, Norrbom et al. 1999, 2008, Korneyev and Dirlbek 2000, Al Dhafer et al. 2012).

***Goniurellia persignata* Freidberg, 1980**

Literature records. Morocco, Eastern region: Defilia, nr. Figuig (Freidberg 1980). Anti Atlas: Tiffoultoute (Norrbom et al. 1999, Heřman and Dirlbek 2006).

New records. Morocco, Rif: Dhar Sbagh Mâasra, 1♀, 12-XII-2015. Anti Atlas: Douar Zaouiet, 1♂, 27-V-2015; Oued Ouarzazate, 1♂, 12-VI-2015 (net sweeping). First record for the Rif.

World distribution. Egypt, China, Crete, Cyprus, Ethiopia, Israel, Saudi Arabia, Sri Lanka, Turkmenistan, Namibia? (Freidberg 1980, Norrbom et al. 1999, Hancock et al. 2001, Merz and Dawah 2005, Herman and Dirlbek 2006).

Genus *SPATHULINA* Rondani, 1856***Spathulina sicula* (Rondani, 1856)**

Literature records. Morocco (Séguy 1930).

New site. Morocco, Rif: Barrage Smir, 1♂, 27-IV-2014 (net sweeping).

World distribution. Spain, Portugal, Italy, Israel, Canary Is. (Norrbom et al. 1999).

Genus *SPHENELLA* Robineau-Desvoidy, 1830

Sphenella marginata (Fallén, 1814)

Literature records. Morocco, Rif: around Tangier (Becker and Stein 1913, Séguay 1930).

New sites. Morocco, Rif: Affluent Tarmast, 1♂, 26-VI-2013; Daya Jbel Zemzem, 5♂♂, 2♀♀, 23-IV-2014; El Malâab, 1♂, 02-II-2015; Oued Maâza (Âachira), 1♀, 10-III-2015; Daya Aïn Jdioui, 1♀, 28-III-2015; Oued Majjou, 1♀, 09-IV-2015; Daya Afrate, 1♂, 1♀, 18-IV-2015 (net sweeping).

World distribution. Egypt, South Africa, Czech Rep., Poland, Portugal, Spain, Sweden, Switzerland, Ukraine, Russia, Madeira, Afghanistan, Canary Islands, China, Iran, Israel, Kazakhstan, Saudi Arabia, Turkey, Turkmenistan, Eritrea, Kenya, Lesotho, Mozambique (Gharajedagi et al. 2012).

Genus *TEPHRITIS* (Latreille, 1804)

Tephritis dioscurea Loew, 1856

Literature records. Morocco, Middle Atlas: El Hajeb (Séguay 1930).

World distribution. Sweden, France, Hungary, Austria, Germany, Switzerland, Russia, Estonia, Latvia, Lithuania, Ukraine, Moldova, Azerbaijan, Armenia, Georgia, Kazakhstan, Turkey, Iran (Foote 1984, Merz 1994, Thompson 1998, Kütük 2005, Zarghani et al. 2010).

Tephritis divisa (Rondani, 1871)

New record. Morocco, Rif: Daya Amsemilil, 1♀, 01-XI-2014 (net sweeping). First record for Morocco and North Africa.

World distribution. Croatia, France, Greece, Israel, Italy, Near East Asia, Spain, Switzerland, Turkey and Ukraine (Norrbom 1999, Korneyev 2003, Merz and Korneyev 2004, Kütük 2005).

Tephritis formosa (Loew, 1844)

Literature record. Morocco, High Atlas: Asni (Séguay 1930).

New records. Morocco, Rif: Oued Abou Bnar, 1♂, 18-V-2014; Oued Sidi Ben Sâada, 1♂, 30-V-2014; Oued Achekrade, 1♂, 31-V-2014; Oued El Kanar, 1♀, 25-IV-2015 (net sweeping). First record for the Rif.

World distribution. Caucasus, Israel, Persia, Russia, Ukraine, Moldova, Azerbaijan, Georgia, Armenia, England, Switzerland, Germany, Iran, Turkey, China (Hendel 1927, Dirlbek and Dirlbek 1971, Giray 1979, Foote 1984, White 1988, Freidberg and Kugler 1989, Merz 1994, Wang 1998).

Tephritis leontodontis De Géer, 1776

Literature record. Morocco (Séguy 1930).

World distribution. Austria, Albania, Belgium, Bulgaria, United Kingdom, Estonia, Ireland, Spain, Italy, Latvia, Lithuania, Netherlands, Germany, Norway, Poland, Russian Federation, Romania, Slovakia, Hungary, Ukraine, Finland, France, United Kingdom, Czech Republic, Sweden, Switzerland (Korneyev 2016).

Tephritis matricariae Loew, 1844

Literature records. Morocco (Séguy 1930).

New sites. Morocco, Rif: Affluent Oued Amsemilil, 1♂, 30-V-2013; El Haouta, 1♀, 30-I-2014; Daya Jbel Zemzem, 1♂, 23-IV-2014; Daya Amsemilil, 1♂, 01-XI-2014; Oued El Hamma, 1♀, 24-IV-2015 (net sweeping).

World distribution. Algeria, Tunisia, Libya, Egypt, England, Netherlands, Austria, Andorra, Greece, Portugal, Spain, Switzerland, Turkey, Iran (Norrbom et al. 1999, Merz 2001, Mohamadzade Namin et al. 2010a).

Tephritis nigricauda Loew, 1856

Literature records. Morocco: Berrechid (Séguy 1930). First record for the Rif.

New sites. Daya Jbel Zemzem, 2♂♂, 23-IV-2014; Oued Maâza (Tarik El Ouasâa), 1♂, 18-IV-2015; Ain El Maounzil, 1♂, 21-IV-2015; Daya Tazia, 1♀, 24-IV-2015; Daya Amsemilil, 1♀, 27-V-2015; Douar Tamakoute, 1♂, 1♀, 25-XII-2015 (net sweeping).

World distribution. Algeria, Egypt, Poland, Portugal, Spain, Iran, Iraq, Jordan, Russia, Syria (Korneyev and Dirlbek 2000, Merz 2001, Gharajedaghi et al. 2012).

Tephritis postica (Loew, 1844)

Literature records. Morocco, Middle Atlas: Volubilis (Heřman and Dirlbek 2006). First record for the Anti Atlas.

New records. Anti Atlas: Ksibat Elhdeb, 1♀, 07-VI-2015 (net sweeping)- 1♂, 1♀, 23-VI-2015- 1♂, 1♀, 26-VI-2015- 1♂, 1♀, 28-VI-2015 (reared); Oued Tinghir, 1♂, 11-VI-2015 (net sweeping).



Figure 5. *Onopordum acanthium* L. host plant of *Tephritis postica*.

Host plant. Flower heads of *Onopordum acanthium* L. (Fig. 5) from which the specimens were reared.

World distribution. Algeria, Tunisia, Spain, France, Czech Rep., Georgia, Germany, Hungary, Poland, Slovakia, Ukraine, Russia, Turkey, Kazakhstan, Kyrgyzstan, China, Iran, Israel, Lebanon (Séguy 1934a, Richter 1970, Merz 1994, 2001, Wang 1998, Klasa 2001, Heřman and Dirlbek 2006, Zarghani et al. 2010, Gharajedaghi et al. 2012, Hancock 2013).

Tephritis praecox (Loew, 1844)

Literature records. Morocco, Rif: Tangier (Heřman and Dirlbek 2006). Middle Atlas: Tizi s'Tkrine (Séguy 1930), Ifrane-Azrou National Park (Heřman and Dirlbek 2006). First record for eastern Morocco.

New sites. Morocco, Rif: Daya El Ânassar, 2♂♂, 24-V-2013; Daya Amsemilil, 1♀, 30-V-2013; Affluent Oued Amsemilil, 1♀, 30-V-2013; Douar Dacheryène, 1♀, 04-IV-2014; Douar Taghbaloute, 1♂, 05-IV-2014; Barrage Nakhla, 1♂, 05-IV-2014; Oued Sahel, 2♂♂, 05-IV-2014; Daya Jbel Zemzem, 4♂♂, 2♀♀, 17-IV-2014- 1♂, 23-IV-2014; Douar Kitane, 1♂, 19-IV-2014; Oued El Hamma, 1♂, 25-IV-2014; Oued El Kebir, 2♂♂, 2♀♀, 25-IV-2014; Aïn El Ma Bared, 1♂, 06-V-2014; Aïn El Malâab, 1♂, 1♀, 17-V-2014; Douar Abou Boubnar (Marabout Sidi Gile), 1♀, 18-V-2014; Maison forestière, 1♂, 1♀, 17-VI-2014; Douar Tizga, 1♀, 25-VI-2014; Oued Aïn Jdioui (Touaret), 1♀, 28-III-2015;

Daya Afrate, 1♂, 18-IV-2015; Oued Jbara, 1♂, 18-IV-2015; Aïn El Malâab, 1♂, 21-IV-2015; Aïn El Maounzil, 11♂♂, 8♀♀, 21-IV-2015; Daya Tazia, 1♂, 24-IV-2015; Oued Jnane Niche, 5♂♂, 4♀♀, 25-IV-2015; Oued Majjou, 1♀, 27-IV-2015; Aïn Tiouila, 1♂, 02-V-2015; Daya Mtahen, 1♂, 1♀, 07-V-2015; Lâazaba, 3♂♂, 1♀, 12-XII-2015; Dhar Sbagh Mâasra, 1♂, 3♀♀, 12-XII-2015; El Hajria, 1♀, 13-XII-2015; Aïn Boharroch, 1♂, 13-XII-2015; Douar Tamakout, 1♀, 25-XII-2015; Douar Ouslaf, 1♂, 2♀♀, 26-XII-2015 (net sweeping). Beni Snassen (Eastern Morocco): Oued Beni Ouaklane, 1♂, 24-XI-2014.

World distribution. Algeria, Spain, Portugal, Switzerland, Great Britain, Austria, Hungary, Madeira, Canary Islands, Syria, Iran, Israel, Iraq, Afghanistan (Séguy 1930, Norrbom et al. 1999, Korneyev and Dirlbek 2000, Smit 2006, Heřman and Dirlbek 2006, Zarghani et al. 2010, Gharajedaghi et al. 2012).

Tephritis pulchra Loew, 1844

Literature records. Morocco (Séguy 1930).

World distribution. North Africa, France, Spain, Austria, Hungary, Italy, Greece, Poland, Iran, Turkey, China (Norrbom et al. 1999, Merz 2001, Freidberg and Kütük 2002, Mohamadzade Namin and Nozari 2011).

Tephritis simplex Loew, 1844

Literature records. Morocco (Séguy 1930).

World distribution. Algeria, Tunisia, Spain, Portugal, Crete, Turkey, Israel (Norrbom et al. 1999, Merz 2001).

Tephritis stictica Loew, 1862

Literature records. Morocco: Rabat (Séguy 1930).

World distribution. Algeria (Séguy 1930), southern Europe (Norrbom et al. 1999).

Tephritis theryi (Séguy 1930)

Literature records. Morocco, High Atlas: Asni (Séguy 1930, Norrbom et al. 1999).

Tephritis vespertina (Loew, 1844)

New record. Morocco, Rif: Daya Mtahen, 1♀, 27-V-2015; Dhar Sbagh Mâasra, 1♂, 1♀, 12-XII-2015 (net sweeping). First record for Morocco.

World distribution. South to North Africa, Andorra, Germany, Hungary, Poland, Portugal, Spain, Switzerland, Ukraine, Turkey (Merz 2001, Hendel 1927, Foote 1984, Norrbom et al. 1999, Merz 1994, Kütük and Özgür 2003).

Genus *TEPHRITOMYIA* Hendel, 1927

Tephritisomyia lauta (Loew, 1869)

Literature records. Morocco, High Atlas: Tachdirt (Séguy 1930); Tizi-n-Tichka (Morgulis 2015). First record for the Rif.

New records. Morocco, Rif: Daya El Birdiyel, 1♂, 27-VI-2013; Daya Amsemlil, 1♀, 01-XI-2014; Lâazaba, 1♂, 12-XII-2015. Anti Atlas: Msidira, 1♂, 18-V-2015; Oued Ouarzazate, 1♂, 6♀♀, 12-VI-2015 (net sweeping).

World distribution. Tunisia, Egypt, Cyprus, Greece, Iran, Iraq, Israel, Lebanon, Syria, Turkey (Freidberg and Kugler 1989, Norrbom et al. 1999, Korneyev and Dirlbek 2000, Mohamadzade Namin and Nozari 2011, Morgulis 2015).

Genus *TRUPANEA* (Schrank, 1795)

Trupanea amoena (Frauenfeld, 1857)

Literature records. Morocco, Middle Atlas: Aïn Leuh (Séguy 1930). First record for the Rif.

New records. Morocco, Rif: Ksar Rimal, 1♂, 05-VI-2013; Oued Jnane niche, 1♂, 14-VI-2013; Affluent Tarmast, 2♀♀, 26-VI-2013; Oued Martil (Tamouda), 1♀, 13-VII-2013; Oued Amsa, 1♂, 13-XII-2013; Oued El Hamma, 2♂♂, 1♀, 25-IV-2014; Oued Boumarouil, 1♀, 10-V-2014; Oued Sidi Yahya Aârab, 1♂, 25-IV-2015; Aïn Tiouila, 1♂, 02-V-2015. Anti Atlas: Oued Massa (Pont Aghbalou), 1♂, 12-V-2015; Centre Sidi Ouassay, 1♂, 12-V-2015; Avant Sidi Binzarne, 2♂♂, 1♀, 12-V-2015; Oued Tisla, 2♀♀, 24-V-2015; Douar Tighrimt, 1♂, 29-V-2015; Oued Draa (Tahtah), 1♂, 03-VI-2015; Jnane Makadir, 1♂, 03-VI-2015; Douar Rggaga, 1♀, 05-VI-2015; Aït Aissa O Brahim, 1♂, 05-VI-2015; Oued Drâa (Ikhf Mezrou), 1♂, 06-VI-2015; Isdaoun, 1♂, 07-VI-2015; Ksibat Elhdeb, 1♀, 07-VI-2015; Oued Tinghir, 2♂♂, 11-VI-2015 (net sweeping).

World distribution. North Africa, Canary Islands, Ceylon, Ethiopia, England, Germany, India, Iran, Israel, Middle Asia, Middle and North Europe, Netherlands, Philippines, Saudi Arabia, Switzerland, Taiwan, Turkey, United Arab Emirates (Hendel 1927, Séguy 1930, Giray 1969, Foote 1984, White 1988, Freidberg and Kugler 1989, Merz 1994, Merz and Korneyev 2004, Merz and Dawah 2005, Merz 2008, Mohamadzade Namin et al. 2010b).

Trupanea guimari (Becker, 1908)

Literature records. Morocco, Middle Atlas: Tizi s'Tkrine, Forest of Azrou (Séguy 1930). High Atlas: Tenfecht (Séguy 1930). First record for the Anti Atlas.

New records. Morocco, Anti Atlas: Centre Sidi Ouassay, 4♂♂, 12-V-2015; Msidira, 1♂, 18-V-2015; Jnane Makadir, 2♂♂, 1♀, 03-VI-2015; Aït Aissa O Brahim, 1♂, 1♀, 05-VI-2015; Ksibat Elhdeb, 1♀, 07-VI-2015 (net sweeping).

World distribution. Algeria, Spain, Canary Islands, Cape Verde Islands (Merz 1992, 1999).

Trupanea stellata (Fuessly, 1775)

Literature records. Morocco, Middle Atlas: Forest of Timelilt (Séguy 1930), Anti Atlas: Goulimine (Séguy 1949). First record for the Rif.

New records. Morocco, Rif: Mizoghar, 1♀, 06-V-2014; Oued Maâza (Tarik El Ouasâa), 1♂, 19-VI-2014; Daya Afrate, 1♀, 18-IV-2015; Aïn El Malâab, 1♀, 21-IV-2015; Oued Tkaraâ, 1♀, 07-V-2015. Anti Atlas: Centre Sidi Ouassay, 1♀, 12-V-2015 (net sweeping).

World distribution. Tunisia, Andorra, Armenia, British Isles, Scandinavia, Czech Rep., Madeira, Poland, Spain, Switzerland, Turkey, Ukraine, Mongolia, China, Canary Islands, Iraq, Israel, Saudi Arabia, United Arab Emirates, Iran, India (Norrbom et al. 1999, Heřman and Dirlbek 2006, David and Ramani 2011, Gharajedaghi et al. 2012, Merz 2011).

Tribe TERELLINI Hendel, 1927

Genus *CHAETORELLIA* Hendel, 1927

Chaetorellia conjuncta (Becker, 1912)

New records. Morocco, Anti Atlas: Airport Sidi Ifni, 5♂♂, 5♀♀, 17-V-2015; Oued Assa: 3♂♂, 2♀♀, 21-V-2015; Oued Sayad, 1♂, 25-V-2015; Oued Foum Zguid (Douar Ouaiftoute), 1♀, 01-VI-2015; Ksibat Elhdeb, 1♂, 07-VI-2015; Oued Ziz (Pont Errachidia), 1♀, 09-VI-2015 (net sweeping)- 1♀, 12-VI-2015- 3♂♂, 23-VI-2015 (reared); Oued Ouarazate, 4♂♂, 2♀♀, 12-VI-2015 (net sweeping)- 1♂, 28-VI-2015- 1♂, 1♀, 01-VII-2015 (reared). First record for Morocco.

Host plants. Flower heads of *Centaurea calcitrapa* L. from which the specimens were reared.

World distribution. Egypt, Afghanistan, Albania, Caucasus, Cyprus, Greece, Hungary, Iraq, Iran, Israel, Jordan, Kazakhstan, Lebanon, Pakistan, Syria, Turkey (White and Marquardt 1989, Norrbom et al. 1999, Korneyev and Dirlbek 2000, Merz 2000).

***Chaetorellia bestia* Hering, 1937**

Literature record. Morocco, High Atlas: Essaouira (Séguy 1930). First record for the Anti Atlas.

New record. Morocco, Anti Atlas: Centre Sidi Ouassay, 2♂♂, 12-V-2015 (net sweeping).

World distribution. Algeria, France, Spain, Italy (Séguy 1930, White and Marquardt 1989, Norrbom et al. 1999).

Genus *TERELLIA* Robineau-Desvoidy, 1830

***Terellia colon* Meigen, 1826**

Literature records. Morocco (Séguy 1930).

World distribution. North Africa, Austria, Azerbaijan, Belgium, Great Britain, Crete, Czech Republic, Denmark, Dodecanese Islands, France, Germany, Georgia, Greece, Hungary, Israel, Italy, Kazakhstan, Kyrgyzstan, Lithuania, Moldova, North Aegean Islands, Poland, Romania, Russia, Sicily, Slovakia, Spain, Sweden, Switzerland, West Siberia, Tajikistan, Netherlands, Turkmenistan, Turkey, Ukraine, Uzbekistan (Foote 1984, Norrbom et al. 1999, Kütük and Özgür 2003, Özgür and Kütük 2003, Merz and Korneyev 2004, Heřman and Dirlbek 2006).

***Terellia fuscicornis* (Loew, 1844)**

Literature records. Morocco (Séguy 1930).

World distribution. North Africa, Crete, Cyprus, France, Greece, Italy, Madeira, Malta, Sardinia, Sicily, Spain, Iran, Israel, Lebanon, Turkey, California, Scotland (Norrbom et al. 1999, Merz 2001, Whittington 2002, Kütük and Yaran 2011, Mohamadzade Namin 2011).

***Terellia longicauda* (Meigen, 1838)**

Literature records. Morocco, High Atlas: Tizi n'Test, Middle Atlas: Aïn Leuh (Séguy 1930, 1934b).

World distribution. Albania, Austria, Great Britain, Bulgaria, Czech Republic, Cyprus, France, Germany, Hungary, Italy, Near East, Poland, Russia, Slovakia, Spain, Switzerland, Ukraine, Turkey (Kütük 2003, Merz and Korneyev 2004).



Figure 6. *Cynara cardunculus* L. host plant of *Terellia* sp. near *longicauda*.

Terellia sp. near *longicauda*

New record. Morocco, Rif: Douar Kitane, 4♂♂, 10-VIII-2016- 2♂♂, 3♀♀, 11-VIII-2016- 1♂, 3♀♀, 12-VIII-2016- 1♀, 13-VIII-2016 (reared). First record for Morocco and North Africa.

Host plants. Flower heads of *Cynara cardunculus* L. (Fig. 6) from which the specimens were reared.

World distribution. Spain, Canary Islands (Merz 1992).

Comments. This is an undescribed, cryptic species of the *Terellia serratulae* L. complex known to infest *Cynara cardunculus* L. in Spain and the Canary Islands. The *serratula* group contains cryptic species, a situation that needs to be resolved, and for which a molecular approach should be adopted (Valery Korneyev, pers. comm.).

Terellia serratulae (Linnaeus, 1758)

Literature records. Morocco, Rif: Tangier (Wiedemann 1824, Becker and Stein 1913).

New sites. Morocco, Rif: Daya Jbel Zemzem, 1♀, 23-IV-2014; Oued Maâza (Tarik El Ouasâa), 2♂♂, 1♀, 18-VI-2014 (net sweeping).

World distribution. Algeria, Tunisia, Albania, Andorra, Armenia, Austria, Azerbaijan, Balearic Islands, Belgium, Great Britain, Bulgaria, China, Corsica, Crete, Cyprus,

Czech Rep., Denmark, Finland, France, Georgia, Germany, Greece, Iran, Iraq, Ireland, Israel, Italy, Kazakhstan, Korea, Lebanon, Malta, Moldova, Mongolia, Netherlands, Norway, Poland, Portugal, Romania, Russia, Sardinia, Sicily, Slovakia, Spain, Sweden, Switzerland, Syria, Turkey, Ukraine (Korneyev and Dirlbek 2000, Han and Kwon 2000, Merz 2001, Heřman and Dirlbek 2006, Kütük and Yaran 2011, Khaghaninia et al. 2012).

***Terellia virens* (Loew, 1846)**

Literature records. Morocco, High Atlas: Jbel Ayachi (White 1989), Tizi-n-Talrhemt (Korneyev et al. 2013). First record for the Anti Atlas.

New record. Morocco, Anti Atlas: Airport Sidi Ifni, 2♂♂, 17-V-2015; Oued Ouarzazate, 5♂♂, 12-VI-2015 (net sweeping)- 1♀, 01-VII-2015 (reared).

Host plants. Flower heads of *Centaurea calcitrapa* L. from which the specimens were reared.

World distribution. Tunisia, Albania, Afghanistan, Austria, Bulgaria, Corsica, Czech Rep., France, Germany, Greece, Hungary, Iran, Iraq, Israel, Italy, Jordan, Kazakhstan, Moldova, Netherlands, Poland, Romania, Russia, Serbia, Sicily, Slovakia, Spain, Switzerland, Turkey, Ukraine, USA (Séguy 1930, Kütük and Yaran 2011, Gharajedaghi et al. 2012, Korneyev et al. 2013).

Subfamily TRYPETINAE (Robineau-Desvoidy, 1830)

Tribe CARPOMYINI Korneyev, 1995

Subtribe CARPOMYINA Norrbom, 1989

Genus *CARPOMYA* Costa, 1854

***Carpomya incompleta* (Becker, 1903)**

New record. Morocco, Anti Atlas: Douar Zaouia, 1♂, 1♀, 11-VI-2015 (net sweeping). First record for Morocco.

World distribution. Italy, Egypt, Ethiopia, Iraq, Israel, Saudi Arabia, Sudan, United Arab Emirates, Burkina Faso, Kenya (Norrbom et al. 1999, Merz 2011, Cope-land 2009, Tankoano et al. 2012).

Tribe TRYPETINI Loew, 1861

Subtribe CHETOSTOMATINA

Genus *CHETOSTOMA* Rondani, 1856

***Chetostoma curvinerve* Rondani, 1856**

New record. Morocco, Rif: Oued El Kelâa, 1♂, 18-XII-2015; Bab El Karne, 1♀, 25-XII-2015 (net sweeping). First record for Morocco.

World distribution. North Africa, Austria, Great Britain, Italy, Netherlands, Portugal, Spain, Switzerland, Uzbekistan, Iran, Israel (Norrbom et al., 1999, Mohamadzade Namin et al. 2010).

Subtribe TRYPETINA Loew, 1861

Genus *EULEIA* Walker, 1835

Euleia heraclei (Linnaeus, 1758)

Literature records. Morocco, Gharb plain: Sidi Slimane (Séguy 1953). First records for the Rif.

New record. Morocco, Rif: Oued Boumarouil, 1♂, 10-V-2014; Aïn El Åakba Larbaâ, 1♂, 18-IV-2015 (net sweeping).

World distribution. Algeria, Madeira, Great Britain, Czech Rep., Estonia, Latvia, Moldova, Poland, Romania, Russia, Spain, Switzerland, Ukraine, Afghanistan, Armenia, Azerbaijan, Georgia, Iran, Israel, Kazakhstan, Kyrgyzstan, Tadjikistan, Turkey, Turkmenistan, Uzbekistan, Japan (Norrbom et al. 1999, Heřman and Dirlbek 2006, Gilasian and Merz 2008).

Euleia marmorea (Fabricius, 1805)

Literature records. Morocco, Rif: Tangier (Séguy 1930, Zimsen 1964).

Acknowledgments

It is our pleasure to warmly thank Professor Valery Korneyev (Schmalhausen Institute of Zoology, Kyiv, Ukraine) for his generous help in confirming the identification of some species and for revising the manuscript.

References

- Al Dhafer HM, Aldryhim YN, Elgharbawy AA, Fadl HH (2012) Insects Associated with Milkweed *Calotropis procera* (Ait.) Ait. in the Ibex Reserve in the Central Region of the Kingdom of Saudi Arabia. The American Entomological Society. Entomological News 122(3): 233–246. <https://doi.org/10.3157/021.122.0304>
- Arambourg Y, Soria F (1961) Note sur quelques Trypetidae de Tunisie. Revue de Pathologie Végétale et de l'Entomologie Agricole France 40: 227–233.
- Becker T, Stein P (1913) Dipteren aus Marokko. Annuaire du Musée Zoologique de l'Académie Impériale de Sciences de Saint-Pétersbourg 18: 62–95.

- Copeland RS (2009) A new species of *Munromyia bezzi* (Diptera: Tephritidae) reared from *Chionanthus battiscombei* (Oleaceae) in northern Kenya. Journal of Natural History 43: 2649–2665. <https://doi.org/10.1080/00222930903216737>
- Copeland RS, White IM, Okumu M, Machera P, Wharton RA (2004) Insects associated with fruits of the *Oleaceae* (Asteridae, Lamiales) in Kenya, with special reference to the Tephritidae (Diptera). In: Evenhuis NL, Kaneshiro KY (Eds) D. Elmo Hardy memorial volume: contributions to the systematics and evolution of Diptera. Bishop Museum Bulletins in Entomology 12: 135–164.
- David KJ, Ramani S (2011) An Illustrated Key to Fruit flies (Diptera: Tephritidae) from Peninsular India and Andaman & Nicobar Islands. Zootaxa 3021: 1–31.
- De Meyer M (2000) Systematic revision of the subgenus *Ceratitis* MacLeay s.s. (Diptera, Tephritidae). Zoological Journal of the Linnean Society 128: 439–467. <http://dx.doi.org/10.1111/j.1096-3642.2000.tb01523.x>
- De Meyer M, Freidberg A (2005) Revision of the fruit fly genus *Capparimyia* (Diptera, Tephritidae). Zoologica Scripta 34: 279–303. <https://doi.org/10.1111/j.1463-6409.2005.00195.x>
- De Meyer M, White IM, Goodger KFM (2013) Notes on the frugivorous fruit fly (Diptera: Tephritidae) fauna of western Africa, with description of a new *Dacus* species. European Journal of Taxonomy 50:1–17. doi: 10.5852/ejt.2013.50
- Dirlbek J, Dirlbek K (1971) Nálezy některých vrtulovitých (Diptera, Trypetidae) z oblasti Krkonoš. (Die Funde einiger Bohrfliegen (Diptera, Trypetidae) aus dem Riesenbirge). Opera Corcontica 7-8: 197–202 (in Czech, German summ.).
- Donati M, Belcari A (2003) A note on insect pests of the caper plant in Jordan, with special reference to *Capparimyia savastani* (Martelli) (Diptera, Tephritidae). Studia Dipterologica 10: 395–400.
- Dirlbek J, Dirlbekova O (1976) Beitrag zur Kenntnis der Bohrfliegen Algeriens (Diptera: Trypetidae). Beiträge zur Entomologie 26: 479–480.
- Evstigneev DA (2011) The fruit flies of the tribe Myopitini (Diptera: Tephritidae) of Ulyanovsk Region (Russia). Ukrainska Entomofaunistyka 2: 21–30
- Foote RH (1984) Family Tephritidae. In: Soos A, Papp L (Eds) Catalogue of Palaearctic Diptera, Vol. 9, Micropezidae - Agromyzidae. Akadémiai Kiadó, Budapest & Elsevier Science Publishers, Amsterdam, 460 pp.
- Freidberg A (1980) A revision of the genus *Goniurellia* Hendel (Diptera: Tephritidae). Journal of the Entomological Society of Southern Africa 43: 257–274.
- Freidberg A, Kugler J (1989) Fauna Palaestina, Insecta IV, Diptera: Tephritidae. The Israel Academy of Sciences and Humanities, Jerusalem, 212 pp.
- Freidberg A, Küyük M (2002) A new species of *Tephritis* from Turkey, with a key to the species of the *Tephritis pulchra* Group. Israel Journal of Zoology 48: 295–311. <https://doi.org/10.1560/H0A3-JJB0-RX53-WMXJ>
- Frey R (1958a) Kanarische Diptera brachycera p.p., von Håkan Lindberg gesammelt. Commentationes Biologicae Societas Scientiarum Fennicae 17(4): 1–63.
- Frey R (1958b) Zur Kenntnis der Diptera brachycera p.p. der Kapverdischen Inseln. Commentationes Biologicae Societas Scientiarum Fennicae 18(4): 1–61.
- Gharali B, Merz B, Lotfalizadeh HD (2005) Report of *Urophora terebrans* (Loew, 1850) (Diptera: Tephritidae) on Centurea from Iran. Entomology and Phytopathology 72(2): 39–39.

- Gilasian E, Merz B (2008) The first report of three genera and fifteen species of Tephritidae (Diptera) from Iran. *Journal of Entomological Society of Iran* 27: 11–14.
- Giray H (1969) Ege bölgesinde yabani otlarda bulunan Trypetidae (Diptera) türleri ile ilgili faunistik araştırmalar. *Ege Üniversitesi Ziraat Fak. Dergisi* 6(1): 71–78.
- Giray H (1979) Türkiye Trypetidae (Diptera) faunasına ait ilk liste. *Türkiye Bitki Koruma Dergisi* 3: 35–46.
- Gharajedaghi Y, Khaghaninia S, Farshbaf Purabad R (2012b) Investigation of the fruit flies (Diptera: Tephritidae) fauna in Ajabshir region (East Azerbaijan Province) with the new record from Iran (Part 2). *Munis Entomology & Zoology* 7(2): 935–945.
- Hancock DL (2013a) A revised checklist of Australian fruit flies (Diptera: Tephritidae). *Australian entomologist* 40 (4): 219–236.
- Hancock DL, Kirk-Spriggs AH, Marais E (2001) An annotated checklist and prvidional atlas of Namibian Tephritidae (Diptera: Schizophora). *Cimbobasia* 17: 41–72.
- Hancock DL, McGuire DJ (2001) New records of Tephritisinae (Diptera: Tephritidae) from Australia and the South Pacific. *Australian Entomologist* 27(4): 101–102.
- Han HY, Kwon YJ (2000) Family Tephritidae. *Insecta Koreana Suppl* 10: 1–113.
- Hendel F (1927) Trypetidae. In: Lindner E (Ed.) *Die Fliegert der Palaarktischen Region* 49: 1–221.
- Hering EM (1937) Neue Bohrfliegen aus der Beckerschen Sammlung. (Dipt.). (16. Beitrag zur Kenntnis der Trypetidae). *Mitteilungen aus dem Zoologischen Museum in Berlin* 22: 244–264.
- Hering EM (1938) Neue palaearktische und exotische Bohrfliegen. 2 I. Beitrag zur Kenntnis der Trypetidae (Dipt.). *Deutsche entomologische Zeitschrift* 2: 397–417.
- Heřman P, Dirlbek J (2006) Contribution to the knowledge of the fruit fly fauna of Algeria, Morocco and Tunisia (Diptera: Tephritidae). In: Kinkorová J (Ed.) *Dipterologica Bohemoslovaca*. 13. *Acta Universitatis Carolinae Biologica* 50(1–2): 53–56.
- Khaghaninia S, Gharajedaghi Y (2012) Faunistic study of the fruit flies (Diptera: Tephritidae) of Kandovan valley with a new record for Iran. *Munis Entomology and Zoology* 7(2): 950–956.
- Khaghaninia S, Gharajedaghi Y, Zarghani E, Farshbaf Purabad R (2012) Some of the fruit flies fauna of Chichekli region with a new report for Iran (Diptera: Tephritidae). *Munis Entomology & Zoology* 7(1): 526–535.
- Klasa A (2001) New for the Polish fauna and rare fruit-flies (Diptera: Tephritidae) with annotated check-list of this family. *Polskie Pismo Entomologiczne* 70: 293–306.
- Korneyev SV (2016) Ревізія роду *Tephritis* (Diptera, Tephritidae) Західної Палеарктики. PhD Thesis. I.I. Schmalhausen Institute of Zoology National Academy of Science of Ukraine, Kiev, 387 pp.
- Korneyev VA (2002) New and little-known Eurasian Dithrycini (Diptera, Tephritidae). *Vestnik Zoologii* 36(3): 3–13.
- Korneyev VA (2003) New and little-known Tephritidae (Diptera, Cyclorrhapha) from Europe. *Vestnik Zoologii* 37(3): 3–12.
- Korneyev VA, Dirlbek J (2000) The fruit flies (Diptera: Tephritidae) of Syria, Jordan and Iraq. *Studia Dipterologica* 7: 463–482.
- Korneyev VA, Evstigneev DA, Karimpour Y, Küyük M, Mohamadzade Namin S, Ömür Koçuncu M, Yaran M (2013) Revision of the *Terellia virens* group (Diptera, Tephritidae) with description of three new species. *Vestnik zoologii* 47(1): 3–25.

- Korneyev VA, Gharali B, Rakhshani H (2011) A review of the genus *Hypenidium* Loew, 1862 (Diptera: Tephritidae). Ukrainska Entomofaunistyka 2(4): 39–46.
- Korneyev VA, White IM (1992) Fruitflies of the genus *Urophora* R.-D. (Diptera, Tephritidae) of East Palaearctic. II. Review of species of the subgenus *Urophora* s.str. (Communication 1). Entomologicheskoye obozreniye 71(3): 688–699. [In Russian. English translation: Entomological Review, Washington 72(5) 1993: 35–47]
- Korneyev VA, White IM (2000) Fruit-flies of the genus *Urophora* R.-D (Diptera, Tephritidae) of East Palaearctic. IV. Conclusion. Entomologicheskoe obozrenie 79: 239–253. [In Russian; English summary; English translation Entomol. Rev Wash 79: 296–309]
- Korneyev VA, White IM (1996) Fruit flies of the genus *Urophora* R.-D. (Diptera: Tephritidae) of Eastern Palaearctic. Entomological Review 76: 499–513.
- Korneyev VA, White IM (1999) Fruit flies of Genus *Urophora* RD. (Diptera: Tephritidae) of East Palaearctic. III. Key to Palaearctic Species. Entomologicheskoe Obozrenie 78(2): 464–482.
- Kütük M (2003) Güney Batı Anadolu bölgesi Tephritidae faunası ve sistematığı üzerine araştırmalar, PhD dissertation, Çukurova University, Adana, 217 pp.
- Kütük M (2005) Two new records of *Tephritis* Latreille, 1804 (Diptera: Tephritidae) from Turkey. Turkish Journal of Zoology 29: 167–170.
- Kütük M, Özgür AF (2003) Faunistical and systematical studies on the genus *Tephritis* Latreille, 1804 (Diptera: Tephritidae) in the South West of Turkey along with new records. Türkiye Entomoloji Dergisi 27: 243–252.
- Kütük M, Yaran M (2011) The fauna and systematic of the genus *Terellia* Rob-Des., 1830 (Diptera: Tephritidae) with a key to the species of *Terellia* in Turkey. Turkish Journal of Zoology 35(4): 509–517.
- Merz B (1992) The fruit flies of the Canary Islands (Diptera: Tephritidae). Entomologica Scandinavica 23: 215–231. <https://doi.org/10.1163/187631292X00308>
- Merz B (1994) Diptera: Tephritidae. Insecta Helvetica Fauna 10: 1–198.
- Merz B (1999) Tephritidae. In: Schumann H, Bahrmann R, Stark A (Eds) Checkliste der Dipteren Deutschlands. Studia Dipterologica 2: 215–218
- Merz B (2000) Additions and corrections to the list of Tephritidae of Hungary (Diptera: Acalyptrata). Folia entomologica hungarica 61: 207–214.
- Merz B (2001) Faunistics of the Tephritidae (Diptera) of the Iberian Peninsula and the Balearic Islands. Bulletin de la Societe Entomologique Suisse 74: 91–98.
- Merz B (2008) Order Diptera, family Tephritidae. In: Van Harten A (Ed.) Arthropod fauna of the UAE. Dar Al Ummah Printing Publishing, Distribution and Advertising, Abu Dhabi, 643–661.
- Merz B (2011) Order Diptera, family Tephritidae, additions and the description of a new species. Arthropod Fauna of the UAE 4: 769–779.
- Merz B, Dawah A (2005) Fruit flies (Diptera, Tephritidae) from Saudi Arabia, with descriptions of a new genus and six new species, Revue Suisse de Zoologie 112(4): 983–1028. <https://doi.org/10.5962/bhl.part.80334>
- Merz B, Korneyev VA (2004) Fauna Europaea: Diptera, Brachycera, Tephritidae. Fauna Europaea version 1.3. <http://www.faunaeur.org> [accessed 25 March 2009]
- Miranda MA, Terrassa J, Miquel M (2008) *Capparimyia savastani* (Martelli, 1911): A New Record of Tephritidae of Economic Significance for Spain Phytoparasitica 36: 247–248.

- Mohamadzade Namin S, Nozari J (2011) The fruit flies (Diptera: Tephritidae) in Kurdistan Province, with new records for Iranian Fauna. Ukrainska Entomofaunistyka 2(4): 47–53.
- Mohamadzade Namin S, Nozari J, Najarpour A (2010a) The Fruit Flies (Diptera: Tephritidae) in the Fauna of Ardabil Province, with New Records for Iran. Ukrainska Entomofaunistyka 1(3): 35–41.
- Mohamadzade Namin S, Nozari J, Rasoulian GR (2010b) The first report of *Goniurellia persignata* Freidberg, 1980 (Diptera: Tephritidae) from Iran. Proceedings of the 19th Iranian Plant Protection Congress, Iranian Research Institute of Plant Protection, Iran 1: 138.
- Morgulis E, Freidberg A, Dorchin N (2015) Phylogenetic Revision of *Tephritisomyia* Hendel (Diptera: Tephritidae), With Description of 14 New Species. Annals of the Entomological Society of America 109(4) 2016: 595–628 <https://doi.org/10.1093/aesa/saw026>
- Munro HK (1984) A Taxonomic Treatise on the Dacidae (Tephritoidea, Diptera) of Africa, Republic of South Africa, Department of Agriculture and Water Supply, Entomology Memoir № 61.
- Norrbom AL, Carroll LE, Thompson RC, White IM, Freidberg A (1999) Systematic database of names. In: Thompson FC (Ed.) Fruit Fly Expert Identification System and Systematic Information Database. Myia 9: 65–251.
- Norrbom AL (2004) Updates to a biosystematic database of world Diptera for Tephritidae throughout 1999. Diptera Data Dissemination Disk (CD-ROM) 2.
- Norrbom AL, Hancock DL (2004) New species and new records of Tephritidae (Diptera) from New Caledonia. Bishop Museum Bulletin in Entomology 12: 67–77.
- Özgür AF, Kütük M (2003) Adana ili meyve sinekleri (Tephritidae: Diptera) faunasının tespiti. Çukurova Üniversitesi Ziraat Fakültesi Dergisi 18(2): 35–44.
- Papachristos AV (2009) Murder by structure: dominance relations and the social structure of gang homicide. American Journal of Sociology 115(1): 74–128. <https://doi.org/10.1086/597791>
- Părvu C, Popescu-Mirceni R, Zaharia R (2006) Faunistic Data on some Dipteran Families (Insecta: Diptera) from Morocco (Results of "Hamada" Expedition 2005) Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa" 49: 271–281.
- Ribera I, Blasco-Zumeta J (1998) Biogeographical links between steppe insects in the Monégros region (Aragón, NE Spain), the Eastern Mediterranean, and Central Asia. Journal of Biogeography 25: 969–986. <https://doi.org/10.1046/j.1365-2699.1998.00226.x>
- Richter VA (1970) [Family Tephritidae (Trypetidae)], In: Key to the insects of the European part of the USSR. Zool. Ins. Acad. Nauk. SSSR, Leningrad, 132–172.
- Séguy E (1930) Contribution à l'étude des Diptères du Maroc. Mémoire de la Société des Sciences Naturelles et Physique du Maroc 24: 1–206.
- Séguy E (1934a) Diptères (Brachycères) (Muscidae, Acalypterae et Scatophagidae). Faune de France 28: 832 pp.
- Séguy E (1934b) Contribution à la connaissance des insectes diptères du Maroc, Notes scientifiques. Terre et Vie 4: 161–162.
- Séguy E (1941) Diptères recueillis par M. Berland dans le Sud Marocain. Annales de la Société Entomologique de France 110: 1–23.
- Séguy E (1949) Diptères de Sud-Marocain (Vallée du Draa) recueillis par M L Berland en 1947-Revue française d'Entomologie 16: 152–161.
- Séguy E (1953) Diptères du Maroc. Encyclopédie Entomologique (B) II. Diptera 11: 77–92.

- Smit JT (2006) The Tephritidae (Diptera) of the Madeiran archipelago, Portugal. In: Merz B (Ed.) Phylogeny, taxonomy, and biology of tephritoid flies (Diptera, Tephritoidea). *Instrumenta Biodiversitatis* 7: 1–274.
- Tankoano PM, Diallo OB, Ouedraogo SN, Some NA, Noula K, Kalinganire A (2012) Inventaire des insectes nuisibles aux fruits des variétés indiennes de *Ziziphus mauritiana* Lam. (Rhamnaceae) au Burkina Faso. Note technique 67: 189–200. www.fruits-journal.org
- Thompson FC (1998) Fruit Fly Expert Identification System and Systematic Information Database. North American Dipterists Society, Backhuys Publishers, Leiden, the Netherlands, 524 pp.
- Wang X (1996) The fruit flies (Diptera: Tephritidae) of the East Asian region. *Acta Zootaxonomica Sinica* 21. Supplement.
- White IM (1988) Tephritid flies: Handbooks for the identification of British insects. Vol. 10, part 5a. Royal Entomological Society of London, 134 pp.
- White IM (1989) A new species of *Terellia* Robineau-Desvoidy associated with *Centaurea solstitialis* L. and a revision of the *Terellia virens* (Loew) species group (Diptera: Tephritidae). *Entomologist's Monthly Magazine* 125: 53–62. <https://doi.org/10.1111/j.1365-3113.1989.tb00289.x>
- White IM (2006) Taxonomy of the Dacina (Diptera: Tephritidae) of Africa and the Middle East. *African Entomology Memoir* 2: 1–156.
- White IM, Elson-Harris MM (1992) Fruit flies of economic significance: their identification and bionomics. CAB International, Wallingford, Oxon, UK and the Australian Center for Agricultural Research, Canberra, Australia, 601 pp.
- White IM, Goodger KFM (2009) African *Dacus* (Diptera: Tephritidae); new species and data, with particular reference to the Tel Aviv University collection. *Zootaxa* 2127: 1–49.
- White IM, Hancock DL (1997) Cabikey to the Indo-Australasian Dacini Fruit Flies. CAB International, Wallingford CDROM.
- White IM, Headrick DH, Norrbom AL, Carroll LE (2000), 33 Glossary. In: Aluja M, Norrbom AL (Eds) *Fruit flies (Tephritidae): phylogeny and evolution of behavior*. CRC Press, London, New York, Washington, 881–924.
- White IM, Korneyev VA (1989) A revision of the Western Palearctic species of *Urophora* Robineau-Desvoidy (Diptera Tephritidae) *Systematic Entomology* 14: 327–374. <https://doi.org/10.1017/S0007485300018459>
- White IM, Marquardt K (1989) A revision of the genus *Chaetorellia* Hendel (Diptera: Tephritidae) including a new species associated with spotted knapweed, *Centaurea maculosa* LAM. (Asteraceae). *Bulletin of Entomological Research* 79: 453–487.
- Whittington AE (2002) *Terellia fuscicornis* (Loew, 1844) (Diptera, Tephritidae) new to Britain. *Entomologist's Monthly Magazine* 138: 119–120.
- Wiedemann CRW (1824) *Munus rectoris* in Academia Christiana Albertina aditurus analecta entomologica ex Museo Regio Haviensi maxime congesta profert iconibusque illustrat. Kiliae [= Kiel], 60 pp. <https://doi.org/10.5962/bhl.title.77322>
- Zarghani E, Khaghaninia S, Pour Abad RF, Gharali B (2010) Two genera and five species as new records for fruit flies fauna of Iran from East Azarbayjan province. *Munis Entomology & Zoology* 5(2): 823–824.
- Zimšen E (1964) The type material of I.C. Fabricius. Munksgaard, Copenhagen, 656 pp.