

Genetic diversity estimates for the *Caenorhabditis* Intervention Testing Program screening panel

Anastasia A Teterina^{1,2}, Anna L Coleman-Hulbert¹, Stephen A Banse¹, John H Willis¹, Viviana I Perez³, Gordon J Lithgow⁴, Monica Driscoll⁵ and Patrick C Phillips^{1§}

¹Institute of Ecology and Evolution, University of Oregon, Eugene, OR, 97403, USA

²Center of Parasitology, Severtsov Institute of Ecology and Evolution RAS, Moscow, Russia

³Division of Aging Biology, National Institute on Aging, Bethesda, MD, 20892, USA

⁴The Buck Institute for Research on Aging, Novato, CA, 94945, USA

⁵Rutgers University, Dept. of Molecular Biology and Biochemistry, Piscataway, NJ, 08854, USA

[§]To whom correspondence should be addressed: pphil@uoregon.edu

Abstract

The *Caenorhabditis* Intervention Testing Program (CITP) was founded on the principle that compounds with positive effects across a genetically diverse test-set should have an increased probability of engaging conserved biochemical pathways with mammalian translational potential. To fulfill its mandate, the CITP uses a genetic diversity panel of *Caenorhabditis* strains for assaying longevity effects of candidate compounds. The panel comprises 22 strains from three different species, collected globally, to achieve inter-population genetic diversity. The three represented species, *C. elegans*, *C. briggsae*, and *C. tropicalis*, are all sequential hermaphrodites, which simplifies experimental procedures while maximizing intra-population homogeneity. Here, we present estimates of the genetic diversity encapsulated by the constituent strains in the panel based on their most recently published and publicly available whole-genome sequences, as well as two newly generated genomic data sets. We observed average genome-wide nucleotide diversity (π) within the *C. elegans* ($1.2e-3$), *C. briggsae* ($7.5e-3$), and *C. tropicalis* strains ($2.6e-3$) greater than estimates for human populations, and comparable to that found in mouse populations. Our analysis supports the assumption that the CITP screening panel encompasses broad genetic diversity, suggesting that lifespan-extending chemicals with efficacy across the panel should be enriched for interventions that function on conserved processes that are shared across genetic backgrounds. While the diversity panel was established by the CITP for studying longevity interventions, the panel may prove useful for the broader research community when seeking broadly efficacious interventions for any phenotype with potential genetic background effects.




Figure 1. Geographic and genetic diversity in the CITP strain panel: (A) The CITP diversity panel represents 22 total nematode strains from three species, *C. elegans* (black circles), *C. briggsae* (orange triangles), and *C. tropicalis* (teal squares). (B) Two-dimensional representation of diversity and relatedness of the strains, LD1 and LD2 represent two latent dimensions. Bright colors indicate the core CITP strains (*C. elegans*: N2-PD1073, JU775, MY16; *C. briggsae*: AF16, HK104, ED3092; and *C. tropicalis*: JU1373, JU1630, QG834), while all other strains in the CITP diversity panel are indicated by muted colors. (C) Nucleotide diversity (π) of the strains used in the CITP diversity panel estimated on non-overlapping 100 kb genomic windows, core strains demonstrated by bright colors, all other strains of the panel in muted colors. The grey boxes represent the mean and the standard deviations.

Description

Model organisms have been fruitful tools for elucidating core biological principles. The power of model organism study, in part, is due to the ability to grow large populations with known genetic makeup. One of the most widely adopted genetic models is the hermaphroditic nematode *Caenorhabditis elegans*. The reproductive style of *C. elegans* makes it particularly easy to generate and maintain large populations of genetically identical individuals. In fact, the control over genetic variability helped make *C. elegans* the first multi-cellular organism to have its entire genome sequenced (*C. elegans* Sequencing Consortium 1998). Despite the success garnered using genetically homogeneous populations, it has become increasingly apparent that many of the phenotypes of interest for study are dependent on genetic background (see Evans *et al.* 2021 for review). Examples of these background-influenced phenotypes range from α -synuclein toxicity (Wang *et al.* 2019), to

behavioral responses to temperature (Stegeman *et al.* 2013), dietary influence on lifespan and reproduction (Stastna *et al.* 2015), and pharmacological efficacy (Lucanic *et al.* 2017). The dependence on genetic background suggests that attempts to identify core biological systems and functionality could benefit from assaying across genetic diversity to identify genetic background-independent phenotypes. One way to achieve this is to use a panel of populations with intra-population homogeneity and inter-population diversity.

With the importance of genetic background effects in mind, the *Caenorhabditis* Intervention Testing Program (CITP) was designed to identify anti-aging and longevity-promoting compound interventions effective in a genetically diverse set of *Caenorhabditis* nematode populations (Lucanic *et al.* 2017). To date, 55 chemical compounds have been tested for reproducible, genetic background-independent effects on longevity (Lucanic *et al.* 2017; Banse *et al.* 2019; Coleman-Hulbert *et al.* 2019; Coleman-Hulbert *et al.* 2020; Morshead *et al.* 2020; Osman *et al.* 2021; Banse *et al.* 2021; Onken *et al.* 2021). The genetic diversity panel used by the CITP is composed of 22 strains from three hermaphroditic species, which facilitates maintenance of intra-population homogeneity. While the full panel is composed of 22 strains, a smaller core sub-panel of nine strains (three from each of three species, *C. elegans* (N2-PD1073, JU775, MY16), *C. briggsae* (AF16, HK104, ED3092), and *C. tropicalis* (JU1373, JU1630, QG843)) is used in initial compound effect characterization (Banse *et al.* 2021). Here, we present estimates of the genetic diversity encapsulated by the constituent strains of the CITP panel, and core sub-panel, based on their most recently published and publicly available whole-genome sequences.

When establishing the genetic diversity panel for compound screening, the CITP sought strains that represented both broad geographic and genetic diversity. The three species represented in the panel are themselves globally distributed, but with ecological restrictions. For example, *C. elegans* is typically isolated from cooler ecological niches than *C. tropicalis* (Kiontke *et al.* 2011; Frézal and Félix 2015; Noble *et al.* 2021), while *C. briggsae* is found in niches that range from cool to warm (Frézal and Félix 2015). The wide species distributions enabled the CITP to assemble a panel of strains isolated worldwide, with representatives from most continents (Figure 1A). We next sought to determine the genetic diversity encapsulated by the panel by collecting genomic data for the strains (see Reagents) and analyzing the genomes for variation within each species (see Software). To visualize the population structure within the panel for each species we used a variational autoencoder approach, popVAE (Battey *et al.* 2021), to project genotypes of the strains on two latent dimensions (Figure 1B). We then determined the nucleotide diversity in the panel. The observed average genome-wide nucleotide diversity (π) among the *C. elegans*, *C. briggsae*, and *C. tropicalis* strains were, respectively, 1.2×10^{-3} , 7.5×10^{-3} , and 2.6×10^{-3} (3.8×10^{-4} , 6.2×10^{-3} , and 2.4×10^{-3} for the nine core CITP strains), which is consistent with previous estimates for those species (Graustein *et al.* 2002; Jovelín *et al.* 2009; Andersen *et al.* 2012; Crombie *et al.* 2019; Noble *et al.* 2021). Figure 1C shows nucleotide diversity estimated on 100 kb windows along the genomes for both the 20 strains in the full panel with available sequencing data, and for the nine strains in the core sub-panel. The estimated level of genetic diversity within these *Caenorhabditis* species is higher than that within human populations (Yu *et al.* 2004; Perry *et al.* 2013; Prado-Martinez *et al.* 2013; Arbiza *et al.* 2014; 1000 Genomes Project Consortium *et al.* 2015) and comparable to that found in mouse populations (Halligan *et al.* 2010; Booker and Keightley 2018). Our analysis supports the assumption that the CITP screening panel encompasses broad genetic diversity, suggesting that lifespan-extending chemicals with efficacy across the panel should be enriched for interventions that function on conserved processes that are shared across genetic backgrounds. While the diversity panel was established by the CITP for studying longevity interventions, the panel may prove useful for the broader research community when seeking broadly efficacious interventions for any phenotype with potential genetic background effects.

Methods

[Request a detailed protocol](#)

To generate estimates of the genetic variability among the strains in the CITP diversity panel we collected publicly available genomic data for 18 of the 22 CITP strains (see Reagents below). Because comparable Illumina-based sequencing was unavailable for N2-PD1073 and ED3092, two strains present in the core-subpanel of nine strains, we generated whole genome data for these two strains using standard protocols. In brief, we used proteinase K to digest whole worms and isolated total genomic DNA using the Genomic DNA Clean & Concentrator kit (Zymo Research). Genomic libraries were then generated using the Nextera XT DNA Library Preparation Kit (Illumina, Inc.). Sequencing was carried out on the Illumina NovaSeq 6000 platform by the Genomics and Cell Characterization Core Facility (GC3F) at the University of Oregon. We then used the two new genomic datasets, along with the 18 available in the NCBI database, to calculate genetic diversity estimates using our software pipeline (described below).

Reagents

While most of the strains in the CITP diversity panel can be obtained from the *Caenorhabditis* Genetics Center (CGC), the *C. elegans* wild-isolates should be ordered from the *Caenorhabditis elegans* Natural Diversity Resource (CeNDR) (Cook *et al.*

2017). Strains unavailable from those sources (e.g., N2-PD1073) can be obtained directly from the CITP upon request. Relevant to the unavailability of PD1073 at the CGC, PD1073 is a close relative of PD1074 which is distributed by the CGC as a wild type reference strain. PD1073, PD1074, and PD1075 are three subclones of the N2-derivative strain VC2010 that were generated in the process of assembling a new N2 reference (“VC2010-1.0”) genome (Yoshimura *et al.* 2019). All strains in the genetic diversity panel and the associated SRA read accession IDs used in this study are listed below:

C. elegans: strain CB4856 (with SRA read accession numbers SRR9322768, SRR9322769, SRR9322775, SRR9322863, SRR9322864), ED3040 (SRR9322720, SRR9322724, SRR9322725, SRR9322726, SRR9322818), JU1088 (SRR9322295, SRR9322300, SRR9322301), JU1652 (SRR9322577, SRR9322578, SRR9322579), JU775 (SRR9322349, SRR9322352, SRR9322355, SRR9322364), MY16 (SRR9322907, SRR9322327, SRR9322913, SRR9322161), QX1211 (SRR9324168, SRR9324217, SRR9323954, SRR9323955);

C. briggsae: AF16 (SRR2002620), HK104 (ERR3063412, ERR3063411, SRR8333803), JU1348 (SRR1793004), JU726 (SRR1792964), NIC20 (SRR1793012), QR25 (SRR1793006);

C. tropicalis: JU1373 (SRR12623131, SRR241785), JU1630 (SRR12623130), NIC58 (SRR12623045, SRR12623063), QG131 (SRR12623047), QG834 (SRR12623046).

No genomic data is publicly available for *C. briggsae* strain JU1264 and *C. tropicalis* NIC122.

Raw reads for *C. elegans* N2-PD1073 and *C. briggsae* ED3092 strains generated in this study were deposited to the SRA database (<https://www.ncbi.nlm.nih.gov/sra>) under the project accession ID PRJNA773598.

The core CITP strains are *C. elegans* N2-PD1073, MY16, and JU775; *C. briggsae* AF16, ED3092, and HK104; *C. tropicalis* JU1373, JU1630, and QG834.

Software

We estimated genetic diversity of the CITP strains using whole-genomic data for 20 out of 22 CITP strains from the panel. Reads from the SRA database were downloaded with SRA-toolkit (the SRA Toolkit Development Team). We evaluated the read quality with FastQC v.0.11.5 (Andrews 2010) and MultiQC v.1.3 (Ewels *et al.* 2016), and filtered and trimmed reads with Skewer v.0.2.2 (Jiang *et al.* 2014). We mapped filtered reads to the *C. elegans*, *C. briggsae*, and *C. tropicalis* genomes obtained from the WormBase database (<https://wormbase.org/>) version WS280 (with accession ID PRJNA13758, PRJNA10731, PRJNA53597, respectively) by BWA-MEM v.0.7.17 (Li 2013) and SAMtools v.1.5 (Li *et al.* 2009), and deduplicated with the Picard tools v.2.17.6 (Broad Institute). We called and filtered variants using GATK v.3.7 (McKenna *et al.* 2010) software and BEDtools v.2.25 (Quinlan and Hall 2010), and estimated the nucleotide diversity using VCFtools v.0.1.15 (Danecek *et al.* 2011), masking repetitive regions, indels, complex variants, and regions with too low or high coverage. We projected genotypes of the CITP strains on two latent dimensions using popVAE (Battey *et al.* 2021), and visualized them in R v.3.5 (R Core team 2018) using packages ggplot2 (Wickham 2016) and ggrepel.

All the tools used for the analysis are publicly available and free, the scripts used to get the results and figures are in the GitHub repository (https://github.com/phillips-lab/CITP_diversity/) and are available under the MIT license. The global map was generated in MATLAB R2021b (MathWorks, Inc.) using the MATLAB Mapping Toolbox. The location coordinates used for the strains are included in the GitHub repository. Figure readability was improved by moving panel relative positions, updating color coding, and improving text aesthetics using Adobe Illustrator 2022 in a manner consistent with image integrity standards.

Acknowledgments: We acknowledge the members of the Phillips labs for helpful discussions. We thank the CITP Advisory Committee, Max Guo and Ronald Kohanski (National Institute on Aging) for extensive discussion. This work was conducted using the resources of the Research Advanced Computing Services at the University of Oregon.

References

- 1000 Genomes Project Consortium, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, Korbel JO, Marchini JL, McCarthy S, McVean GA, Abecasis GR. 2015. A global reference for human genetic variation. *Nature* 526: 68-74. PMID: 26432245.
- Andersen EC, Gerke JP, Shapiro JA, Crissman JR, Ghosh R, Bloom JS, Félix MA, Kruglyak L. 2012. Chromosome-scale selective sweeps shape *Caenorhabditis elegans* genomic diversity. *Nat Genet* 44: 285-90. PMID: 22286215.
- Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. [Computer Software]
- Arbiza L, Gottipati S, Siepel A, Keinan A. 2014. Contrasting X-linked and autosomal diversity across 14 human populations. *Am J Hum Genet* 94: 827-44. PMID: 24836452.

- Banse SA, Lucanic M, Sedore CA, Coleman-Hulbert AL, Plummer WT, Chen E, Kish JL, Hall D, Onken B, Presley MP, Jones EG, Blue BW, Garrett T, Abbott M, Xue J, Guo S, Johnson E, Foulger AC, Chamoli M, Falkowski R, Melentijevic I, Harinath G, Huynh P, Patel S, Edgar D, Jarrett CM, Guo M, Kapahi P, Lithgow GJ, Driscoll M, Phillips PC. 2019. Automated lifespan determination across *Caenorhabditis* strains and species reveals assay-specific effects of chemical interventions. *Geroscience* 41: 945-960. PMID: 31820364.
- Banse SA, Sedore CA, Johnson E, Coleman-Hulbert AL, Onken B, Hall D, Jackson EG, Huynh P, Foulger AC, Guo S, Garrett T, Xue J, Inman D, Morshead ML, Plummer WT, Chen E, Bhamik D, Chen MK, Harinath G, Chamoli M, Quinn RP, Falkowski R, Edgar D, Schmidt MO, Lucanic M, Guo M, Driscoll M, Lithgow GJ, Phillips PC. 2021. Antioxidants green tea extract and nordihydroguaiaretic acid confer species and strain specific lifespan and health effects in *Caenorhabditis* nematodes. *bioRxiv preprint*. DOI: DOI:10.1101/2021.11.09.464847
- Bathey CJ, Coffing GC, Kern AD. 2021. Visualizing population structure with variational autoencoders. *G3 (Bethesda)* 11: jkaa036. PMID: 33561250.
- Booker TR, Keightley PD. 2018. Understanding the Factors That Shape Patterns of Nucleotide Diversity in the House Mouse Genome. *Mol Biol Evol* 35: 2971-2988. PMID: 30295866.
- C. elegans* Sequencing Consortium. 1998. Genome sequence of the nematode *C. elegans*: a platform for investigating biology. *Science* 282: 2012-8. PMID: 9851916.
- Coleman-Hulbert AL, Johnson E, Sedore CA, Banse SA, Guo M, Driscoll M, Lithgow GJ, Phillips PC. 2019. *Caenorhabditis* Intervention Testing Program: the tyrosine kinase inhibitor imatinib mesylate does not extend lifespan in nematodes. *microPublication Biology* 10.17912/micropub.biology.000131. PMID: 32010883.
- Coleman-Hulbert AL, Johnson E, Sedore CA, Banse SA, Guo M, Driscoll M, Lithgow GJ, Phillips PC. 2020. *Caenorhabditis* Intervention Testing Program: the creatine analog β -guanidinopropionic acid does not extend lifespan in nematodes. *microPublication Biology* 10.17912/micropub.biology.000207 PMID: 31998863.
- Cook DE, Zdraljevic S, Roberts JP, Andersen EC. 2017. CeNDR, the *Caenorhabditis elegans* natural diversity resource. *Nucleic Acids Res* 45: D650-D657. PMID: 27701074.
- Crombie TA, Zdraljevic S, Cook DE, Tanny RE, Brady SC, Wang Y, Evans KS, Hahnel S, Lee D, Rodriguez BC, Zhang G, van der Zwagg J, Kiontke K, Andersen EC. 2019. Deep sampling of Hawaiian *Caenorhabditis elegans* reveals high genetic diversity and admixture with global populations. *Elife* 8: e80465. PMID: 31793880.
- Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT, Sherry ST, McVean G, Durbin R, 1000 Genomes Project Analysis Group. 2011. The variant call format and VCFtools. *Bioinformatics* 27: 2156-8. PMID: 21653522.
- Evans KS, van Wijk MH, McGrath PT, Andersen EC, Sterken MG. 2021. From QTL to gene: *C. elegans* facilitates discoveries of the genetic mechanisms underlying natural variation. *Trends Genet* 37: 933-947. PMID: 34229867.
- Ewels P, Magnusson M, Lundin S, Källér M. 2016. MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics* 32: 3047-8. PMID: 27312411.
- Frézal L, Félix MA. 2015. *C. elegans* outside the Petri dish. *Elife* 4: e05849. PMID: 25822066.
- Graustein A, Gaspar JM, Walters JR, Palopoli MF. 2002. Levels of DNA polymorphism vary with mating system in the nematode genus *Caenorhabditis*. *Genetics* 161: 99-107. PMID: 12019226.
- Halligan DL, Oliver F, Eyre-Walker A, Harr B, Keightley PD. 2010. Evidence for pervasive adaptive protein evolution in wild mice. *PLoS Genet* 6: e1000825. PMID: 20107605.
- Jiang H, Lei R, Ding SW, Zhu S. 2014. Skewer: a fast and accurate adapter trimmer for next-generation sequencing paired-end reads. *BMC Bioinformatics* 15: 182. PMID: 24925680.
- Jovelin R, Dunham JP, Sung FS, Phillips PC. 2009. High nucleotide divergence in developmental regulatory genes contrasts with the structural elements of olfactory pathways in *Caenorhabditis*. *Genetics* 181: 1387-97. PMID: 19001295.
- Kiontke KC, Félix MA, Ailion M, Rockman MV, Braendle C, Pénigault JB, Fitch DH. 2011. A phylogeny and molecular barcodes for *Caenorhabditis*, with numerous new species from rotting fruits. *BMC Evol Biol* 11: 339. PMID: 22103856.
- Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv:1303.3997 [q-bio.GN]*.
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* 25: 2078-9. PMID:

19505943.

Lucanic M, Plummer WT, Chen E, Harke J, Foulger AC, Onken B, Coleman-Hulbert AL, Dumas KJ, Guo S, Johnson E, Bhaumik D, Xue J, Crist AB, Presley MP, Harinath G, Sedore CA, Chamoli M, Kamat S, Chen MK, Angeli S, Chang C, Willis JH, Edgar D, Royal MA, Chao EA, Patel S, Garrett T, Ibanez-Ventoso C, Hope J, Kish JL, Guo M, Lithgow GJ, Driscoll M, Phillips PC. 2017. Impact of genetic background and experimental reproducibility on identifying chemical compounds with robust longevity effects. *Nat Commun* 8: 14256. PMID: 28220799.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D, Gabriel S, Daly M, DePristo MA. 2010. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res* 20: 1297-303. PMID: 20644199.

Morshead ML, Sedore CA, Jones EG, Hall D, Plummer WT, Garrett T, Lucanic M, Guo M, Driscoll M, Phillips PC, Lithgow G. 2020. *Caenorhabditis* Intervention Testing Program: the farnesoid X receptor agonist obeticholic acid does not robustly extend lifespan in nematodes. *microPublication Biology* 10.17912/micropub.biology.000257. PMID: 32550518.

Noble LM, Yuen J, Stevens L, Moya N, Persaud R, Moscatelli M, Jackson JL, Zhang G, Chitrakar R, Baugh LR, Braendle C, Andersen EC, Seidel HS, Rockman MV. 2021. Selfing is the safest sex for *Caenorhabditis tropicalis*. *Elife* 10: e62587. PMID: 33427200.

Onken B, Sedore CA, Coleman-Hulbert AL, Hall D, Johnson E, Jones EG, Banse SA, Huynh P, Guo S, Xue J, Chen E, Harinath G, Foulger AC, Chao EA, Hope J, Bhaumik D, Plummer T, Inman D, Morshead M, Guo M, Lithgow GJ, Phillips PC, Driscoll M. 2021. Metformin treatment of diverse *Caenorhabditis* species reveals the importance of genetic background in longevity and healthspan extension outcomes. *Aging Cell* 1: e13488. PMID: 34837316.

Osman HC, Sedore CA, Jackson EG, Battistoni ET, Hall D, Foulger A, Lucanic M, Guo M, Driscoll M, Phillips P, Lithgow GJ. 2021. *Caenorhabditis* Intervention Testing Program: the herbicide diuron does not robustly extend lifespan in nematodes. *microPublication Biology* 10.17912/micropub.biology.000448. PMID: 34585102.

Perry GH, Louis EE Jr, Ratan A, Bedoya-Reina OC, Burhans RC, Lei R, Johnson SE, Schuster SC, Miller W. 2013. Aye-aye population genomic analyses highlight an important center of endemism in northern Madagascar. *Proc Natl Acad Sci U S A* 110: 5823-8. PMID: 23530231.

Prado-Martinez J, Sudmant PH, Kidd JM, Li H, Kelley JL, Lorente-Galdos B, Veeramah KR, Woerner AE, O'Connor TD, Santpere G, Cagan A, Theunert C, Casals F, Laayouni H, Munch K, Hobolth A, Halager AE, Malig M, Hernandez-Rodriguez J, Hernando-Herraez I, Prüfer K, Pybus M, Johnstone L, Lachmann M, Alkan C, Twigg D, Petit N, Baker C, Hormozdiari F, Fernandez-Callejo M, Dabad M, Wilson ML, Stevison L, Campubí C, Carvalho T, Ruiz-Herrera A, Vives L, Mele M, Abello T, Kondova I, Bontrop RE, Pusey A, Lankester F, Kiyang JA, Bergl RA, Lonsdorf E, Myers S, Ventura M, Gagneux P, Comas D, Siegmund H, Blanc J, Agueda-Calpena L, Gut M, Fulton L, Tishkoff SA, Mullikin JC, Wilson RK, Gut IG, Gonder MK, Ryder OA, Hahn BH, Navarro A, Akey JM, Bertranpetit J, Reich D, Mailund T, Schierup MH, Hvilsom C, Andrés AM, Wall JD, Bustamante CD, Hammer MF, Eichler EE, Marques-Bonet T. 2013. Great ape genetic diversity and population history. *Nature* 499: 471-5. PMID: 23823723.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* 26: 841-2. PMID: 20110278.

R Core Team. 2018. R: A language and environment for statistical computing.

Stastna JJ, Snoek LB, Kammenga JE, Harvey SC. 2015. Genotype-dependent lifespan effects in peptone deprived *Caenorhabditis elegans*. *Sci Rep* 5: 16259. PMID: 26539794.

Stegeman GW, de Mesquita MB, Ryu WS, Cutter AD. 2013. Temperature-dependent behaviours are genetically variable in the nematode *Caenorhabditis briggsae*. *J Exp Biol* 216: 850-8. PMID: 23155083.

Wang YA, Snoek BL, Sterken MG, Riksen JAG, Stastna JJ, Kammenga JE, Harvey SC. 2019. Genetic background modifies phenotypic and transcriptional responses in a *C. elegans* model of α -synuclein toxicity. *BMC Genomics* 20: 232. PMID: 30894116.

Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York. ISBN 978-3-319-24277-4. DOI: 10.1007/978-3-319-24277-4

Yoshimura J, Ichikawa K, Shoura MJ, Artiles KL, Gabdank I, Wahba L, Smith CL, Edgley ML, Rougvie AE, Fire AZ, Morishita S, Schwarz EM. 2019. Recompleting the *Caenorhabditis elegans* genome. *Genome Res* 29: 1009-1022. PMID: 31123080.

1/27/2022 - Open Access

Yu N, Jensen-Seaman MI, Chemnick L, Ryder O, Li WH. 2004. Nucleotide diversity in gorillas. *Genetics* 166: 1375-83. PMID: 15082556.

Funding: This work was supported by funding from National Institutes of Health grants (U01 AG045844, U01 AG045864, U01 AG045829, U24 AG056052, and R35 GM131838)

Author Contributions: Anastasia A Teterina: Data curation, Formal analysis, Methodology, Writing - review and editing, Visualization, Software, Funding acquisition. Anna L Coleman-Hulbert: Conceptualization, Project administration, Writing - original draft, Writing - review and editing, Visualization. Stephen A Banse: Conceptualization, Project administration, Writing - original draft, Writing - review and editing, Visualization. John H Willis: Investigation, Writing - review and editing. Viviana I Perez: Funding acquisition, Writing - review and editing, Conceptualization. Gordon J Lithgow: Conceptualization, Funding acquisition, Writing - review and editing. Monica Driscoll: Conceptualization, Funding acquisition, Writing - review and editing. Patrick C Phillips: Conceptualization, Project administration, Supervision, Writing - review and editing, Funding acquisition.

Reviewed By: Erik Andersen

History: Received November 22, 2021 **Revision received** December 15, 2021 **Accepted** January 11, 2022 **Published** January 27, 2022

Copyright: © 2022 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Teterina, AA; Coleman-Hulbert, AL; Banse, SA; Willis, JH; Perez, VI; Lithgow, GJ; Driscoll, M; Phillips, PC (2022). Genetic diversity estimates for the *Caenorhabditis* Intervention Testing Program screening panel. *microPublication Biology*. <https://doi.org/10.17912/micropub.biology.000518>