

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.jfda-online.com

Original Article

Beverage-induced enhanced bioavailability of carbamazepine and its consequent effect on antiepileptic activity and toxicity



Sanmoy Karmakar ^{a,b,*}, Sreerupa Biswas ^a, Rammohan Bera ^a,
Samiran Mondal ^c, Amit Kundu ^a, Md Asif Ali ^a, Tuhinadri Sen ^{a,b}

^a Division of Pharmacology, Department of Pharmaceutical Technology, Jadavpur University, Kolkata 700032, India

^b School of Natural Product Studies, Jadavpur University, 188 Raja S.C. Mallick Road, Kolkata 700032, India

^c Animal Health Centre, Purulia, West Bengal, India

ARTICLE INFO

Article history:

Received 14 April 2014

Received in revised form

14 July 2014

Accepted 16 July 2014

Available online 3 December 2014

Keywords:

bioavailability

black tea

carbamazepine

grapefruit juice

HPLC

ABSTRACT

The present study was undertaken to investigate the food–drug interaction of carbamazepine (CBZ). Common fruit juices [grapefruit juice (GFJ), lime juice (LJ)], known to inhibit the enzyme cytochrome P450 3A4 (CYP3A4), and some widely consumed beverages [milk (M), black tea (BT)] were involved in this study in the presence of CBZ, as might happen during clinical therapy. The effects of the beverages on the pharmacokinetics and drug-induced toxicity of CBZ was observed after concomitant administration for a period of 28 days. Accordingly, the influence of altered bioavailability of CBZ on its antiepileptic activity was investigated. A significant shift in the C_{max} as well as T_{max} of CBZ was observed in the presence of LJ and GFJ. This increase in bioavailability significantly enhanced hepatotoxicity and delayed the onset of tremor and piloerection against pentylene tetrazole (PTZ)-induced seizure in experimental animals. However, increased toxicity of CBZ was found to be absent with BT. Thus, from our observation, LJ or GFJ in the presence of CBZ significantly increased the bioavailability of CBZ, which might lead to increased toxicity and antiepileptic activity of the drug.

Copyright © 2014, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan

LLC. Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

1. Introduction

Antipsychotics and antidepressants are clinically important drugs often used for chronic treatment thereby rendering adverse effects and aspects of drug–drug or food–drug interactions of these drugs to serious clinical vigilance [1]. Carbamazepine (CBZ), a drug with a comparatively narrow

therapeutic index, is a concern among health care professionals because it may result in clinically significant drug interactions. Therefore, bioavailability studies in the presence of other drugs and food are becoming increasingly necessary to avoid toxic effects [2] or clinical failure. It has been observed that inhibition of cytochrome P450 (CYP) enzymes is the most common mechanism that produces serious and potentially life-threatening drug interactions [3]. CBZ undergoes

* Corresponding author. Division of Pharmacology, Department of Pharmaceutical Technology, Jadavpur University, 188 Raja S.C. Mallick Road, Kolkata 700032, India.

E-mail address: sanmoykarmakar@gmail.com (S. Karmakar).

<http://dx.doi.org/10.1016/j.jfda.2014.07.012>

1021-9498/Copyright © 2014, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan LLC. Open access under [CC BY-NC-ND license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

biotransformation by CYP3A4 into carbamazepine-10,11-epoxide [4] and as a consequence of CYP inhibition or induction, plasma concentrations of CBZ may reach toxic or sub-therapeutic levels.

Similar studies are available on incidents of increased bioavailability and enhanced pharmacodynamic effects following concomitant administration of citrus juice and calcium channel blockers such as felodipine and nifedipine [5].

In the present study, we investigated the *in vivo* pharmacokinetic interaction and pharmacodynamic effects of CBZ with commonly consumed beverages including lime juice (*Citrus aurantifolia*; LJ), grapefruit juice (*Citrus paradise*; GFJ), milk (M), and black tea (*Camellia sinensis*; BT) in experimental animals.

2. Methods

2.1. Chemicals and reagents

A CBZ suspension (Tegretol; Novartis Pharmaceuticals, Hyderabad, India), acetonitrile [high performance liquid chromatography (HPLC) grade; Sigma Aldrich, Maharashtra, India], potassium dihydrogen phosphate (Merck Pvt. Ltd, Mumbai, India), ortho-phosphoric acid (Merck Pvt. Ltd, Mumbai, India), methanol (HPLC grade, Sigma Aldrich, Maharashtra, India), and HPLC grade water (resistivity of 18.2 MU cm) generated from a Milli Q water purification system (Elix; Milli Q A10 Academic, Molsheim, France) were used throughout the analysis. Biochemical kits and pentylene tetrazole (PTZ) were obtained from Merck Pvt. Ltd. (Mumbai, India) Crude CBZ and valdecoxib were obtained as gifts from Cosmas Pharmacls (Ludhiana, Punjab, India).

2.2. Animal husbandry and maintenance

Healthy adult male Wistar rats weighing 170–180 g and male Swiss albino mice of 25–30 g, procured from TAAB Biostudy Services, 67/1B, Ibrahimpur Road, Kolkata 700032, India, were used for the study. The animals were grouped and housed in wire cages with not more than six animals per cage in a controlled environment (12 h light–dark cycle, temperature of $25 \pm 2^\circ\text{C}$ and $50 \pm 20\%$ relative humidity). During the period of study, the animals had free access to standard dry pellet diet (Nutrilab Rodent; Provimi) and water *ad libitum*. The study was conducted in accordance with the Institutional Ethical Committee (constituted under the Guidelines Committee for the Purpose of Control and Supervision of Experiments on Animals, Reg. No. 367).

2.3. Preparation of beverages

LJ (*Citrus aurantifolia*), GFJ (*Citrus paradise*), and M were obtained from local commercial sources. Juice was obtained by squeezing the edible portion of the fruits and then filtered. BT (*Camellia sinensis*) extract was prepared by soaking 2 g of black tea leaves (Lipton; Hindustan Unilever Ltd.) in 10 mL of boiling water followed by filtration [6]. Beverages were administered at a dose of 10 mL/kg.

2.4. High performance liquid chromatography conditions

The HPLC system consisted of an LC-20A Dvp pump (Shimadzu, Kyoto, Japan), a Shimadzu UV absorbance detector, a Hamilton syringe, and a Shimadzu SDP-20Avp system controller. The system was equipped with a Luna 5 μ C18 column (250 mm \times 4.6 mm, 5 μ m; Phenomenex, Torrance, CA, USA), preceded by a precolumn. The isocratic mobile phase comprised a mixture of 100 mM potassium dihydrogen orthophosphate (KH_2PO_4 , pH 3.2) and acetonitrile in a ratio of 60:40 v/v was delivered at a flow rate of 0.8 mL/min at 25°C . The UV detector was set at 245 nm and the volume of injection was 20 μ L. The column was equilibrated for at least 20 minutes with the mobile phase flowing through the system prior to the injection of the drug standards. The run time was set at 15 minutes with the system operating at air-conditioned temperature (20°C).

2.5. Grouping and dosing of animals for pharmacokinetic study

The maximum tolerated dose of CBZ in humans was extrapolated to a rat dose [7]. Rats were randomly grouped into nine ($n = 9$). CBZ at a dose of 105.70 mg/kg was administered orally (equivalent to 17.14 mg/kg in humans). Group I received CBZ and water (W) whereas Groups II, IV, VI, and VIII received CBZ 30 minutes after a dose of LJ, GFJ, M, and BT, respectively. Groups III, V, VII, and IX received LJ, GFJ, M, and BT, respectively.

2.6. Serum biochemistry

Serum samples collected on Day 0, Day 2, Day 15, and Day 29 were analyzed for serum glutamic pyruvic transaminase (SGPT), serum glutamic oxaloacetic transaminase (SGOT), alkaline phosphate (ALP), total protein (TP), albumin (ALB), albumin/globulin ratio (A/G), blood urea nitrogen (BUN), and creatinine (CR) using a Microlab-300 auto-analyzer.

2.7. Histopathological study

On Day 29, the animals were sacrificed by cervical decapitation under anesthesia. The liver and kidney tissues were fixed in neutral phosphate-buffered 10% formalin solution and kept in a fridge until the tissues were prepared for histological examination. The tissues were processed, embedded in paraffin, and sectioned at 3–5 μ m. The sections were examined after staining with hematoxylin and eosin (H&E).

2.8. Pharmacokinetic experiment

On Day 1 and Day 28, each animal of Groups I, II, IV, VI, and VIII was anesthetized and blood was collected from the retro-orbital plexus or tail vein into tubes containing EDTA at time points of 0 hours, 0.25 hours, 0.50 hours, 1 hour, 3 hours, 6 hours, and 8 hours. The samples were immediately centrifuged at 1150g at 15°C for 5 minutes and the plasma was separated. The plasma sample collected was stored at -20°C until analysis.

The standard stock solution of analyte was prepared by dissolving the accurately weighed CBZ in dimethyl sulfoxide (DMSO) to get a final concentration of 2 mg/mL. The working stock solutions for the calibration curve were 4.88 µg/mL, 9.77 µg/mL, 19.53 µg/mL, 39.06 µg/mL, 78.125 µg/mL, 156.25 µg/mL, 312.5 µg/mL, 625 µg/mL, 1250 µg/mL, and 2500 µg/mL in methanol. A 10-point standard curve was prepared by spiking 2 µL of the working stock to 98 µL of blank plasma to get final concentrations of 0.1 µg/mL, 0.2 µg/mL, 0.4 µg/mL, 0.8 µg/mL, 1.56 µg/mL, 3.125 µg/mL, 6.25 µg/mL, 12.5 µg/mL, 25 µg/mL, and 50 µg/mL, respectively. The internal standard (Valdecoxib, 20 µg/mL) solution was made in the 100% acetonitrile that had been used to precipitate the protein samples. All stocks were stored in polypropylene tube vials at 2–8°C. The plasma sample aliquot was taken in a microcentrifuge tube and then direct precipitation of the matrix was performed by adding ice-cold acetonitrile (3 × the sample solution) containing the internal solution. The mixture was vortexed in a cyclomixer for 5 minutes and centrifuged at 1500g for 10 minutes at 15°C; 120 µL of the clear supernatant was collected and 20 µL of it was used for HPLC analysis.

2.9. Pharmacodynamic analysis/PTZ-induced convulsion

Male Swiss albino mice were grouped into six ($n = 9$). Group I received PTZ at a dose of 60 mg/kg intraperitoneally (i.p.) [8] and Group II received CBZ and PTZ. Groups III, IV, V, and VI were administered CBZ and PTZ along with LJ, GFJ, M, and BT, respectively. Group II received PTZ 30 minutes after administration of CBZ, and Groups III, IV, V, and VI received PTZ 1.5 hours after CBZ. This was in accordance with the T_{max} obtained from the plasma concentration time profile of rats. The dose of CBZ was 52.86 mg/kg per os (orally, p.o.; equivalent to 4.29 mg/kg in humans). The onset of tremor, piloerection, and convulsions, the number of convulsions, the recovery from the convulsive phase, and the subsequent mortality were recorded for experimental animals as well as the control group.

2.10. Data analysis

Results were expressed as mean ± standard error of the mean (SEM; $n = 9$). Statistical analyses were performed with one-way analysis of variance followed by the post-hoc Dunnett's test; $p < 0.05$ was considered to be statistically significant. To determine the toxicity and pharmacokinetic parameters of CBZ, the concentration–time data were analyzed by the noncompartmental analysis method. The area under the plasma concentration–time curve (AUC) from 0 hours to 8 hours after administration was calculated by the linear trapezoidal method. Mean plasma concentration–time profiles of CBZ per dose group were obtained by averaging the individual concentrations per sampling time point and by plotting the mean concentrations versus the sampling time point. The C_{max} was defined as the highest observed concentration of CBZ in plasma. The T_{max} was defined as the time to reach the C_{max} . Half-lives were determined from rate constants using the relationship: $t_{1/2} = \ln 2/k$.

3. Results

3.1. Serum biochemistry

SGOT, SGPT, CR, and BUN were significantly increased whereas TP was significantly decreased in rats belonging to Groups II and IV. There was a significant elevation in SGPT and CR in rats belonging to Group VI. However, it was observed that significant changes in serum biochemistry parameters were higher in Groups II and IV in comparison to Group VI. Thus, LJ and GFJ caused a greater change in parameters than M. No significant changes were observed in other groups. The data for Day 0 and Day 29 are represented in Table 1.

3.2. Morphologic pathology

Photomicrographs of the liver (Fig. 1) and kidney (Fig. 2) samples showed normal architecture in Group I (CBZ + W) rats. Significant toxic changes were observed in liver and kidney of rats treated with CBZ along with LJ, GFJ, and M. In Group II (CBZ + LJ), histological study of the liver revealed fatty changes, pyknotic nucleus, nuclear vacuolation, central vein congestion, peri-biliary infiltration, and widening of the sinusoidal space. Histological study of the kidney revealed pyknotic necrosis of the nucleus, necrosis of the tubular epithelium, and necrosis of the epithelial cell lining of the glomerulus. In Group IV (CBZ + GFJ), histopathology of the liver showed hepatocyte and hepatic plate disruption, hemorrhage from the central vein, necrotic and degenerative changes, focal fibrosis, and fatty changes in the peri-biliary area. Histological study of the kidney expressed focal necrosed glomeruli, hemorrhage, focal epithelial swelling, Hyalinization of the focal tubule, degenerative changes of the epithelium in the distal convoluted tubule, and swelling. In Group VI (CBZ + M), liver tissues showed biliary hyperplasia, eosinophilia, necrotic changes, granular cytoplasm, cell disruption, and fatty changes whereas kidney tissues revealed hemorrhage, protein cast, coagulative necrosis, and severe hemorrhage. There were no significant histopathological changes in the liver and kidney tissues in Group VIII (CBZ + BT).

3.3. Pharmacokinetic analysis

Significant increases ($p < 0.05$) in the T_{max} and AUC_{0-8} of CBZ in rats belonging to Groups II, IV, VI, and VIII were observed in comparison to rats belonging to Group I, both on Day 1 and Day 28 (Table 2), whereas the C_{max} values of Groups II and IV were significantly increased on Day 1. Therefore, beverages increased the bioavailability of CBZ (AUC increased) when both were administered p.o. concomitantly. However, when we compared the C_{max} , T_{max} , and AUC_{0-8} of Day 1 with the corresponding C_{max} , T_{max} , and AUC_{0-8} of Day 28 of the same group, we observed that there was a decrease in C_{max} and AUC_{0-8} and an increase in T_{max} . The plasma concentration time profiles for Day 1 and Day 28 are represented in Figs. 3 and 4, respectively.

Table 1 – Biochemical analysis of blood serum after administration of CBZ with W, LJ, GFJ, M, or BT.

Parameter	Day 0					Day 29				
	CBZ + W	CBZ + LJ	CBZ + GFJ	CBZ + M	CBZ + BT	CBZ + W	CBZ + LJ	CBZ + GFJ	CBZ + M	CBZ + BT
SGOT	130.33 ± 17.01	131.25 ± 17.61	131.11 ± 17.65	136.67 ± 18.40	132.44 ± 18.49	141.11 ± 18.41	286.52 ± 40.90*	270.00 ± 38.45*	156.89 ± 23.4	129.44 ± 15.95
SGPT	51.89 ± 3.91	55.89 ± 7.25	45.22 ± 5.82	42.78 ± 5.58	49.89 ± 3.74	56.56 ± 4.65	138.52 ± 18.80*	83.67 ± 9.37*	72.11 ± 10.78*	49.22 ± 4.26
ALP	31.33 ± 3.83	30.78 ± 3.94	29.44 ± 3.42	28.22 ± 2.98	33.89 ± 5.48	33.33 ± 4.06	45.78 ± 7.81	41.44 ± 5.98	36.56 ± 4.86	32.89 ± 5.04
TP	7.42 ± 1.05	7.40 ± 0.72	7.51 ± 0.93	8.00 ± 0.83	7.54 ± 0.79	6.84 ± 0.84	4.74 ± 0.43*	4.71 ± 0.63*	6.67 ± 0.91	7.88 ± 1.38
ALB	3.84 ± 0.46	3.90 ± 0.52	3.42 ± 0.47	3.73 ± 0.42	3.73 ± 0.48	3.52 ± 0.39	2.97 ± 0.24	2.72 ± 0.40	3.02 ± 0.22	3.90 ± 0.64
A/G	1.45 ± 0.26	1.53 ± 0.18	1.43 ± 0.27	1.34 ± 0.19	1.48 ± 0.26	1.37 ± 0.28	2.05 ± 0.27	1.67 ± 0.29	1.56 ± 0.15	1.55 ± 0.25
CR	0.35 ± 0.04	0.34 ± 0.05	0.36 ± 0.04	0.31 ± 0.04	0.30 ± 0.04	0.38 ± 0.06	2.58 ± 0.43*	2.57 ± 0.04*	2.10 ± 0.28*	0.31 ± 0.05
BUN	15.48 ± 2.54	15.96 ± 1.59	17.62 ± 2.56	13.36 ± 1.34	17.62 ± 1.85	16.39 ± 1.98	24.64 ± 2.94*	25.86 ± 3.05*	18.66 ± 2.29	18.54 ± 1.98

Data are presented as mean ± SEM (n = 9).

*p < 0.05 versus CBZ + W group.

A/G = albumin/globulin ratio; ALB = albumin; ALP = alkaline phosphatase; BT = black tea; BUN = blood urea nitrogen; CBZ = carbamazepine; CR = creatinine; GFJ = grapefruit juice; LJ = lime juice; M = milk; SEM = standard error of the mean; SGOT = serum glutamic oxaloacetic transaminase; SGPT = serum glutamic pyruvic transaminase; TP = total protein; W = water.

3.4. PTZ-induced convulsions in mice

The mice of Groups III–VI showed significant delayed onset of tremor and piloerection, and episodes of convulsion were absent when compared with Group I. The disappearance of signs and symptoms of convulsion were also significantly faster in Groups III–VI than in Group I. No mortality was observed in Groups III–VI. The anticonvulsive effect against PTZ-induced convulsion was found to be absent in Group II (CBZ + PTZ). However, the increased bioavailability of CBZ in Groups III–VI probably resulted in the anticonvulsive effect of CBZ against PTZ-induced convulsion at the same dose. Data relating to the effect of the beverages on the anticonvulsive effect of CBZ are represented in Table 3.

4. Discussion

CYP enzymes are important for the metabolism of drugs and xenobiotics. This is particularly important for drugs with a narrow therapeutic index that are generally advocated for chronic treatment where the change in bioavailability may eventually affect efficacy and toxicity [9–11]. It has also been documented that such drugs often present interaction with some commonly consumed foodstuffs having relevance to involvement of similar CYP enzyme isoforms [12]. CBZ is one such drug; it is used for epilepsy on a long-term basis with reported clinically relevant liver toxicity [13]. During long-term drug therapy, restrictions regarding food are not convenient as far as patient compliance is concerned. Accordingly, this study concerning the interaction of CBZ with common beverages (LJ, GFJ, BT, and M) was undertaken. CBZ is a known substrate of CYP3A4 [14] whereas the beverages LJ and GFJ are known inhibitors of the same isoform [15,16]. In our study, beverages were co-administered on a chronic basis along with CBZ and some general indicators of toxic manifestations relevant to CBZ were monitored in experimental animals. It was found that LJ and GFJ in combination with CBZ resulted in significant toxicity with particular reference to reduction of body weight, food and water intake, and increases in hepatic markers namely SGPT, SGOT, TP, CR, and BUN. Toxicity was also confirmed through histopathological studies of kidney and liver samples in the affected groups of animals. Histological study of the liver revealed fatty changes, pyknotic nucleus, nuclear vacuolation, central vein congestion, peri-biliary infiltration, and widening of the sinusoidal space while that of the kidney revealed focal necrosed glomeruli, hemorrhage, focal epithelial swelling, hyelination of the focal tubule, degenerative changes of the epithelium in the distal convoluted tubule, and swelling. The observed toxicity complies with the reported toxic manifestation of CBZ that might have occurred due to the reduced activity of CYP3A4, eventually leading to increased bioavailability of CBZ in the presence of LJ or GFJ. [It has been reported that pomegranate juice increases the bioavailability of CBZ in rats [12].]

Accordingly, an attempt was made to justify the above apprehension where the bioavailability of CBZ in the presence of beverages, namely LJ, GFJ, M, and BT, was quantitatively analyzed using HPLC. Our observation indicated a significant shift in the C_{max} as well as T_{max} of CBZ in the presence of LJ or




Fig. 1 – Representative histopathological picture of the male rat liver (H&E, 10 \times). (A) CBZ and W: (a) central vein and (b) normal cells. (B) CBZ and LJ: (a) fatty changes; (b) pyknotic nucleus; (c) nuclear vacuolation; (d) widening of the sinusoidal space; and (e) central vein congestion. (C) CBZ and GFJ: (a) hepatocyte and hepatic plate disruption; (b) hemorrhage from the central vein; and (c) necrotic and degenerative changes. (D) CBZ and M: (a) biliary hyperplasia; (b) increased eosinophilia; and (c) necrotic changes. Fatty changes are present in the entire area. (E) CBZ and BT leaf extract. BT = black tea; CBZ = carbamazepine; GFJ = grapefruit juice; H&E = hematoxylin and eosin; LJ = lime juice; M = milk; W = water.

GFJ, although the maximum shift was found with LJ (at least in our experimental conditions).

Attempts were also made to investigate the anticonvulsant effects owing to the observed increase in bioavailability of CBZ in the presence of LJ or GFJ. Accordingly, the antiepileptic activity of the said drug against PTZ-induced seizure in experimental animals was measured. It was observed that in the presence of LJ or GFJ the onset of tremor and piloerection was

significantly delayed, the disappearance of signs and symptoms of seizures was significantly shortened, and episodes of seizures and even mortality were found to be completely absent as compared to the control group of animals.

In this context, it is interesting to consider that similar effects on CBZ were also observed with BT or M, although these beverages are not reported to be significantly potent CYP3A4 inhibitors; however, a few reports are available in support of




Fig. 2 – Representative histopathological picture of the male rat kidney. (A) CBZ and W (H&E, 10×). (B) CBZ and LJ (H&E 40×): (a) pyknotic nucleus of the necrotic cell and (b) necrosis of the tubular epithelium. (C) CBZ and GFJ (H&E 40×): (a) focal epithelial swelling; (b) hyelinization of the focal tubule; (c) degenerative changes of the epithelium in the distal convoluted tubule; and (d) swelling. (D) CBZ and M (H&E 40×): (a) hemorrhage; (b) protein cast; and (c) coagulative necrosis. (E) CBZ and BT leaf extract (H&E 40×). BT = black tea; CBZ = carbamazepine; GFJ = grapefruit juice; H&E = hematoxylin and eosin; LJ = lime juice; M = milk; W = water.

this theory [17,18]. It is important to mention that the solubility of CBZ is known to increase with an increase in pH, which might be a possible reason for enhanced bioavailability of CBZ when concomitantly administered with M [19,20]. Interestingly

enough, the increased bioavailability-related toxicity of CBZ was found to be absent with BT, which might have some relation to the *in vivo* antioxidant property of BT [21,22] while similar protection was found to be absent in M. Therefore,

Table 2 – Pharmacokinetic parameters of CBZ after administration of CBZ with W, LJ, GFJ, M, or BT on Day 1 and Day 28.

		Pharmacokinetic parameters of CBZ				
		Group I	Group II	Group IV	Group VI	Group VIII
		CBZ + W	CBZ + LJ	CBZ + GFJ	CBZ + M	CBZ + BT
C_{max} (µg/mL)	Day 1	26.40 ± 2.54	32.26 ± 2.84*	30.98 ± 2.49*	28.97 ± 3.16	28.13 ± 3.44
	Day 28	21.88 ± 2.59	18.47 ± 3.15	17.01 ± 2.46	17.49 ± 1.88	17.05 ± 1.67
T_{max} (h)	Day 1	0.5 ± 0.00	1.5 ± 0.00	1.5 ± 0.00	1.5 ± 0.00	1.5 ± 0.00
	Day 28	0.25 ± 0.00	3 ± 0.00	3 ± 0.00	3 ± 0.00	3 ± 0.00
AUC_{0-8}	Day 1	147.79 ± 4.69	185.50 ± 5.31*	177.16 ± 3.88*	173.2 ± 4.62*	168.75 ± 5.02*
	Day 28	72.35 ± 4.36	96.83 ± 5.16*	92.84 ± 4.36*	87.28 ± 3.84*	90.56 ± 4.44*

Data are presented as mean ± SEM (n = 9).

*p < 0.05 versus CBZ + W group.

AUC = area under the plasma concentration–time curve; BT = black tea; CBZ = carbamazepine; GFJ = grapefruit juice; LJ = lime juice; M = milk; SEM = standard error of the mean; W = water.

among the various combinations of beverages and CBZ used in this study, BT appears to be comparatively safe.

Thus, from our observation, the presence of LJ or GFJ along with CBZ leads to increased bioavailability of CBZ. This in turn may increase the probability of liver and kidney toxicity of the drug in a dose that is otherwise expected to be safe.

In summary, this study demonstrates that concomitant administration of beverages such as GFJ, LJ, M, and BT along with a narrow therapeutic index drug like CBZ may result in an increased bioavailability where subsequent monitoring of toxicity to the liver and kidney will be of clinical importance.




Fig. 3 – Plasma concentration time profile of rats treated with CBZ and W, LJ, GFJ, M, or BT on Day 1. BT = black tea; CBZ = carbamazepine; GFJ = grapefruit juice; LJ = lime juice; M = milk; W = water.




Fig. 4 – Plasma concentration time profile of rats treated with CBZ and W, LJ, GFJ, M, or BT on Day 28. BT = black tea; CBZ = carbamazepine; GFJ = grapefruit juice; LJ = lime juice; M = milk; W = water.

Table 3 – The effect of beverages on the anticonvulsive effect of CBZ against PTZ-induced convulsions.

Group	Onset of tremor (min)	Onset of piloerection (min)	Onset of convulsions (min)	Number of convulsions	Recovery from convulsive phase (min)	Mortality (%)
I (PTZ)	0.83 ± 0.09	0.52 ± 0.08	12.667 ± 1.36	3.67 ± 0.38	73.25 ± 8.45	11.11
II (CBZ + W + PTZ)	1.40 ± 0.21	1.24 ± 0.16	11.56 ± 0.87	3.33 ± 0.33	68.63 ± 6.89	11.11
III (CBZ + LJ + PTZ)	6.90 ± 0.98*	6.09 ± 0.86*	—	0*	38.56 ± 4.29*	0
IV (CBZ + GFJ + PTZ)	2.94 ± 0.43*	2.64 ± 0.29*	—	0*	45.00 ± 6.01*	0
V (CBZ + M + PTZ)	2.77 ± 0.27*	2.72 ± 0.38*	—	0*	43.67 ± 5.88*	0
VI (CBZ + BT + PTZ)	2.60 ± 0.18*	1.73 ± 0.22*	—	0*	48.44 ± 6.30*	0

Data are presented as mean ± SEM (n = 9).

*p < 0.05 versus Group I.

BT = black tea; CBZ = carbamazepine; GFJ = grapefruit juice; LJ = lime juice; M = milk; PTZ = pentylene tetrazole; SEM = standard error of the mean; W = water.

Conflicts of interest

The authors declare that there are no conflicts of interest.

Acknowledgments

We would like to extend our sincere thanks to All India Council for Technical Education – Research Promotion Scheme (AICTE-RPS), the Government of India, University Grants Commission – Department of Special Assistance (UGC DSA), and University Grants Commission – Universities with Potential for Excellence Phase II (UGC UPE-II) for providing the financial support needed for this study.

REFERENCES

- Tanaka E, Hisawa S. Clinically significant pharmacokinetic drug interactions with psychoactive drugs: antidepressants and antipsychotics and the cytochrome P450 system. *J Clin Pharm Ther* 1999;24:7–16.
- Morrow JL, Routledge PA. Poisoning by anticonvulsants. *Adverse Drug React Acute Poisoning Rev* 1989;8:97–109.
- Johnson MD, Newkirk G, White JR. Clinically significant drug interactions. *Postgrad Med* 1999;105:93–195.
- Kerr BM, Thummel KE, Wurden CJ, Klein SM, Kroetz DL, Gonzalez FJ, Levy RH. Human liver carbamazepine metabolism: role of CYP3A4 and CYP2C8 in 10, 11-epoxide formation. *Biochem Pharmacol* 1994;47:1969–79.
- Bailey DG, Spence JD, Munoz C, Arnold JM. Interaction of citrus juices with felodipine and nifedipine. *Lancet* 1991;337:268–9.
- Nag Chaudhuri AK, Karmakar S, Roy D, Pal S, Pal M, Sen T. Antiinflammatory activity of Indian Black tea (Sikkim Variety). *Pharmacol Res* 2005;51:169–75.
- Shannon RS, Minakshi N, Nihal A. Dose translation from animal to human studies revisited. *FASEB J* 2007;22:659–61.
- Mizoguchi H, Nakade J, Tachibana M, Ibi D, Someya E, Koike H, Kamei H, Nabeshima T, Itohara S, Takuma K, Sawada M, Sato J, Yamada K. Matrix metalloproteinase-9 contributes to kindled seizure development in pentylene tetrazole treated mice by converting pro-BDNF to mature BDNF in the hippocampus. *J Neurosci* 2011;31:12963–71.
- Apfelbaum JD, Caravati EM, Kerns 2nd WP, Bossart PJ, Larsen G. Cardiovascular effects of carbamazepine toxicity. *Ann Emerg Med* 1995;25:631–5.
- Spiller HA, Krenzelok EP. Carbamazepine overdose. A prospective study of serum concentrations and toxicity. *J Toxicol Clin Toxicol* 1990;28:445–58.
- Tibbals J. Acute toxic reactions to carbamazepine: clinical effects and serum concentration. *J Paediatr* 1992;121:295–9.
- Hidaka M, Okumura M, Fujita K, Ogikubo T, Yamasaki K, Iwakiri T, Setoguchi N, Arimori K. Effects of pomegranate juice on human cytochrome P450 3A (CYP3A) and carbamazepine pharmacokinetics in rats. *Drug Metab Dispos* 2005;33:644–8.
- Gram L, Bentsen KD. Hepatic toxicity of antiepileptic drugs: a review. *Acta Neurol Scand Suppl* 1983;97:81–90.
- Kang P, Liao M, Wester MR, Leeder JS, Pearce RE, Correia MA. CYP3A4-mediated carbamazepine (CBZ) metabolism: Formation of a covalent CBZ-CYP3A4 adduct and alteration of the enzyme kinetic profile. *Drug Metab Dispos* 2008;36:490–9.
- Bailey DG, Dresser GK, Bend JR. Bergamottin, lime juice and red wine as inhibitors of cytochrome P450 3A4 activity: comparison with grapefruit juice. *Clin Pharmacol Ther* 2003;273:529–37.
- Paine MF, Criss AB, Watkin PB. Two major grapefruit juice components differ in time to onset of intestinal CYP3A4 inhibition. *J Pharmacol Exp Ther* 2005;312:1151–60.
- Kazuhiro K, Lee R, Xu H, Harper P, Ito S. Effects of human milk and formula on expression of MDR1, MRP2, and CYP3A in the C2BBE1 intestinal cell model. *Clin Pharmacol Ther* 2004;75. <http://dx.doi.org/10.1016/j.clpt.2003.11.191>.
- Mancuso C, Barone E. Therapeutic use of tea derivatives: all that glitters is not gold. *Blood* 2009;114:2359–60.
- Javadzadeh Y1, Mohammadi A, Khoei NS, Nokhodchi A. Improvement of physicochemical properties of carbamazepine by recrystallization at different pH values. *Acta Pharm* 2009;59:187–97.
- Rao NGR, Patel T, Kumar R. Development of fast dissolving carbamazepine tablets: effect of functionality of superdisintegrants. *Int J Pharm Sci Nanotech* 2010;3:824–33.
- Benzie IF, Szeto YT. Total antioxidant capacity of teas by the ferric reducing/antioxidant power assay. *J Agric Food Chem* 1999;47:633–6.
- Pecorari M, Villaño D, Testa MF, Schmid M, Serafini M. Biomarkers of antioxidant status following ingestion of green tea at different polyphenol concentrations and antioxidant capacity in human volunteers. *Mol Nutr Food Res* 2010;54:S278–83.