

Three new species of *Sesioctonus* Viereck (Hymenoptera, Braconidae, Agathidinae) from Peru

Lidia Sulca^{1,†}, Michael J. Sharkey^{2,‡}

1 Natural History Museum, University of San Marcos, Lima, Peru **2** Department of Entomology, University of Kentucky, Lexington, Kentucky, 40502, USA

† [urn:lsid:zoobank.org/author:E8FE9278-2683-46D3-A6B8-DD7E6037AFBD](https://doi.org/urn:lsid:zoobank.org/author:E8FE9278-2683-46D3-A6B8-DD7E6037AFBD)

‡ [urn:lsid:zoobank.org/author:77B8EC3A-442C-4A7A-AF85-A31C27E257F2](https://doi.org/urn:lsid:zoobank.org/author:77B8EC3A-442C-4A7A-AF85-A31C27E257F2)

Corresponding author: Michael J. Sharkey (msharkey@uky.edu)

Academic editor: C. van Achterberg | Received 15 March 2012 | Accepted 15 May 2012 | Published 21 May 2012

[urn:lsid:zoobank.org/pub:0345EA97-A583-4CAB-9AE8-D8463A6D063B](https://doi.org/urn:lsid:zoobank.org/pub:0345EA97-A583-4CAB-9AE8-D8463A6D063B)

Citation: Sulca L, Sharkey MJ (2012) Three new species of *Sesioctonus* Viereck (Hymenoptera, Braconidae, Agathidinae) from Peru. ZooKeys 196: 1–10. doi: 10.3897/zookeys.196.3086

Abstract

Three new species of *Sesioctonus* (Braconidae: Agathidinae) are described and illustrated, i.e., *Sesioctonus huggerti*, *S. wayquecha*, and *S. bina*. Two new Peruvian species records for *Sesioctonus* are reported: *S. longinoi* and *S. diazi*. A revised key to all known species of *Sesioctonus* is presented.

Keywords

Insecta, taxonomy, biodiversity, parasitoid

Introduction

Sesioctonus Viereck, 1912 is a Neotropical genus of Agathidinae for which the biology is largely unknown, only *S. parathyridis* is recorded as a larval parasitoid of *Parathyris perspicilla* Stall (Lepidoptera: Arctiidae) (Viereck 1912). Briceño (2003) revised the species of *Sesioctonus* and included twenty six species and Sharkey and Briceño (2005) described five new species. Here we describe three new species from Peru and two additional species are reported for the first time from Peru.

Methods

Morphological terminology follows that of Sharkey and Wharton (1997). Figures in this paper that are followed by the letter ‘B’ refer to those in Briceño (2003). The species descriptions are of the holotypes with variation given in parenthesis.

Unless otherwise stated specimens are deposited in the Natural History Museum, University of San Marcos, Lima, Peru (MUSM), with duplicates deposited in the Hymenoptera Institute Collection at the University of Kentucky, USA (HIC).

Results and discussion

Diagnosis

Members of *Sesioctonus* are restricted to the Neotropical realm of the New World and may be distinguished from all other agathidine braconids with the following combination of characters: Mesoscutum smooth, lacking notauli; tarsal claws simple, lacking a basal claw; hind coxal cavities open, sharing a common opening with the metasomal foramen.

Key to *Sesioctonus* species of the world, modified from Sharkey and Briceño (2003)

- | | | |
|------|--|------------------------------|
| 1 | Occipital tubercles present (Figs. 16B–18B) | 2 |
| – | Occipital tubercles absent. (Figs. 19B) | 17 |
| 2(1) | Epicnemial carina straight medially or absent, not indented at midline, between fore coxae) (Figs. 4B, 23B)..... | 3 |
| – | Epicnemial carina bilobed medially (indented at midline, between fore coxae) (Figs. 3B, 22B) | 6 |
| 3(2) | Epicnemial carina complete laterally (Figs. 3B, 22B)..... | 4 |
| – | Epicnemial carina incomplete or absent laterally (Fig. 23B) | 5 |
| 4 | (3) Interantennal space with longitudinal rounded keel; face without median longitudinal carinae | <i>S. garciai</i> Briceño |
| – | Interantennal space lacking longitudinal keel; face with median longitudinal carinae | <i>S. huggertii</i> sp. n. |
| 5(3) | Face with median longitudinal carina (Fig. 13B)..... | <i>S. acrolophus</i> Briceño |
| – | Face without median longitudinal carina (similar to Figs. 12B, 14B) | <i>S. analogus</i> Briceño |
| 6(3) | Mid coxa color variable, but not completely melanic | 7 |
| – | Mid coxa completely melanic..... | 10 |
| 7(6) | Fore wing banded from base: yellow, black, yellow, black... <i>S. chaconi</i> Briceño | |
| – | Fore wing infusate (melanic) | 8 |
| 8(7) | Fore tibia with spines; mid femur yellowish orange..... | 9 |

- Fore tibia without spines; mid femur melanic
..... ***S. longinoi* (part) Sharkey & Briceño**
- 9(8) Median longitudinal carinae of propodeum absent, median areola of metanotum with lateral carinae not meeting posteriorly, subpronope triangular
..... ***S. peruviensis* Briceño**
- Median longitudinal carinae of propodeum present, median areola of metanotum with lateral carinae meeting posteriorly, subpronope oval ***S. bina* sp. n.**
- 10(6) Longitudinal carina(e) of scutellar depression present and fore wing banded from base: yellow, black, yellow, black ***S. venezuelensis* Briceño**
- Longitudinal carina(e) of scutellar depression absent **and/or** fore wing not banded **11**
- 11(10) Mesoscutum black; median areola of metanotum with longitudinal rugosities (Fig. 29B); median tergite of first metasomal segment without pair of lateral longitudinal carina (similar to Fig. 34B); fore wing (RS+M)a vein complete (Fig. 10aB) ***S. kompsos* Briceño**
- Mesoscutum yellowish orange, or if black then not combining other characters **12**
- 12(11) Mesoscutum melanic **13**
- Mesoscutum yellowish orange **14**
- 13(12) Fore wing infusate with large hyaline spot; metasoma reddish brown except last few segments melanic ***S. brasiliensis* Briceño**
- Fore wing either infusate without hyaline spot or hyaline basally and infusate apically; metasoma yellowish orange ***S. dichromus* Briceño**
- 14(12) Median longitudinal carina of propodeum present and complete
..... ***S. ariasi* Briceño**
- Median longitudinal carina of propodeum absent or incomplete **15**
- 15(14) Subpronope triangular, three sides almost equal (Fig. 1B); fore wing 3RSa vein absent (Fig. 10B) ***S. boliviensis* Briceño**
- Subpronope more oval-shaped, weakly triangular with vertical sides longer than dorsal side (Fig. 2B); fore wing 3RSa vein present (Fig. 9B) **16**
- 16(15) Fore wing banded from base: yellow, black, yellow, black ***S. diazi* Briceño**
- Fore wing infusate (melanic) ***S. longinoi* (part) Sharkey & Briceño**
- 17(1) Occiput excavated (similar to Figs. 16B–18B) ***S. eumenetes* Briceño**
- Occiput not excavated (Fig. 19B) **18**
- 18(17) Interantennal space without sharp longitudinal keel **19**
- Interantennal space with sharp longitudinal keel (Fig. 11B) **34**
- 19(18) Basal sterna of metasoma chalk-white **20**
- Basal sterna of metasoma not chalk-white, rather melanic or yellowish orange **22**
- 20(19) Head orange (Fig. 1dB) ***S. susanai* Sharkey & Briceño**
- Head black **20**
- 21(20) Fore and hind coxa pale yellow (Fig. 1bB) ... ***S. stephaniai* Sharkey & Briceño**

–	Fore and hind coxa melanic (Fig. 1aB).....	<i>S. philipi</i> Sharkey & Briceño
22(19)	Median areola of metanotum with lateral carinae meeting posteriorly (Figs. 25B, 26B)	23
–	Median areola of metanotum with lateral carinae absent or, if present, not meeting posteriorly (Figs. 27B, 28B).....	32
23(22)	Epicnemial carina present (Figs. 3B, 4B).....	24
–	Epicnemial carina absent	28
24(23)	Epicnemial carina complete laterally (Fig. 3B)	25
–	Epicnemial carina incomplete laterally (Fig. 4B)	30
25(24)	Hind tibia entirely melanic	<i>S. amazonensis</i> Briceño
–	Hind tibia mostly yellowish orange.....	26
26(25)	Propodeum with central areola absent.....	27
–	Propodeum with central areola present	<i>S. areolatus</i> Briceño
27(26)	Antenna with more than 29 flagellomeres; interantennal space with rounded longitudinal keel (similar to Fig. 12B); hind tibia yellowish orange in basal half, melanic apically.....	<i>S. miyayensis</i> Briceño
–	Antenna with less than 28 flagellomeres; interantennal space without longitudinal keel; hind tibia mostly yellowish orange, melanic apically	<i>S. clavijoi</i> Briceño
28(23)	Scutellar depression with longitudinal carinae; body color yellow, white, and black (Fig. 1cB)	<i>S. torresi</i> Sharkey & Briceño
–	Scutellar depression without longitudinal carinae; body color yellowish orange and black.....	29
29(28)	(RS+M)a vein of fore wing complete, median tergite of first metasomal segment with pair of lateral longitudinal carinae.....	<i>S. ammosakron</i> Briceño
–	(RS+M)a vein fore wing incomplete, median tergite of first metasomal segment without pair of lateral longitudinal carinae	<i>S. wayquecha</i> sp. n.
30 (24)	Epicnemial carina straight medially (between fore coxae) (Fig. 4B); body length less than 3mm.....	<i>S. dominicus</i> Briceño
–	Epicnemial carina bilobed medially (indented at midline, between fore coxae) (Fig. 3B); body length more than 3mm	31
31(30)	Fore wing (RS+M)a vein complete (Fig. 10aB)	<i>S. armandoi</i> Briceño
–	Fore wing (RS+M)a vein incomplete (Fig. 9aB)	<i>S. biospleres</i> Briceño
32(22)	Epicnemial carina present, complete or incomplete laterally (Figs. 3B, 4B)	33
–	Epicnemial carina completely absent.....	<i>S. chrestos</i> Briceño
33(32)	Fore wing banded, yellow, black, yellow, black; labial palpus 3-segmented	<i>S. galeos</i> Briceño
–	Fore wing infusate; labial palpus 4-segmented	<i>S. theskelos</i> Briceño
34(18)	Third and fourth labial palpomeres not fused; first metasomal median tergite with depression posteriad spiracle (Figs. 36B, 37B).....	<i>S. grandis</i> Briceño
–	Third and fourth labial palpomeres fused, first metasomal median tergite with or without depression posteriad spiracle.....	35

- 35(36) First metasomal median tergite with depression posteriad spiracle (similar to Figs. 3, 36).....***S. qui* Briceno**
 – First metasomal median tergite without depression posteriad spiracle
***S. parathyridis* Viereck**

New Species Descriptions

***Sesioctonus huggerti* Sulca & Sharkey, sp. n.**

urn:lsid:zoobank.org:act:A198E0BC-7DFE-42CB-B5AD-9DF7E63597E6

http://species-id.net/wiki/Sesioctonus_huggerti

Figure 1

Diagnosis. Distinguished from all other known species of *Sesioctonus* by the following suite of characters: Interantennal space lacking longitudinal keel, epicnemial carinae straight medially.

Description. ♀ *Length.* Length of body, excluding ovipositor, 5 mm.

Head. Flagellum with 30 flagellomeres. Interantennal space lacking longitudinal keel. Antennal sockets moderately excavated. Face with median longitudinal carina. Gena not expanded posteroventrally. Occipital tubercles present. Occiput not excavated. Mandible concave. Outer tooth of mandible not longer than inner tooth. Maxillary palpus with 4 palpomeres. Third and fourth labial palpomeres not fused. *Mesosoma.* Subpronope elongate-oval. Longitudinal carinae of scutellar depression absent. Scutellum convex. Median areola of metanotum smooth, with median longitudinal carina, and with lateral carinae present and not meeting posteriorly. Propodeum convex. Median longitudinal carina of propodeum absent. Epicnemial carina complete, sharp, straight medially (between fore coxae). Hind femur 6 times as long as wide. (RS+M)a vein of fore wing incomplete. 3RSa vein of fore wing absent. 2–1A vein of hind wing not tubular. Cub vein of hind wing not tubular. *Metasoma.* Median tergite of first metasomal segment without pair of lateral longitudinal carinae. Hind wing with 4 hamuli. First metasomal median tergite without depression posteriad spiracle. Length/width ratio of first metasomal median tergite 0.63. Ovipositor 4 mm.

Color. Head melanic. Maxillary palpomeres melanic. Labial palpomeres melanic. Pronotum melanic. Mesoscutum yellowish orange. Scutellum yellowish orange. Metanotum yellowish orange. Propodeum melanic. Propleuron melanic. Mesopleuron yellowish orange. Metapleuron melanic. Fore coxa melanic. Fore trochanter melanic. Fore trochantellus melanic. Fore femur melanic. Fore tibia melanic. Fore tarsus melanic. Mid coxa melanic. Mid trochanter melanic. Mid trochantellus melanic. Midfemur melanic. Mid tibia melanic. Mid tarsus melanic. Hind coxa melanic. Hind trochanter melanic. Hind trochantellus melanic. Hind femur melanic. Hind tibia melanic. Hind tarsus melanic. Fore wing entirely infusate. Stigma melanic. Hind wing entirely infusate. First metasomal tergum melanic. Second metasomal tergum me-




Figure 1. *Sesiioctonus huggerti*. **1a** dorsal habitus **1b** lateral habitus **1c** wings **1d** propodeum and first metasomal segment.

lanic. Third metasomal tergum melanic. Fourth metasomal tergum melanic. Fifth to eighth metasomal terga melanic. Ovipositor yellowish orange.

♂ Unknown.

Etymology. Named in honor of the late Lars Huggert who collected the type specimen.

Holotype. PERU, Madre de Dios, Puerto Maldonado, 6–11.i.1984, L. Huggert Leg. (Canadian National Collection).

Distribution. Known only from the type locality in Peru.

***Sesiioctonus wayquecha* Sulca & Sharkey, sp. n.**

urn:lsid:zoobank.org:act:19BD24A0-162D-405A-8BF0-5CDA62C5FE86

http://species-id.net/wiki/Sesiioctonus_wayquecha

Figure 2 a,b,c,d

Diagnosis. Distinguished from all other known species of *Sesiioctonus* by the following suite of characters: occipital tubercles absent, epicnemial carina completely absent, antennal socket not excavated, gena moderately expanded posteroventrally.

Description. ♀ *Length.* Length of body, excluding ovipositor, 4.3–5.5 mm.

Head. Flagellum with 31 flagellomeres. Interantennal space lacking longitudinal keel. Antennal sockets not excavated. Face without median longitudinal carina. Gena moderately expanded posteroventrally. Occipital tubercles absent. Occiput not excavated. Mandible concave. Outer tooth of mandible not longer than inner tooth. Maxillary palpus with 5 palpomeres. Third and fourth labial palpomeres not fused. *Mesosoma.* Subpronope elongate-oval. Longitudinal carinae of scutellar depression absent. Scutellum convex. Median areola of metanotum smooth, without median longitudinal carina, and with lateral carinae present and meeting posteriorly. Propodeum convex. Median longitudinal carina of propodeum absent. Epicnemial carina completely absent. Fore tibial spines present. Mid tibia with 7 spines. Hind tibia with 15 spines. Hind femur 3.3–4 times as long as wide. (RS+M)a vein of fore wing complete. 3RSa




Figure 2. *Sesioctonus wayquecha*. **2a** lateral habitus **2b** wings **2c** dorsal scutellum and propodeum **2d** anterior head **2e** dorsal habitus.

vein of fore wing absent. 2–1A vein of hind wing not tubular. Cub vein of hind wing not tubular. *Metasoma*. Median tergite of first metasomal segment without pair of lateral longitudinal carinae. Hind wing with 4 hamuli. First metasomal median tergite without depression posteriad spiracle. Length/width ratio of first metasomal median tergite 0.63. Ovipositor 0.5–0.6 mm.

Color. Head melanic. Antenna melanic. Maxillary palpomeres yellowish orange. Labial palpomeres yellowish orange. Pronotum melanic. Mesoscutum melanic. Scutellum melanic. Metanotum melanic. Propodeum mostly yellowish orange with melanic spots. Propleuron mostly melanic with yellowish orange areas. Mesopleuron melanic. Meta-pleuron yellowish orange. Fore coxa yellowish orange. Fore trochanter yellowish orange. Fore trochantellus yellowish orange. Fore femur yellowish orange. Fore tibia melanic with yellowish orange ends. Fore tarsus melanic. Mid coxa yellowish orange. Mid trochanter yellowish orange. Mid trochantellus yellowish orange. Mid femur yellowish orange. Mid tibia melanic. Mid tarsus melanic. Hind coxa yellowish orange. Hind trochanter yellowish orange. Hind trochantellus yellowish orange. Hind femur yellowish orange. Hind tibia melanic with a yellow orange apical spot. Hind tarsus melanic. Fore wing entirely infusate. Stigma melanic. Hind wing entirely infusate. First metasomal tergum yellowish orange. Second metasomal tergum yellowish orange. Third metasomal tergum yellowish orange. Fourth metasomal tergum yellowish orange but median tergum melanic. Fifth to eighth metasomal terga melanic. Ovipositor yellowish orange.

♂. As in the female (above).

Etymology. Named after the type locality, Wayquecha which means ‘brother’ in Quechua.

Holotype. PERU. ♀, Cusco, Wayquecha, 13°11'21"S, 71°35'04"W 2837m, 6–20.x.2007, C. Castillo. Leg.

Paratypes: PERU: Cusco: 2♀♀, Wayquecha, 13°11'21"S, 71°35'4"W, 2837m, Malaise, 20.x.2007, C. Castillo Leg.; 3♀♀, 1♂, Wayquecha, 13°10'31"S, 71°34' 53"W, 2692m, Malaise, 10.ix.2007, C. Castillo Leg.; ♀, Wayquecha 13°11'S, 71°35'W, 2800m, sweep, 12.ix.2007, C. Castillo Leg.; ♂ Wayquecha, 13°10'31"S, 71°34' 53", 2692m, Malaise, 22.x.2007, C. Castillo Leg.

Distribution. Known only from one locality in Peru.

***Sesioctonus bina* Sulca & Sharkey, sp. n.**

urn:lsid:zoobank.org:act:5AE5EEED-ACF8-47D4-8189-531FFDBB1209

http://species-id.net/wiki/Sesioctonus_bina

Figure 3

Diagnosis. Distinguished from all other known species of *Sesioctonus* by the following suite of characters: occiput not excavated, subpronope oval, median tergite of first metasomal segment with pair of lateral longitudinal carinae.

Description. ♀ *Length.* Length of body, excluding ovipositor, 3.35 mm. Flagellum broken after flagellomere 28. Interantennal space with longitudinal rounded keel. Antennal sockets moderately excavated. Face without median longitudinal carina. Gena not expanded posteroventrally. Occipital tubercles present. Occiput not excavated. Mandible concave. Outer tooth of mandible not longer than inner tooth. Maxillary palpus with 4 palpomeres. *Mesosoma.* Subpronope oval. Longitudinal carinae of scutellar depression absent. Scutellum convex. Median areola of metanotum smooth, without median longitudinal carina, and with lateral carinae present and meeting posteriorly. Propodeum convex. Median longitudinal carina of propodeum present. Epicnemial carina complete, blunt, bilobed medially (between fore coxae). Fore tibial spines present. Mid tibia with 3 spines. Hind tibia with 12 spines. Hind femur 4 times as long as wide. (RS+M)a vein of fore wing incomplete. 3RSa vein of fore wing present. 2–1A vein of hind wing tubular. Cub vein of hind wing not tubular. *Metasoma.* Median tergite of first metasomal segment with pair of lateral longitudinal carinae. Hind wing with 3 hamuli. First metasomal median tergite without depression posteriad spiracle. Length width ratio of first metasomal median tergite 0.5. Ovipositor 1.68 mm.

Color. Head black and yellowish orange. Antenna melanic. Maxillary palpomeres yellowish orange. Labial palpomeres yellowish orange. Pronotum melanic. Mesoscutum melanic. Scutellum melanic. Metanotum melanic. Propodeum melanic. Propleuron melanic. Mesopleuron melanic. Metapleuron melanic. Fore coxa yellowish orange. Fore trochanter yellowish orange. Fore trochantellus yellowish orange. Fore femur yellowish orange. Fore tibia yellowish orange. Fore tarsus mostly yellowish orange, but apical tarsomere melanic. Mid coxa yellowish orange. Mid trochanter yellowish orange. Mid




Figure 3. *Sesioctonus bina*. **3a** lateral head and mesosoma **3b** lateral habitus **3c** wings **3d** propodeum and first metasomal segment.

trochantellus yellowish orange. Mid femur yellowish orange. Mid tibia yellowish orange in basal half, melanic apically, or yellowish orange basally, otherwise melanic. Mid tarsus melanic. Hind coxa melanic. Hind trochanter melanic. Hind trochantellus melanic. Hind femur melanic. Hind tibia melanic in basal and apical third, yellowish orange medially. Hind tarsus melanic. Fore wing entirely infusate. Stigma melanic. Hind wing entirely infusate. First metasomal tergum melanic. Second metasomal tergum yellowish orange but median tergite melanic. Third metasomal tergum melanic. Fourth metasomal tergum melanic. Fifth to eighth metasomal terga melanic. Ovipositor yellowish orange.

♂ unknown

Etymology. Bina means ‘wasp’ in Shipibo, an indigenous language of the Peruvian Amazon.

Holotype. ♀, PERU, Cusco, Rocotal, 16.ix.2007, Sweep, C. Castillo Leg.

New Peruvian Distribution Records

Sesioctonus longinoi

♀, Cusco, Cosñipata valley, San Pedro, 13°03'23"S, 71°32'55"W, 1520m, Malaise, 7.i.2009, C. Castillo. leg. ♀, Cusco, San Pedro, 13°03'23"S, 71°32'55"W, 1520m, Malaise, C. Castillo. leg.

Sesioctonus diazi

♀, Cusco, Reserva Comunal AmaraKaeri, Rio Azul, 12°49,8'24"S, 71°05'55"W, 507m, 11.x.2010. C. Castillo. leg. ♀, Loreto, Alto Nanay, Albarenga north, 18M 0533605 9645694, 142m, C. Castillo leg.

References

- Briceño GR (2003) Taxonomic revision of the Genus *Sesioctonus* Viereck (Hymenoptera: Braconidae: Agathidinae). *Journal of Hymenoptera Research*, 12 (2): 238–271.
- Sharkey MJ, Wharton RA (1997) Morphology and terminology. In: Wharton RA, Marsh PM, Sharkey MJ (Eds), *Manual of the New World genera of the family Braconidae* (Hymenoptera). Special publication of the International Society of Hymenopterists, Number 1. Washington D.C., 19–38.
- Sharkey MJ, Briceño R (2005) Review of *Sesioctonus* Viereck (Hymenoptera: Braconidae: Agathidinae) from Colombia with descriptions of five new species. *Zootaxa* 1002: 45–58.