Original Article Open Access

# Does Multi-morbidity Mediate the Effect of Socioeconomics on Self-rated Health? Cross-country Differences

Shervin Assari<sup>1,2</sup>, Maryam Moghani Lankarani<sup>1,2</sup>

Department of Psychiatry, School of Medicine, <sup>2</sup>Center for Research on Ethnicity, Culture and Health, School of Public Health, University of Michigan, Michigan, USA

#### Correspondence to:

Dr. Shervin Assari, Department of Psychiatry, School of Medicine, Michigan, 4250 Plymouth Rd., Ann Arbor, MI 48109-2700, USA. E-mail: assari@umich.edu

How to cite this article: Assari S, Moghani Lankarani M. Does multi-morbidity mediate the effect of socioeconomics on self-rated health? cross-country differences. Int J Prev Med 2015;6:85.

# **ABSTRACT**

**Background:** This study explored cross-country differences in how multi-morbidity explains the effects of socioeconomic characteristics on self-rated health.

**Methods:** The study borrowed data from the Research on Early Life and Aging Trends and Effects. Participants were 44,530 individuals (age >65 years) who were sampled from 15 countries (i.e. United States, China, India, Russia, Costa Rica, Puerto Rico, Mexico, Argentina, Barbados, Brazil, Chile, Cuba, Uruguay, Ghana and South Africa). Multi-morbidity was measured as number of chronic medical conditions. In Model I, main effects of socioeconomic factors on self-rated health were calculated using country-specific logistic regressions. In Model II, number of chronic conditions were also added to the models to find changes in coefficients for demographic and socioeconomic factors.


**Results:** In the United States, number of chronic medical conditions explained the effect of income on subjective health. In Puerto Rico, number of chronic medical conditions explained the effect of marital status on subjective health. In Costa Rica, Argentina, Barbados, Cuba, and Uruguay, number of chronic medical conditions explained gender disparities in subjective health. In China, Mexico, Brazil, Russia, Chile, India, Ghana and South Africa, number of chronic medical conditions did not explain the effect of demographic or socioeconomic factors on subjective health.

**Conclusions:** Multi-morbidity explains the effect of demographic and socioeconomic factors on subjective health in some but not other countries. Further research is needed.

Keywords: Chronic medical conditions, physical health, sociodemographics, subjective health

# INTRODUCTION

Research has consistently shown cross-country differences in subjective and objective measures of health and



well-being. [1-6] World Values Survey, European Values Study, Eurobarometer, and Latino-barometer, have shown self-rated health, physical health, life expectancy, and all-cause mortality vary across countries. [7-10] What is, however, not known is cross-country differences in factors that influence health and well-being.

Socioeconomic status (SES) is associated with health and well-being. [11] High social status provides multiple benefits for individuals through a higher access to financial and material resources. [12] Unfortunately, most of our knowledge about the mechanisms of the effects of SES on health and well-being has originated from

studies conducted within a single country, [13,14] thus cross-country comparison of mechanisms for the effects of SES on health is still needed. [15-18]

Gender is a demographic and social factor with major implications for health and well-being.[19] Compared to men, women tend to report a higher number of chronic conditions and also poorer self-reported health.[19,20] Pinquart and Sörensen listed four reasons for gender differences in subjective measures of health. First, women may have lower material resources due to gender inequities and gendered social power. Gendered labor market may result in lower stable employment among women.[11] Among those in the labor market, women's pensions are lower than men's. [21,22] Women more frequently live in poverty than older men.<sup>[23]</sup> Women also have lower access to health care resources while also requiring more care in later life than men.<sup>[24]</sup> Due to gender difference in longevity, a larger part of women's life is spent with illness and disabilities.[25] Third, older women are more likely to be widowed than older men. [23] In the United States, nearly four times as many older women than men live alone. [24] Finally, due to gender differences in response sets, women may have a higher tendency to report negative feelings and emotions.[26]

Multi-morbidity - defined as multiple chronic medical conditions-is an important predictor of subjective health,[11] as people with higher number of chronic medical conditions report lower physical and mental health-related quality of life (HRQL).[27-29] In one study, chronic medical conditions had a consistent dose-dependent effect on all aspects of well-being, namely physical functioning, role limitation due to physical health problems, bodily pain, general health, vitality, social functioning, role limitation due to emotional problems and mental health.[30] Another study in the United States showed that any additional chronic medical condition leads to 3-4 decrements of mental quality of life decline.[31] Another study among 21,133 showed an association between the presence of one chronic condition and lower well-being across all subjective health domains including physical function, fatigue, pain, emotional distress, and social function. The presence of two or more morbidities was associated with a larger decrement in HRQL, compared to a single condition.<sup>[32]</sup> Another large study of adults showed that after adjustments for SES, and health behaviors (i.e., smoking, alcohol consumption, and physical activity), people with 3 or more chronic medical conditions are more likely to report poor general health, mental distress, physical distress, and activity limitations compared to individuals who have one or two chronic conditions.[33,34]

Despite our knowledge about cross-country differences in health and well-being, [7-10] it is not clear if countries are also different in the mechanisms by which demographic (i.e., age and gender), and socioeconomic (e.g., income and education) influence health, and if chronic medical conditions explain some of health disparities associated with low SES. [1-6] Research on Early Life and Aging Trends and Effects (RELATE) has provided a unique opportunity for cross-country studies on socioeconomic, chronic medical conditions and subjective health among the elderly. The RELATE is a cross-national survey in 15 countries located in North America, South America, Asia, and Africa. [35,36]

Using RELATE data, this study compared 15 countries for the mediating effect of multi-morbidity (defined as number of chronic medical conditions) on the effects of demographic and socioeconomic characteristics on self rated health.

# **METHODS**

# Study design and participants

Data came from the RELATE, a study composed of the following national surveys: (1) China Health and Nutrition Study (CHNS), (2) Chinese Longitudinal Healthy Longevity Survey, (3) Costa Rican Study of Longevity and Healthy Aging, (4) Puerto Rican Elderly: Health Conditions, (5) Study of Aging Survey on Health and Well-being of Elders, (6) WHO Study on Global Ageing and Adult Health, and (7) Wisconsin Longitudinal Study. [35,36]

We included 44,530 elderly individuals (age >65 years). Participants were sampled in the following 15 countries: Puerto Rico (n=3,102,7.0%), Costa Rica (n=2,374,5.3%), China (n=2,2034,49.5%), India (n=2069,4.6%), Mexico (n=2,031,4.6%), Ghana (n=1,923,4.3%), Russia (n=1,860,4.2%), Brazil (n=1633,3.7%), Cuba (n=1,345,3.0%), South Africa (n=1,333,3.0%) Barbados (n=1,160,2.6%), Uruguay (n=1,066,2.4%), Chile (n=958,2.2%), United States (n=894,2.0%), Argentina (n=748,1.7%).

From the participating countries, Barbados, Puerto Rico and the United States represent high-income countries; Argentina, Cuba, Uruguay, Chile, Costa Rica, Brazil, Mexico, and Russia represent upper middle-income countries; China and India represent lower middle-income countries; and Ghana represents a low-income country. Thus, the RELATE project represented countries from a diverse range in national income levels.

### Measures

Demographic characteristics

The study collected data on age (continuous variable) and gender (dichotomous variable).

## Socioeconomic characteristics

The study also measured the following socioeconomic factors: Education level (four-level categorical variable of no schooling, primary to elementary, secondary to intermediate, and higher), household income (continuous variable), and marital status (dichotomous variable of married and other statuses).

# Multi-morbidity

We measured multi-morbidity as number of chronic medical conditions. Chronic medical conditions were measured based on self-reported physician diagnosis of seven conditions (i.e., hypertension, cancer, pulmonary disease, heart disease, diabetes, stroke, and arthritis). The multi-morbidity score potentially ranged from 0 to 7, with a higher score indicative of the higher number of chronic medical conditions. The agreement between self-report and physician diagnosis of chronic medical conditions has been shown to be high (kappa: 0.74–0.92). [37]

#### Main outcome

The outcome was subjective health (self-rated health), which was measured using a single-item. Although it was measured using a five Likert scale (i.e., very bad health, bad, moderate, good, very good), we collapsed the scale to a dichotomous outcome: Poor health (very bad health, bad health) versus good health (moderate health, good health, very good health).

Single-item measures of subjective health and well-being have been frequently used and are well accepted. [38-46] Test-retest reliability for single-item is high for brief time intervals, with a range from 0.7 to 0.8. [42] Validation studies have consistently shown surprisingly high correlations between single-item indicators and multi-item measures of perceived health. [42] Single-item measures of self-rated health also have high predictive validity for mortality above and beyond traditional risk factors. [47]

# **Data analysis**

Statistical analysis was conducted using SPSS 20.0 for Windows (IBM Inc. Armonk, NY). We used country-specific logistic regressions to determine if the effects of demographic (i.e., age and gender) and socioeconomic (i.e., education, income, and marital status) factors vary across countries or not.

We also tested if number of chronic medical conditions differently explains the links between demographic and socioeconomic factors and subjective health. Although most country-specific surveys had sampling weights, the current study did not apply sampling weights, because sampling weights were not applicable to data from United States (Wisconsin) and China (CHNS).

# **Mediation analysis**

For this purpose, we used hierarchical regression approach. In Model I, only demographic and socioeconomic factors were included. In Model II, we also entered number of

chronic physical conditions. Changes in the regression coefficients of demographic and socioeconomic factors from significant to nonsignificant association was defined as full mediation. [48]

# **RESULTS**

Table 1 shows that mean age, education, income, and number of chronic medical conditions were significantly different across countries.

#### Model I

Based on Model I, high age was associated with better subjective health in China, Costa Rica and Argentina. High age was associated with poor subjective health in Barbados, India, South Africa and Russia. Age and subjective health were not significantly associated In Puerto Rico, United States, Mexico, Brazil, Chile, Cuba and Uruguay [Table 2].

Female gender was associated with worse subjective health in China, Costa Rica, Puerto Rico, Barbados, Cuba and Uruguay. Gender was not associated with subjective health in other countries [Table 2].

In the US, Ghana and South Africa, education was not associated with subjective health High education was associated with high subjective health in all other countries [Table 2].

Surprisingly, high income was associated with poor subjective health in Ghana. In Argentina, Chile, Cuba, Uruguay and South Africa, income was not associated with subjective health. In other countries, high-income was associated with better subjective health [Table 2].

Being currently married was associated with better subjective health in Mexico. Being currently married was associated with worse subjective health in Costa Rica, Puerto Rico, and Brazil. Marital status was not associated with subjective health in other countries [Table 2].

#### **Model II**

The number of chronic conditions explained the association between gender and subjective health in Costa Rica, Argentina, Barbados, Cuba, Uruguay. In these 5 countries, the significant association between gender and subjective health became nonsignificant after introducing the number of chronic conditions to the model [Table 3].

In the US, number of chronic conditions explained the association between income and subjective health. That said, the association between income and subjective health that was significant in the previous model did not remain significant after entering number of conditions to the model [Table 3].

In Puerto Rico, number of conditions explained the association between marital status and subjective health. That said, the association between marital status and

Table 1: Comparison of demographic and socioeconomic factors among participants in each country

	China	Costa Rica		United States	Mexico	Argentina	Barbados	Brazil	Chile	Cuba	Uruguay	India	Ghana	South Africa	Russia	P*
Age	82.99	78.94	76.25	66.15	74.78	73.96	75.54	76.63	74.84	75.99	73.96	73.39	75.05	73.81	74.32	< 0.001
	(11.74)	(9.11)	(7.72)	(0.52)	(6.83)	(6.01)	(7.01)	(6.77)	(6.81)	(7.5)	(6.15)	(6.12)	(7.22)	(6.52)	(5.98)	
Education	1.55	1.94	2.55	0.78	1.83	2.53	2.15	1.74	2.05	2.34	2.3	1.48	1.31	1.59	2.99	< 0.001
	(0.86)	(0.75)	(0.93)	(2.06)	(0.98)	(0.76)	(0.63)	(0.75)	(1)	(0.73)	(0.92)	(1.15)	(1.1)	(1.46)	(0.72)	
Income	5.02	0.5	5.96	24.54	12.02	3.13	9.49	3.83	280.04	1.48	44.07	16.37	148.51	11.96	67.91	< 0.001
	(9.07)	(1.25)	(9.34)	(35.92)	(32.9)	(5.35)	(32.48)	(7.68)	(261.74)	(5.71)	(74.19)	(43.11)	(259.31)	(38.87)	(51.78)	
Number	0.93	0.82	1.22	1.15	0.88	1.11	0.98	1.02	1.07	1.28	1.05	0.66	0.47	0.83	2	< 0.001
of CMCs	(1.57)	(0.91)	(1.02)	(1.02)	(0.9)	(0.91)	(0.9)	(0.98)	(0.96)	(1.04)	(0.94)	(0.89)	(0.73)	(0.96)	(1.27)	

\*ANOVA, CMCs=Chronic medical conditions

subjective health that was significant in the absence of number of chronic conditions in the previous model did not stay significant after entering this variable to the model. Interestingly, after entering number of conditions to the model, the nonsignificant association between age and subjective health became statistically significant. This finding suggests that number of conditions may have a suppressor effect on age – subjective health association [Table 3].

In China, Mexico, Brazil, Russia, Chile, India, Ghana, and South Africa, number of chronic medical conditions did not fully explain the associations between demographic and SES and subjective health. In other words, in these countries, the associations between demographic and SES factors and subjective health were above and beyond number of chronic medical conditions [Table 3].

# **DISCUSSION**

This study documented major cross-country differences in the role of multi-morbidity - defined as number of chronic medical conditions - in explaining the associations between demographic and socioeconomic characteristics and subjective health. In Costa Rica, Argentina, Barbados, Cuba, and Uruguay, number of chronic medical conditions explain the association between gender and subjective health. In the United States number of chronic medical conditions explained the association between income and subjective health. In Puerto Rico, they explain the association between marital status and subjective health. In China, Mexico, Brazil, Russia, Chile, India, Ghana and South Africa, number of conditions do not explain the associations between SES and subjective health. These findings suggest that mechanisms for the effects of demographic and socioeconomic factors on self rated health may vary across countries.

Women reported worse self rated health only in six countries (i.e. China, Costa Rica, Puerto Rico, Barbados, Cuba and Uruguay), and in Costa Rica, Argentina, Barbados, Cuba, and Uruguay, chronic conditions explained such an association. Literature has consistently shown gender differences in health and well-being. [19] Although some researchers have attributed the effect of gender on perceived health to income, [21-23] and marital status, [23,24] the effect of gender on self rated health was above and beyond these SES factors in our study. Research has shown that women report higher number of chronic conditions and poorer health, [19] and we found that higher rate of multi-morbidity may be the reason female gender is associated with poor self-rated health, at least in some countries in South America.

Multi-morbidity had a suppressor effect on the association between age and self rated health in Puerto Rico. That is, only after controlling for number of chronic conditions, the association between age and self rated health became significant. This finding has important implications for future research on age effect on subjective health in Puerto Rico. The inclusion of a suppressor to a regression equation is known to enhances the relationship between subjective health and age in this country by removing the unwanted variance from the predictor variable. [49] In other words, in Puerto Rico, studies that wish to test the association between age and well-being need to control for number of conditions not as a mediator, but as a suppressor.

With an exception of only three countries (i.e. United States, Ghana and South Africa), education was associated with a better subjective health. United States, Ghana, and South Africa were the three countries where education did not independently predict self rated health. In some countries, the effect of education on health and well-being may be due to income or marital status, [50] which were both controlled in our study.

In nine countries including the United States, high income was associated with better self rated health. Surprisingly, high income was found to be associated with poor subjective health in Ghana. In five other countries, income was not associated with self rated health. Why income does not have a protective effect on perceived health in Ghana, Argentina, Chile, Cuba, Uruguay, and South Africa needs more research.

Table 2: Results of model I on the associations between demographics and socioeconomics and self-rated health

	В	SE	OR	95% CI for OR		
				Lower	Upper	
China						
Age	-0.020	0.002	0.981	0.977	0.984	
Female	0.129	0.040	10.138	10.052	10.23	
Currently married	0.070	0.044	10.072	0.984	10.16	
Education level	-0.250	0.026	0.779	0.741	0.819	
Income	-0.022	0.003	0.979	0.972	0.985	
Costa Rica						
Age	-0.014	0.005	0.986	0.976	0.997	
Female	0.193	0.098	10.213	10.001	10.469	
Currently married	0.255	0.104	10.290	10.052	10.58	
Education level	-0.351	0.072	0.704	0.611	0.811	
Income	-0.141	0.064	0.869	0.766	0.986	
Puerto Rico	0.111	0.001	0.000	0.700	0.000	
Age	-0.007	0.007	0.993	0.980	10.00	
Female	0.492	0.101	10.636	10.343	10.99	
Currently married	0.239	0.106	10.270	10.032	10.56	
Education level	-0.490	0.058	0.612	0.546	0.687	
Income	-0.023	0.007	0.977	0.964	0.991	
United States	0.025	0.007	0.377	0.304	0.551	
Age	0.031	0.277	10.032	0.599	10.77	
Female	0.111	0.337	10.117	0.577	20.16	
Currently married	-0.516	0.361	0.597	0.294	10.21	
Education level	-0.726	0.627	0.484	0.234	10.65	
Income	-0.720 $-0.020$	0.027	0.484	0.142	10.00	
Mexico Mexico	-0.020	0.010	0.900	0.901	10.00	
	0.002	0.000	0.007	0.001	10.01	
Age Female	-0.003 0.052	0.008 0.111	0.997 10.054	0.981		
				0.848	10.31	
Currently married	-0.194	0.110	0.824	0.663	10.02	
Education level	-0.261	0.066	0.770	0.676	0.878	
Income	-0.011	0.003	0.989	0.983	0.996	
Argentina	0.000	0.045	0.004	0.000	0.004	
Age	-0.036	0.015	0.964	0.936	0.994	
Female	0.382	0.197	10.466	0.996	20.15	
Currently married	0.026	0.191	10.026	0.705	10.49	
Education level	-0.725	0.126	0.484	0.378	0.620	
Income	-0.029	0.023	0.971	0.928	10.01	
Barbados						
Age	0.029	0.010	10.030	10.010	10.05	
Female	0.374	0.145	10.453	10.094	10.93	
Currently married	-0.185	0.154	0.831	0.614	10.12	
Education level	-0.315	0.117	0.730	0.581	0.918	
Income	-0.005	0.003	0.995	0.990	10.000	
Brazil						
Age	-0.002	0.008	0.998	0.983	10.01	

Contd...

Age, gender, income, education, chronic medical conditions, and perceived health have complex inter-relations, and

Table 2: Contd

Table 2: Contd					
Female	0.142	0.115	10.153	0.920	10.444
Currently married	0.334	0.115	10.397	10.114	10.752
<b>Education level</b>	-0.196	0.073	0.822	0.713	0.948
Income	-0.035	0.009	0.965	0.949	0.982
Chile					
Age	0.001	0.011	10.001	0.980	10.022
Female	0.185	0.160	10.203	0.879	10.648
Currently married	-0.110	0.161	0.896	0.653	10.228
<b>Education level</b>	-0.295	0.075	0.744	0.643	0.862
Income	0.000	0.000	10.000	0.999	10.000
Cuba					
Age	-0.010	0.009	0.990	0.972	10.008
Female	0.456	0.141	10.577	10.197	20.078
Currently married	0.087	0.149	10.091	0.815	10.461
<b>Education level</b>	-0.265	0.091	0.767	0.642	0.917
Income	-0.018	0.012	0.982	0.960	10.005
Uruguay					
Age	0.013	0.011	10.013	0.990	10.035
Female	0.341	0.158	10.406	10.032	10.916
Currently married	0.102	0.154	10.107	0.818	10.498
<b>Education level</b>	-0.467	0.084	0.627	0.532	0.738
Income	0.000	0.001	10.000	0.998	10.002
India					
Age	0.042	0.008	10.043	10.026	10.060
Female	0.048	0.124	10.049	0.822	10.339
Currently married	-0.057	0.121	0.945	0.745	10.198
Education level	-0.223	0.069	0.800	0.699	0.917
Income	-0.009	0.003	0.991	0.985	0.996
Ghana					
Age	0.053	0.009	10.054	10.036	10.073
Female	0.118	0.180	10.126	0.790	10.603
Currently married	-0.104	0.179	0.902	0.635	10.280
Education level	-0.111	0.082	0.895	0.761	10.051
Income	0.001	0.000	10.001	10.000	10.001
South Africa					
Age	0.034	0.012	10.035	10.012	10.059
Female	0.188	0.196	10.206	0.822	10.771
Currently married	-0.093	0.192	0.911	0.626	10.327
Education level	-0.036	0.057	0.965	0.864	10.078
Income	-0.008	0.005	0.992	0.983	10.002
Russia					
Age	0.056	0.011	10.058	10.035	10.080
Female	0.234	0.143	10.263	0.954	10.672
Currently married	-0.178	0.137	0.837	0.640	10.094
Education level	-0.253	0.086	0.776	0.656	0.918
Income	-0.003	0.001	0.997	0.994	0.999
( I=( ontidence interval	( )U = ( )ddc rot	10 CF-C+00	dard arror		

CI=Confidence interval, OR=Odds ratio, SE=Standard error

these links may vary based on country. Dynamics of social power associated with age, gender, income, education vary from one to another country. The degree by which income, education, gender, age, and other demographic and social

Table 3: Results of Model II on the associations between demographics, socioeconomics, and number of chronic medical conditions and self-rated health

	В	SE	OR	95% CI for OR		
				Lower	Upper	
China						
Age	-0.010	0.002	0.990	0.986	0.994	
Female	0.141	0.043	1.151	1.057	1.254	
Currently married	0.066	0.048	1.069	0.973	1.173	
Education level	-0.217	0.029	0.805	0.760	0.852	
Income	-0.020	0.004	0.980	0.973	0.987	
Number of CMCs	0.136	0.011	1.146	1.122	1.171	
Costa Rica						
Age	-0.013	0.006	0.987	0.976	0.998	
Female	0.079	0.102	1.082	0.886	1.322	
Currently married	0.221	0.108	1.247	1.009	1.540	
Education level	-0.375	0.075	0.687	0.593	0.797	
Income	-0.123	0.062	0.884	0.782	0.999	
Number of CMCs	0.440	0.054	1.552	1.396	1.726	
Puerto Rico	0.110	0.001	11002	1.000	11720	
Age	-0.015	0.007	0.985	0.972	0.999	
Female	0.327	0.106	1.386	1.125	1.708	
Currently married	0.155	0.112	1.168	0.938	1.453	
Education level	-0.487	0.062	0.615	0.544	0.694	
Income	-0.024	0.002	0.977	0.963	0.991	
Number of CMCs	0.802	0.058	2.230	1.992	2.496	
United States	0.002	0.030	2.230	1.332	2.430	
Age	0.220	0.281	1.246	0.718	2.163	
Female	-0.090	0.453	0.914	0.716	2.703	
Currently married	-0.030 $-0.272$	0.433	0.762	0.376	1.964	
Education level	-0.272 -0.941	1.061	0.702	0.230	3.125	
Income	-0.941 -0.018	0.015	0.983	0.049	1.011	
Number of CMCs	0.667	0.013	1.949	1.352	2.808	
Mexico	0.007	0.107	1.545	1.332	2.000	
	0.006	0.000	0.004	0.070	1 010	
Age	-0.006	0.008	0.994	0.979	1.010	
Female	-0.023	0.113	0.977	0.783	1.220	
Currently married	-0.194	0.112	0.824	0.662	1.025	
Education level	-0.269	0.067	0.764	0.669	0.872	
Income	-0.011	0.003	0.990	0.983	0.996	
Number of CMCs	0.242	0.057	1.274	1.140	1.424	
Argentina	0.007	0.010	0.000	0.000	0.00	
Age	-0.037	0.016	0.963	0.933	0.995	
Female	0.241	0.212	1.273	0.839	1.930	
Currently married	0.026	0.207	1.027	0.685	1.540	
Education level	-0.658	0.133	0.518	0.399	0.671	
Income	-0.044	0.028	0.957	0.906	1.011	
Number of CMCs	0.888	0.110	2.431	1.959	3.018	
Barbados						
Age	0.030	0.011	1.031	1.009	1.052	
Female	0.151	0.156	1.163	0.856	1.579	
Currently married	-0.158	0.165	0.854	0.618	1.179	
<b>Education level</b>	-0.301	0.125	0.740	0.579	0.946	

Table 3: Contd...

lable 3: Contd					
Income	-0.005	0.003	0.995	0.990	1.001
Number of CMCs	0.825	0.091	2.281	1.908	2.727
Brazil					
Age	-0.003	0.008	0.997	0.981	1.014
Female	0.032	0.122	1.032	0.812	1.311
Currently married	0.305	0.122	1.357	1.068	1.723
Education level	-0.213	0.077	0.808	0.695	0.940
Income	-0.035	0.009	0.966	0.949	0.983
Number of CMCs	0.656	0.062	1.927	1.707	2.175
Chile					
Age	0.008	0.011	0.992	0.971	1.014
Female	0.015	0.168	1.015	0.731	1.411
Currently married	-0.099	0.167	0.906	0.653	1.258
Education level	-0.310	0.077	0.734	0.631	0.854
Income	0.000	0.000	1.000	0.999	1.001
Number of CMCs	0.665	0.089	1.944	1.633	2.314
Cuba					
Age	-0.013	0.010	0.987	0.968	1.006
Female	0.051	0.154	1.052	0.779	1.422
Currently married	0.004	0.159	1.004	0.734	1.372
Education level	-0.275	0.097	0.759	0.628	0.918
Income	-0.018	0.015	0.982	0.954	1.012
Number of CMCs	0.924	0.082	2.519	2.145	2.958
Uruguay					
Age	0.009	0.012	1.009	0.985	1.034
Female	0.112	0.174	1.118	0.795	1.574
Currently married	-0.063	0.170	0.939	0.673	1.310
Education level	-0.493	0.090	0.611	0.512	0.729
Income	0.000	0.001	1.000	0.997	1.002
Number of CMCs	0.963	0.089	2.620	2.200	3.119
India					
Age	0.041	0.008	1.042	1.025	1.059
Female	0.010	0.125	1.010	0.790	1.291
Currently married	-0.075	0.122	0.928	0.730	1.178
Education level	-0.298	0.072	0.742	0.645	0.854
Income	-0.010	0.003	0.990	0.984	0.996
Number of CMCs	0.345	0.058	1.413	1.260	1.584
Ghana					
Age	0.053	0.009	1.054	1.036	1.073
Female	0.068	0.182	1.070	0.749	1.530
Currently married	-0.118	0.179	0.889	0.625	1.263
Education level	-0.131	0.083	0.877	0.745	1.032
Income	0.001	0.000	1.001	1.000	1.001
Number of CMCs	0.160	0.085	1.173	0.992	1.387
South Africa					
Age	0.040	0.012	1.041	1.016	1.065
Female	0.060	0.206	1.062	0.709	1.590
Currently married	-0.172	0.201	0.842	0.568	1.248
<b>Education level</b>	-0.062	0.059	0.940	0.838	1.055
Income	-0.008	0.005	0.992	0.982	1.002
Number of CMCs	0.458	0.081	1.580	1.348	1.853

Contd... Contd...

Table 3: Contd...

	В	SE	OR	95% CI for OR		
				Lower	Upper	
Russia						
Age	0.053	0.011	1.055	1.032	1.078	
Female	0.072	0.152	1.074	0.798	1.447	
Currently married	-0.126	0.144	0.882	0.664	1.171	
<b>Education level</b>	-0.290	0.091	0.748	0.626	0.894	
Income	-0.005	0.002	0.995	0.992	0.998	
Number of CMCs	0.561	0.054	1.753	1.576	1.949	

CI=Confidence interval, OR=Odds ratio, SE=Standard error

factors shape populations' access to resources may vary from one to another country. Thus, countries may differ on how social and health factors confound, mediate, or suppress the effects of demographic and socioeconomic factors on health and well-being. There is empirical evidence suggesting that the complex inter-relation between socioeconomics, chronic medical conditions, and subjective health vary across countries. [16-18,46]

Interestingly, number of chronic medical conditions did not fully explain the effect of age on subjective health in any of the 15 countries. Age and subjective health were differently linked across countries, as well. In three countries (i.e. China, Costa Rica and Argentina), high age was associated with better subjective health, while in four countries (i.e. Barbados, India, South Africa and Russia), high age was associated with low subjective health. Interestingly, in seven other countries (i.e. Puerto Rico, United States, Mexico, Brazil, Chile, Cuba and Uruguay), there was no linear association between age and subjective health. There are studies reporting a positive net effect of age on well-being among the elderly. [51,52]

The number of chronic medical conditions explained the association between marital status and subjective health in Puerto Rico. Being currently married was associated with better subjective health in Mexico. The same status was linked to worse subjective health in three other countries (i.e. Costa Rica, Puerto Rico, and Brazil). Marital status was not associated with subjective health in the other 11 countries. The two main hypotheses explaining health effects of marital status are marriage protection and marriage selection. [53-55] None of these theories, however, have hypothesized being married as a risk factor for health. More research is needed to find out why married people do worse in Costa Rica, Puerto Rico, and Brazil.

# Limitations

There are a number of limitations to consider. First, the study was a cross-sectional study, and causative associations are not plausible. In addition, cross – country differences in the validity of chronic medical conditions measurement and even subjective health cannot be

ruled out. The study did not measure mental health as a predictor of self-rated health. Exclusion of individuals with a mental disorder limits generalizability of the findings to the populations. The study also ignores the type of chronic conditions, as only number of conditions were included in the models.

#### **CONCLUSIONS**

To conclude, our study suggests cross-country differences in the way number of chronic medical conditions may explain the effects of demographic and socioeconomic factors on subjective health.

#### **ACKNOWLEDGEMENTS**

Publication of this manuscript was possible with the support of Department of Psychiatry and School of Public Health, University of Michigan to Shervin Assari. Publicly available data were used for this analysis. Data were downloaded from ICPSR. Research on Early Life and Aging Trends and Effects was funded by United States, Department of Health and Human Services, National Institutes of Health, and National Institute on Aging (K25AG027239) to Dr. Mary McEniry, University of Michigan. According to the following website (http://www.icpsr. umich.edu/icpsrweb/DSDR/studies/34241), these data are freely available for the public.

Received: 04 Jan 14 Accepted: 03 Jan 15

# Published: 03 Sep 15 REFERENCES

 Zborowski M. Cultural components in response to pain. J Soc Issues 1952;8:16-30.

- Zola IK. Culture and symptoms An analysis of patients' presenting complaints. Am Sociol Rev 1966;31:615-30.
- Fabrega H Jr. The study of disease in relation to culture. Behav Sci 1972;17:183-203.
- Kleinman A, Kleinman J. Somatization: The interconnections in Chinese society among culture, depressive experiences, and the meaning of pain. Berkeley (CA): University of California Press; 1985.
- Gureje O, Simon GE, Ustun TB, Goldberg DP. Somatization in cross-cultural perspective: A World Health Organization study in primary care. Am J Psychiatry 1997;154:989-95.
- Gureje O, Ustün TB, Simon GE. The syndrome of hypochondriasis: A cross-national study in primary care. Psychol Med 1997;27:1001-10.
- Easterlin RA, McVey LA, Switek M, Sawangfa O, Zweig JS. The happiness-income paradox revisited. Proc Natl Acad Sci U S A 2010;107:22463-8.
- Jen MH, Sund ER, Johnston R, Jones K. Trustful societies, trustful individuals, and health: An analysis of self-rated health and social trust using the World Value Survey. Health Place 2010;16:1022-9.
- Jen MH, Jones K, Johnston R. Global variations in health: Evaluating Wilkinson's income inequality hypothesis using the World Values Survey. Soc Sci Med 2009;68:643-53.
- Kim D, Kawachi I, Hoorn SV, Ezzati M. Is inequality at the heart of it? Cross-country associations of income inequality with cardiovascular diseases and risk factors. Soc Sci Med 2008;66:1719-32.
- Pinquart M, Sörensen S. Influences of socioeconomic status, social network, and competence on psychological well-being in the elderly. Psychol Aging 2000;15:187-224.
- Anderson C, Kraus MW, Galinsky AD, Keltner D. The local-ladder effect: Social status and subjective well-being. Psychol Sci 2012;23:764-71.

- Diener E, Suh EM, Lucas RE, Smith HE. Subjective well-being: Three decades of progress. Psychol Bull 1999;125:276-302.
- Mousavi M, Shiani M, Mohammadi MA, Sadjadi H, Tabatabaee F, Assari S. Life satisfaction in Iran; A national representative study. Sci Res Essays 2011:6:1839-44.
- Assari S. Cross-country variation in additive effects of socio-economics, health behaviors, and comorbidities on subjective health of patients with diabetes.
   Diabetes Metab Disord 2014;13:36.
- Assari S, Lankarani RM, Lankarani MM. Correction: Cross-country differences in the association between diabetes and disability. J Diabetes Metab Disord 2014;13:73.
- Assari S. Cross-country differences in the additive effects of socioeconomics, health behaviors and medical comorbidities on disability among older adults with heart disease. J Tehran Univ Heart Cent 2015;10:24-33.
- Assari S, Moghani Lankarani M. Association between heart disease and subjective health in ten north, middle, and South American countries. Int J Travel Med Glob Health 2014;2:141-7.
- Olsson A, Hasselgren M, Hagquist C, Janson S. The association between medical conditions and gender, well-being, psychosomatic complaints as well as school adaptability. Acta Paediatr 2013;102:550-5.
- Jette A. Disability trends and transitions. In: Binstock R, George LK, editors. Handbook of Aging and the Social Sciences. 4th ed. San Diego, CA: Academic Press; 1996. p. 94-117.
- Golombok S, Fivush R. Gender Development. New York: Cambridge University Press: 1994.
- Moen P. Gender, age, and the life course. In: Binstock RH, George LK, editors. Handbook of Aging and the Social Sciences. 4th ed. San Diego, CA: Academic Press; 1996. p. 171-87.
- 23. Arber S, Ginn J. Women and aging. Rev Clin Gerontol 1994;4:349-58.
- Hobbs FB, Damon BL. 65+in the United States. Washington, DC: U.S. Government Printing Office; 1996.
- Katz S, Branch LG, Branson MH, Papisdero JA, Beck JC, Greer DS. Active life expectancy. N Engl J Med 1983;309:1218-24.
- Phillips DL, Segal BE. Sexual status and psychiatric symptoms. Am Soc Rev 1969:34:58-72.
- Azarkeivan A, Hajibeigi B, Alavian SM, Lankarani MM, Assari S. Associates
  of poor physical and mental health-related quality of life in beta
  thalassemia-major/intermedia. J Res Med Sci 2009;14:349-55.
- Alishiri GH, Bayat N, Fathi Ashtiani A, Tavallaii SA, Assari S, Moharamzad Y. Logistic regression models for predicting physical and mental health-related quality of life in rheumatoid arthritis patients. Mod Rheumatol 2008;18:601-8.
- Khedmat H, Karami GR, Pourfarziani V, Assari S, Rezailashkajani M, Naghizadeh MM. A logistic regression model for predicting health-related quality of life in kidney transplant recipients. Transplant Proc 2007;39:917-22.
- Crouchley K, Daly A. Chronic Disease and Quality of Life in Western Australia. Department of Health, Western Australia; 2007. Available from: http://www.health.wa.gov.au/publications/documents/Chronic%20Disease.pdf. [Last accessed on 2015 Jan].
- Bayliss M, Rendas-Baum R, White MK, Maruish M, Bjorner J, Tunis SL. Health-related quality of life (HRQL) for individuals with self-reported chronic physical and/or mental health conditions: Panel survey of an adult sample in the United States. Health Qual Life Outcomes 2012;10:154.
- Rothrock NE, Hays RD, Spritzer K, Yount SE, Riley W, Cella D. Relative to the general US population, chronic diseases are associated with poorer health-related quality of life as measured by the Patient-Reported Outcomes Measurement Information System (PROMIS). J Clin Epidemiol 2010;63:1195-204.
- Centers for Disease Control and Prevention: Behavioral Risk Factor Surveillance System Survey Data; 2007. Available from: http://www.cdc.gov/

- brfss/annual\_data/annual\_2007.htm. [Last accessed on 2014 Apr].
- Chen HY, Baumgardner DJ, Rice JP. Health-related quality of life among adults with multiple chronic conditions in the United States, Behavioral Risk Factor Surveillance System, 2007. Prev Chronic Dis 2011;8:A09.
- McEniry M. Research on Early Life and Aging Trends and Effects (RELATE):
   A Cross-National Study. ICPSR34241-v1. Ann Arbor, Ml: Inter-university
   Consortium for Political and Social Research [distributor]; 2013. Available
   from: http://www.icpsr.umich.edu/icpsrweb/DSDR/studies/34241. [ Last
   accessed on 2014 Apr].
- McEniry M, Moen S, McDermott J. Methods Report on the Compilation of the RELATE Cross-National Data on Older Adults from 20 Low, Middle and High Income Countries. Ann Arbor, MI: University of Michigan; 2013.
- Baumeister H, Kriston L, Bengel J, Härter M. High agreement of self-report and physician-diagnosed somatic conditions yields limited bias in examining mental-physical comorbidity. J Clin Epidemiol 2010;63:558-65.
- Andrews FM. Social indicators of perceived life quality. Soc Indic Res 1974;1:279-99.
- Andrews FM, Crandall R.The validity of measures of self-reported well-being. Soc Indic Res 1976;3:1-19.
- Knäuper B, Turner PA. Measuring health: Improving the validity of health assessments. Qual Life Res 2003;12 Suppl 1:81-9.
- 41. Verbrugge LM.A global disability indicator. J Aging Stud 1997;11:337-62.
- McDowell I. Measuring Health: A Guide to Rating Scales and Questionnaires.
   3<sup>rd</sup> ed. New York: Oxford University Press; 2006.
- Bélanger A, Berthelot JM, Guimond E, Houle CA. Head-to-head comparison of two generic health status measures in the household population: McMaster Health Utilities Index (Mark3) and the EQ-5D. Ottawa, Canada: Statistics Canada Report; 2002. p. 1-62.
- McDowell I. Measures of self-perceived well-being. J Psychosom Res 2010;69:69-79.
- Hunt MO.The individual, society, or both? A comparison of black, Latino, and white beliefs about the causes of poverty. Soc Forces 1996;75:293-332.
- Assari S. Race and Ethnicity, Religion Involvement, Church-based Social Support and Subjective Health in United States: A Case of Moderated Mediation. Int J Prev Med 2013;4:208-17.
- Idler EL, Benyamini Y. Self-rated health and mortality: A review of twenty-seven community studies. | Health Soc Behav 1997;38:21-37.
- Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. J Pers Soc Psychol 1986;51:1173-82.
- MacKinnon DP, Krull JL, Lockwood CM. Equivalence of the mediation, confounding and suppression effect. Prev Sci 2000;1:173-81.
- Kaljee LM, Chen X. Social capital and risk and protective behaviors: A global health perspective. Adolesc Health Med Ther 2011;2011:113-122.
- Cheng ST.Age and subjective well-being revisited: A discrepancy perspective. Psychol Aging 2004;19:409-15.
- Horley J, Lavery JJ. Subjective well-being and age. Social Indicators Research 1995;34:275-82
- Waldron I, Hughes ME, Brooks TL. Marriage protection and marriage selection – Prospective evidence for reciprocal effects of marital status and health. Soc Sci Med 1996;43:113-23.
- Waldron I, Weiss CC, Hughes ME. Marital status effects on health: Are there differences between never married women and divorced and separated women? Soc Sci Med 1997;45:1387-97.
- Waldron I, Hughes ME, Brooks TL. Marriage protection and marriage selection – prospective evidence for reciprocal effects of marital status and health. Soc Sci Med 1996;43:113-23.

Source of Support: Nil, Conflict of Interest: None declared.