

Long noncoding RNA H19 mediated the chemosensitivity of breast cancer cells via Wnt pathway and EMT process

Hongli Gao¹

Guijun Hao¹

Yue Sun²

Liye Li³

Yukun Wang²

¹Department of Oncology, The Fourth People's Hospital of Jinan, Jinan, Shandong 250031, People's Republic of China; ²Department of Internal Medicine, ³Department of Surgery (Breast Surgery), Affiliated Hospital of Shandong Academy of Medical Sciences, Jinan, Shandong 250031, People's Republic of China

Background: Breast cancer is still one of the major public health burdens worldwide, although there is tremendous progress in early diagnosis and treatment of breast cancer. Tamoxifen was one of the most popular endocrine therapies for early-stage estrogen receptor (ER) + breast cancer patients. However, a high incidence of drug resistance develops along with poor prognosis in breast cancer. Currently, long noncoding RNAs (lncRNAs) are emerging and are well suited to play a role in the development of cancer and tamoxifen resistance. However, there is little reported about the relationship of breast cancer resistance to tamoxifen and lncRNA H19. Here, we validated that lncRNA H19 was highly expressed in breast cancer especially in patients with late stage (III and IV), compared to normal tissues and early stage cancers (I and II).

Methods: Quantitative polymerase chain reaction (qPCR) was utilized for comparison of lncRNA H19 expression level in breast cancers with different stages. qPCR and Western blotting were used to detect gene and protein, respectively.

Results: We found that lncRNA H19 expression level manipulated breast cancer cell proliferation both in parental breast cancer cell lines and tamoxifen-resistant cell lines. Knockdown of lncRNA H19 elevated tamoxifen sensitivity for promoting cell growth and inhibiting apoptosis in tamoxifen-resistant breast cancer cells. Moreover, knockdown of H19 inhibited Wnt pathway and epithelia–mesenchymal transition in tamoxifen-resistance breast cancer cells.

Conclusion: Taken together, the results of this study provided the evidence for H19 in regulating tamoxifen-resistant breast cancer and might provide novel options in the future treatment of tamoxifen-resistance breast cancer patients.

Keywords: lncRNA H19, breast cancer, tamoxifen-resistance, EMT, Wnt pathway

Background

Breast cancer is the most common invasive cancer in women which causes approximately 4 million deaths in worldwide each year.¹ For decades, classification of breast cancer stages was based on the tumor size and the metastasis of cancer to lymph nodes or other parts.² Recently, molecular classification, such as Her-2, estrogen receptor (ER), and progesterone receptor, was developed. It has great advances for personalized treatment of breast cancer with better outcomes than conventional classification.^{3–5} Tamoxifen is a nonsteroidal antiestrogen and as an adjuvant endocrine therapy has been extensively used for early ER positive and node negative breast cancer treatment. However, tamoxifen resistance can result in breast cancer recurrence and mortality.⁶ Therefore, it is essential to explore molecular targets involved in tamoxifen resistance of breast cancer.

It was well known that long noncoding RNA (lncRNA) involved in diabetes, neurological diseases, and cancers.^{7–9} H19 was the first identified lncRNA in human breast

Correspondence: Yukun Wang
Department of Internal Medicine,
Affiliated Hospital of Shandong Academy
of Medical Sciences, No 38 Wuyingshan
Road, Jinan, Shandong 250031, People's
Republic of China
Tel +86 531 5862 8612
Email dr_yukunwang@sina.com

cancer, and the expression of H19 is higher in ER positive cells.¹⁰ Additionally, in ER-negative MDA-MB-231 cell line, highly proliferation ability of these cells was notably with ectopic overexpression of H19. The function of lncRNA in proliferation, invasion, and metastasis have become a hot spot of research.

Several studies have preliminarily revealed that lncRNA mediates the metastasis of tumor by regulating the epithelia–mesenchymal transition (EMT) process of tumor cells. Hu et al screened a series of high levels of lncRNAs in MCF10A (MCF10A/Twist) with high expression of twisted protein (TWIST) and proved that dysregulation lncRNAs was involved in the EMT process by regulating the activation of the target gene and the corresponding pathways of WNT and MAPK, inducing cell invasion and metastasis.¹¹ It was reported that that H19 can lead to de-repression of β -catenin, which eventually activated Wnt/ β -catenin pathway and hence potentiated osteogenesis.¹² Moreover, H19 mediates methotrexate resistance in colorectal cancer through activating Wnt/ β -catenin pathway.¹³ lncRNA aquaporin can promote the migration and metastasis of EMT in intestinal cancer cells.¹⁴ Zhou et al demonstrated H19 regulated EMT and mesenchymal–epithelial transition on breast cancer.¹⁵ Therefore, expression level of lncRNAs plays an important role in Wnt pathway and EMT in tumor cells.

The aim of the present study was to identify new mechanisms underlying tamoxifen resistance in order to lay the foundation for therapies that could have greater efficacy. We focused on the mechanism of lncRNA H19 for drug sensitivity regulation in tamoxifen-resistance breast cancer. Taken together, this article might give a new hint for further clinical treatment of tamoxifen-resistant breast cancer patients.

Patients and methods

Patients' recruitment and tissue sample collection

A total of 30 pairs of fresh breast cancer tissues and adjacent normal tissues from 30 treatment-naïve breast cancer patients were collected in two Chinese hospitals (The Fourth People's Hospital of Jinan and Affiliated Hospital of Shandong Academy of Medical Sciences). The clinical information including tumor size, pathological stage, and lymphatic metastasis was collected and reviewed. For all the patients, sample were sampling intraoperatively and immediately frozen in liquid nitrogen with RNALater (Qiagen NV, Venlo, the Netherlands) and then stored at -80°C for further experimental using. This study was approved by the institutional Review Board for Human Research of The

Fourth People's Hospital of Jinan and the Affiliated Hospital of Shandong Academy of Medical Sciences. All recruited patients provided written informed consents.

Cell culture

Human breast cancer cell lines (MCF-7 and SK-BR-3) were obtained from Stem Cell Bank, Chinese Academy of Sciences (Shanghai, China). The tamoxifen-resistant MCF-7-R and SK-BR-3-R cell lines were developed by 70 days selection of MCF-7 and SK-BR-3 cells co-cultured with 800 nM 4-hydroxytamoxifen (4OH-TAM, Sigma-Aldrich Co., St Louis, MO, USA). The cells were cultured in RPMI-1640 (Sigma-Aldrich Co.) supplemented with 10% charcoal–dextran-stripped fetal bovine serum (Thermo Fisher Scientific, Waltham, MA, USA) and penicillin (100 U/mL, Sigma-Aldrich Co.) and streptomycin (100 U/mL, Sigma-Aldrich Co.) at 37°C with 5% CO_2 in a humidified cell culture incubator (Sanyo Electric Co., Ltd., Osaka, Japan).

Cell transfection

siRNA specific for H19 was purchased from Invitrogen (Thermo Fisher Scientific). To minimize nonspecific effects of interfering RNAs, non-targeting control siRNA, also obtained from Invitrogen, was used as negative control (NC). The siRNA transfection reagent (Invitrogen) was used according to the manufacturer's protocol. Briefly, 3×10^4 cells in media without antibiotic were plated in 6-well plates and then transfected with 100 nM siRNA for H19 or NC when cells were 70% confluence the plate well at 24 hours using lipofectamine 2000 reagent (Thermo Fisher Scientific). The cells were sampled at indicated time point for the different experiments with indicated purposes such as cell proliferation, invasion, and protein expression. For the validation of chemosensitivity experiment, cells with 24 hours siRNA transfection and then treated with tamoxifen in multiple concentrations and analyzed the cell proliferation and apoptosis rate.

Cell counting kit 8 assay

According to the manufacture instruction (Beyotime Institute of Biotechnology, Haimen, People's Republic of China), the procedure briefly described as following, MCF-7, MCF-7-R, SK-BR-3, or SK-BR-3R cells at the logarithmic growth phase were plated in 96-well at the density of 5,000 cells/well and cultured for 24 hours. And then, siRNA or NC were added with transfection reagents and cultured for another 72 hours. Cell counting kit solution (20 μL /well) was added to each well, and the cells were further incubated

for another half hours. In the validation of chemosensitivity, following the transfection of siRNA or NC, cells were cultured with tamoxifen with dose escalation from 0 to 100 μM (0, 0.01, 0.1, 1, 10, and 100 μM). The living cell number was measured by the OD value at 450 nm wavelength using microplate reader.

Transwell cell migration and invasion assay

For in vitro cell migration and invasion assay, the published on-line available protocol was used. Briefly as following: 100 μL cells (3×10^6 cells/well) transfected with H19 siRNA or NC were plated on the top of the membrane in a transwell inserted into the 24-well plate. When the cells settled down, 600 μL RPMI-1640 media with 20% FBS was added into the bottom of the lower chamber in a 24-well plate. Then, 48 hours later, cells that did not migrate from the top of the membrane were removed carefully and the migrated cells were fixed and stained with crystal violet. The cells in different fields of view were counted under microscope and using the average sum of cells to analyze the migration and invasion ability.

Flow cytometry analysis of apoptotic cells

Apoptosis was demonstrated by Annexin V-FITC and propidium iodide (PI) (Thermo Fisher Scientific) according to the manufacturer's protocol. Briefly, 1×10^6 cells were plated in 100 mm dishes (Corning Incorporated, Corning, NY, USA) and then transfected with H19 or mock vector. Then, 72 hours later, the cells were lysed and collected following twice washing with cold PBS and re-suspended with 500 μL binding buffer. And then cells were stained with Annexin-FITC and PI with another further incubated for 5 minutes at room temperature before running the flow cytometry. For each group, the samples were triplicate.

Real-time PCR analysis

For indicated treatments, cells were collected, and total RNA was isolated from these cells or tumor tissues from patients using PicoPure™ RNA Isolation Kit (Arcturus, Sunnyvale, CA, USA) according to the manufacturer's instructions. First, 2 μg of mRNA was transcribed into cDNA using SuperScript III RNase H Reverse Transcriptase (Thermo Fisher Scientific). Following the reverse transcription reactions, real-time PCR was performed with Taqman (Thermo Fisher Scientific) primers for lncRNA H19 and GAPDH and Vii™ 7 system (Bio-Rad Laboratories, Hercules, CA,

USA) was used to run the qPCR procedure according the manufacturers instruments. Based on the real-time PCR results of Ct number, the expression of mRNA levels was calculated using the $2^{-\Delta\Delta\text{Ct}}$ method and normalized to an internal reference gene GAPDH.

Western blot

Cells were collected at 72 hours with the treatment of H19 siRNA or NC and lysed in RIPA buffer (Sigma-Aldrich Co.). Protein concentration was quantified using BCA Protein Assay Kit (Beyotime). Then, 30 μg of total protein from each group was loaded and resolved with 10% Tris-SDS gel. After the electrophoresis, the gel was electro-transferred to polyvinylidene fluoride membranes (Thermo Fisher Scientific). For the following blocking and antibody incubation, iBind kit (Thermo Fisher Scientific) was used according the manufactures' instrument. The primary antibodies: anti-B-catenin (Cell Signaling Technology, Inc., Danvers, MA, USA) (1:300), anti-E-Cadherin (Cell Signaling) (1:300) anti-vimentin (Santa Cruz Biotechnology Inc., Dallas, TX, USA) (1:1,000), anti-GAPDH (Santa Cruz Biotechnology Inc.) (1:1,000), and anti-Histone (Santa Cruz Biotechnology Inc.) (1:1,000). Horseradish peroxidase-conjugated anti-rabbit and anti-mouse, and anti-goat IgG was used as the secondary antibody (Santa Cruz Biotechnology Inc.) (1:3,000). The signals were detected with super sensitive reagent (Thermo Fisher Scientific).

Statistical analysis

All statistical analysis was done with two-tailed Student's *t*-test available in GraphPad Prism7 (GraphPad Software, Inc., La Jolla, CA, USA). In all assays, one-way ANOVA was used to determine the difference between groups and the *P*-value <0.01 was considered statistically significant difference. All experiments were performed in triplicate and all value was presented as the mean \pm SD.

Results

lncRNA H19 has been reported to express in multiple cancers and takes a key role in tumorigenesis and metastasis.¹⁶ We first analyzed H19 mRNA level in breast cancer tissues and normal adjacent tissues from the 30 patients. RT-qPCR demonstrated that H19 mRNA was much higher in breast cancer tissues compared to normal tissues (Figure 1A, $P < 0.01$). Due to the fact that lncRNA H19 was associated with tumor metastasis, we investigated the correlation of lncRNA H19 level and tumor stage. As shown in Figure 1B, patients at late stage (stage III and IV) expressed relatively higher level




Figure 1 Comparison of lncRNA H19 expression level in breast cancers with different TNM stage.

Notes: Total RNA was extracted from both tumor cells and normal tissue cells of 30 enrolled breast cancer patients, and the H19 expression level was analyzed by RT-qPCR. **(A)** The relative fold change of H19 expression in breast cancer cells compared to normal adjacent normal tissues ($P < 0.01$). **(B)** Comparison of H19 expression of breast cancer patient with different stage ($P < 0.01$).

of H19 compared to patients at early stage cancers (stage I and II) ($P < 0.01$). Taken together, these results demonstrated that highly expressed lncRNA H19 might be correlated with metastatic breast cancer.

lncRNA H19 expression level manipulated breast cancer cell proliferation

In order to investigate the relationship of lncRNA H19 and tamoxifen-resistance breast cancer cells, we detected the level of H19 in tamoxifen-resistant breast cancer cell

lines. Results revealed that the level of lncRNA H19 in tamoxifen-resistance cells (MCF-7-R and SK-BR-3-R) was much higher than normal breast cancer cells (MCF-7 and SK-BR-3) ($P < 0.01$, Figure 2A). Next, to focus on the association of H19 and breast cancer cell proliferation, we assessed the impact of H19 on cell proliferation using cell counting kit-8 (CCK8). As shown in Figure 2B, both MCF-7-R and SK-BR-3-R acquired tamoxifen resistance. In addition, compared with control group, MCF-7 and SK-BR-3 cells transfected with H19 displayed faster growth, especially 48 hours and 72 hours after transfection




Figure 2 (Continued)




Figure 2 lncRNA H19 affected breast cancer cell proliferation.

Notes: (A) H19 expression level in parental cells (MCF-7 and SK-BR-3) and tamoxifen-resistance cells (MCF-7-R, SK-BR-3-R) ($P < 0.01$; $*P < 0.01$ vs the MCF-7 group; $^{\#}P < 0.01$ vs the SK-BR-3 group). (B) Parental cells (MCF-7 and SK-BR-3) and tamoxifen-resistance cells (MCF-7-R, SK-BR-3-R) were treated with 4OH-TAM for 72 hours (range from 0 to 10 μM). (C and D) Cell proliferation measurement using CCK8 after H19 or mock vector transfection in parental cells (MCF-7 and SK-BR-3, $P < 0.01$; [C] $*P < 0.01$ vs the vector group; [D] $*P < 0.01$ vs the NC group). (E and F) Cell proliferation measurement using CCK8 after siRNA of H19 transfection in tamoxifen-resistant cells (MCF-7-R and SK-BR-3-R, $P < 0.01$; [E] $*P < 0.01$ vs the vector group; [F] $*P < 0.05$ vs the NC group). (G) Cell proliferation measurement using EdU staining after siRNA of H19 transfection in tamoxifen-resistant cells (10 \times).

Abbreviation: NC, negative control.

(Figure 2C and D, $P < 0.01$). Additionally, when we use siRNA of H19 to suppress H19 level of MCF-7-R and SK-BR-3-R cells, the cell proliferation was inhibited (Figure 2E and F). This observation was further confirmed by EdU

staining, since siRNA H19-treated group contained less EdU-positive cells compared with NC group (Figure 2G), suggesting the strong correlation of lncRNA H19 level and cell proliferation.

Knockdown of lncRNA H19 improved tamoxifen sensitivity in terms of cell apoptosis in tamoxifen-resistant breast cancer cells

To investigate the contribution of lncRNA H19 to tamoxifen-resistance in breast cancer cells, we transfected H19 into tamoxifen-sensitive parental control cells (MCF-7 or SK-BR-3), as well as H19 siRNA into tamoxifen-resistant cells (MCF-7-R or SK-BR-3-R). All these cells were exposed to tamoxifen treatment. Under the same concentration of tamoxifen, sensitive cell lines displayed higher cell survival than resistant cell line (Figure 3A and B). We also observed that elevated level of H19 increased cell survival rate in sensitive cells, while knocking down of H19 decreased cell survival rate in resistant cells ($P < 0.01$). This indicated that in the presence of tamoxifen, suppression of lncRNA H19 could ameliorate the resistance to tamoxifen in tamoxifen-resistant cell lines.

Next, we performed FACS assay to investigate the effect of H19 on cell apoptosis. Results showed that transfection of H19 can inhibit apoptosis in MCF-7 parental control cells, and knockdown of H19 by siRNA can ameliorate the low level of apoptosis status in tamoxifen-resistant cells (MCF-7-R) (Figure 3C and D). This observation was further confirmed in SK-BR-3 and SK-BR-3-R cells (Figure 3E and F). These findings indicated that knockdown of lncRNA H19 could improve tamoxifen sensitivity in tamoxifen-resistant breast cancer cells.

Knockdown of lncRNA H19 improved tamoxifen sensitivity in terms of cell invasion in tamoxifen-resistant breast cancer cells

We performed Transwell experiments to interrogate the effect of lncRNA H19 on breast cancer cell invasion in




Figure 3 (Continued)




Figure 3 lncRNA H19 expression manipulates the tamoxifen sensitivity of breast cancer cells.

Notes: lncRNA H19 mimic was used to transfect breast cancer cells to increase H19 expression level and H19 siRNA was used to knockdown H19 expression level in breast cancer cell lines. Then, 72 hours after transfection, tamoxifen with dose escalation were added to the media as an anti-tumor agent. After 48 hours co-culture (120 hours post transfection), the proliferation of the cells was determined by CCK8 kit and the apoptotic cells were observed by performing Annexin V and PI double staining. **(A)** Cell survival rate of MCF-7 (* $P < 0.01$ compared to mock vector or NC group). **(B)** Cell survival rate of SK-BR-3 (* $P < 0.01$ compared to mock vector or NC group). **(C and D)** Effect of lncRNA H19 expression manipulation on cell apoptosis in MCF-7 and MCF-7-R measured by FACS. **(E and F)** Effect of lncRNA H19 expression manipulation on cell apoptosis in SK-BR-3 and SK-BR-3-R measured by FACS. * $P < 0.01$ vs the SK-BR-3 vector group; # $P < 0.01$ vs the SK-BR-3-R vector group.

Abbreviation: NC, negative control.

the presence of tamoxifen (1 μ M). Results showed that transfection of lncRNA H19 enhanced MCF-7 invasion to two folds compared to vector control. Tamoxifen-resistant MCF-7 cells with H19 knockdown displayed less invasion capability compared to control group under tamoxifen treatment ($P < 0.01$, Figure 4A and B). This finding was further confirmed in SK-BR-3 and SK-BR-3-R cells ($P < 0.01$, Figure 4C and D). This indicated that knockdown of lncRNA H19 improved tamoxifen sensitivity in terms of cell invasion in tamoxifen-resistant breast cancer cells.

Knockdown of H19 inhibited Wnt pathway in tamoxifen-resistance breast cancer cells

As lncRNA H19 is thought to participate in the breast cancer cell metastasis and Wnt pathway has been shown to play a critical role in promoting metastasis.¹⁵ Thus, we investigated the correlation of H19 and Wnt pathway. β -Catenin plays a central role mediating the canonical Wnt/ β -catenin signaling pathway. We found that knockdown of H19 by siRNA transfection can promote the translocation of β -catenin from nuclear to cytoplasm in tamoxifen-resistant cells (Figure 5A). The re-localization of β -catenin prompted us to investigate

Wnt-dependent transcription by using the Dual-luciferase reporter system and TOP/FOP reporter assays system. Upon knock-down of H19, the TOP/FOP ratio decreased to about 50% when compared to the control in both MCF-7-R and SK-BR-3-R cells (Figure 5B), indicating that loss of H19 inhibited Wnt-dependent transcription, in line with the loss of nuclear staining of beta-catenin observed in Figure 5A. We also measured the protein level of β -catenin, and results showed that knockdown of H19 reduced the expression of β -catenin in both MCF-7-R and SK-BR-3-R cells (Figure 5C and D).

R-Spondin (RSPO) proteins are a family of secreted molecules that strongly potentiate Wnt/ β -catenin signaling. We found that RSPO1 can reverse the β -catenin decrease induced by H19 knockdown in both two tamoxifen-resistant cell lines. (Figure 5E–G). In terms of cell proliferation, results showed that RSPO1 can block the inhibition of cell proliferation induced by H19 knockdown, both in MCF-7-R and SK-BR-3-R cells (Figure 5H and I). Taken together, these findings suggested that in tamoxifen-resistant cells, Wnt/beta-catenin signal pathway acted as a downstream of H19, and inhibition of H19 level can reduce the activity of Wnt/beta-catenin signal pathway and cell proliferation.




Figure 4 Effect of lncRNA H19 expression manipulation on breast cancer cell migration.

Notes: (A and B) Cell migration after transfection of H19 or H19 siRNA in MCF-7 and MCF-7-R cells (20 \times , $P < 0.01$; * $P < 0.01$ vs the MCF-7 vector group; # $P < 0.01$ vs the MCF-7-R NC group). (C and D) Cell migration after transfection of H19 or H19 siRNA in SK-BR-3 and SK-BR-3-R cells (20 \times , $P < 0.01$; * $P < 0.01$ vs the SK-BR-3 vector group; # $P < 0.01$ vs the SK-BR-3-R NC group).

Abbreviation: NC, negative control.

Knockdown of H19 inhibited EMT in tamoxifen-resistance breast cancer cells

EMT is an important process in tumor development. EMT is essential for developmental processes and has also been shown to occur in wound healing, in organ fibrosis, and in the initiation of metastasis in cancer progression. Wnt/ β -catenin signal pathway is an inducer of EMT and plays an important role in EMT, which displayed an upregulation of N-cadherin and downregulation of E-cadherin and vimentin. In this study, we found that knockdown of H19 by siRNA transfection can significantly reduce the expression of N-cadherin, as well as increase E-cadherin and vimentin level, both in MCF-7-R and SK-BR-3-R cells (Figure 6A–D). This observation indicated that knockdown of H19 can inhibit EMT in tamoxifen-resistance breast cancer cells.

Discussion

Breast cancer is the most common malignancy in women in worldwide and multiple advanced therapies have been developed. Tamoxifen is the widespread treatment for patients with ER positive breast cancer. Drug resistance is still a main obstacle preventing successful breast cancer treatment.^{17,18} Therefore, better understanding of the biological mechanisms for the acquired resistance to tamoxifen might suggest novel direction to overcome tamoxifen resistance.

lncRNA H19 is highly expressed in majority of human cancers and its overexpression is often correlated with tumor metastasis.^{19,20} Several studies about lncRNA H19 on drug resistance reported that H19 can reduce the sensitivity of breast cancer to paclitaxel via inactivation of BIK/NOXA, which were two important pro-apoptotic genes.²¹ Moreover,




Figure 5 The H19 knockdown deregulated Wnt pathway in tamoxifen-resistant breast cancer cells.

Notes: (A) H19 siRNA transfection promoted the translocation of β -catenin from nuclear to cytoplasm in tamoxifen-resistant cells. Red staining indicates β -catenin, blue means DAPI, and purple presents the merged color of β -catenin and DAPI. (B) Dual-luciferase reporter system and TOP/FOP reporter assays to presents the Wnt pathway transcription activity. (C) H19 knockdown decreased β -catenin protein level both in MCF-7-R and SK-BR-3-R cells. (D) Quantification of β -catenin protein level after H19 knockdown. (E–G) β -catenin protein level response in the presence of H19 knockdown and R-spondin I. (H and I) The response of cell proliferation in the presence of H19 knockdown and R-spondin I. * $P < 0.01$ vs the NC group; # $P < 0.01$ vs the R-spondin I group.

Abbreviation: NC, negative control.




Figure 6 The H19 knockdown de-regulated EMT in tamoxifen-resistant breast cancer cells.

Notes: (A) H19 knockdown lead to increase of E-cadherin and decrease od N-cadherin and vimentin. (B–D) Quantification of protein level in response to H19 knockdown. * $P < 0.01$ vs the MCF-7-R NC group; # $P < 0.01$ vs the SK-BR-3-R NC group.

Abbreviation: NC, negative control.

H19 inactivated p53 via physically interacts with p53 in gastric cancer cells and finally regulated the cell cycle.¹⁶ Stable expression of H19 significantly promotes EMT progression and accelerates in vivo and in vitro tumor growth in gastric cancer. H19 was an important factor contributing to drug resistance in breast cancer. lncRNA H19 overexpressed in various types of tumor cells such as bladder cancer, liver cancer, colon cancer, and breast cancer.^{11,22,23} Adriaenssens found that the expression of H19 increased in 72.5% breast cancer tissues and overexpression of an ectopic H19 gene enhances the tumorigenic properties of breast cancer cells.²⁴ Berteaux et al also found that over expression of H19 in breast cancer is the reason for promoting cell proliferation.²⁵ Similar to these studies, the expression of lncRNA H19 in breast cancer cells was investigated, and the underlying possible

molecular mechanism of H19 overexpression regulated tamoxifen resistant was interrogated. It was found that the expression level of H19 in tamoxifen-resistant breast cancer cells was significantly higher than that of tamoxifen-sensitive breast cancer cells.

More and more studies have shown that lncRNA H19 is involved in the development of tumor invasion and metastasis. In bladder cancer cells, the high expression of lncRNA H19 promotes the expression of the transcription factor EZH2, down the expression of E-cadherin, and then enhances the invasion ability of the cells in the hepatoma cells.²⁶ lncRNA H19 is expressed by upregulation of Slug to induce the tumor metastasis.²⁷ The above research results have certain reference significance for this study. The migration and invasion ability of tumor cells is a manifestation of

their malignant biological behavior and is also one of the main factors that affect tumor progression and prognosis. The metastatic risk of breast cancer is high because of the failure of endocrine therapy, and the expression of H19 in tamoxifen-resistant breast cancer is significantly improved, and our results also suggests that H19 is associated with TAM resistance in breast cancer. Therefore, we further study the effect of H19 on the metastasis of TAM breast cancer and the underlying involved pathways. We found that knockdown of lncRNA H19 can ameliorate tamoxifen resistance and promote cell apoptosis and inhibit cell proliferation in tamoxifen-resistant breast cancer cells. Moreover, knockdown of H19 can inhibited Wnt pathway and EMT in tamoxifen-resistance breast cancer cells. Therefore, in this study, we provide evidence for a novel mechanism that is employed by the lncRNA H19 to ameliorate tamoxifen resistance in breast cancer cells. Due to the crucial role of H19 in the amelioration of drug resistance, it may be used as a useful biomarkers and potential therapeutic target for breast cancer patients.

Over the last decade, researchers put efforts to illuminate the complex network of ER transcriptional effects such as ER/ERK/MAPK pathway and stress-activated protein kinase/c-junNH2 terminal kinase pathway and suggested that alterations of the elements in these signaling pathway play an important role in resistance to tamoxifen in breast cancer treatment.^{28–30} Moreover, with the development of molecular research, several critical importance targets of breast cancer cells have been identified, such as HER2, other receptor tyrosine kinases, and components of the PI3K/AKT/mTOR and Raf/MEK/ERK pathways.^{31,32} In the current study, H19 overexpression may be related to the poor prognosis of tamoxifen-resistance breast cancer. Therefore, we present a rationale for dual inhibition of these pathways in the treatment of breast cancer patients with acquired resistance.

Taken together, the reduction of H19 expression level inhibits the expression of transcription factors related to the Wnt pathway and EMT, thus inhibiting the invasion ability of tamoxifen-resistance breast cancer, suggesting that H19 may be related to the poor prognosis of tamoxifen-resistance breast cancer, and is expected to be a drug target for the treatment of metastatic breast cancer after tamoxifen-resistance. This provided a theoretical basis for further studies about targeted drugs investigation.

Disclosure

The authors report no conflicts of interest in this work.

References

- Bartsch R, Bergen E. ASCO 2017: highlights in breast cancer. *Memo*. 2017;10(4):228–232.
- Agarwala V, Choudhary N, Gupta S. A risk-benefit assessment approach to selection of adjuvant chemotherapy in elderly patients with early breast cancer: a mini review. *Indian J Med Paediatr Oncol*. 2017;38(4):526–534.
- Başaran GA, Twelves C, Diéras V, Cortés J, Awada A. Ongoing unmet needs in treating estrogen receptor-positive/HER2-negative metastatic breast cancer. *Cancer Treat Rev*. 2018;63:144–155.
- Calhoun BC, Collins LC. Predictive markers in breast cancer: an update on ER and HER2 testing and reporting. *Semin Diagn Pathol*. 2015;32(5):362–369.
- Lee A, Djamgoz MBA. Triple negative breast cancer: emerging therapeutic modalities and novel combination therapies. *Cancer Treat Rev*. 2018;62:110–122.
- Wu Y, Zhang Z, Cenciarini ME, et al. Tamoxifen resistance in breast cancer is regulated by the EZH2-ER α -GREB1 transcriptional axis. *Cancer Res*. 2018;78(3):671–684.
- Huang X, Luo YL, Mao YS, Ji JL, Ji J. The link between long noncoding RNAs and depression. *Prog Neuropsychopharmacol Biol Psychiatry*. 2017;73:73–78.
- Mirza AH, Kaur S, Pociot F. Long non-coding RNAs as novel players in β cell function and type 1 diabetes. *Hum Genomics*. 2017;11(1):17.
- Luo Q, Chen Y. Long noncoding RNAs and Alzheimer's disease. *Clin Interv Aging*. 2016;11:867–872.
- Collette J, Le Bourhis X, Adriaenssens E. Regulation of human breast cancer by the long non-coding RNA H19. *Int J Mol Sci*. 2017;18(11):2319.
- Hu P, Yang J, Hou Y, et al. LncRNA expression signatures of twist-induced epithelial-to-mesenchymal transition in MCF10A cells. *Cell Signal*. 2014;26(1):83–93.
- Liang WC, Fu WM, Wang YB, et al. H19 activates Wnt signaling and promotes osteoblast differentiation by functioning as a competing endogenous RNA. *Sci Rep*. 2016;6:20121.
- Wu KF, Liang WC, Feng L, et al. H19 mediates methotrexate resistance in colorectal cancer through activating Wnt/ β -catenin pathway. *Exp Cell Res*. 2017;350(2):312–317.
- Chen J, Wang T, Zhou YC, et al. Aquaporin 3 promotes epithelial-mesenchymal transition in gastric cancer. *J Exp Clin Cancer Res*. 2014;33:38.
- Zhou W, Ye XL, Xu J, et al. The lncRNA H19 mediates breast cancer cell plasticity during EMT and MET plasticity by differentially sponging miR-200b/c and let-7b. *Sci Signal*. 2017;10(483):eaak9557.
- Li T, Mo X, Fu L, Xiao B, Guo J. Molecular mechanisms of long non-coding RNAs on gastric cancer. *Oncotarget*. 2016;7(8):8601–8612.
- Fedele P, Ciccarese M, Surico G, Ciniéri S. An update on first line therapies for metastatic breast cancer. *Expert Opin Pharmacother*. 2018;19(3):243–252.
- Gonzalez-Angulo AM, Morales-Vasquez F, Hortobagyi GN. Overview of resistance to systemic therapy in patients with breast cancer. *Adv Exp Med Biol*. 2007;608:1–22.
- Li CX, Li HG, Huang LT, et al. H19 lncRNA regulates keratinocyte differentiation by targeting miR-130b-3p. *Cell Death Dis*. 2017;8(11):e3174.
- Wei Y, Liu Z, Fang J. H19 functions as a competing endogenous RNA to regulate human epidermal growth factor receptor expression by sequestering let-7c in gastric cancer. *Mol Med Rep*. 2018;17(2):2600–2606.
- Si X, Zang R, Zhang E, et al. LncRNA H19 confers chemoresistance in ER α -positive breast cancer through epigenetic silencing of the proapoptotic gene BIK. *Oncotarget*. 2016;7(49):81452–81462.
- Yu M, Bardia A, Wittner BS, et al. Circulating breast tumor cells exhibit dynamic changes in epithelial and mesenchymal composition. *Science*. 2013;339(6119):580–584.

23. Gunasinghe NP, Wells A, Thompson EW, Hugo HJ. Mesenchymal-epithelial transition (MET) as a mechanism for metastatic colonisation in breast cancer. *Cancer Metastasis Rev.* 2012;31(3-4):469-478.
24. Lottin S, Adriaenssens E, Dupressoir T, et al. Overexpression of an ectopic H19 gene enhances the tumorigenic properties of breast cancer cells. *Carcinogenesis.* 2002;23(11):1885-1895.
25. Ohana P, Kopf E, Bibi O, et al. The expression of the H19 gene and its function in human bladder carcinoma cell lines. *FEBS Lett.* 1999;454(1-2): 81-84.
26. Luo M, Li Z, Wang W, Zeng Y, Liu Z, Qiu J. Long non-coding RNA H19 increases bladder cancer metastasis by associating with EZH2 and inhibiting E-cadherin expression. *Cancer Lett.* 2013;333(2): 213-221.
27. Matouk IJ, Raveh E, Abu-Lail R, et al. Oncofetal H19 RNA promotes tumor metastasis. *Biochim Biophys Acta.* 2014;1843(7):1414-1426.
28. Ring A, Dowsett M. Mechanisms of tamoxifen resistance. *Endocr Relat Cancer.* 2004;11(4):643-658.
29. Yamaguchi N, Nakayama Y, Yamaguchi N. Down-regulation of Forkhead box protein A1 (FOXA1) leads to cancer stem cell-like properties in tamoxifen-resistant breast cancer cells through induction of interleukin-6. *J Biol Chem.* 2017;292(20):8136-8148.
30. Chen S, Yao F, Xiao Q, et al. EZH2 inhibition sensitizes tamoxifen-resistant breast cancer cells through cell cycle regulation. *Mol Med Rep.* 2018;17(2):2642-2650.
31. Saini KS, Loi S, de Azambuja E, et al. Targeting the PI3K/AKT/mTOR and Raf/MEK/ERK pathways in the treatment of breast cancer. *Cancer Treat Rev.* 2013;39(8):935-946.
32. Herter-Sprue GS, Greulich H, Wong KK. Activating mutations in ERBB2 and their impact on diagnostics and treatment. *Front Oncol.* 2013;3:86.

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress